

✔ Congratulations! You passed!

Grade
received 100%

Latest Submission
Grade 100%

To pass 70% or
higher

Go to next item

1.

1 / 1 point

Gradient descent is an algorithm for finding values of parameters w and b that minimize the cost function J .

repeat until convergence {

$$w = w - \alpha \frac{\partial}{\partial w} J(w, b)$$

$$b = b - \alpha \frac{\partial}{\partial b} J(w, b)$$

When $\frac{\partial J(w, b)}{\partial w}$ is a negative number (less than zero), what happens to w after one update step?

- ☐ It is not possible to tell if w will increase or decrease.
- ☐ w decreases
- ☒ w increases.
- ☐ w stays the same

✔ Correct

The learning rate is always a positive number, so if you take W minus a negative number, you end up with a new value for W that is larger (more positive).

2.

1 / 1 point

For linear regression, what is the update step for parameter b ?

☒ $b = b - \alpha \frac{1}{m} \sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)})$

☐ $b = b - \alpha \frac{1}{m} \sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)}) x^{(i)}$

✔ Correct

The update step is $b = b - \alpha \frac{\partial J(w, b)}{\partial b}$ where $\frac{\partial J(w, b)}{\partial b}$ can be computed with this expression:
 $\sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)})$