

# Conformal Self Mappings

Jaemin Oh

KAIST

October 6, 2020

# Contents

# conformal map

- ▶  $U, V$  : open subsets of  $\mathbb{C}$

# conformal map

- ▶  $U, V$  : open subsets of  $\mathbb{C}$
- ▶  $f$  is a function of  $U$  into  $V$

# conformal map

- ▶  $U, V$  : open subsets of  $\mathbb{C}$
- ▶  $f$  is a function of  $U$  into  $V$
- ▶  $f$  is conformal if  $f$  is bijective and holomorphic.

# conformal map

- ▶  $U, V$  : open subsets of  $\mathbb{C}$
- ▶  $f$  is a function of  $U$  into  $V$
- ▶  $f$  is conformal if  $f$  is bijective and holomorphic.
- ▶ conformal = biholomorphic

# conformal map

- ▶  $U, V$  : open subsets of  $\mathbb{C}$
- ▶  $f$  is a function of  $U$  into  $V$
- ▶  $f$  is conformal if  $f$  is bijective and holomorphic.
- ▶ conformal = biholomorphic
- ▶ If  $h$  is holomorphic function of  $U$ , and  $U$  is somewhat complicated, by considering  $h \circ f$ , we can change the domain of  $h$ .

# characterizing conformal self mapping of $\mathbb{C}$

- ▶ natural example:  $az + b$  for  $a \neq 0$



# characterizing conformal self mapping of $\mathbb{C}$

- ▶ natural example:  $az + b$  for  $a \neq 0$
- ▶ In fact, above form is all of them.

# characterizing conformal self mapping of $\mathbb{C}$

- ▶ natural example:  $az + b$  for  $a \neq 0$
- ▶ In fact, above form is all of them.
- ▶ Note that we are considering not just entire function.  
Conformal self mappings of  $\mathbb{C}$  has more condition than entire function.

## Lemma 6.1.2.

If  $f : \mathbb{C} \rightarrow \mathbb{C}$  is a conformal then  $\lim_{|z| \rightarrow \infty} |f(z)| = \infty$ .

## proof of lemma 6.1.2.

- Fix  $M$ . We want to show existence of  $N$  such that  $f(\{z : |z| > N\}) \subset \{w : |w| > M\}$ .

## proof of lemma 6.1.2.

- ▶ Fix  $M$ . We want to show existence of  $N$  such that  $f(\{z : |z| > N\}) \subset \{w : |w| > M\}$ .
- ▶ Above is equivalent to  $\{z : |z| > N\} \subset f^{-1}(\{w : |w| > M\})$  since  $f$  is bijective.

## proof of lemma 6.1.2.

- ▶ Fix  $M$ . We want to show existence of  $N$  such that  $f(\{z : |z| > N\}) \subset \{w : |w| > M\}$ .
- ▶ Above is equivalent to  $\{z : |z| > N\} \subset f^{-1}(\{w : |w| > M\})$  since  $f$  is bijective.
- ▶ Above is equivalent to  $\{z : |z| \leq N\} \supset f^{-1}(\{w : |w| \leq M\})$  by taking complement.

## proof of lemma 6.1.2.

- ▶ Fix  $M$ . We want to show existence of  $N$  such that  $f(\{z : |z| > N\}) \subset \{w : |w| > M\}$ .
- ▶ Above is equivalent to  $\{z : |z| > N\} \subset f^{-1}(\{w : |w| > M\})$  since  $f$  is bijective.
- ▶ Above is equivalent to  $\{z : |z| \leq N\} \supset f^{-1}(\{w : |w| \leq M\})$  by taking complement.
- ▶ Existence of  $N$  is clear since RHS of above is compact( $\Rightarrow$  closed and bounded).

# characterizing conformal self mapping of $\mathbb{C}$

## lemma 6.1.3

$f$  is a conformal self mapping of  $\mathbb{C}$ . Then there are  $B, D > 0$  such that  $|z| > D$  implies  $|f(z)| < B|z|$ .

## proof of lemma 6.1.3.

- ▶ There is  $C$  such that  $|z| > C$  implies  $|f(z)| > 1$ .



## proof of lemma 6.1.3.

- ▶ There is  $C$  such that  $|z| > C$  implies  $|f(z)| > 1$ .
- ▶ Define  $g(z) = 1/f\left(\frac{1}{z}\right)$  for  $z \in D'(0, \frac{1}{C})$ .

## proof of lemma 6.1.3.

- ▶ There is  $C$  such that  $|z| > C$  implies  $|f(z)| > 1$ .
- ▶ Define  $g(z) = 1/f\left(\frac{1}{z}\right)$  for  $z \in D'(0, \frac{1}{C})$ .
- ▶ As  $z \rightarrow 0$ ,  $g \rightarrow 0$ . So  $g$  has removable singularity at 0.

## proof of lemma 6.1.3.

- ▶ There is  $C$  such that  $|z| > C$  implies  $|f(z)| > 1$ .
- ▶ Define  $g(z) = 1/f\left(\frac{1}{z}\right)$  for  $z \in D'(0, \frac{1}{C})$ .
- ▶ As  $z \rightarrow 0$ ,  $g \rightarrow 0$ . So  $g$  has removable singularity at 0.
- ▶  $g'(0) \neq 0$  since  $g$  is injection since  $f$  is injection.

## proof of lemma 6.1.3.

- ▶ There is  $C$  such that  $|z| > C$  implies  $|f(z)| > 1$ .
- ▶ Define  $g(z) = 1/f\left(\frac{1}{z}\right)$  for  $z \in D'(0, \frac{1}{C})$ .
- ▶ As  $z \rightarrow 0$ ,  $g \rightarrow 0$ . So  $g$  has removable singularity at 0.
- ▶  $g'(0) \neq 0$  since  $g$  is injection since  $f$  is injection.
- ▶ Near 0,  $\left|\frac{g(z)}{z}\right| > \frac{1}{B}$ .

## proof of lemma 6.1.3.

- ▶ There is  $C$  such that  $|z| > C$  implies  $|f(z)| > 1$ .
- ▶ Define  $g(z) = 1/f\left(\frac{1}{z}\right)$  for  $z \in D'(0, \frac{1}{C})$ .
- ▶ As  $z \rightarrow 0$ ,  $g \rightarrow 0$ . So  $g$  has removable singularity at 0.
- ▶  $g'(0) \neq 0$  since  $g$  is injection since  $f$  is injection.
- ▶ Near 0,  $\left|\frac{g(z)}{z}\right| > \frac{1}{B}$ .
- ▶ Near 0,  $\left|f\left(\frac{1}{z}\right)\right| < \frac{B}{|z|}$ .

## proof of lemma 6.1.3.

- ▶ There is  $C$  such that  $|z| > C$  implies  $|f(z)| > 1$ .
- ▶ Define  $g(z) = 1/f\left(\frac{1}{z}\right)$  for  $z \in D'(0, \frac{1}{C})$ .
- ▶ As  $z \rightarrow 0$ ,  $g \rightarrow 0$ . So  $g$  has removable singularity at 0.
- ▶  $g'(0) \neq 0$  since  $g$  is injection since  $f$  is injection.
- ▶ Near 0,  $\left|\frac{g(z)}{z}\right| > \frac{1}{B}$ .
- ▶ Near 0,  $\left|f\left(\frac{1}{z}\right)\right| < \frac{B}{|z|}$ .
- ▶  $z \mapsto \frac{1}{z}$  leads the conclusion.

## characterizing

- ▶ Consider  $|f^{(n)}(0)| \leq \frac{n!}{r^n} \sup_{w \in \partial D(0,r)} |f(w)|$ .

## characterizing

- ▶ Consider  $|f^{(n)}(0)| \leq \frac{n!}{r^n} \sup_{w \in \partial D(0,r)} |f(w)|$ .
- ▶ For  $r > D$ , supremum above  $\leq Br$  by lemma 6.1.3.



## characterizing

- ▶ Consider  $|f^{(n)}(0)| \leq \frac{n!}{r^n} \sup_{w \in \partial D(0,r)} |f(w)|$ .
- ▶ For  $r > D$ , supremum above  $\leq Br$  by lemma 6.1.3.
- ▶ If  $n > 1$ , by letting  $r \rightarrow \infty$ ,  $n$ -th derivative of  $f$  at 0 must be zero.

## characterizing

- ▶ Consider  $|f^{(n)}(0)| \leq \frac{n!}{r^n} \sup_{w \in \partial D(0,r)} |f(w)|$ .
- ▶ For  $r > D$ , supremum above  $\leq Br$  by lemma 6.1.3.
- ▶ If  $n > 1$ , by letting  $r \rightarrow \infty$ ,  $n$ -th derivative of  $f$  at 0 must be zero.
- ▶ Therefore  $f$  must be polynomial of degree at most 1.

## characterizing

- ▶ Consider  $|f^{(n)}(0)| \leq \frac{n!}{r^n} \sup_{w \in \partial D(0,r)} |f(w)|$ .
- ▶ For  $r > D$ , supremum above  $\leq Br$  by lemma 6.1.3.
- ▶ If  $n > 1$ , by letting  $r \rightarrow \infty$ ,  $n$ -th derivative of  $f$  at 0 must be zero.
- ▶ Therefore  $f$  must be polynomial of degree at most 1.
- ▶ But  $f$  must be nonconstant. So  $f(z) = az + b$  for  $a \neq 0$ .

# characterizing

- ▶ Consider  $|f^{(n)}(0)| \leq \frac{n!}{r^n} \sup_{w \in \partial D(0,r)} |f(w)|$ .
- ▶ For  $r > D$ , supremum above  $\leq Br$  by lemma 6.1.3.
- ▶ If  $n > 1$ , by letting  $r \rightarrow \infty$ ,  $n$ -th derivative of  $f$  at 0 must be zero.
- ▶ Therefore  $f$  must be polynomial of degree at most 1.
- ▶ But  $f$  must be nonconstant. So  $f(z) = az + b$  for  $a \neq 0$ .
- ▶ We are characterized conformal self mappings of  $\mathbb{C}$ .

## remark

- ▶  $h$  is holomorphic on  $\{z : |z| > \alpha\}$  and  $\lim_{|z| \rightarrow \infty} |h(z)| = \infty$ .

## remark

- ▶  $h$  is holomorphic on  $\{z : |z| > \alpha\}$  and  $\lim_{|z| \rightarrow \infty} |h(z)| = \infty$ .
- ▶ By same procedure in proof of lemma 6.1.3, we can conclude that there are  $B, D > 0$  such that  $|z| > D \Rightarrow |h(z)| < B|z|^n$  for some  $n$ .

## remark

- ▶  $h$  is holomorphic on  $\{z : |z| > \alpha\}$  and  $\lim_{|z| \rightarrow \infty} |h(z)| = \infty$ .
- ▶ By same procedure in proof of lemma 6.1.3, we can conclude that there are  $B, D > 0$  such that  $|z| > D \Rightarrow |h(z)| < B|z|^n$  for some  $n$ .
- ▶ Why  $n$ ? Because we cannot say  $g'(0) \neq 0$ . But,  $g^{(n)}(0) \neq 0$  for some  $n$  since  $g$  is nonconstant since  $h$  is nonconstant.

## remark

- ▶  $h$  is holomorphic on  $\{z : |z| > \alpha\}$  and  $\lim_{|z| \rightarrow \infty} |h(z)| = \infty$ .
- ▶ By same procedure in proof of lemma 6.1.3, we can conclude that there are  $B, D > 0$  such that  $|z| > D \Rightarrow |h(z)| < B|z|^n$  for some  $n$ .
- ▶ Why  $n$ ? Because we cannot say  $g'(0) \neq 0$ . But,  $g^{(n)}(0) \neq 0$  for some  $n$  since  $g$  is nonconstant since  $h$  is nonconstant.
- ▶ Note that entire function  $\varphi$  which satisfies  $\lim_{|z| \rightarrow \infty} |\varphi(z)| = \infty$  must be polynomial.



# characterizing conformal self mapping of unit disc

- ▶ natural example : rotation ( $f(z) = wz$  for  $|w| = 1$ )

# characterizing conformal self mapping of unit disc

- ▶ natural example : rotation ( $f(z) = wz$  for  $|w| = 1$ )
- ▶ In fact, above form is all of them which fixes origin.

# characterizing conformal self mapping of unit disc

- ▶ natural example : rotation ( $f(z) = wz$  for  $|w| = 1$ )
- ▶ In fact, above form is all of them which fixes origin.

## lemma 6.2.1.

$f : D \rightarrow D$  is biholomorphic which fixes origin iff  $f(z) = wz$  for  $|w| = 1$



## proof of lemma 6.2.1.

- ▶  $g = f^{-1}$ . Then both of  $f, g$  are fixing origin.

## proof of lemma 6.2.1.

- ▶  $g = f^{-1}$ . Then both of  $f, g$  are fixing origin.
- ▶ Schwarz lemma says  $|f'(0)|$  and  $|g'(0)|$  are  $\leq 1$ .

## proof of lemma 6.2.1.

- ▶  $g = f^{-1}$ . Then both of  $f, g$  are fixing origin.
- ▶ Schwarz lemma says  $|f'(0)|$  and  $|g'(0)|$  are  $\leq 1$ .
- ▶ Chain rule says  $f'(0)g'(0) = 1$ . This leads  $|f'(0)| = |g'(0)| = 1$ .

## proof of lemma 6.2.1.

- ▶  $g = f^{-1}$ . Then both of  $f, g$  are fixing origin.
- ▶ Schwarz lemma says  $|f'(0)|$  and  $|g'(0)|$  are  $\leq 1$ .
- ▶ Chain rule says  $f'(0)g'(0) = 1$ . This leads  $|f'(0)| = |g'(0)| = 1$ .
- ▶ Uniqueness of Schwarz lemma tells us that  $f(z) = f'(0)z$ .

# Mobius transformation

- ▶  $\varphi_a(z) = \frac{z-a}{1-\bar{a}z}$  for  $|a| < 1$  is called Mobius transformation.



# Mobius transformation

- ▶  $\varphi_a(z) = \frac{z-a}{1-\bar{a}z}$  for  $|a| < 1$  is called Mobius transformation.
- ▶ Theorem 5.5.2 says  $\varphi_a$  is conformal self mapping of unit disc. So we take it for granted.

# Mobius transformation

- ▶  $\varphi_a(z) = \frac{z-a}{1-\bar{a}z}$  for  $|a| < 1$  is called Mobius transformation.
- ▶ Theorem 5.5.2 says  $\varphi_a$  is conformal self mapping of unit disc. So we take it for granted.

## theorem 6.2.3.

$f$  is conformal self mapping of unit disc. Then  $f(z) = w\varphi_a(z)$  for some  $|a| < 1$  and  $|w| = 1$ .



## proof of theorem 6.2.3.

- ▶  $f(0) = b$ . Let  $g = \varphi_b \circ f$ . Then  $g$  fixes origin.

## proof of theorem 6.2.3.

- ▶  $f(0) = b$ . Let  $g = \varphi_b \circ f$ . Then  $g$  fixes origin.
- ▶  $g(z) = wz$  for some  $|w| = 1$ . Namely,  $f(z) = \varphi_b^{-1}(wz)$ .

## proof of theorem 6.2.3.

- ▶  $f(0) = b$ . Let  $g = \varphi_b \circ f$ . Then  $g$  fixes origin.
- ▶  $g(z) = wz$  for some  $|w| = 1$ . Namely,  $f(z) = \varphi_b^{-1}(wz)$ .
- ▶ But  $\varphi_b^{-1} = \varphi_{-b}$ .

## proof of theorem 6.2.3.

- ▶  $f(0) = b$ . Let  $g = \varphi_b \circ f$ . Then  $g$  fixes origin.
- ▶  $g(z) = wz$  for some  $|w| = 1$ . Namely,  $f(z) = \varphi_b^{-1}(wz)$ .
- ▶ But  $\varphi_b^{-1} = \varphi_{-b}$ .
- ▶  $f(z) = \frac{wz+b}{1+\overline{b}wz}$

## proof of theorem 6.2.3.

- ▶  $f(0) = b$ . Let  $g = \varphi_b \circ f$ . Then  $g$  fixes origin.
- ▶  $g(z) = wz$  for some  $|w| = 1$ . Namely,  $f(z) = \varphi_b^{-1}(wz)$ .
- ▶ But  $\varphi_b^{-1} = \varphi_{-b}$ .
- ▶  $f(z) = \frac{wz+b}{1+\overline{b}wz}$
- ▶ Simple calculation leads  $f(z) = w\varphi_{-bw^{-1}}(z)$ .

## proof of theorem 6.2.3.

- ▶  $f(0) = b$ . Let  $g = \varphi_b \circ f$ . Then  $g$  fixes origin.
- ▶  $g(z) = wz$  for some  $|w| = 1$ . Namely,  $f(z) = \varphi_b^{-1}(wz)$ .
- ▶ But  $\varphi_b^{-1} = \varphi_{-b}$ .
- ▶  $f(z) = \frac{wz+b}{1+\overline{b}wz}$
- ▶ Simple calculation leads  $f(z) = w\varphi_{-bw^{-1}}(z)$ .
- ▶ Take  $a = -bw^{-1}$ .



# preliminaries of linear fractional transformation

- ▶ Riemann sphere is  $\mathbb{C} \cup \infty \cong S^2$  by stereographic projection.

# preliminaries of linear fractional transformation

- ▶ Riemann sphere is  $\mathbb{C} \cup \infty \cong S^2$  by stereographic projection.
- ▶  $p_i \rightarrow p_0$  in R-sphere is equivalent to  $\pi^{-1}(p_i) \rightarrow \pi^{-1}(p_0)$  in  $S^2$ .

## preliminaries of linear fractional transformation

- ▶ Riemann sphere is  $\mathbb{C} \cup \infty \cong S^2$  by stereographic projection.
- ▶  $p_i \rightarrow p_0$  in R-sphere is equivalent to  $\pi^{-1}(p_i) \rightarrow \pi^{-1}(p_0)$  in  $S^2$ .
- ▶ Note that image of north-pole in  $S^2$  under projection is  $\infty$ .

# preliminaries of linear fractional transformation

- ▶ Riemann sphere is  $\mathbb{C} \cup \infty \cong S^2$  by stereographic projection.
- ▶  $p_i \rightarrow p_0$  in R-sphere is equivalent to  $\pi^{-1}(p_i) \rightarrow \pi^{-1}(p_0)$  in  $S^2$ .
- ▶ Note that image of north-pole in  $S^2$  under projection is  $\infty$ .
- ▶ Also, above definition of limit in extended plane is congruent to definition using metric.

# linear fractional transformation

- ▶ Let  $ad - bc \neq 0$ ,  $a, b, c, d \in \mathbb{C}$ .  $f(z) = \frac{az+b}{cz+d}$  is called linear fractional transformation if it satisfies two more conditions.

# linear fractional transformation

- ▶ Let  $ad - bc \neq 0$ ,  $a, b, c, d \in \mathbb{C}$ .  $f(z) = \frac{az+b}{cz+d}$  is called linear fractional transformation if it satisfies two more conditions.
- ▶ If  $c = 0$ ,  $f(\infty) = \infty$ . In this case,  $f$  is a linear map.

# linear fractional transformation

- ▶ Let  $ad - bc \neq 0$ ,  $a, b, c, d \in \mathbb{C}$ .  $f(z) = \frac{az+b}{cz+d}$  is called linear fractional transformation if it satisfies two more conditions.
- ▶ If  $c = 0$ ,  $f(\infty) = \infty$ . In this case,  $f$  is a linear map.
- ▶ If  $c \neq 0$ ,  $f(-\frac{d}{c}) = \infty$ ,  $f(\infty) = \frac{a}{c}$ .

# linear fractional transformation

- ▶ Let  $ad - bc \neq 0$ ,  $a, b, c, d \in \mathbb{C}$ .  $f(z) = \frac{az+b}{cz+d}$  is called linear fractional transformation if it satisfies two more conditions.
- ▶ If  $c = 0$ ,  $f(\infty) = \infty$ . In this case,  $f$  is a linear map.
- ▶ If  $c \neq 0$ ,  $f(-\frac{d}{c}) = \infty$ ,  $f(\infty) = \frac{a}{c}$ .
- ▶ Note that  $f(p_i) \rightarrow f(p_0)$  when  $p_i \rightarrow p_0$  for all  $p_0 \in \mathbb{C} \cup \infty$ .



# linear fractional transformation

- ▶ Let  $ad - bc \neq 0$ ,  $a, b, c, d \in \mathbb{C}$ .  $f(z) = \frac{az+b}{cz+d}$  is called linear fractional transformation if it satisfies two more conditions.
- ▶ If  $c = 0$ ,  $f(\infty) = \infty$ . In this case,  $f$  is a linear map.
- ▶ If  $c \neq 0$ ,  $f(-\frac{d}{c}) = \infty$ ,  $f(\infty) = \frac{a}{c}$ .
- ▶ Note that  $f(p_i) \rightarrow f(p_0)$  when  $p_i \rightarrow p_0$  for all  $p_0 \in \mathbb{C} \cup \infty$ .
- ▶ Above says continuity of  $f$  on extended plane.