

Andy's math/science background information

J. Andrew Fingerhut (andy.fingerhut@gmail.com)

June 1, 2025

Contents

1	Introduction	1
2	Notation	1
2.1	Maxwell's Equations	1
2.1.1	Questions about Maxwell's Equations	2
3	Gradient	3
4	Galilean relativity	3
5	Experiments	4
6	Questions	4
7	Index of A History of the Theories of Aether and Electricity	4
7.1	Definitions of some key terms	8
7.2	Noteworthy events in aether vs. action at a distance	9

1 Introduction

This document is a place to write up notes on background in math that help in learning science.

2 Notation

\mathbb{R}^3 - The set of all points in 3-dimensional space, i.e. where each point is specified by a sequence of 3 real-valued coordinates.

∇ - Called “nabla”, and often called “del”. Some history of it can be found on the Nabla Symbol page [8].

∇f - Gradient of a scalar function $f : \mathbb{R}^3 \rightarrow \mathbb{R}$.

$\text{grad} f$ - another way to write the gradient of f , ∇f

TODO: Any 3Blue1Brown-quality YouTube videos showing examples and definition of gradient, divergence, and curl? If so, give links to them below in appropriate sections.

2.1 Maxwell's Equations

In partial differential form:

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0} \quad \text{Gauss's Law} \quad (1)$$

$$\nabla \cdot \mathbf{B} = 0 \quad \text{Gauss's Law for magnetism} \quad (2)$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad \text{Maxwell-Faraday Equation (Faraday's law of induction)} \quad (3)$$

$$\nabla \times \mathbf{B} = \mu_0(\mathbf{J} + \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}) \quad \text{Ampère-Maxwell law} \quad (4)$$

where:

- \mathbf{E} is the electric field
- \mathbf{B} is the magnetic field
- ρ is the electric charge density
- \mathbf{J} is the current density
- ϵ_0 is the vacuum permittivity
- μ_0 is the vacuum permeability

Statement of the equations in English:

Gauss's Law (equation (1)) says that electric fields are “produced” radially outward from volumes with positive electric charge, and “consumed” radially into volumes with negative electric charge.

Gauss's Law for Magnetism (equation (??)) says that there are no magnetic monopoles, i.e. no sources or sinks for net magnetic field. All magnetic flux “swirls around” in loops.

2.1.1 Questions about Maxwell's Equations

Questions about Gauss's Law TODO: Is Gauss's Law a strict generalization of Coulomb's Law? In what ways? One way is that Gauss's Law gives the electric field generated by a distribution of charge, whereas Coulomb's Law gives the force between two point charges. So applying to a continuous distribution of charge is one generalization, and what electric field is created by that distribution of charge is another (as long as you then combine Gauss's Law with the Lorentz force law in order to calculate the forces).

TODO: It seems that Gauss's Law, if you take ρ and \mathbf{E} as functions of time, gives an electric field that changes instantaneously everywhere when the charge distribution ρ changes. There does not appear to be anything related to retarded positions of charge in that equation anywhere. How does the finite speed of electric field propagation enter into Maxwell's equations?

Doing a Google search on the question “is Gauss's law consistent with relativistic effects” gives answers that it *is* consistent with special relativity, including the following paragraphs:

- Maxwell's equations are relativistic: Gauss's law is one of Maxwell's equations. These equations, which describe classical electromagnetism, are inherently Lorentz invariant. This means they hold true in all inertial reference frames, consistent with the principle of relativity, which states that the laws of physics are the same for all observers in uniform motion relative to one another.
- Lorentz covariance: Maxwell's equations, and thus Gauss's law, can be written in a covariant form that is explicitly consistent with Lorentz transformations. These transformations describe how physical quantities change when moving from one inertial frame to another, incorporating relativistic effects like time dilation and length contraction.

Questions about Gauss's Law for Magnetism Figures for electromagnetic waves that are the propagation of light in a vacuum show \mathbf{E} and \mathbf{B} fields as perpendicular to each other, and each a sinusoid with the same phase as each other. These figures must be dramatic simplifications of the full \mathbf{E} and \mathbf{B} fields, e.g. perhaps they show at one instant in time all \mathbf{E} and \mathbf{B} fields only along the line that is the “center” of the ray of light? In particular, I am wondering how they look in all of space, such that it is clear that their divergence is 0 at all times, at all positions in space. It might be instructive to have a visual representation, perhaps an animation, somehow showing what the \mathbf{E} field is at all points in space at one particular instant of time. The \mathbf{B} field has a very similar shape. And if you know what the shape of these fields are at one instant of time, then you know that they merely “shift” in the direction of light propagation over time at speed c , without changing shape.

Is there a way to write them that has only ρ without \mathbf{J} ? If one emphasizes the view that all current is due to moving charge, is there a way to write Maxwell's equations in a way that makes this clear? For example, is there a way to write it with some single function x that somehow represents both ρ and \mathbf{J} ?

I realize that it is possible to have non-zero \mathbf{J} even when ρ is a constant over time, e.g. a wire carrying a constant current in a loop. Here charge is moving, but the density of the charge at any one point in

space remains the same over time, because in any given sub-volume within the wire, as much charge is entering that volume per unit time as is leaving per unit time.

Answer from Google to this search phrase: “are there versions of Maxwell’s equations that combine charge density and current density”

Yes, Maxwell’s equations can be formulated in a way that combines charge density and current density into a single entity called the four-current. This formulation is part of the covariant formulation of electromagnetism, which is a more elegant and relativistic way to express Maxwell’s equations.

More details available on the Wikipedia page “Covariant formulation of classical electromagnetism” [6].

Since this is a four-vector, it has just as much “information” as the scalar field ρ plus the vector field J . It makes sense that if someone had found a way to combine ρ and J into a single time-varying scalar field, they would have done so long ago.

3 Gradient

Gradient has been generalized to many coordinate systems other than the 3-dimensional Cartesian coordinate system \mathbb{R}^3 , but I will focus on \mathbb{R}^3 . The Wikipedia page on Gradient [7] is not too bad for me, as long as I skim over the parts that generalize it to other coordinate systems.

Griffiths [3] Section 1.2.2 “Gradient” is good at giving the definition and some useful examples and properties of the gradient. He defines ∇T this way, where $\hat{\mathbf{x}}$, $\hat{\mathbf{y}}$, and $\hat{\mathbf{z}}$ are the unit vectors in the direction of the three coordinate axes:

$$\nabla T \equiv \frac{\partial T}{\partial x} \hat{\mathbf{x}} + \frac{\partial T}{\partial y} \hat{\mathbf{y}} + \frac{\partial T}{\partial z} \hat{\mathbf{z}}$$

where:

- $T : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a scalar function that is continuous and differentiable

Some properties of the gradient:

- $\nabla T(x, y, z)$ is the gradient evaluated at a position given by (x, y, z) . The vector points in a direction that function T increases most quickly.
- The function T can be approximated at points \mathbf{r} near $\mathbf{r}_0 = (x, y, z)$ by the linear function $T(\mathbf{r}_0) + (\mathbf{r} - \mathbf{r}_0) \cdot \nabla T(\mathbf{r}_0)$
- The instantaneous rate of change of T in direction \mathbf{u} (a unit vector) from \mathbf{r}_0 is $\mathbf{u} \cdot \nabla T(\mathbf{r}_0)$. Note that it is always 0 in a direction perpendicular to $\nabla T(\mathbf{r}_0)$, and the negative of the magnitude of $\nabla T(\mathbf{r}_0)$ in the opposite direction.
- Consider a “level set”, i.e. a surface defined by all of the points \mathbf{r} where $T(\mathbf{r}) = c$ for some constant c . Then ∇T evaluated at any point on that surface, is normal to the surface.

4 Galilean relativity

In classical Newtonian mechanics, physical laws are invariant under Galilean relativity, i.e. any laws that hold in one inertial frame also hold in all other inertial frames, i.e. those that move with a constant velocity relative to the first frame.

This applies to Newton’s three laws of motion, for example.

It does not apply to measurements of quantities that depend upon velocity, e.g. an object’s momentum or kinetic energy.

However, the properties that momentum is conserved, and energy is conserved, is preserved across inertial frames, even if the absolute values of the kinetic energy and momentum differ between frames.

If there are forces that are velocity-dependent, e.g. a frictional force resisting an object traveling through a medium like air or water that is a function of velocity relative to that medium, then as long as that force is always written as a function of the relative velocity between the object and the medium, then note that the measurements of such relative velocities are the same in different inertial frames.

For other examples and some mathematical derivations, see [2].

See this video [1] that gives a similar mathematical demonstration that Newton’s second law is invariant under Galilean relativity, but Maxwell’s equations are *not*.

5 Experiments

- 1848-49: Fizeau’s measurement of the speed of light in air [4]
- 1851: Fizeau’s experiment to measure the relative speed of light in a moving medium (water) [5]

6 Questions

- Historically, how was aberration of light distinguished/separated from other effects like parallax?

7 Index of A History of the Theories of Aether and Electricity

All page numbers are of the form “I(number)” for a page number in Volume I, or “II(number)” for a page in Volume II.

Done for I, pp. 1-93

- aberration of light, adumbration by Pardies: I22
- aberration of light, discovery by Bradley: I94
- action at a distance: I5, I28, I30-31, I36, I48, I50, I51, I59
- action, as in principle of least action: I97
- Aepinus, Franz Ulrich: I51
- aether: I5, I13, I19, I20, I30, I95, I98
- Ampère, André-Marie: I83
- ammunitionm, finite, argument against corpuscular theory of light: I97
- atomism: I13
- Bartholin, Erasmus: I25
- Berzelius, Jöns Jacob: I78
- Biot, Jean-Baptiste: I82
- birefringence, discovered 1669: I25
- Bologna stone: I20
- Bradley, James: I94
- Carlisle, Anthony: I73
- Cavendish, Henry: I54
- Cavendish, Charles: I54
- colors of light, Descartes’ theory: I9
- colors of light, Hooke’s theory: I16
- colors of light, Newton’s theory: I17
- colors of thin plates: I13, I21, I102
- conductor, origin of name: I42
- corpuscular theory of light: I10, I20, I23, I31, I97
- corpuscular theory of light, argument against: I97, I100
- Coulomb, Augustin: I57

- Davy, Humphry: I74
- de Maricourt: see entry alphabetized as “Maricourt”
- Desaguliers, Jean Théophile: I41
- diamond emitting light when rubbed or heated: I14
- dielectric: I52
- diffraction: I14, I20
- du Fay: see entry alphabetized as “Fay”
- epistemological rationalism: I7
- electric charge, conservation of: I47
- electric charge, induction of: I52
- electric current effect on magnets: I81-83
- electric current effect on magnets, and vice versa: I83
- electric current carrying wires attract or repel: I83
- electric current through human body, used as sensory evidence: I45, I55, I67, I71, I73
- electric current through wire, similarity with flow of heat through wire: I91
- electric fluid: I42
- electric fluid, one-fluid theory: I50
- electric fluid, two-fluid theory: I58
- electric force, how it varies with distance: I53, I54, I57
- electric fluid in conductor is all at surface at equilibrium: I51, I53, I61
- electric pile: I71-73
- electric pile, experiments varying the strength of: I89
- electrolysis of water and other chemicals, both by continuous current and frictional electricity: I74
- electrical conducting power of different materials and shapes, generated by frictional electricity: I55
- electrical conducting power of different materials, lengths, cross-sectional areas, and temperatures, generated by electric piles: I90
- electrification, discovery of two kinds by du Fay in 1733: I44
- electrification only at the surface, discovery in 1729: I43, I49
- electrical emanations: I36
- electricity, discoveries by Gilbert, and naming: I35
- electricity, discovery of transfer between electrified bodies: I42
- electricity, ordinary matter (without electric fluid) repels other ordinary matter: I52
- electricity, resinous: I44
- electricity, theory of electric fluid that electrified bodies have a surplus or deficiency of when electrified: I46, I49
- electricity, vitreous: I44

- electrodynamics, origin of name: I85
- electrodynamics, Ampère's theories and mathematical analysis of: I83-88
- electrostatic mathematics by Poisson: I61-62
- extraordinary ray: see birefringence
- du Fay, Charles-François: I43
- fits of easy transmission and easy reflection, theory of, about light: I22
- Franklin, Benjamin: I45, I47
- frogs in electrical experiments: I67-69
- Galvani, Luigi: I67
- galvanism, i.e. electric current caused by touching of different metals: I70
- galvanism, proof tha it was electrical in nature: I71
- Gilbert, William: I34
- Gravesande: see entry alphabetized as "s Gravesande"
- Gray, Stephen: I41
- Green, George: I65
- Grene's theorem: I65
- gravity: I28, I31
- heat: I20, I38
- heat, from focused light: I23
- heat vs. light, heat absorbed by glass but not light: I40
- heat vs. light, heat transmitted by different colors of light: I99
- Herschel, Sir William: I99
- Huygens' principle of light wave propagation: I24
- induction of electric charge: see electric charge, induction of
- infrared light: I99
- interference of waves, applied to water and light: I101
- interference of light waves, applied to colors of thin plates: I102
- inter-phenomena: I30
- Leyden phial, Leyden jar: I45
- magnet, use as a compass, as early as 1186 in Europe: I33
- magnet, Earth is a magnet: I34
- magnetic fluid, two-fluid theory: I57, I59
- magnetic force between poles decreases as square of distance: I56, I59
- magnetic moment: I63
- magnetic poles, discovery and naming: I33
- magnetism, discoveries by Gilbert: I35

- magnetism, distinguishing behavior from gravity: I55
- magnetism, created by lightning: I80-81
- de Maricourt, Pierre, also known as Peregrinus: I33
- de Maupertuit, Peirre-Louis-Moreau: I96
- mechanical philosophy: I6
- Michell, John: I56
- van Musschenbroek, Pieter: I45
- Neckam, Alexander: I33
- Newton's rings: I13, I102
- Nicholson, William: I73
- Oersted, Hans Christian: I80
- Ohm, Georg Simon: I90
- Ohm's law: I92
- opacity: I25
- plenum: I5, I13
- Poisson, Siméon Denis: I60
- polarized light: I19, I27
- poles of magnets: see magnetic poles
- potential: I65
- Priestley, Joseph: I53
- principle of least time: I12
- reflection of light: I10, I24
- refraction of light: I10, I15, I17, I20, I25, I95, I101
- resinous: see electricity, resinous
- Ritter, Johann Wilhelm: I74, I76
- Roemer, Olaf: I22
- 's Gravesande, Wilhelm Jacob: I37
- Savart, Félix: I82
- Snell, Willebrord: I10
- sound as periodic vibrational waves: I21
- speed of light propagation: I22
- thermo-electric circuit: I89
- Torricelli: I24
- ultraviolet light: I100
- wave theory of light: I14, I23
- Wilcke, Johan Carl: I51

- vacuum, light traveling through: I24
- van Musschenbroek: see entry alphabetized as “Musschenbroek”
- vitreous: see electricity, vitreous
- void: I5, I13
- Volta, Alessandro: I69
- voltaic polarization: I76
- Watson, William: I45
- wave theory of light: I100-101
- waves, interference of: see “interference”
- Young, Thomas: I100

7.1 Definitions of some key terms

aberration of light

diffraction of light Diffraction of light is the bending of light waves as they pass around the edges of an obstacle or through a narrow opening. This bending causes the light to spread out and illuminate areas where a shadow would otherwise be expected. Diffraction is a wave phenomenon, meaning it’s a characteristic behavior of waves, including light, sound, and water waves.

Bending Around Obstacles: When light encounters an obstacle, like the edge of a barrier or a small opening, it doesn’t just travel in a straight line. Instead, the light waves spread out and bend around the obstacle, a phenomenon known as diffraction.

Interference: Diffraction often occurs alongside interference, where waves combine, either constructively (creating brighter areas) or destructively (creating darker areas). This combination of diffraction and interference is responsible for the characteristic patterns observed in diffraction experiments.

Wavelength Dependence: The extent of diffraction depends on the wavelength of the light and the size of the opening or obstacle. When the wavelength of the light is comparable to the size of the opening or obstacle, diffraction is more pronounced.

Examples: A classic example is the single-slit experiment where light passes through a narrow opening and creates a diffraction pattern on a screen behind it. Another example is the silver lining seen around clouds in the sky, caused by sunlight diffracting through water droplets or ice crystals.

diffusion of light Light diffusion is the process where light scatters or spreads out in various directions, rather than traveling in a straight line. This happens when light encounters a medium or surface with irregularities, causing it to scatter at different angles. This scattering results in a softer, more even light, reducing harsh shadows and glare.

Scattering: When light hits a rough or uneven surface (like a matte surface or a translucent material), it scatters in multiple directions instead of reflecting at a single angle.

Softening: This scattering effect diffuses the light, making it appear softer and more gentle.

Reduced glare: By scattering the light, diffusion reduces the intensity of the light source, minimizing glare and harsh shadows.

Examples:

- Transmission diffusion: Light passing through a frosted glass or plastic panel.
- Reflection diffusion: Light bouncing off a whiteboard or textured wall.
- Softboxes and diffusers: Professional tools like softboxes and diffusion panels are used in photography and filmmaking to soften and spread out light.
- Overcast days: The clouds act as a natural diffuser, scattering sunlight and creating soft, even light.

dispersion of light Dispersion of light is the phenomenon where white light, upon passing through a medium like a prism, separates into its constituent colors (like in a rainbow). This happens because different wavelengths (colors) of light bend at slightly different angles when they enter and exit the medium. When light passes from one medium to another (like from air to glass), it bends (refracts). The amount of bending depends on the wavelength of the light. Shorter wavelengths (like violet and blue) bend more than longer wavelengths (like red and orange).

induction of electric charge

induction of electric current

interference

polarization of light

reflection

refraction Refraction of light is the bending of light as it passes from one medium to another, caused by a change in the speed of light. This bending occurs because light travels at different speeds in different materials. For example, light travels slower in water than in air, causing it to bend when entering or exiting water.

7.2 Noteworthy events in aether vs. action at a distance

Vol. I, p. 28:

Newton claimed nothing more for his discovery than that it provided the necessary instrument for mathematical prediction, and he pointed out that it did not touch on the question of the mechanism of gravity. As to this, he conjectured that the density of the aether might vary from place to place, and that bodies might tend to move from the denser parts of the medium toward the rarer; but whether this were the true explanation or not, at any rate, he said, to suppose 'that one body may act upon another at a distance through a vacuum, without the mediation of anything else, ... is to me so great an absurdity, that I believe no man, who has in philosophical matters a competent faculty for thinking, can ever fall into.'

According to Wikipedia [9], Newton wrote this in 1692 in a letter to Richard Bentley, 5 years after *Principia* was first published in 1687. I do not know if Newton said anything similar in *Principia* itself. Very likely he did not, or else it would be more widely quoted on this topic.

Vol. I, p. 30:

The rejection of the inverse-square law of gravitation by the French Cartesians antagonised the younger disciples of Newton to such an extent that the latter hardened into opposition not only to the vortices but to the whole body of Cartesian notions, including the aether. In the second edition (1713) of the *Principia*, there is a preface written by Roger Cotes (1682-1716), in which the Newtonian law of action at a distance is championed as being the only formulation of the facts of experience which does not introduce unverifiable and useless suppositions.

Vol. I, p. 31:

When the eighteenth-century natural philosophers found by experience that the Newtonian law was marvellously powerful, yielding formulae by which practically every observable motion in the solar system could be predicted, while on the other hand the search for an explanation of inter-phenomena led to no practical results, opinion set in favour of Cotes's attitude, which came to prevail widely; and in the middle of the century R. G. Boscovich (1711-87) ... attempted to account for all known physical effects in terms of action at a distance between point particles.

References

- [1] History of Special Relativity (Part 1) - Galilean Invariance & Maxwell's Equations, 2018. URL <https://www.youtube.com/watch?app=desktop&v=P79Ur8LVcy8>.
- [2] Galilean Transformation, 2025. URL https://qiweb.tudelft.nl/mecharela/galilean_transformation.
- [3] David J. Griffiths. *Introduction to Electrodynamics*. Pearson Education, 4th edition, 2013.
- [4] Wikipedia. Fizeau's measurement of the speed of light in air, 1849. URL https://en.wikipedia.org/wiki/Fizeau%27s_measurement_of_the_speed_of_light_in_air.
- [5] Wikipedia. Fizeau experiment, 1851. URL https://en.wikipedia.org/wiki/Fizeau_experiment.
- [6] Wikipedia. Covariant formulation of classical electromagnetism, 2025. URL https://en.wikipedia.org/wiki/Covariant_formulation_of_classical_electromagnetism.
- [7] Wikipedia. Gradient, 2025. URL <https://en.wikipedia.org/wiki/Gradient>.
- [8] Wikipedia. Nabla Symbol, 2025. URL https://en.wikipedia.org/wiki/Nabla_symbol.
- [9] Wikipedia. Newton's law of universal gravitation, 2025. URL https://en.wikipedia.org/wiki/Newton%27s_law_of_universal_gravitation.