

Aim:

Design an algorithm and implement using a C program which finds the **sum** of the **infinite series**

$$1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots,$$

Print the result as shown in the example.

Sample Input and Output:

Enter the value of x and n: 4 5
sum = 3.666667

Source Code:

infinite.c

```
#include<stdio.h>
#include<math.h>
int main()
{
    int x,n,m,i=0,fact=1;
    float k,sum;
    printf("Enter the value of x and n: ");
    scanf("%d %d",&x,&n);
    while(i<=n)
    {
        if(i%2==0)
        {
            fact=1;
            for(m=1;m<i+1;m++)
            {
                fact=fact*m;
            }
            k=pow(x,i)/fact;
        }
        if(i%4!=0)
        {
            fact=1;
            for(m=1;m<=i;m++)
            {
                fact=fact*m;
            }
            k=- (pow(x,i))/fact;
        }
        sum=sum+k;
        i=i+2;
    }
    printf("sum = %f\n",sum);
}
```

Test Case - 1
User Output
Enter the value of x and n: 4 5
sum = 3.666667

Test Case - 2
User Output
Enter the value of x and n: 12 5
sum = 793.000000