

Queues with a Dynamic Schedule

John Gilbertson

October 2016

Background

Static Schedules

Dynamic Schedules

Schedule Comparison

Simulation Studies

Background

Queues with Scheduled Arrivals

-



Assumptions

- Single server
- Service times are independent exponential RVs with mean μ
- All customers arrive punctually
- Total number of customers is fixed

Static vs. Dynamic



Static Schedules

Objective Function

- Denote the expected waiting time of customer i by w_i

$$\mathbb{E}[\text{total customers' waiting time}] = \sum_{i=1}^n w_i$$


- Denote the interarrival time between customer i and customer $i + 1$ by x_i

$$\mathbb{E}[\text{server availability time}] = \sum_{i=1}^{n-1} x_i + w_n + \mu$$

- Objective is to minimise a linear combination of these times

$$\phi(\mathbf{x}) = (1 - \gamma) \sum_{i=1}^n w_i + \gamma \left(\sum_{i=1}^{n-1} x_i + w_n + \mu \right)$$

Model for 15 Customers



- **Dome-shape:** increase for first customers, remain constant, then decrease for last few customers
- As relative cost of server availability time (γ) increases, customers arrive earlier

Dynamic Schedules

Markov Decision Process

-

Erlang Distribution

- Waiting time of customer in position $r + 1$

$$X \sim \text{Erlang}(r, \mu)$$

- Distribution function:

$$F(a; r) = \mathbb{P}(X \leq x) = \begin{cases} 0 & \text{for } x = 0 \\ 1 - \sum_{i=0}^{r-1} \frac{1}{i!} \left(\frac{x}{\mu}\right)^i e^{-\frac{x}{\mu}} & \text{for } x > 0 \end{cases}$$

- Conditional expectation:

$$\mathbb{E}[X \mid X \leq a] = \mu r \times \frac{F(a; r+1)}{F(a; r)}$$

- Suppose $Y \sim \text{Exp}(\mu)$ independent of X

$$\mathbb{E}[X \mid X \leq a, X + Y > a] = \frac{ar}{r+1}$$

Schedule Comparison


Expected Cost Comparison



Simulation Studies




Schedule Cost



■

Customer Arrival Times



Customer Waiting Times

