

Direct Tests of Black Hole Accretion Rate Prescriptions: I. Bondi Accretion at Different Scales

JAMES AGOSTINO,¹ MING-YI LIN,¹ NATASHA JONES,¹ ANNE M. MEDLING,¹
LORETO BARCOS-MUÑOZ,^{2,3} DANIEL ANGLÉS-ALCÁZAR,⁴ CLAUDIO RICCI,^{5,6} GEORGE C. PRIVON,^{7,8,9}
VIVIAN U,^{10,11} PAUL TORREY,^{12,13,14} PHILIP F. HOPKINS,¹⁵ AND CLAIRE MAX¹⁶

¹*Ritter Astrophysical Research Center and Department of Physics & Astronomy, University of Toledo, Toledo, OH 43606, USA*

²*National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA, 22903, USA*

³*Department of Astronomy, University of Virginia, 530 McCormick Road, Charlottesville, VA, 22903, USA*

⁴*Department of Physics, University of Connecticut, 196 Auditorium Road, U-3046, Storrs, CT 06269, USA*

⁵*Instituto de Estudios Astrofísicos, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Avenida Ejercito Libertador 441, Santiago, Chile*

⁶*Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China*

⁷*National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903*

⁸*Department of Astronomy, University of Florida, P.O. Box 112055, Gainesville, FL 32611, USA*

⁹*Department of Astronomy, University of Virginia, 530 McCormick Road, Charlottesville, VA 22904, USA*

¹⁰*4129 Frederick Reines Hall, Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA*

¹¹*IPAC, California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125, USA*

¹²*Department of Astronomy, University of Virginia, 530 McCormick Road, Charlottesville, VA 22904*

¹³*Virginia Institute for Theoretical Astronomy, University of Virginia, Charlottesville, VA 22904, USA*

¹⁴*The NSF-Simons AI Institute for Cosmic Origins, USA*

¹⁵*TAPIR, Mailcode 350-17, California Institute of Technology, Pasadena, CA 91125, USA*

¹⁶PLACEHOLDER

ABSTRACT

We present spatially resolved parsec-scale measurements of nuclear conditions (gas density and kinetic temperature) relevant for black hole accretion rate predictions in the Seyfert 2 galaxy, NGC 1068. We inject these parameters into the prescription for a Bondi-like accretion model, then compare the resulting accretion rate prediction to the empirical accretion rate derived from hard X-ray observations. Cosmological simulations have spatial resolution ranging from ~ 10 pc to \sim kpc scales, and so for reasonable comparison we test these accretion rate predictions in pixel-sized radial steps out to 500 pc. Compared to warm H₂ gas, CO gas is the dominant mass carrier close to the SMBH. We find that the Bondi accretion rate (\dot{M}_{Bondi}) of cold molecular gas alone (measured using CO) overestimates the true accretion rate by up to 13 dex in a small aperture ($r \lesssim 5$ pc) around the black hole, and by at least 9 dex inside large apertures ($r \lesssim 500$ pc). These results are the first in a series of direct tests of accretion rate prescriptions, and they suggest that using a Bondi accretion formalism to model supermassive black hole accretion in Seyfert 2 galaxies may lead to overestimated accretion rates in simulations.

1. INTRODUCTION

Supermassive black holes (SMBHs), despite their small gravitational radius of the sphere

of influence ($1\sim 100$ pc), are thought to be a key piece of the connection between pc and kpc scales of galaxy evolution. Observations of galaxies with active galactic nuclei (AGN) have shown both directly and indirectly that AGN can inject energy into their surrounding environments, which can ultimately quench or in some cases trigger star formation (see [Fabian 2012](#) for an observational review; [Mercedes-Feliz et al. \(2023\)](#) for a simulated example).

Active galactic nuclei do not only reside in the central part of galaxies, they may also significantly impact several global properties of galaxies and their surrounding inter-galactic media, allowing us to indirectly infer their influence on those observables. Relationships between black hole mass and global galaxy properties, like the velocity dispersion of stars in the galactic bulge, have been well-calibrated and show tight correlations (see [Kormendy & Ho 2013](#); [McConnell & Ma 2013](#) for reviews). These correlations suggest that AGN radiative feedback, which in part depends on black hole mass, may leave an imprint on bulge stellar velocity dispersion (see [Ferrarese & Merritt 2000](#); [Gebhardt et al. 2000](#) for seminal studies) but fueling regulation ([Escala 2007](#); [Chen et al. 2013](#); [Anglés-Alcázar et al. 2013](#); [Anglés-Alcázar et al. 2017](#)) and non-causal mass averaging through mergers ([Peng 2007](#); [Hirschmann et al. 2010](#); [Jahnke & Macciò 2011](#)) have also been proposed as plausible drivers of black hole-galaxy scaling relations. Star formation in massive halos is suppressed (e.g. in [Behroozi et al. 2013](#); [Torrey et al. 2014](#)), which could be caused by heating of the interstellar medium (ISM) from AGN feedback. In the high energy regime, a discrepancy is found between the observed and expected correlations between X-ray luminosities and temperatures of gas in the intra-cluster medium (called the L_X -T relation, see [Mushotzky 1984](#); [Markevitch 1998](#)). This discrepancy suggests that gas in the intra-cluster medium evolves dif-

ferently from dark matter; energetics input by host AGN could be a factor as to why.

Alongside indirect cases of the impact of AGN feedback on galaxy formation histories, the direct effects of AGN on the ISM have been observed for decades. Since more than 100 years ago ([M87](#); [Curtis 1918](#)) radio jets powered by a central SMBH have been seen to extend up to ~ 0.9 Mpc outside from their host galaxies (e.g. Centaurus A; [Burns et al. 1983](#)). Outflows driven by these SMBHs have been observed in the process of depleting the ISM at outflow rates of $700 M_\odot \text{ yr}^{-1}$ (e.g. in Mrk 231; [Feruglio et al. 2010](#)). NGC 1068, which is the test case in the rest of this Paper has a complex and well studied AGN-driven outflow that has been observed to impact its ISM on sub-kpc scales (e.g. [Wilson & Ulvestad 1983](#); [Müller-Sánchez et al. 2011](#); [García-Burillo et al. 2014](#); [Saito et al. 2022](#); [Hviding et al. 2023](#); [Holden & Tadhunter 2023](#); [Gallimore & Impellizzeri 2023](#); [Mutie et al. 2024](#); [Hagiwara et al. 2024](#)).

The energy output of an AGN is driven by mass accretion onto its accretion disk, fueled by inflows in the nuclei of galaxies. This gas then accelerates to speeds of up to $> 0.1c$ in the accretion disk, and that disk can power radiative outflows. These energetics interact with the ISM, with the effects of which we call feedback. The direct observational feedback can be classified as two mechanisms: radiative (quasar mode) or kinematic (radio mode) ([Fabian 2012](#)). In the quasar mode, occurring when the black hole accretes mass quickly, photons from the accretion disk couple to the ISM, transferring momentum in a powerful jet. In radio mode, accretion onto the disk is slower, and the primary feedback mechanism is in the form of collimated radio jets that typically appear narrower than quasar-mode jets (see [Cielo et al. 2018](#) for a simulated comparison between the feedback of the two modes). Both modes can drive outflows, but the quasar-mode is thought to start the quenching

process (the spatial extent of which grows over time) and then the radio-mode maintains that quenched state (see Fabian 2012 and Morganti 2017 for reviews).

As simulations of galaxy evolution have been informed by increasingly detailed observations, theorists have begun to study the physical mechanisms that drive AGN feedback and how that feedback impacts the simulated host galaxies. Dubois et al. (2013a) (see also Dubois et al. 2013b; Taylor & Kobayashi 2015) examined how AGN jets impact cold gas and transform blue, disk galaxies into red ellipticals. Building on these studies, Rosas-Guevara et al. (2015), who simulated accretion in galaxies of varied halo mass, find that in galaxies with M_{Halo} above $10^{11.5} M_{\odot}$, as was observed in Behroozi et al. (2013), star formation is suppressed by AGN feedback. Valentini et al. (2020) perform a suite of cosmological simulations in which they couple AGN feedback to different phases of the ISM. They find, in part, that energy output from the AGN as feedback must couple with both the cold and hot phases in order to avoid excessive SMBH growth.

As is seen in both observations and simulations, global galaxy properties can be affected by accretion-dependent feedback. Theorists have attempted to model the physical processes causing those properties to change. Williamson et al. (2020) perform radiation hydrodynamics modeling of the 1-100 pc scales in a nuclear region of a simulated AGN host. They demonstrate that increasingly polar winds are produced when anisotropic radiation from the AGN shifts the mass distribution of the outflow originating from the AGN. Meenakshi et al. (2022) simulated the direct interaction between AGN jet-induced outflows on 2 kpc scale and the ISM and found shocked emission fronts in the ISM that could be responsible for stunting star formation. On $r < 1$ pc scale, Wada et al. (2023) were able to induce radiation-driven dusty out-

flows which impact the ISM as they continue on their outward paths. Tying the small and large scales together has been an ongoing challenge.

Due to computational constraints, large-scale cosmological simulations that can model hundreds of Mpc^3 at a time are not able to directly resolve the physical processes that drive gas accretion at $<< 1$ pc scales where accretion takes place, and so sub-grid prescriptions for black hole accretion and its subsequent feedback must be adopted.¹ The ‘sub-grid’ is defined as the region below the gridded resolution of the simulation. Unfortunately, there is no unified model for these sub-grid physics, and different studies use different accretion prescriptions. The most commonly applied prescription is the one described in Bondi (1952), often referred to as the Bondi accretion prescription. The equation for the mass accretion rate follows the form:

$$\dot{M}_{\text{Bondi}} = \frac{4\pi G^2 M_{\text{BH}}^2 \rho}{(c_s^2 + v_{\text{rel}}^2)^{3/2}} \quad (1)$$

where G is the gravitational constant, M_{BH} is the mass of the black hole, ρ is the gas density, c_s is the sound speed, and v_{rel} is the relative velocity of the gas. In the pure Bondi case, the gas is assumed to be stationary relative to the galactic potential, so v_{rel} is zero. This model is theoretically predicated on gas free-falling onto the SMBH once it reaches the Bondi radius, $R_{\text{Bondi}} = 2GM_{\text{BH}}/c_s^2$. The Bondi radius is where the escape velocity of the SMBH (based on its mass) equals the sound speed of the gas in the nuclear region. The physical scale of the Bondi radius is typically of order 0.1-300 pc if we assume c_s of 400 km s^{-1} and SMBH mass range of $10^6 \sim 10^9 M_{\odot}$. Some large scale cosmological simulation suites that use a pure

¹ Although there is also much variation in AGN feedback prescriptions, this program will focus on discussing the accretion rate prescriptions, on which all feedback depends.

Bondi prescription to account for SMBH accretion, including *MassiveBlack-II* (Khandai et al. 2015) and *IllustrisTNG* (Weinberger et al. 2017; Pillepich et al. 2018a).

Physically, the issue with the Bondi accretion formalism is that it ignores both the angular momentum of the gas and interactions due to self-gravity between the gas, stellar, and dark matter components, which is only appropriate in the case of hot, virialized gas (Hobbs et al. 2012; Negri & Volonteri 2017; Anglés-Alcázar et al. 2021). Recent studies have shown that gas and other accreting material still has angular momentum inside what may be the Bondi radius, particularly in gas-rich mergers or galaxies with Seyfert AGN (e.g. in Davies et al. 2004; Hicks et al. 2013; Medling et al. 2014; Lin et al. 2016), and so Bondi accretion timescales may be much shorter than in reality where angular momentum delays accretion. Feedback from the AGN in such models self-regulates this rapid growth (Anglés-Alcázar et al. 2015).

Because of the M_{BH}^2 dependence of accretion rate in Bondi accretion prescriptions, low mass BH seed growth is suppressed such that BHs do not grow quick enough to match their expected mass at corresponding redshifts. To account for this discrepancy, some large-scale cosmological simulation suites adjust the accretion physics by using modified versions of Bondi accretion. The prescription in the *Illustris* (the predecessor to *IllustrisTNG*; Vogelsberger et al. 2013; Genel et al. 2014) and *Magneticum Pathfinder* hydrodynamical simulation suites (Hirschmann et al. 2014; Bocquet et al. 2016; Dolag et al. 2016) modify Bondi by multiplying Equation 1 by a constant (unitless) ‘boost’ factor α (following the prescription of Springel et al. 2005; Di Matteo et al. 2005; Springel & Hernquist 2005). The boost factor is used to account for the volume average of the Bondi-rates for both the cold and hot phases in the simulations and typically has a value = 100. Another large-scale cosmologi-

cal model, *Horizon-AGN* (Dubois et al. 2016), uses an α similar to *Illustris* and *Magneticum*, but instead of a constant value, their boost factor (following the prescription from Booth & Schaye 2009; see also Dubois et al. 2012) is $\alpha = (\rho/\rho_0)^2$ or $\alpha = 1$ for densities above and below the threshold for star formation respectively. *EAGLE* (Schaye et al. 2015) uses a pure Bondi prescription alongside the heuristic correction from Rosas-Guevara et al. (2015) to account for variable angular momentum of accreting gas. Another approach, used by the large-scale *Romulus* suite (Tremmel et al. 2017) is to adjust the Bondi accretion rate depending on the motion of the simulated gas particles. In *Romulus*, if the smallest relative velocity (which they equate to v_{bulk} , the bulk motion of the gas) of the gas particle closest to the SMBH is faster than the rotational velocity of the gas, they replace the relative velocity of the SMBH (in Equation 1) with v_{bulk} and multiply the Bondi rate by a density-dependent boost factor similar to *Horizon-AGN*.

Bondi or Bondi-like accretion prescriptions are the most commonly used, but theorists have also designed accretion prescriptions with very different underlying physics. One large-scale simulation ($[100 h^{-1} \text{ Mpc}]^3$ volume) suite that in part uses one of these prescriptions is *SIMBA* (Davé et al. 2019a). In *SIMBA*, pure Bondi accretion is still applied for hot gas accretion where, as we mentioned, it is most appropriate. But, they then apply a torque-limited accretion formalism for the cold gas where instabilities in the disk drive mass inflow (Hopkins & Quataert 2011; Anglés-Alcázar et al. 2017).

Understanding if and in which cases different sub-grid prescriptions are accurately estimating accretion rates onto the black holes of galaxies is critically important to cosmological simulations and conclusions drawn from them. Without an accurate prescription for accretion over time, simulations cannot accurately implement

the impact of AGN feedback, and as such may have incorrect outcomes with regards to galaxy formation and evolution. Depending on the assumed accretion prescription, simulations find that BH scaling relations are driven either by feedback efficiency (in Bondi-like models), or accretion efficiency (in a torque-driven model; see Anglés-Alcázar et al. 2021 for further discussion). Theorists’ conclusions on which physics drive the co-evolution between BH mass and global galaxy properties is directly dependent on which accretion model is implemented. Determining which accretion formalism is most appropriate in which circumstances is critical to understanding BH-galaxy co-evolution in our Universe.

Observationally testing how black hole accretion rate prescriptions perform has only become possible in recent times. In this study, in the prototypical Seyfert 2 galaxy NGC 1068, we directly measure the parameters that go into Bondi accretion, ρ_{gas} and c_s , on physical scales ranging from 2-500 pc. We then plug these measured parameters into the pure Bondi accretion prescription as a function of radius to mimic what a simulation at that resolution would estimate for the black hole accretion rate. Finally, we test these predicted Bondi accretion rates against empirically derived accretion rates using hard (14-195 keV) X-ray data from The Burst Alert Telescope (BAT) AGN Spectroscopic Survey (BASS) (Ricci et al. 2017). The BAT instrument (Barthelmy et al. 2005; Krimm et al. 2013) on *Swift* (Gehrels et al. 2004) is a hard X-ray detector that surveys the entire sky, reporting X-ray sources to within 1-4 arcmin accuracy.

In this work, we use cosmological parameters of $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.28$, and $\Omega_\Lambda = 0.72$ (Hinshaw et al. 2009). To calculate spatial scales and luminosity distance to NGC 1068 we use Ned Wright’s Cosmology Calculator (Wright 2006).

2. NUCLEAR STRUCTURE OF NGC 1068

NGC 1068 is one of the most-studied prototypical Seyfert galaxies, and as such a wealth of information has already been published about its nuclear structure. The studies described here are not an exhaustive list, but are included to provide context relevant to our analysis.

Under our 2.2 pc resolution, NGC 1068 hosts a water maser that is thought to originate from the accretion disk on much smaller ($<0.1 \text{ pc}$) scales. Greenhill et al. (1996) observed the maser with very long baseline interferometry (VLBI) using both the Very Long Baseline Array and Very Large Array to achieve 0.65 pc resolution. They used the velocity gradient of the maser emission to infer a rotational velocity of the gas, and in turn constrain M_{BH} . Kumar (1999) modeled the 0.65-1.1 pc disk from which the maser emission is thought to be ejected from. The clumps in their disk model interact with each other, leading to eventual accretion onto the SMBH.

On slightly larger scales with near and mid infrared interferometry, multiple authors were able to resolve a two-component dusty torus (Jaffe et al. 2004; Raban et al. 2009). One component is smaller and more elongated ($1.35 \times 0.45 \text{ pc}$) in size than the other ($3 \times 4 \text{ pc}$). In the nucleus of Circinus, another Seyfert 2 AGN, Tristram et al. (2014) also found a two-component dusty torus. Images like these that showed structure inconsistent with the prior, observationally-defined, Type 2 classification of these galaxies (unless foreground extinction was applied) fundamentally challenged the AGN unification model (Antonucci 1993).

Gómez Rosas et al. (2022) used sub-pc resolution observations of NGC 1068 taken with the MATISSE/ESO/VLTI interferometer between 3 and 13 μm to map the dust temperature distribution of the dust observed in the previously mentioned studies. They confirm an optically thick pc scale dusty structure and a second, less




Figure 1. Nuclear region of NGC 1068 in the CO(2-1) flux (left, from ALMA), $2.2\mu\text{m}$ line + continuum (middle), and the continuum subtracted rovibrational H_2 1-0 S(1) transition (right), described in Section 3. The CO(2-1) moment 0 map is masked below $3\times\text{rms}$ and the red box in the CO(2-1) moment map represents the field of view of the NIR data. The very small, red ellipse in the bottom left represents the beam size of the ALMA data (41×30 mas). All three images show flux peaks at the AGN’s location and both the CO and H_2 maps have enhanced emission in the CND ring. The red circle in the H_2 1-0 S(1) moment map represents the aperture in which T_{kin} is calculated in Figure 3.

378 optically thick disk that extends to at least 10
 379 pc. [García-Burillo et al. \(2019\)](#) (who in part
 380 use the same ALMA data as we describe in Sec-
 381 tion 3.2) find a 14 pc CO(2-1) nuclear disk with
 382 a PA (~ 110 - 140 deg) aligned with the water
 383 maser disk PA (140 deg). Also in [García-Burillo](#)
 384 [et al. \(2019\)](#), they observe the CND, which as
 385 can be seen in Figure 1, has a gas deficit inside
 386 the outer ring in its central ~ 130 pc region.

387 To resolve the kinematics of the 10 pc in-
 388 ner disk (often referred to as the torus) and
 389 outer ring, [Imanishi et al. \(2020\)](#) observe both of
 390 these scales using the bright (relative to CO(2-
 391 1)) HCN $J=(3-2)$ and HCO $^+$ $J=(3-2)$ tran-
 392 sitions with ALMA at 1.4 pc resolution. They
 393 find that the torus as observed with these dense
 394 gas tracers rotates in the opposite direction with
 395 respect to the outer ring. This is particularly
 396 surprising because the water maser emission is
 397 rotating in the same direction as the outer ring
 398 rather than the torus it is physically closer to
 399 (see Figure 1 of [Imanishi et al. 2020](#)). In the

400 work of [García-Burillo et al. \(2019\)](#), the authors
 401 find that a “significant part” of the observed
 402 counter-rotation in CO(2-1) can be attributed
 403 to a northern AGN-driven wind. To make
 404 a more robust determination though, [García-](#)
 405 [Burillo et al. \(2019\)](#) say that higher resolution
 406 data is required so that the outflowing compo-
 407 nent can be better disentangled from the rotat-
 408 ing component.

409 Outflows originating from the AGN can serve
 410 to regulate black hole accretion, and NGC 1068
 411 hosts a complex outflow in the NE direction,
 412 perpendicular to the nuclear disk. The largest
 413 outflow component is seen as the radio jet (e.g.
 414 in [Gallimore et al. 1996](#)). [Mutie et al. \(2024\)](#)
 415 present higher resolution (~ 4 pc) *e*-MERLIN
 416 5 GHz data along with archival VLA 10 GHz,
 417 and VLA 21 GHz images of the jet. These
 418 images together show not only the central jet
 419 emission, but also detail in the larger scale bow
 420 shock, >200 pc from the SMBH in the same
 421 NE direction, which exhibits direct evidence of

the AGN’s impact on the ISM. The impact of the jet on the ISM is studied in part in both [Hviding et al. \(2023\)](#) and [Holden & Tadhunter \(2023\)](#), who both show evidence for gas ionization consistent with shock ionization or radiation-bounded AGN-photoionization along the outflow’s path on 160 pc to kpc scale. [García-Burillo et al. \(2014\)](#) show that the CO kinematics on distances 50 to 400 pc are spatially correlated with the radio jet, evidence that the AGN is influencing even the cold ISM. ALMA CO(6-5) observations from [Gallimore et al. \(2016\)](#) show that this molecular outflow originates within 2 pc from the SMBH, and has velocities relative to systemic of about 400 km s⁻¹.

3. NGC 1068 OBSERVATIONS

For NGC 1068, we made use of <3 pc scale resolution both in the near infrared (NIR) with Keck/OSIRIS+AO (adaptive optics; PI Medling), and in the sub-mm with ALMA archival data (PI García-Burillo).

3.1. *Keck/OSIRIS K-band Integral Field Spectroscopy*

The first of two sets of data we are using in this project is a set of high resolution integral field unit (IFU) Keck/OSIRIS+AO (OH-Suppressing InfraRed Imaging Spectrograph, [Larkin et al. 2006](#)) integrations, for which we mosaic all frames into a single data cube. These observations were taken with the Kbb filter (broad-band K between 1.965 - 2.381 μm) with the 35 mas pixel⁻¹ plate scale on 2018 December 28th, 2019 January 22nd, and 2019 October 7th for a total exposure time of 6120 seconds (51 frames, 120 seconds each). Weather impacted observations on 2019 October 7th, during which the laser guiding system was also not working. We used the galaxy nucleus as the natural guide star in NGS mode, and as the tip/tilt star in LGS mode. AO corrections in those frames without the laser produced larger

point spread functions with full-width at half-maximum (FWHM) values between 3 and 5 pixels compared to ~ 2 with the laser on other nights. We reduced the Keck/OSIRIS+AO observations using the OSIRIS Data Reduction Pipeline (OSIRISDRP, [Lyke et al. 2017](#); [Lockhart et al. 2019](#)) version 4.2.0, which we use to extract a spectrum for each spatial pixel, assemble the spectra into a cube, and mosaic the 51 total frames together to form the final image, which has a 0.17” point spread function (PSF) FWHM. Flux calibration was applied for each night before final mosaicking.

The resulting mosaic reveals a strong K-Band continuum (particularly near the AGN) and H₂ 1-0 rovibrational emission (S(0), $\lambda_{\text{rest}} = 2.2235\mu\text{m}$; S(1), $\lambda_{\text{rest}} = 2.1218\mu\text{m}$; S(2)), $\lambda_{\text{rest}} = 2.0338\mu\text{m}$. These line + continuum and continuum-subtracted H₂ 1-0 S(1) maps are shown in the middle and right panels of Figure 1 respectively. The line + continuum map was made using the Cube Analysis and Rendering Tool for Astronomy (CARTA, [Comrie et al. 2021](#)) and the continuum subtracted H₂ 1-0 S(1) map was made using QFitsView ([Ott 2012](#)). Both images show peaks of emission on or near the position of the central engine, and NGC 1068’s circumnuclear disk (CND) ring can be seen in the H₂ map.

3.2. *ALMA Band 6 Long-baseline Interferometry*

We chose the highest resolution CO J=(2-1) (hereafter CO(2-1)) available on the ALMA archive that shows strong emission (PI García-Burillo, Project code 2016.1.00232.S; see also [García-Burillo et al. 2019](#)). We retrieved the CO(2-1) spectral cube product from the ALMA archive, which has a rms of 0.25 mJy over 20 km s⁻¹, and was imaged using a Briggs ([Briggs 1995](#)) robust value of 0, resulting in a beam size of 41 \times 30 mas. We then used this spectral cube with the image cube analysis tools in CARTA ([Comrie et al. 2021](#)) to create a moment

0 (flux) map of the CO(2-1) emission. Figure 1 (left) shows this CO(2-1) moment 0 map which is masked below $3 \times \text{rms}$ and is used for our analysis in Section 4. Like in the warm H_2 observations, both the AGN and CND ring are bright sources in CO(2-1).

4. PRESCRIPTION PARAMETERS

In this pilot study, we examine the performance of the most commonly used accretion prescription for black hole growth in NGC 1068. The Bondi accretion formalism with a relative velocity of zero (also known as Bondi-Hoyle, or Bondi-Hoyle-Lyttleton e.g. Hoyle & Lyttleton 1939; Bondi & Hoyle 1944; Bondi 1952) follows the form:

$$\dot{M}_{\text{Bondi}} = 4\pi G^2 M_{\text{BH}}^2 \rho c_s^{-3} \quad (2)$$

where G is the gravitational constant, M_{BH} is the mass of the black hole, ρ is the gas density and c_s is the sound speed.

Bondi accretion is predicated on a spherically symmetric, non-self-gravitating gas distribution in which the gas inside the Bondi radius has no angular momentum. While this kind of environment may not be most appropriate for describing all galaxy nuclei (see Section 6 for additional information), Bondi accretion is a simple analytical prescription that can be applied inside a sphere of any radius, which makes it convenient as a sub-grid prescription.

In this subsection, we outline the methods for measuring each free parameter in the Bondi prescription using the available high resolution data from Section 3.

4.1. Parameter 1: black hole mass

Greenhill et al. (1996) imaged NGC 1068's water maser emission at a 0.65 pc scale using very long baseline interferometry. From the rotation curve of the water maser emission, they found the enclosed mass within that radius to be $\sim 1 \times 10^7 M_\odot$ (with uncertainty of order unity). Another study by Lodato & Bertin (2003) derive

a smaller black hole mass of $\sim 8 \pm 0.3 \times 10^6 M_\odot$ in a self-gravitating accretion disk model that matches the Greenhill et al. (1996) and Greenhill & Gwinn (1997) observations well. The Lodato & Bertin (2003) model corrects for non-Keplerian motion in the velocity profile of the water maser emission, but this could be an over-correction. In fact, other studies have found that the disk rotation may still be dominated by the black hole (Imanishi et al. 2018). Albeit with a worse fit to the velocities from the maser emission, Lodato & Bertin (2003) also fit a Keplerian rotation model, which has a best fit black hole mass of $\sim 1.5 \pm 0.02 \times 10^7 M_\odot$. In the absence of clear evidence in favor of one of the newer modeling schemes, we adopt the Greenhill et al. (1996) value of $M_{\text{BH}} = \sim 1 \times 10^7 M_\odot$ as an intermediate M_{BH} measurement.

4.2. Parameter 2: gas density

4.2.1. Choice of volume element

To measure the gas density, we first must define our volume element. In cosmological simulations, typically, a fixed number of gas particles exist inside a spherical region with radius r centered on the location of the SMBH. This volume makes up the black hole kernel, in which the accretion physics are prescribed. Although studies like the ones discussed in Section 2 and Vollmer et al. (2022) have shown that the ~ 10 pc cold gas distribution is more disk-like, we opt to use a sphere of volume $V = \frac{4}{3}\pi r^3$ centered on the AGN for which we vary the radius with the goal of mimicking the spherical radial aperture that simulations typically use to evaluate Bondi accretion.

4.2.2. Cold gas mass

To measure the molecular gas (H_2 and He) mass inside the sphere, we use the CO(2-1) data described in Section 3.2. To obtain a molecular gas mass, we utilize the conversion factor α_{CO} . The exact value of α_{CO} depends on several factors including the size scale and environment




Figure 2. *Left:* Moment 0 map of CO(2-1) in NGC 1068, with flux density values masked below $3 \times \text{rms}$. The red circle indicates the location and size of the 100 pc aperture (centered on the AGN) corresponding to the red dot in the right panel, which has $M_{\text{enc}, \text{H}_2 + \text{He}} = 4.09 \pm 1.49 \times 10^8 M_\odot$. The small, white ellipse in the bottom left represents the beam size of the ALMA data (41×30 mas). *Right:* Integrated mass profile inside the radial aperture. Details on the conversion to molecular gas mass can be found in Section 4.2.

over which the CO flux is measured. The picture is further complicated by the distinction between $\alpha_{\text{CO}(1-0)}$ and $\alpha_{\text{CO}(2-1)}$, where the difference is dictated by the ratio between the line luminosity of the two rotational transitions: r_{21} ($r_{21} = L'_{\text{CO}(2-1)} / L'_{\text{CO}(1-0)}$), which depends on the temperature of the gas. In this work, we follow the same α_{CO} methodology as in García-Burillo et al. (2019) who use the Milky Way $\alpha_{\text{CO}(1-0)} = 4.3 \pm 1.29 M_\odot (\text{K km s}^{-1} \text{pc}^2)^{-1}$ recommended by Bolatto et al. (2013). We use $\alpha_{\text{CO}(1-0)}$ in conjunction with the averaged line intensity ratios for NGC 1068's northern and southern CND regions (because the CND ring contains the majority of the nuclear gas mass): $r_{21} = 2.2 \pm 0.4$, from Viti et al. (2014) to calculate a final

$$\alpha_{\text{CO}(2-1)} = \frac{\alpha_{\text{CO}(1-0)}}{r_{21}} = \frac{4.3 \pm 1.29}{2.2 \pm 0.4} = 1.95 \pm 0.73 M_\odot (\text{K km s}^{-1} \text{pc}^2)^{-1}. \quad (3)$$

The outflowing components of NGC 1068 may have a lower $\alpha_{\text{CO}(1-0)}$, but we expect the Milky Way value to be closer to the average for the purpose of measuring integrated enclosed masses, especially at larger r . The outflow's impact on our gas mass measurement is expected

to be small as there is not much CO(2-1) emission between the AGN and CND ring, and the CND ring itself does not visually appear disturbed along the path of the outflow. $\alpha_{\text{CO}(2-1)}$ is then multiplied by the sum of the flux density inside a circular aperture of radius r , to match our spherical geometry. The enclosed mass profile is shown alongside a snapshot of the aperture geometry in Figure 2.

García-Burillo et al. (2019), who center their $r = 200$ pc aperture measurement on the center of the CND ring, find a molecular ($\text{H}_2 + \text{helium}$) gas mass of $\approx 1.4 \times 10^8 M_\odot$. We measure molecular gas mass within the same aperture (using CARTA to measure flux) and find $1.3 \pm 0.5 \times 10^8 M_\odot$ ($1.4 \pm 0.5 \times 10^8 M_\odot$ if centered on the AGN), both of which are consistent with the García-Burillo et al. (2019) measurement. For comparison to another nearby Seyfert 2, in the nuclear region of Circinus, using the warm gas tracer H_2 1-0 S(1), Müller Sánchez et al. (2006) find the total molecular gas mass to be $1.7 \times 10^7 M_\odot$ within $0.8''$ (52 pc). Integrated inside the same physical distance from the SMBH in NGC 1068, we find a molecular gas mass of $8.8 \pm 3.2 \times 10^7 M_\odot$, higher by almost 1 dex.

To convert enclosed mass to density we divide by the volume of the sphere with radius r (see Section 4.2.1) with r defined by our circular aperture size used for measuring mass. In this sphere with $r = 100$ pc centered on the AGN as shown in Figure 2 (left), we find a molecular gas mass density of $93.3 \pm 71.1 \text{ M}_\odot \text{pc}^{-3}$.

4.2.3. Warm H_2 gas mass

We also calculate an enclosed mass using the warm H_2 gas measured from the NIR data, following Equation 6 of Storch-Bergmann et al. (2009), which uses the line flux of the H_2 1-0 S(1) rovibrational transition at $\lambda_{\text{rest}} = 2.1218 \text{ }\mu\text{m}$. In NGC 1068, Martins et al. (2010) used the NASA 3-m Infrared Telescope Facility (IRTF) and found a nuclear (slit $1'' \times 2''$) extinction $E(\text{B-V})$ of 1.13 (from their Table 4). Assuming the standard extinction law of Cardelli et al. (1989) with $R_v = 3.1$, the extinction A_v ($A_v = R_v \times E(\text{B-V})$) is ~ 3.5 . Based on $A_k \sim A_v/10$ (Howarth 1983), the extinction-corrected H_2 gas mass inside $r < 1.7''$ (111 pc) is $\sim 68 \text{ M}_\odot$, which is about 1.38 times the observed value. The warm H_2 mass is inconsequential compared to the CO-derived value of $4.08 \pm 1.49 \times 10^8 \text{ M}_\odot$ in the same region.

One reason that the warm gas measurement in this subsection may be so small is due to the radiative environment in NGC 1068's nucleus. Under local thermal equilibrium (LTE, where the energy distribution can be described by a single number locally) conditions, the H_2 emission can be excited by the equilibrium value for temperatures $T \approx 1000 \text{ K}$ (Davies et al. 2005). To reach such high excitation temperatures in NGC 1068, H_2 emission lines can be excited through several mechanisms, as described below:

(1) *UV fluorescence*: This excitation mechanism dominates in photodissociation regions (PDRs). Far-ultraviolet (FUV, $\lambda > 912 \text{ }\text{\AA}$) radiation pumps the molecule into electronically excited states, leading to subsequent cascades that

emit fluorescent emission (Wakelam et al. 2017). This mechanism is dominant in Seyfert 1 galaxies (Davies et al. 2005). Although NGC 1068 is classified as a Seyfert 2 galaxy and is expected to have less FUV radiation, the HST/FOC UV image shows a bright nucleus with polarization (Barnouin et al. 2023) within our OSIRIS field of view (FOV).

(2) *Shocks and outflows*: Veilleux et al. (1997) suggest that shocks associated with nuclear outflows are a likely heating source for H_2 in many Seyfert 2 galaxies. May & Steiner (2017) analyzed VLT/SINFONI and Gemini/NIFS data with a larger FOV covering the entire CNB and proposed that the CNB could be an expanding bubble.

(3) *X-ray heating from the AGN*: X-ray emission can penetrate deeply into regions that are opaque to UV photons and influence H_2 excitation (Matt et al. 1997). All of these mechanisms can contribute to H_2 emission.

We measure the H_2 1-0 S(1) extinction-corrected intrinsic flux ($F_{\text{intrinsic}} = F_{\text{observed}} \times 10^{(0.4A_k)}$) and directly convert it to the warm H_2 gas mass. Due to the rectangular FOV, only an aperture radius of $< 0.3''$ is fully contained within the OSIRIS FOV, suggesting that H_2 emission at radii $> 0.3''$ is incomplete.

4.3. Parameter 3: sound speed of the gas

The final parameter required in the Bondi accretion formalism is the sound speed of the gas. The sound speed for an ideal gas is:

$$c_s = \sqrt{\frac{\gamma k_B T_K}{\mu m_p}} \quad (4)$$

where γ is the adiabatic index (1, as the gas is assumed to be isothermal in each sub-region), k_B is the Boltzmann constant $1.381 \times 10^{-16} \text{ erg K}^{-1}$, T_K is the temperature of the gas (K), and μ is the mean molecular weight of the gas, which is 2.7 since we assume the molecular gas is H_2 , 10% helium, and trace metals, and m_p is the

mass of a proton (kg). All but the temperature in this case are constants.

For the temperature of the molecular gas, we use two methods: one using CO rotation diagrams (cold gas), and another using an excitation diagram for the molecular H₂ (warm gas) from our Keck/OSIRIS+AO NIR data.

4.3.1. CO-derived c_s

For a temperature from CO transitions we refer to the work of Viti et al. (2014) who infer the temperature of the gas in the CND of NGC 1068 by using CO rotation diagrams. This method assumes that the gas is in LTE, and that the observations are mostly in the Rayleigh-Jeans regime where the intensity of the radiation is proportional to the temperature. This temperature is also known as the ‘rotational temperature’ and is equal to the kinetic temperature if all CO levels are thermalized (Goldsmith & Langer 1999). Because of these assumptions, this temperature should be considered a lower limit, which translates to an upper limit on our final accretion rate because $\dot{M}_{\text{Bondi}} \propto c_s^{-3}$. For the central region of NGC 1068, Viti et al. (2014) find a temperature of 50 ± 5 – 7 K via the CO rotation diagram method (see Section 3.1.1. of their work for more details). Plugging that and the other constants into Equation 4, we find that the speed of sound in the cold molecular gas phase is $391.0 \pm 135.4 \text{ m s}^{-1}$.

4.3.2. H₂-derived c_s

As shown in Section 4.2.3, warm H₂ is also present in NGC 1068’s nuclear regions, so we also consider the sound speed for this component of the ISM. To measure the temperature which we then use in the c_s calculation, we use the H₂ 1-0 S(0), S(1), and S(2) rovibrational line fluxes in the Keck/OSIRIS NIR data described in Section 3.1. Assuming the H₂ gas is in LTE, the H₂ excitation temperature is equal to the kinetic temperature. Figure 3(a) shows the H₂ excitation diagram, which is the column density

in the upper level of each transition normalized by its statistical weight (N_u/g_u) as a function of energy of the level as a temperature (E_u). The best-fit slope of this relationship is related to T_K as $\frac{N_u}{g_u} \propto e^{(-\frac{h\nu}{kT_K})}$ in the LTE description of energy level populations (see pages 322, 327 of Wilson et al. 2013). Solving for T_K then yields

$$-\frac{1}{T_K} \propto \frac{\ln \frac{N_u}{g_u}}{\frac{E_u}{k}}.$$

Because we have spatially resolved data for these H₂ lines, we can derive kinetic temperatures from 12-111 pc and apply them at the matched distances in the accretion rate prediction. While the Keck/OSIRIS+AO data has a higher resolution than 6 pc, the H₂ 1-0 S(1) and S(2) lines are not detected in a $r \leq 6$ pc (0.1”) aperture. Fluxes of the lines are measured using the line fitting tool in QFitsView (Ott 2012), which we use to fit the continuum and one Gaussian component to the integrated (within a region circular region with radius r) spectrum. Figure 3(b) shows the range of excitation temperatures as a function of radius. T_{kin} ranges from 678-2261 K, and peaks at $r \leq 85$ pc where $T_{\text{kin}} = 2261^{+3683}_{-1631}$ K. High temperatures may be caused by the influence of the PDR (Section 4.2.2 describes observations of this for NGC 1068), which is found to increase the H₂ 1-0 S(1) emission by up to 70% in the some luminous infrared galaxies (Davies et al. 2000; Davies et al. 2003). Using Equation 4 (with a mean molecular weight of H₂ only) results in H₂ sound speeds between 1440-2629 m s⁻¹, peaking at $r = 85$ pc.

5. RESULTS: \dot{M}_{Bondi} VS. $\dot{M}_{\text{X-RAY}}$

5.1. Calculating \dot{M}_{Bondi}

Now that we have calculated each parameter for the Bondi accretion prescription in Section 4, we are ready to estimate a Bondi accretion rate. Because our parameters are spatially resolved, we calculate accretion rate as a function of radial distance r representing a simulated resolution:




Figure 3. (*Left:*) Column density in the upper level of each H₂ 1-0 S(0), S(1), and S(2) transition as a function of energy level of that transition as a temperature (K) inside a $r = 21$ pc circular aperture centered on the AGN as shown in Figure 1. The best fit slope (using linear regression), as described in Section 4.3.2, is the temperature of the gas in that region if we assume LTE. (*Right:*) T_{kin} estimated as in the excitation diagram on the left but instead inside circular apertures matching the methods of Section 4.2 from 0.2" to 1.7" in steps of 0.1". The mean of the derived temperatures is 1612^{+2840}_{-1216} K.

$$\dot{M}_{\text{Bondi}}(r) = 4\pi G^2 M_{\text{BH}}^2 \rho(\leq r) c_s(\leq r)^{-3}. \quad (5)$$

where M_{BH} is in kg, ρ is in kg m^{-3} and c_s is in m s^{-1} . Figure 4 shows the Bondi accretion rate for the cold and warm derived cases as a function of radius. The Bondi accretion rate derived from the cold gas component ranges between about 10^9 (higher than the M_{BH} of NGC 1068's SMBH) and $10^6 M_{\odot} \text{ yr}^{-1}$. As the enclosed mass found in Section 4.2 for the warm H₂ gas component in $r < 170$ pc is small ($68 M_{\odot}$), and the temperature gradient is high (678-2261 K, see Section 4.3.2) relative to the values found for the cold gas component, the resulting Bondi accretion rates are much smaller (between about 10^{-2} and $3 M_{\odot} \text{ yr}^{-1}$) for the warm gas. These results suggest that the cold gas is the dominant carrier of mass accretion on $r < 170$ pc scales. Table 1 shows a range of precise values for both the cold and warm Bondi accretion rates.

5.2. Calculating X-ray accretion rates

To understand how well the Bondi accretion formalism compares to the real accretion rate, we compare it to the X-ray derived accretion

rate. To calculate an accretion rate from X-ray measurements, we use *Swift*/BAT data from the BAT AGN Spectroscopic Survey (BASS, Ricci et al. 2017). They present intrinsic luminosities in the 14-195 keV band, which we use alongside the bolometric correction, Equation 17 in Gupta et al. (2024):

$$\log(\kappa_{14-195}) = (0.13 \pm 0.04) \times \log(\lambda_{\text{Edd}}) + (1.04 \pm 0.05) \quad (6)$$

to calculate bolometric luminosity. Because Ricci et al. (2017) measure a neutral column density of $\log N_{\text{H}} = 25.0 \text{ cm}^{-2}$ in NGC 1068 and the X-ray continuum might not be well estimated when the emission is dominated by reprocessed radiation in environments like this, we conservatively estimate uncertainty on the input intrinsic 14-195 keV luminosity to be ± 0.4 dex. We then use that bolometric luminosity in the equation from Netzer & Trakhtenbrot (2014), $L_{\text{bol}} = \eta \dot{M} c^2$, solving for \dot{M} where η is the unitless mass-to-radiation conversion efficiency that depends on the spin of the black hole. For




Figure 4. X-ray and (Cold-derived) Bondi accretion rates as a function of radius over which nuclear parameters are measured. Color bars and ranges displayed to the right denote uncertainties on each X-ray measurement. Regardless of which gas component is used to estimate \dot{M}_{Bondi} , the Bondi prescription overestimates \dot{M}_{BH} by orders of magnitude. For the cold gas case, which represents the majority of gas available for accretion in NGC 1068, Bondi overpredicts the accretion rate by between 9 and 13 orders of magnitude.

stationary, retrograde disk, and maximally rotating SMBHs respectively, the values for η are 0.057, 0.038, and 0.32 (Netzer & Trakhtenbrot 2014). For NGC 1068, we find $\dot{M}_{\text{X-ray}}$ values equal to $1.51 \pm 0.81 \times 10^{-3} \text{ M}_{\odot} \text{ yr}^{-1}$ (stationary SMBH), $2.26 \pm 1.21 \times 10^{-3} \text{ M}_{\odot} \text{ yr}^{-1}$ (retrograde accretion disk), and $2.69 \pm 1.43 \times 10^{-4} \text{ M}_{\odot} \text{ yr}^{-1}$ (maximally spinning SMBH). As shown in Figure 4 and Table 1, \dot{M}_{Bondi} overestimates the accretion rate by several orders of magnitude in the warm gas case to up to 13 orders of magnitude in the cold gas case in small aperture radii. In Section 6 we discuss the implications of such

a discrepancy with respect to cosmological simulations.

Vollmer et al. (2022) used the IR-derived bolometric luminosity for the AGN in NGC 1068 from Vollmer et al. (2018) to calculate $\dot{M}_{\text{BH}} \sim L_{\text{bol}} / (0.1c^2) \sim 0.05 \text{ M}_{\odot} \text{ yr}^{-1}$. They calculate a mass accretion rate onto their modeled accretion disk for NGC 1068 to be $2 \times 10^{-3} \text{ M}_{\odot} \text{ yr}^{-1}$ ($\eta = 0.1$), which is in agreement with our $\dot{M}_{\text{X-ray}}$ values.

6. DISCUSSION: RESULTS IN THE CONTEXT OF SIMULATIONS

Method	Accretion rate ($M_{\odot} \text{ yr}^{-1}$)						
X-rays ($\epsilon = 0.038$)	$2.26 \pm 1.21 \times 10^{-3}$						
X-rays ($\epsilon = 0.057$)	$1.51 \pm 0.81 \times 10^{-3}$						
X-rays ($\epsilon = 0.32$)	$2.69 \pm 1.43 \times 10^{-4}$						
Bondi ($< r$)	5 pc	10 pc	25 pc	50 pc	100 pc	200 pc	500 pc
($T_{\text{Kin}} = 50 \text{ K}$)	$1.86 \pm 2.05 \times 10^9$	$4.87 \pm 5.37 \times 10^8$	$7.16 \pm 7.89 \times 10^7$	$5.69 \pm 6.27 \times 10^7$	$3.87 \pm 0.95 \times 10^7$	$9.54 \pm 1.05 \times 10^6$	$8.75 \pm 9.64 \times 10^6$
($T_{\text{Kin}} = 678\text{--}2261 \text{ K}$)	*	*	$2.85^{+10.07}_{-7.54}$	$1.73^{+5.53}_{-4.89} \times 10^{-1}$	$1.76^{+8.23}_{-4.18} \times 10^{-2}$	*	*

Table 1. Accretion rate measurements and estimates for rates derived from X-ray luminosities and the pure Bondi accretion prescription inside various radii. In NGC 1068, where the cold gas phase makes up the bulk of the gas mass in the nucleus, the cold gas derived Bondi accretion estimate outpaces the X-ray derived accretion rates by between 9 and 13 orders of magnitude. *H₂ 1-0 S(1) and S(2) lines required to calculate temperature for the Bondi calculation are not detected in this aperture.

To inform theorists on which accretion prescriptions in their simulations are best to use and when, we have designed our measurements to fit in the practical context of those simulations. Large scale cosmological simulations must use sub-grid physics for accretion because of computing constraints. Some examples of hydrodynamical galaxy evolution simulations that use spherically symmetric, Bondi or Bondi-like black hole accretion formalisms are *Illustris*/*IllustrisTNG* (Genel et al. 2014; Vogelsberger et al. 2014; Pillepich et al. 2018b), *Magneticum Pathfinder* (Hirschmann et al. 2014; Bocquet et al. 2016; Dolag et al. 2016), *MassiveBlack-II* (Khandai et al. 2015), *Eagle* (Schaye et al. 2015), *Horizon-AGN* (Dubois et al. 2016), *Romulus* (Tremmel et al. 2017), and *SIMBA* (Davé et al. 2019b, uses Bondi for hot gas only). The resolution of the hydrodynamical gas cells in which these sub-grid physics is typically close to 1 kpc. In the highest resolution zoom-in simulations, the spherical radius in which particle calculations are made is approximately 10 pc (Wetzel et al. 2023). Hyper-refinement simulations (e.g. Anglés-Alcázar et al. 2021; Hopkins et al. 2024), where gas resolution elements are dynamically split to reach high resolution can reach spatial scales smaller than 10 pc, but these simulations can only be

practically run for short periods of cosmic time due to computing constraints.

Because we have spatially-resolved measurements, we are able to examine the performance of Bondi accretion at a range of spatial scales. For the Bondi accretion rate derived from warm gas we are limited by the field of view of OSIRIS ($0.56 \times 2.24''$ with our observational setup), but the ALMA data extends to over 500 pc away from the SMBH.

Table 1 shows the Bondi accretion rates estimated at radii between 5-500 pc as calculated in Section 5, and the X-ray accretion rates as calculated in Section 5.2, which are all plotted together in Figure 4. At all aperture radii, regardless of whether we are estimating \dot{M}_{Bondi} using the cold or warm gas component, the parameterized Bondi accretion rate exceeds the X-ray derived accretion rate (by 1 or more dex in the warm gas case and by 9 or more dex in the cold gas case).

This is, perhaps, not a surprising result. Past studies have hinted towards Bondi accretion overestimating the real accretion rate. Di Matteo et al. (2000) found that luminosities calculated using estimated Bondi accretion rates for six black holes with masses of $0.22\text{--}5.2 \times 10^9 M_{\odot}$ determined in Magorrian et al. (1998) were 4-6 orders of magnitude higher than the real luminosities of the galaxy nuclei. Hopkins

et al. (2016) model SMBH accretion in a gas-rich nuclear disk in a massive simulated galaxy with 0.1 pc resolution. In their study, applying a pure Bondi accretion formalism resulted in an accretion rate $\sim 10^8$ times higher than the luminosity-derived accretion rate native to their simulation. Their pure Bondi accretion rate ($\sim 10^7 M_\odot \text{yr}^{-1}$), agrees with our cold-gas derived pure Bondi accretion rate between approximately 25 and 200 pc in NGC 1068. Near the SMBH, pure Bondi accretion ignores the possibility that gas particles may have angular momentum. The gas in the simulation used in Hopkins et al. (2016) is primarily cold and is supported by angular momentum rather than radiation pressure. Observations show that especially in gas-rich galaxies that naturally host molecular torii, the $r < 100$ pc cold gas reservoir is large, has significant angular momentum, and is the primary candidate for black hole accretion fueling (Davies et al. 2004; Hicks et al. 2013; Medling et al. 2014; Lin et al. 2016; Gaspari et al. 2015). Ignoring the angular momentum of the cold gas is likely the primary cause of the overestimate that Bondi accretion makes in Di Matteo et al. (2000), Hopkins et al. (2016), and in this work.

If NGC 1068 is typical, these results suggest that the usage of pure Bondi accretion is likely to struggle to accurately predict real black hole accretion rates. From our example, in the cold gas estimate, which represents the majority of the mass available for accretion, the Bondi accretion prediction dramatically (by up to 13 orders of magnitude) overpredicts the true accretion rate. Understanding the physical mechanisms that drive accretion on the sub-grid scales in galactic nuclei can inform the future development of accretion prescriptions. The Bondi prescription allows particles to fall directly onto the BH inside the Bondi radius, but our results suggest that angular momentum plays an important role in some nuclei.

7. CONCLUSIONS AND FUTURE EXPANSION OF THIS PROJECT

In this study we estimate a Bondi accretion rate as a function of radius for NGC 1068 using two different molecular gas tracers, and compare the result to the direct accretion rate derived from hard X-ray luminosity of the AGN. Compared to warm H_2 gas, CO gas is the dominant mass carrier close to the SMBH. Following this, the cold gas derived Bondi accretion rate estimate outpaces the X-ray derived value by more than 9 orders of magnitude at all aperture sizes.

This paper is a pilot for a wider study of AGN and accretion prescriptions. Direct probes of sub-grid accretion prescriptions may, as our sample expands, help identify which physical processes dominate accretion on a variety of spatial scales, and in turn provide recommendations for appropriate sub-grid prescriptions to describe them. The results in this work support previous evidence that in high resolution cosmological simulations, applying a Bondi accretion prescription can lead to large overestimates of \dot{M}_{BH} and therefore large overestimates of AGN feedback, which in turn impacts the global galaxy evolutionary track. We note that this is a test for a specific Seyfert 2 AGN. To make more robust recommendations about the application of the Bondi accretion prescription for sub-grid accretion physics we must directly test Bondi on more galaxies.

8. ACKNOWLEDGEMENTS

The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community; we are privileged to be guests on your sacred mountain. We wish to pay respect to the Atacameño community of the Chajnantor Plateau, whose traditional home now also includes the ALMA observatory. This work makes

use of the following data from ALMA: project
 2016.1.00232.S (PI García-Burillo). ALMA is
 a partnership of ESO (representing its mem-
 ber states), NSF (USA) and NINS (Japan),
 together with NRC (Canada) and NSC and
 ASIAA (Taiwan) and KASI (Republic of Ko-
 rea), in cooperation with the Republic of Chile.
 The Joint ALMA Observatory is operated by
 ESO, AUI/NRAO and NAOJ. The National
 Radio Astronomy Observatory is a facility of
 the National Science Foundation operated un-
 der cooperative agreement by Associated Uni-
 versities, Inc. Some of the data presented herein
 were obtained at the W. M. Keck Observa-
 tory, which is operated as a scientific partner-
 ship among the California Institute of Tech-
 nology, the University of California and the
 National Aeronautics and Space Administra-
 tion. The Observatory was made possible by the
 generous financial support of the W. M. Keck
 Foundation. The authors also wish to thank
 the W.M. Keck Observatory staff for their ef-
 forts on the OSIRIS+AO instrumentation. JA,
 AMM, M-YL, and NJ acknowledge support
 from NSF CAREER grant number 2239807 and
 Cottrell Scholar Award CS-CSA-2024-092 from
 the Research Corporation for Science Advance-
 ment. PT acknowledges support from NSF-
 AST 2346977 and the NSF-Simons AI Insti-
 tute for Cosmic Origins which is supported
 by the National Science Foundation under Co-
 operative Agreement 2421782 and the Simons
 Foundation award MPS-AI-00010515. D.A.A.
 acknowledges support from NSF grant AST-
 2108944 and CAREER award AST-2442788,
 NASA grant ATP23-0156, STScI grants JWST-
 GO-01712.009-A, JWST-AR-04357.001-A, and
 JWST-AR-05366.005-A, an Alfred P. Sloan Re-
 search Fellowship, and Cottrell Scholar Award
 CS-CSA-2023-028 by the Research Corporation
 for Science Advancement.

Software: Astropy (Astropy Collaboration
 et al. 2013, 2018, 2022), Matplotlib (Hunter
 2007), NumPy (Harris et al. 2020).

REFERENCES

- 1064 Anglés-Alcázar, D., Davé, R., Faucher-Giguère,
 1065 C.-A., Özel, F., & Hopkins, P. F. 2017,
 1066 MNRAS, 464, 2840
 1067 Anglés-Alcázar, D., Özel, F., & Davé, R. 2013,
 1068 ApJ, 770, 5
 1069 Anglés-Alcázar, D., Özel, F., Davé, R., Katz, N.,
 1070 Kollmeier, J. A., & Oppenheimer, B. D. 2015,
 1071 ApJ, 800, 127
 1072 Anglés-Alcázar, D., et al. 2021, ApJ, 917, 53
 1073 Antonucci, R. 1993, ARA&A, 31, 473
 1074 Astropy Collaboration et al. 2013, A&A, 558, A33
 1075 —. 2018, AJ, 156, 123
 1076 —. 2022, ApJ, 935, 167
 1077 Barnouin, T., Marin, F., Lopez-Rodriguez, E.,
 1078 Huber, L., & Kishimoto, M. 2023, A&A, 678,
 1079 A143
 1080 Barthelmy, S. D., et al. 2005, SSRv, 120, 143
 1081 Behroozi, P. S., Wechsler, R. H., & Conroy, C.
 1082 2013, ApJ, 770, 57
 1083 Bocquet, S., Saro, A., Dolag, K., & Mohr, J. J.
 1084 2016, MNRAS, 456, 2361
 1085 Bolatto, A. D., Wolfire, M., & Leroy, A. K. 2013,
 1086 ARA&A, 51, 207
 1087 Bondi, H. 1952, MNRAS, 112, 195
 1088 Bondi, H., & Hoyle, F. 1944, MNRAS, 104, 273
 1089 Booth, C. M., & Schaye, J. 2009, MNRAS, 398, 53
 1090 Briggs, D. S. 1995, in American Astronomical
 1091 Society Meeting Abstracts, Vol. 187, American
 1092 Astronomical Society Meeting Abstracts, 112.02
 1093 Burns, J. O., Feigelson, E. D., & Schreier, E. J.
 1094 1983, ApJ, 273, 128
 1095 Cardelli, J. A., Clayton, G. C., & Mathis, J. S.
 1096 1989, ApJ, 345, 245
 1097 Chen, C.-T. J., et al. 2013, ApJ, 773, 3
 1098 Cielo, S., Bieri, R., Volonteri, M., Wagner, A. Y.,
 1099 & Dubois, Y. 2018, MNRAS, 477, 1336
 1100 Comrie, A., et al. 2021, CARTA: The Cube
 1101 Analysis and Rendering Tool for Astronomy
 1102 Curtis, H. D. 1918, Publications of Lick
 1103 Observatory, 13, 9
 1104 Davé, R., Anglés-Alcázar, D., Narayanan, D., Li,
 1105 Q., Rafieferantsoa, M. H., & Appleby, S. 2019a,
 1106 MNRAS, 486, 2827
 1107 —. 2019b, MNRAS, 486, 2827
 1108 Davies, R., Ward, M., & Sugai, H. 2000, ApJ, 535,
 1109 735
 1110 Davies, R. I., Sternberg, A., Lehnert, M., &
 1111 Tacconi-Garman, L. E. 2003, ApJ, 597, 907
 1112 Davies, R. I., Sternberg, A., Lehnert, M. D., &
 1113 Tacconi-Garman, L. E. 2005, ApJ, 633, 105
 1114 Davies, R. I., Tacconi, L. J., & Genzel, R. 2004,
 1115 ApJ, 613, 781
 1116 Di Matteo, T., Quataert, E., Allen, S. W.,
 1117 Narayan, R., & Fabian, A. C. 2000, MNRAS,
 1118 311, 507
 1119 Di Matteo, T., Springel, V., & Hernquist, L. 2005,
 1120 Nature, 433, 604
 1121 Dolag, K., Komatsu, E., & Sunyaev, R. 2016,
 1122 MNRAS, 463, 1797
 1123 Dubois, Y., Devriendt, J., Slyz, A., & Teyssier, R.
 1124 2012, MNRAS, 420, 2662
 1125 Dubois, Y., Gavazzi, R., Peirani, S., & Silk, J.
 1126 2013a, MNRAS, 433, 3297
 1127 Dubois, Y., Peirani, S., Pichon, C., Devriendt, J.,
 1128 Gavazzi, R., Welker, C., & Volonteri, M. 2016,
 1129 MNRAS, 463, 3948
 1130 Dubois, Y., Pichon, C., Devriendt, J., Silk, J.,
 1131 Haehnelt, M., Kimm, T., & Slyz, A. 2013b,
 1132 MNRAS, 428, 2885
 1133 Escala, A. 2007, ApJ, 671, 1264
 1134 Fabian, A. C. 2012, ARA&A, 50, 455
 1135 Ferrarese, L., & Merritt, D. 2000, ApJL, 539, L9
 1136 Feruglio, C., Maiolino, R., Piconcelli, E., Menci,
 1137 N., Aussel, H., Lamastra, A., & Fiore, F. 2010,
 1138 A&A, 518, L155
 1139 Gallimore, J. F., Baum, S. A., O’Dea, C. P., &
 1140 Pedlar, A. 1996, ApJ, 458, 136
 1141 Gallimore, J. F., & Impellizzeri, C. M. V. 2023,
 1142 ApJ, 951, 109
 1143 Gallimore, J. F., et al. 2016, ApJL, 829, L7
 1144 Gámez Rosas, V., et al. 2022, Nature, 602, 403
 1145 García-Burillo, S., et al. 2014, A&A, 567, A125
 1146 —. 2019, A&A, 632, A61
 1147 Gaspari, M., Brighenti, F., & Temi, P. 2015,
 1148 A&A, 579, A62
 1149 Gebhardt, K., et al. 2000, ApJL, 539, L13
 1150 Gehrels, N., et al. 2004, ApJ, 611, 1005
 1151 Genel, S., et al. 2014, MNRAS, 445, 175
 1152 Goldsmith, P. F., & Langer, W. D. 1999, ApJ,
 1153 517, 209
 1154 Greenhill, L. J., & Gwinn, C. R. 1997, Ap&SS,
 1155 248, 261
 1156 Greenhill, L. J., Gwinn, C. R., Antonucci, R., &
 1157 Barvainis, R. 1996, ApJL, 472, L21
 1158 Gupta, K. K., et al. 2024, A&A, 691, A203

- 1159 Hagiwara, Y., Baan, W. A., Imanishi, M., &
 1160 Diamond, P. 2024, *MNRAS*, 528, 3668
 1161 Harris, C. R., et al. 2020, *Nature*, 585, 357
 1162 Hicks, E. K. S., Davies, R. I., Maciejewski, W.,
 1163 Emsellem, E., Malkan, M. A., Dumas, G.,
 1164 Müller-Sánchez, F., & Rivers, A. 2013, *ApJ*,
 1165 768, 107
 1166 Hinshaw, G., et al. 2009, *ApJS*, 180, 225
 1167 Hirschmann, M., Dolag, K., Saro, A., Bachmann,
 1168 L., Borgani, S., & Burkert, A. 2014, *MNRAS*,
 1169 442, 2304
 1170 Hirschmann, M., Khochfar, S., Burkert, A., Naab,
 1171 T., Genel, S., & Somerville, R. S. 2010,
 1172 *MNRAS*, 407, 1016
 1173 Hobbs, A., Power, C., Nayakshin, S., & King,
 1174 A. R. 2012, *MNRAS*, 421, 3443
 1175 Holden, L. R., & Tadhunter, C. N. 2023, *MNRAS*,
 1176 524, 886
 1177 Hopkins, P. F., & Quataert, E. 2011, *MNRAS*,
 1178 415, 1027
 1179 Hopkins, P. F., Torrey, P., Faucher-Giguère,
 1180 C.-A., Quataert, E., & Murray, N. 2016,
 1181 *MNRAS*, 458, 816
 1182 Hopkins, P. F., et al. 2024, *The Open Journal of*
 1183 *Astrophysics*, 7, 18
 1184 Howarth, I. D. 1983, *MNRAS*, 203, 301
 1185 Hoyle, F., & Lyttleton, R. A. 1939, *Proceedings of*
 1186 *the Cambridge Philosophical Society*, 35, 405
 1187 Hunter, J. D. 2007, *Computing in Science &*
 1188 *Engineering*, 9, 90
 1189 Hviding, R. E., Hickox, R. C., Väisänen, P.,
 1190 Ramphul, R., & Hainline, K. N. 2023, *AJ*, 166,
 1191 111
 1192 Imanishi, M., Nakanishi, K., Izumi, T., & Wada,
 1193 K. 2018, *ApJL*, 853, L25
 1194 Imanishi, M., et al. 2020, *ApJ*, 902, 99
 1195 Jaffe, W., et al. 2004, *Nature*, 429, 47
 1196 Jahnke, K., & Macciò, A. V. 2011, *ApJ*, 734, 92
 1197 Khandai, N., Di Matteo, T., Croft, R., Wilkins,
 1198 S., Feng, Y., Tucker, E., DeGraf, C., & Liu,
 1199 M.-S. 2015, *MNRAS*, 450, 1349
 1200 Kormendy, J., & Ho, L. C. 2013, *ARA&A*, 51, 511
 1201 Krimm, H. A., et al. 2013, *ApJS*, 209, 14
 1202 Kumar, P. 1999, *ApJ*, 519, 599
 1203 Larkin, J., et al. 2006, in *Society of Photo-Optical*
 1204 *Instrumentation Engineers (SPIE) Conference*
 1205 *Series*, Vol. 6269, *Ground-based and Airborne*
 1206 *Instrumentation for Astronomy*, ed. I. S.
 1207 McLean & M. Iye, 62691A
 1208 Lin, M.-Y., et al. 2016, *MNRAS*, 458, 1375
 1209 Lockhart, K. E., et al. 2019, *AJ*, 157, 75
 1210 Lodato, G., & Bertin, G. 2003, *A&A*, 398, 517
 1211 Lyke, J., et al. 2017, *OSIRIS Toolbox:*
 1212 *OH-Suppressing InfraRed Imaging*
 1213 *Spectrograph pipeline*, *Astrophysics Source*
 1214 *Code Library*, record ascl:1710.021
 1215 Magorrian, J., et al. 1998, *AJ*, 115, 2285
 1216 Markevitch, M. 1998, *ApJ*, 504, 27
 1217 Martins, L. P., Rodríguez-Ardila, A., de Souza,
 1218 R., & Gruenwald, R. 2010, *MNRAS*, 406, 2168
 1219 Matt, G., et al. 1997, *A&A*, 325, L13
 1220 May, D., & Steiner, J. E. 2017, *MNRAS*, 469, 994
 1221 McConnell, N. J., & Ma, C.-P. 2013, *ApJ*, 764, 184
 1222 Medling, A. M., et al. 2014, *ApJ*, 784, 70
 1223 Meenakshi, M., et al. 2022, *MNRAS*, 516, 766
 1224 Mercedes-Feliz, J., et al. 2023, *MNRAS*, 524, 3446
 1225 Morganti, R. 2017, *Frontiers in Astronomy and*
 1226 *Space Sciences*, 4, 42
 1227 Müller Sánchez, F., Davies, R. I., Eisenhauer, F.,
 1228 Tacconi, L. J., Genzel, R., & Sternberg, A.
 1229 2006, *A&A*, 454, 481
 1230 Müller-Sánchez, F., Prieto, M. A., Hicks, E. K. S.,
 1231 Vives-Arias, H., Davies, R. I., Malkan, M.,
 1232 Tacconi, L. J., & Genzel, R. 2011, *ApJ*, 739, 69
 1233 Mushotzky, R. F. 1984, *Physica Scripta Volume*
 1234 *T*, 7, 157
 1235 Mutie, I. M., et al. 2024, *MNRAS*, 527, 11756
 1236 Negri, A., & Volonteri, M. 2017, *MNRAS*, 467,
 1237 3475
 1238 Netzer, H., & Trakhtenbrot, B. 2014, *MNRAS*,
 1239 438, 672
 1240 Ott, T. 2012, *QFitsView: FITS file viewer*,
 1241 *Astrophysics Source Code Library*, record
 1242 ascl:1210.019
 1243 Peng, C. Y. 2007, *ApJ*, 671, 1098
 1244 Pillepich, A., et al. 2018a, *MNRAS*, 473, 4077
 1245 —. 2018b, *MNRAS*, 473, 4077
 1246 Raban, D., Jaffe, W., Röttgering, H.,
 1247 Meisenheimer, K., & Tristram, K. R. W. 2009,
 1248 *MNRAS*, 394, 1325
 1249 Ricci, C., et al. 2017, *ApJS*, 233, 17
 1250 Rosas-Guevara, Y. M., et al. 2015, *MNRAS*, 454,
 1251 1038
 1252 Saito, T., et al. 2022, *ApJ*, 935, 155
 1253 Schaye, J., et al. 2015, *MNRAS*, 446, 521
 1254 Springel, V., Di Matteo, T., & Hernquist, L. 2005,
 1255 *MNRAS*, 361, 776
 1256 Springel, V., & Hernquist, L. 2005, *ApJL*, 622, L9

- 1257 Storch-Bergmann, T., McGregor, P. J., Riffel,
 1258 R. A., Simões Lopes, R., Beck, T., & Dopita,
 1259 M. 2009, MNRAS, 394, 1148
 1260 Taylor, P., & Kobayashi, C. 2015, MNRAS, 448,
 1261 1835
 1262 Torrey, P., Vogelsberger, M., Genel, S., Sijacki,
 1263 D., Springel, V., & Hernquist, L. 2014,
 1264 MNRAS, 438, 1985
 1265 Tremmel, M., Karcher, M., Governato, F.,
 1266 Volonteri, M., Quinn, T. R., Pontzen, A.,
 1267 Anderson, L., & Bellovary, J. 2017, MNRAS,
 1268 470, 1121
 1269 Tristram, K. R. W., Burtscher, L., Jaffe, W.,
 1270 Meisenheimer, K., Hönig, S. F., Kishimoto, M.,
 1271 Schartmann, M., & Weigelt, G. 2014, A&A,
 1272 563, A82
 1273 Valentini, M., et al. 2020, MNRAS, 491, 2779
 1274 Veilleux, S., Goodrich, R. W., & Hill, G. J. 1997,
 1275 ApJ, 477, 631
 1276 Viti, S., et al. 2014, A&A, 570, A28
 1277 Vogelsberger, M., Genel, S., Sijacki, D., Torrey,
 1278 P., Springel, V., & Hernquist, L. 2013, MNRAS,
 1279 436, 3031
 1280 Vogelsberger, M., et al. 2014, MNRAS, 444, 1518
 1281 Vollmer, B., Schartmann, M., Burtscher, L.,
 1282 Marin, F., Hönig, S., Davies, R., & Goosmann,
 1283 R. 2018, A&A, 615, A164
 1284 Vollmer, B., et al. 2022, A&A, 665, A102
 1285 Wada, K., Kudoh, Y., & Nagao, T. 2023,
 1286 MNRAS, 526, 2717
 1287 Wakelam, V., et al. 2017, Molecular Astrophysics,
 1288 9, 1
 1289 Weinberger, R., et al. 2017, MNRAS, 465, 3291
 1290 Wetzel, A., et al. 2023, ApJS, 265, 44
 1291 Williamson, D., Hönig, S., & Venanzi, M. 2020,
 1292 ApJ, 897, 26
 1293 Wilson, A. S., & Ulvestad, J. S. 1983, ApJ, 275, 8
 1294 Wilson, T. L., Rohlfs, K., & Hüttemeister, S.
 1295 2013, Tools of Radio Astronomy
 1296 Wright, E. L. 2006, PASP, 118, 1711