

Finitistic dimension conjecture

Jacob Fjeld Grevstad

2020

Abstract

FDC yo! This is abstract!

Contents

Notation	iii
Introduction	1
1 The homological conjectures	1
1.1 Implications	3
1.2 Wedderburn correspondence	6
2 Recollement	10
2.1 Triangular matrix rings and vertex removal	13
3 Contravariant finiteness	19
4 Repdimension	27
4.1 The Igusa-Todorov function	31
4.2 Stably hereditary algebras	34
4.3 Special biserial algebras	37
5 Vanishing radical powers	37
6 Monomial algebras	38
7 Unbounded derived category	42
8 Summary	43
9 Dual conjectures	45
10 Personal appendix	46

Notation

k a field Λ findim alg, J radical

$\text{mod } \Lambda$ finite dimensional (left-)modules, $\text{Mod } \Lambda$ all (left-)modules.

${}_{\Gamma}M_{\Lambda}$ is a $\Gamma - \Lambda$ -bimodule. Left Γ -module, right Λ -module

Quiver/path algebra. Multiplication is written right to left

$\text{Hom}_{\mathcal{C}}(M, N)$ can be written as $\mathcal{C}(M, N)$ or sometimes simply (M, N)

$D : \text{mod } \Lambda \rightarrow \text{mod } \Lambda^{\text{op}}$ is the duality $DM = \text{Hom}(M, k)$

$\mathcal{D}^b(\Lambda)$ bounded derived category, K^b , $K^{+,b}$, $K^{-,b}$, \mathcal{D} , etc. $X^{\geq n}$ hard truncation, $[i]$ shift

All functors are additive

$I(M)$ injective envelope, P_M^0 projective cover, $\cdots \rightarrow P_M^1 \rightarrow P_M^0$ projective resolution. $\Omega^n M$ syzygy.

pd projective dimension

Introduction

This is an introduction

1 The homological conjectures

Finitistic Dimension Conjecture (FDC)

Definition 1.1 (Finitistic dimension). For a finite dimensional algebra Λ the *finitistic dimension* of Λ , denoted $\text{findim}(\Lambda)$ is defined by

$$\text{findim}(\Lambda) = \{\text{pd } M : M \in \text{mod } \Lambda, \text{pd } M < \infty\}.$$

Conjecture 1 (Finitistic dimension conjecture). *For a finite dimensional algebra the finitistic dimension is always finite.*

$$\text{findim}(\Lambda) < \infty$$

Watanabe Tilting Conjecture (WTC)

Definition 1.2 (Watanabe tilting). Let T be a in $\text{mod } \Lambda$ for a finite dimensional algebra Λ . Then T is *Watanabe tilting* if

- i) We have that $\text{Ext}^n(T, T) = 0$ for all $n > 0$.
- ii) There is an exact sequence

$$\eta: 0 \rightarrow \Lambda \rightarrow T_0 \rightarrow T_1 \rightarrow \cdots$$

where T_i is in $\text{add } T$.

- iii) The sequence $\text{Hom}(\eta, T)$ is exact. Equivalently we have $\text{Ext}^1(\text{Ker } f, T) = 0$ for every f in η .

Watanabe tilting generalizes the idea of a tilting module. A watanabe tilting module is a tilting module if it has finite projective dimension and η is bounded. The Watanabe tilting conjecture states that this last restriction is superfluous.

Conjecture 2 (Watanabe tilting conjecture). *If T is Watanabe tilting and has finite projective dimension then T is a tilting module. In other words we can choose η to be bounded.*

Gorenstein Symmetry Conjecture (GSC)

Conjecture 3 (Gorenstein symmetry conjecture). *If Λ is a finite dimensional algebra the injective dimension of Λ as a left module is finite if and only if the projective dimension of $D\Lambda_\Lambda$ is finite.*

The conjecture describes a sort of symmetry between the projective and injective modules. Equivalently we could formulate the conjecture as Λ having finite injective dimension as a left module if and only if it has finite injective dimension as a right module.

Vanishing Conjecture (VC)

If Λ is a finite dimensional algebra we denote by $K^b(\text{inj } \Lambda)$ the homotopy category of bounded complexes of injectives. The category $K^{+,b}(\text{inj } \Lambda)$ is the homotopy category of complexes of injectives that are bounded below, and bounded in homology. There is an equivalence of categories between $K^{+,b}$ and the bounded derived category $\mathcal{D}^b(\Lambda)$. This allows us to consider $K^b(\text{inj } \Lambda)$ as a subcategory of $\mathcal{D}^b(\Lambda)$. Using this we define the perpendicular subcategory

$$K^b(\text{inj } \Lambda)^\perp = \{X \in \mathcal{D}^b(\Lambda) : \text{Hom}(I, X) = 0 \text{ for all } I \in K^b(\text{inj } \Lambda)\}.$$

The vanishing conjecture then states that this subcategory is 0.

Conjecture 4 (Vanishing conjecture). *If Λ is a finite dimensional algebra then $K^b(\text{inj } \Lambda)^\perp = 0$.*

Nunke Condition (NuC)

Conjecture 5 (Nunke condition). *If $X \neq 0$ is a non-zero module over a finite dimensional algebra Λ , then there is an $n \geq 0$ such that $\text{Ext}^n(D\Lambda, X) \neq 0$.*

Strong Nakayama Conjecture (SNC)

The strong Nakayama condition is a slight weakening of the Nunke condition.

Conjecture 6 (Strong Nakayama conjecture). *If S is a simple module over a finite dimensional algebra Λ , then there is an $n \geq 0$ such that $\text{Ext}^n(D\Lambda, S) \neq 0$.*

Auslander–Reiten Conjecture (ARC)

Conjecture 7 (Auslander–Reiten conjecture). *Let Λ be finite dimensional algebra. If M is a module over Λ such that $\text{Ext}^n(M, M \oplus \Lambda) = 0$ for all $n > 0$, then M is projective.*

Note that if we replace M by $M \oplus \Lambda$ then we get the equivalent formulation: If M is a generator with $\text{Ext}^n(M, M) = 0$ for all $n > 0$, then M is projective.

Nakayama Conjecture (NC)

Definition 1.3 (Dominant dimension). Let Λ be a finite dimensional algebra, and let

$$0 \longrightarrow \Lambda \longrightarrow I^0 \longrightarrow I^1 \longrightarrow \dots$$

be the minimal injective resolution of Λ . Then the *dominated dimension* of Λ is

$$\text{domdim}(\Lambda) = \inf\{n \mid I^n \text{ is not projective}\}.$$

Conjecture 8 (Nakayama conjecture). *If Λ has infinite dominant dimension, then Λ is self injective.*

1.1 Implications

The homological conjectures are related in the way presented in the diagram below.

$$\begin{array}{ccccccc} \text{FDC} & \implies & \text{WTC} & \implies & \text{GSC} & & \\ \Downarrow & & & & & & \\ \text{VC} & \implies & \text{NuC} & \implies & \text{SNC} & \implies & \text{ARC} \implies \text{NC} \end{array}$$

The remainder of this section is used to prove these implications.

Theorem 1.4. *[Hap93, 1.2] The finitistic dimension conjecture implies the vanishing conjecture.*

Proof. Let $I^\bullet \in K^b(\text{inj } \Lambda)^\perp$ be non-zero. Since $\mathcal{D}^b(\Lambda) \cong K^{+,b}(\text{inj } \Lambda)$ we may assume I^\bullet is a complex of injectives, and WLOG we may assume it concentrated in degrees $i \geq 0$, and that $d^0: I^0 \rightarrow I^1$ is not split mono. Since

if its concentrated in degrees $i \geq k$ we can just shift it, and if d^0 is split mono then replacing I^0 by 0, and I^1 by I^1/I^0 gives a homotopic complex.

$\text{Hom}(D\Lambda, I^\bullet)$ is in $\text{add Hom}(D\Lambda, D\Lambda) = \text{add } \Lambda$ so $\text{Hom}(D\Lambda, I^\bullet)$ is a complex of projectives.

$$\begin{array}{ccccc} 0 & \longrightarrow & D\Lambda & \longrightarrow & 0 \\ \downarrow & & \downarrow f & & \downarrow \\ I^{i-1} & \xrightarrow{d^{i-1}} & I^i & \xrightarrow{d^i} & I^{i+1} \end{array}$$

Since I^\bullet is in $K^b(\text{inj } \Lambda)^\perp$ and $D\Lambda$ is in $K^b(\text{inj } \Lambda)$, whenever $d^i f = 0$, f^\bullet is homotopic to 0. Meaning f factors through d^{i-1} . This means that $\text{Hom}(D\Lambda, I^\bullet)$ is an exact complex. Further since $\text{Hom}(D\Lambda, -)$ is an equivalence between $\text{inj } \Lambda$ and $\text{proj } \Lambda$ we have that $\text{Hom}(D\Lambda, d^0)$ is not split mono.

$\text{Cok Hom}(D\Lambda, d^i)$ has a projective resolution of length i . This resolution is the direct sum of its minimal resolution and an acyclic bounded complex of projectives. Since bounded acyclic complexes of projectives are split and $\text{Hom}(D\Lambda, d^0)$ is not, we must have that the minimal resolution has length i , and so $\text{findim}(\Lambda) = \infty$. \square

Theorem 1.5. [Hap93, 1.2] *The vanishing conjecture implies the Nunke condition.*

Proof. Assume there is an $X \neq 0$ with $\text{Ext}^i(D\Lambda, X) = 0$ for all $i \geq 0$. Then X considered as a stalk complex is in $K^b(\text{inj } \Lambda)^\perp$. Proceed by induction: If $I[-i] \in K^b(\text{inj } \Lambda)$ is a stalk complex then $\mathcal{D}^b(I[-i], X) = \text{Ext}^i(I, X)$. This is 0 because $D\Lambda$ is the sum of the indecomposable injectives.

Let $I \in K^b(\text{inj } \Lambda)$ be a complex of width n . WLOG assume I concentrated in degrees $0 \leq i \leq n-1$. Then

$$I^{>0} \rightarrow I \rightarrow I^0 \rightarrow I^{>0}[1]$$

is a triangle, and $I^{>0}$ has width $n-1$. Taking the long exact sequence in $\mathcal{D}^b(-, X)$ it follows that $\mathcal{D}^b(I, X) = 0$. \square

Theorem 1.6. *The Watamatsu tilting conjecture implies the Gorenstein symmetry conjecture.*

Proof. $D(\Lambda_\Lambda)$ is watamatsu tilting. WTC then gives us that if $D\Lambda$ has finite projective dimension then Λ has a finite coresolution by modules in $\text{add } D\Lambda$. In other words Λ has finite injective dimension.

For the other direction assume ${}_{\Lambda}\Lambda$ has finite injective dimension. Then $D({}_{\Lambda}\Lambda)$ has finite projective dimension, so WTC gives us that Λ_{Λ} has finite injective dimension. Which means $D(\Lambda_{\Lambda})$ has finite projective dimension. \square

Proposition 1.7. *The Auslander–Reiten conjecture is equivalent to M a generator with $\text{Ext}^n(M, M) = 0$ for $n > 0$ implies M projective.*

Do I need this, or is it obvious?

Proof. Assume ARC and that M satisfies the hypothesis. Then since M is a generator Λ is in $\text{add } M$ and thus $\text{Ext}^n(M, \Lambda) = 0$. So $\text{Ext}^n(M, M \oplus \Lambda) = 0$ and M is projective.

For the other direction Assume M satisfies $\text{Ext}^n(M, M \oplus \Lambda) = 0$. Then $\text{Ext}^n(M \oplus \Lambda, M \oplus \Lambda) = 0$, so $M \oplus \Lambda$ is projective, which means that M is projective. \square

Theorem 1.8. *The strong Nakayama conjecture implies the Auslander–Reiten conjecture.*

Proof. $\text{Ext}^i(D\Lambda, S) = \text{Ext}^i(DS, \Lambda)$, so SNC means that for every simple there is an i such that $\text{Ext}^i(S, \Lambda) \neq 0$.

Assume M is a nonprojective generator such that $\text{Ext}^n(M, M) = 0$ for all $n > 0$. Let Γ be $\text{End}(M)^{\text{op}}$, and let

$$0 \longrightarrow M \longrightarrow I_0 \longrightarrow I_1 \longrightarrow \cdots$$

be an injective resolution of M . Since $\text{Ext}^n(M, M) = 0$ when we apply $(M, -) := \text{Hom}(M, -)$ we get an exact sequence.

$$0 \longrightarrow \Gamma \longrightarrow (M, I_0) \longrightarrow (M, I_1) \longrightarrow \cdots$$

By Proposition 1.11 this is an injective resolution of Γ .

prop cited before stated

Since M is a non-projective generator it has every indecomposable projective as a summand and a nonprojective summand. So M has more indecomposable summands than Λ which means that Γ has more indecomposable projectives than Λ . It follows that Γ also has more injectives and thus has an injective not on the form (M, I) . Let Q be such an injective and let S be its socle. Then $\text{Hom}_{\Gamma}(S, (M, I_i)) = 0$ for all i , so $\text{Ext}^i(S, \Gamma) = 0$ for all i . Thus Γ does not satisfy SNC. \square

The next proposition requires part of the theory of Wedderburn projectives. The relevant theory is proven in Section 1.2 below.

Theorem 1.9. *The Auslander–Reiten conjecture implies the Nakayama conjecture.*

Proof. Assume Γ has dominant dimension ∞ , but is not self injective. Let

$$0 \longrightarrow \Gamma \longrightarrow I_0 \longrightarrow I_1$$

be an injective copresentation of Γ . Let P be the sum of the projective covers of all nonisomorphic simple modules in the socle of I_0 . Then by Proposition 1.14 we have that P is Wedderburn projective.

Let $\Lambda = \text{End}(P)^{\text{op}}$ and let $M = \text{Hom}(P, \Gamma)$. Then M is a nonprojective generator, we want to show that $\text{Ext}^{>0}(M, M) = 0$.

We have functors $(M, -): \text{mod } \Lambda \rightarrow \text{mod } \Gamma$ and $(P, -): \text{mod } \Gamma \rightarrow \text{mod } \Lambda$. By Proposition 1.11 $(M, -)$ is fully faithful and $(P, -) \circ (M, -) = \text{id}_\Lambda$.

Let $0 \rightarrow M \rightarrow Q_0 \rightarrow Q_1$ be an injective copresentation of M . Applying $(M, -)$ we get an injective copresentation of Γ . We conclude that all the projective-injective modules are in the essential image of $(M, -)$.

In other words if I^\bullet is the minimal injective resolution of Γ then $Q^\bullet := (P, I^\bullet)$ is the minimal injective resolution of M , and $(M, Q^\bullet) = I^\bullet$. This means that (M, Q^\bullet) is exact away from 0, so $\text{Ext}^{>0}(M, M) = 0$.

But then M is a nonprojective generator with $\text{Ext}^{>0}(M, M) = 0$, so Λ does not satisfy ARC. \square

Proposition 1.10. *[AR75] $\text{SNC} \Rightarrow \text{NC}$*

Proof. $\text{Ext}(D\Lambda, S) = \text{Ext}(DS, \Lambda)$. $\text{Ext}(DS, \Lambda)$ being nonzero means $I(DS)$ appears in the injective resolution of Λ . If all injectives appear in the resolution and the dominant dimension is infinity then all injectives are projective. Thus Λ is self injective. \square

This is unnecessary, but maybe nice to include?

1.2 Wedderburn correspondence

Proposition 1.11. *Let Λ be an artin algebra and M a generator. Let $\Gamma = \text{End}(M)^{\text{op}}$ and $P = (M, \Lambda)$. Then we have the following:*

- i) We have an isomorphism of ring $\text{End}_\Gamma(P)^{\text{op}} \cong \Lambda$ and an isomorphism of Λ -modules $(P_\Lambda, \Gamma) \cong M$.
- ii) The composition $(P, -) \circ (M, -)$ is the identity on $\text{mod } \Lambda$.

Proof.

- i) By Yoneda lemma we have an equivalence $(M, -): \text{add } M \rightarrow \text{add}(M, M) = \text{proj } \Gamma$. Since M is a generator Λ is in $\text{add } M$. So

$$\text{End}(P) = ((M, \Lambda), (M, \Lambda)) = \text{End}(\Lambda) = \Lambda^{\text{op}}$$

and

$$(P, \Gamma) = ((M, \Lambda), (M, M)) = (\Lambda, M) = M.$$

- ii) Let X be a Λ -module. Since $\text{add } M$ has only a finite number of indecomposables it is functorially finite. So we can take an M -resolution of X .

$$\cdots \rightarrow M_1 \rightarrow M_0 \rightarrow X \rightarrow 0$$

Since $\text{add } M$ contains the projectives this is exact. Applying $(M, -)$ we get a projective resolution of (M, X) . Since (M, X) is determined by its projective resolution and X is determined by its M -resolution we need only show that $(P, -) \circ (M, -)$ is the identity on $\text{add } M$.

Then again by Yoneda lemma $(P, (M, M')) = (\Lambda, M') = M'$.

□

Proposition 1.12. *Let M be a module and I an injective module. If the projective cover of the socle of I is a direct summand of M , then (M, I) is an injective $\Gamma := \text{End}(M)^{\text{op}}$ -module.*

Proof. Let $J \leq \Gamma$ be a left ideal and let $\psi: J \rightarrow (M, I)$ be any Γ -linear map. By Lemma 10.3 it is enough to show that ψ factors through Γ . Assume J is generated by f_i . If we can find $\gamma: M \rightarrow I$ such that $\gamma \circ f_i = \psi(f_i)$ then we would get our factorization by mapping $1 \in \Gamma$ to γ .

$$\begin{array}{ccc} \bigoplus M & & \\ \downarrow \sum f_i & \searrow \sum \psi(f_i) & \\ M & \xrightarrow{\quad \gamma \quad} & I \end{array}$$

Next we want to show that the kernel of $\sum \psi(f_i)$ contains the kernel of $\sum f_i$. To see this let K be the kernel of $\sum f_i$ and let K' be the kernel of $\sum \psi(f_i)$. If K' does not contain K , then $Q := K/K' \cap K$ is a nonzero module that is mapped injectively into I . So the socle of Q is a summand of the socle of I . Then by assumption the projective cover of the socle of Q is a direct summand of M . By the lifting property of projectives we get a map $M \rightarrow K$ such that the composition with $\sum \psi(f_i)$ is non-zero.

Let a_i be the composition $M \longrightarrow K \hookrightarrow \bigoplus M \xrightarrow{\pi_i} M$. Then we get $\sum f_i \circ a_i = 0$. Applying ψ we get $\sum \psi(f_i) \circ a_i = 0$, which gives a contradiction. Thus K' contains K .

Using this we get the following commutative diagram:

$$\begin{array}{ccc}
 \bigoplus M & & \\
 \downarrow & \searrow \sum \psi(f) & \\
 (\bigoplus M)/K & \xrightarrow{\quad} & I \\
 \downarrow & \nearrow \exists \gamma & \\
 M & &
 \end{array}
 \quad \text{with a curved arrow } \sum f_i \text{ from } \bigoplus M \text{ to } M.$$

Since I is injective it lifts monomorphisms so we know that γ exists. Thus (M, I) is an injective Γ -module. \square

Definition 1.13 (Wedderburn projective). Let Γ be an artin algebra and P a finitely generated projective. Let $\Lambda = \text{End}(P)^{\text{op}}$ and $M = (P, \Gamma)$. P is said to be *Wedderburn projective* if $\text{End}(M)^{\text{op}} = \Gamma$.

Proposition 1.14. *If P contains the projective cover of all simple modules that appear in the socle of an injective copresentation of Γ , then P is Wedderburn projective.*

To prove this we first need the next proposition as a lemma.

Proposition 1.15. *Let P be a projective Γ -module, and let $\Lambda = \text{End}(P)^{\text{op}}$. Then $(P, -): \text{mod } \Gamma \rightarrow \text{mod } \Lambda$ is fully faithful on $\text{add } I(P/JP)$.*

Proof. We want to show that the map $\text{Hom}_{\Gamma}(I, I') \rightarrow \text{Hom}_{\Lambda}((P, I), (P, I'))$ is an isomorphism. Let's first show injectivity. Let $f: I \rightarrow I'$ be a non-zero map. Then the socle of $\text{Im } f$ is a semisimple submodule of I' , so it is in $\text{add } P/JP$. Then there exists a nonzero map from P to $\text{Im } f$. Since P is projective this lifts to a map $\hat{f}: P \rightarrow I$. Then $f \circ \hat{f}$ is non-zero, so $\text{Hom}_{\Gamma}(I, I') \rightarrow \text{Hom}_{\Lambda}((P, I), (P, I'))$ is injective.

The argument for surjectivity is similar to that for Proposition 1.12. Let $\psi: (P, I) \rightarrow (P, I')$ be a Λ -linear map. Let $f_i: P \rightarrow I$ generate (P, I) as a Λ -module. Consider the diagram

$$\begin{array}{ccc} \oplus P & \xrightarrow{\sum f_i} & I \\ & \searrow \scriptstyle \Sigma \psi(f_i) & \downarrow \scriptstyle ? \\ & & I' \end{array}$$

We wish to show that there is a map at $?$ completing the diagram. We first show that $K' := \ker \sum \psi(f_i)$ contains $K := \ker \sum f_i$. Assume for the sake of contradiction that it does not. Then $Q := K/K' \cap K$ is mapped injectively into I' by $\sum \psi(f_i)$. So the socle of Q is in $\text{add } P/JP$, and we have a non-zero map $P \rightarrow Q$.

Since P is projective this extends to a map $P \rightarrow K$. Let a_i be the compositions $P \longrightarrow K \longrightarrow \oplus P \xrightarrow{\pi_i} P$. Then clearly $\sum f_i \circ a_i = 0$, but $\sum \psi(f_i) \circ a_i$ is non-zero. Since ψ is Λ -linear this is a contradiction, so K' contains K .

Then we get an induced diagram

$$\begin{array}{ccc} \oplus P & & \\ \downarrow & & \\ (\oplus P)/K & \xrightarrow{\sum f_i} & I \\ & \searrow \scriptstyle \Sigma \psi(f_i) & \downarrow \scriptstyle \exists \\ & & I' \end{array}$$

Now because I' is injective we know that there is a lift, and so $\text{Hom}_\Gamma(I, I') \rightarrow \text{Hom}_\Lambda((P, I), (P, I'))$ is surjective, and thus an isomorphism. \square

Corollary 1.15.1. *Proposition 1.14*

Proof. Let $\Gamma \rightarrow I_0 \rightarrow I_1$ be a minimal injective presentation of Γ . Then by Proposition 1.12 we have that $(P, I_0) \rightarrow (P, I_1)$ is an injective presentation of (P, Γ) . The proposition gives us that $(P, -)$ is fully faithful on I_0 and I_1 . Since the endomorphisms of Γ are exactly endomorphisms of $I_0 \rightarrow I_1$ up to homotopy this means that

$$\Gamma^{\text{op}} = \text{End}_\Gamma(\Gamma) = \text{End}_\Lambda((P, \Gamma))$$

So P is Wedderburn projective. \square

2 Recollement

Definition 2.1 (Recollement). A *recollement* between triangulated categories \mathcal{T}' , \mathcal{T} and \mathcal{T}'' is a collection of six functors satisfying:

$$\begin{array}{ccccc}
 & i^* & & j_! & \\
 & \perp & & \perp & \\
 \mathcal{T}' & \xleftarrow{i_* = i_!} & \mathcal{T} & \xleftarrow{j^! = j^*} & \mathcal{T}'' \\
 & \perp & & \perp & \\
 & i^! & & j_* &
 \end{array}$$

- (i) All functors are exact/triangulated
- (ii) $j^* i_* = 0$
- (iii) $i^* i_* \cong i^! i_! \cong \text{id}_{\mathcal{T}'}$ (induced by unit/counit)
- (iv) $j^! j_! \cong j^* j_* \cong \text{id}_{\mathcal{T}''}$
- (v) For every $X \in \mathcal{T}$ we have the following distinguished triangles:

$$j_! j^! X \xrightarrow{\varepsilon} X \xrightarrow{\eta} i_* i^* X \longrightarrow j_! j^! X[1]$$

$$i_! i^! X \xrightarrow{\varepsilon} X \xrightarrow{\eta} j_* j^* X \longrightarrow i_! i^! X[1]$$

Note that (iii) and (iv) are equivalent to i_* , $j_!$, and j_* being fully faithful.

We are specifically interested in recollements when the triangulated categories in question are (bounded) derived categories of finite dimensional algebras.

Lemma 2.2. Let $\mathcal{D}^b(\Lambda') \xrightleftharpoons[i_*]{i^*} \mathcal{D}^b(\Lambda)$ be exact functors with an adjoint pair (i^*, i_*) . Then i^* preserves bounded projective complexes and i_* preserves bounded injective complexes.

Proof. The bounded projective complexes can be characterized as the complexes P such that for any complex Y there is an integer t_Y such that $\text{Hom}(P, Y[t]) = 0$ for $t \geq t_Y$.

cite or
out-
line the
proof?

Let P be a bounded complex of projectives in $\mathcal{D}^b(\Lambda)$. Then we want to show that i^*P is as well. Let Y be any complex in $\mathcal{D}^b(\Lambda')$. Then $\mathcal{D}^b(\Lambda')(i^*P, Y[t]) = \mathcal{D}^b(\Lambda)(P, i_*Y[t])$, so since P is a bounded complex of projectives there is t_Y such that this vanishes for $t \geq t_Y$.

Hom-set
notation

The statement for injectives is exactly dual. \square

Lemma 2.3. *Let $\mathcal{D}^b(\Lambda') \xrightleftharpoons[i^!]{i^*} \mathcal{D}^b(\Lambda)$ be exact functors with adjoint pairs (i^*, i_*) and $(i_*, i^!)$. Then the homology of i_*X is uniformly bounded for $X \in \text{mod } \Lambda'$ considered as a complex concentrated in degree 0. I.e. there is an r , independent of X , such that $H^j(i_*X) = 0$ for $j \notin (-r, r)$.*

Proof. We first prove that there is an r' , independent of X , such that $H^j(i_*X) = 0$ for $j \geq r'$. Let P be $i^*\Lambda \in \mathcal{D}^b(\Lambda')$. Then by Lemma 2.2 P is a bounded complex of projectives.

Thus there is an r' such that $P^{-j} = 0$ for $j \geq r'$. Then

$$\mathcal{D}^b(\Lambda')(P, X[j]) = \mathcal{D}^b(\Lambda)(\Lambda, i_*X[j]) = H^j(i_*X) = 0$$

for $j \geq r'$ and any Λ' -module X , when considered as a complex concentrated in degree 0.

Next we prove that there is an r'' such that $H^{-j}(i_*X) = 0$ for $j \geq r''$. The argument is completely dual. Let I be $i^!D\Lambda \in \mathcal{D}^b(\Lambda') = K^{+,b}(\text{inj } \Lambda')$. Then again by Lemma 2.2 I is a bounded complex of injectives.

Thus there is an r'' such that $I^j = 0$ for $j \geq r''$. Then

$$\mathcal{D}^b(\Lambda')(X, I[j]) = \mathcal{D}^b(\Lambda)(i_*X, D\Lambda[j]) = H^{-j}(i_*X) = 0$$

for $j \geq r''$ and any Λ' -module X , when considered as a complex concentrated in degree 0.

Letting r be the maximum of r' and r'' we get that $H^j(X)$ is zero outside of $(-r, r)$. \square

Theorem 2.4. [Hap93, 3.3] *Given a recollement between bounded derived categories*

$$\begin{array}{ccccc}
 & i^* & & j^! & \\
 & \downarrow & & \downarrow & \\
 \mathcal{D}^b(\Lambda') & \xrightarrow{i_* = i_!} & \mathcal{D}^b(\Lambda) & \xrightarrow{j^! = j^*} & \mathcal{D}^b(\Lambda''), \\
 & \uparrow & & \uparrow & \\
 & i^! & & j_* &
 \end{array}$$

then we have that $\text{findim}(\Lambda) < \infty \iff \text{findim}(\Lambda') < \infty$ and $\text{findim}(\Lambda'') < \infty$.

Proof. Assume $\text{findim}(\Lambda) < \infty$. We begin by showing that $\text{findim}(\Lambda') < \infty$.

Let $T = \Lambda' / \text{rad } \Lambda'$. Then the projective dimension of X is the largest t for which $\text{Ext}^t(X, T) \neq 0$. Let X be a module in $\text{mod } \Lambda'$ with finite projective dimension. We consider X as a complex concentrated in degree 0. Then since X is isomorphic to its projective resolution, by Lemma 2.2 $i_* X$ is a bounded complex of projectives. Say:

$$i_* X = 0 \rightarrow P^{-s} \rightarrow \dots \rightarrow P^{s'} \rightarrow 0$$

By Lemma 2.3 we know there is an r independent of X such that $H^{-j}(X) = 0$ for $j \geq r$. Truncating $i_* X$ at $-r$ gives a projective resolution of $\ker d_{i_* X}^{-r}$. Since $\text{findim}(\Lambda) < \infty$ this means that $s \leq r + \text{findim}(\Lambda)$.

Since $i_* T$ is in $\mathcal{D}^b(\Lambda)$ it is a bounded complex, in particular there is a t_0 such that $i_* T^t = 0$ for $t \geq t_0$. Then by the bounds above $\mathcal{D}^b(\Lambda)(i_* X, i_* T[t]) = 0$ for $t \geq t_0 + s \geq t_0 + r + \text{findim}(\Lambda)$. Since i_* is fully faithful this equals $\mathcal{D}^b(\Lambda')(X, T[t])$, and so $\text{findim}(\Lambda') \leq t_0 + r + \text{findim}(\Lambda)$. In particular it is finite.

The proof for $\text{findim}(\Lambda'')$ is the same, just replacing i_* with $j_!$.

For the converse assume Λ' and Λ'' both have finite finitistic dimension. Let $T = \Lambda / \text{rad } \Lambda$, and X be a Λ -module with finite projective dimension, and consider both modules as a complex concentrated in degree 0. By Definition 2.1v we have distinguished triangles:

$$j_! j^! X \longrightarrow X \longrightarrow i_* i^* X \longrightarrow j_! j^! X[1]$$

$$i_! i^! T \longrightarrow T \longrightarrow j_* j^* T \longrightarrow i_! i^! T[1]$$

Let $(-, -)_m := \mathcal{D}^b(\Lambda)(-, -[m])$, and $X_j := j_! j^! X$, $X_i := i_* i^* X$, $T_i := i_! i^! T$, $T_j = j_* j^* T$. Then we have long exact sequences:

$$\cdots \longrightarrow (X, T_i)_m \longrightarrow (X, T)_m \longrightarrow (X, T_j)_m \longrightarrow (X, T_i)_{m+1} \longrightarrow \cdots$$

$$\cdots \longrightarrow (X_i, T_i)_m \longrightarrow (X, T_i)_m \longrightarrow (X_j, T_i)_m \longrightarrow (X_i, T_i)_{m+1} \longrightarrow \cdots$$

$$\cdots \longrightarrow (X_i, T_j)_m \longrightarrow (X, T_j)_m \longrightarrow (X_j, T_j)_m \longrightarrow (X_i, T_j)_{m+1} \longrightarrow \cdots$$

We have

$$(X_i, T_j)_m = (i_* i^* X, j_* j^* T)_m = (j^* i_* i^* X, j^* T)_m = 0$$

and

$$(X_j, T_i)_m = (j_! j^! X, i_! i^! T)_m = (j^! X, j^! i_! i^! T)_m = 0$$

which combined with long exact sequences gives us that $(X_i, T_i)_m = (X, T_i)_m$ and $(X_j, T_j)_m = (X, T_j)_m$. If we can show that $(X_i, T_i)_m$ and $(X_j, T_j)_m$ are bounded, then $(X, T_i)_m$ and $(X, T_j)_m$, and consequently $(X, T)_m$ would be bounded. Which would give a bound on the projective dimension of X .

We start by bounding $(X, T_i)_m = (X_i, T_i)_m$. First note that

$$(X_i, T_i)_m = (i_* i^* X, i_! i^! T)_m = (i^* i_* i^* X, i^! T)_m = (i^* X, i^! T)_m$$

Since X has finite projective dimension we can think of it as a bounded complex of projectives. Then by Lemma 2.2 $i^* X$ is as well. By the second half of Lemma 2.3 (using (i^*, i_*) instead of $(i_*, i^!)$) we have that there is an r such that $H^{-j}(i^* X) = 0$ for all $j \geq r$. This means that thinking of $i^* X$ as a complex of projectives, it is 0 in degree $-t$ for all $t \geq r + \text{pd ker } d_{i^* X}^{-r}$, in particular it is 0 for all $t \geq r + \text{findim}(\Lambda')$. Since $i^! T$ is a bounded complex, it has an upper bound, say t_0 . Thus $(i^* X, i^! T)_m = 0$ for all $m \geq t_0 + r + \text{findim}(\Lambda')$.

The bound on $(X, T_j)_m$ is similar, using the finitistic dimension of Λ'' . Taking the maximum of these two bounds we get a bound on $(X, T)_m$, which gives a bound on the projective dimension independent of X , hence a bound on $\text{findim}(\Lambda)$. \square

2.1 Triangular matrix rings and vertex removal

Something somethign triangular.

Definition 2.5 (Comma category). Let \mathcal{A} and \mathcal{B} be categories and let $F: \mathcal{A} \rightarrow \mathcal{B}$ be a functor. Then the *comma category* (F, \mathcal{B}) has as objects triplets (A, B, f) with $A \in \mathcal{A}$, $B \in \mathcal{B}$, and $f: FA \rightarrow B$ a morphism in \mathcal{B} . The morphisms are pairs $(\alpha, \beta): (A, B, f) \rightarrow (A', B', f')$ with $\alpha: A \rightarrow A'$ and $\beta: B \rightarrow B'$ such that the following diagram commutes:

$$\begin{array}{ccc} FA & \xrightarrow{f} & B \\ F\alpha \downarrow & & \downarrow \beta \\ FA' & \xrightarrow{f'} & B'. \end{array}$$

Proposition 2.6. *If \mathcal{A} and \mathcal{B} are abelian categories and F is right exact, then the comma category (F, \mathcal{B}) is abelian.*

Proof. We need to show that (F, \mathcal{B}) has kernels, has cokernels, and that image equals coimage. First we show that it contains kernels. Let $(\alpha, \beta): (A, B, f) \rightarrow (A', B', f')$ be a morphism in the comma category. Then we have a diagram:

$$\begin{array}{ccccccc} F \ker \alpha & \xrightarrow{F\iota_\alpha} & FA & \xrightarrow{F\alpha} & FA' \\ \downarrow \theta & & \downarrow f & & \downarrow f' \\ 0 & \longrightarrow & \ker \beta & \xrightarrow{\iota_\beta} & B & \xrightarrow{\beta} & B' \end{array}$$

Since $\beta f F \iota_\alpha = f' F \alpha F \iota_\alpha = 0$ there is a unique θ making the diagram commute. I claim the kernel of (α, β) is $(\ker \alpha, \ker \beta, \theta)$. Indeed if (α', β') is any map such that $(\alpha, \beta) \circ (\alpha', \beta') = 0$ then $\alpha \alpha' = 0$ and $\beta \beta' = 0$ so both α' and β' factor uniquely through ι_α and ι_β .

$$\begin{array}{ccccc} FA'' & \xrightarrow{\alpha''} & F \ker \alpha & \xrightarrow{F\iota_\alpha} & FA \\ \downarrow f'' & & \downarrow \theta & & \downarrow f \\ B'' & \xrightarrow{\beta''} & \ker \beta & \xrightarrow{\iota_\beta} & B \end{array}$$

The only thing left to verify is that the left square commutes. This follows from the outer rectangle commuting, and that ι_β is a monomorphism.

Showing that cokernels exists is similar, but relies on F being right exact. The construction is completely dual, but to verify commutativity at the end instead of using that ι_β is mono we must use that $F\pi_\alpha: FA' \rightarrow F \text{Cok } \alpha$ is an epimorphism. This follows from F being right exact. I leave the details to the reader.

or do I?

Now the image equaling the coimage follows from \mathcal{A} and \mathcal{B} being abelian, and the way we constructed the kernels and cokernels. \square

For the rest of this section we assume F is a right exact functor between abelian categories so that the comma category is abelian. We also assume \mathcal{A} and \mathcal{B} has enough projectives, whenever we mention projective objects. In particular we are interested in the case when \mathcal{A} and \mathcal{B} are module categories over finite dimensional algebras.

Definition 2.7. For \mathcal{A} and \mathcal{B} abelian categories and F right exact we define the following functors:

$$\begin{aligned} U: (F, \mathcal{B}) &\longrightarrow \mathcal{A} \times \mathcal{B} & T: \mathcal{A} \times \mathcal{B} &\longrightarrow (F, \mathcal{B}) \\ (A, B, f) &\longmapsto (A, B) & (A, B) &\longmapsto (A, B \oplus FA, FA \hookrightarrow FA \oplus B) \\ (\alpha, \beta) &\longmapsto (\alpha, \beta) & (\alpha, \beta) &\longmapsto (\alpha, F\alpha \oplus \beta) \end{aligned}$$

$$\begin{aligned} C: (F, \mathcal{B}) &\longrightarrow \mathcal{A} \times \mathcal{B} & Z: \mathcal{A} \times \mathcal{B} &\longrightarrow (F, \mathcal{B}) \\ (A, B, f) &\longmapsto (A, \text{Cok } f) & (A, B) &\longmapsto (A, B, 0) \\ (\alpha, \beta) &\longmapsto (\alpha, \hat{\beta}) & (\alpha, \beta) &\longmapsto (\alpha, \beta) \end{aligned}$$

Proposition 2.8. *With the definitions above U and Z become exact functors.*

Proof. Using the characterization of exact sequences shown in Proposition 2.6 a short exact sequence in (F, \mathcal{B}) is a commutative diagram

$$\begin{array}{ccccccc} FA'' & \xrightarrow{F\alpha'} & FA & \xrightarrow{F\alpha} & FA' & \longrightarrow & 0 \\ \downarrow f'' & & \downarrow f & & \downarrow f' & & \\ 0 & \longrightarrow & B'' & \xrightarrow{\beta'} & B & \xrightarrow{\beta} & B' \longrightarrow 0 \end{array}$$

such that the sequences

$$\begin{aligned} 0 &\longrightarrow A'' \xrightarrow{\alpha'} A \xrightarrow{\alpha} A' \longrightarrow 0 \\ 0 &\longrightarrow B'' \xrightarrow{\beta'} B \xrightarrow{\beta} B' \longrightarrow 0 \end{aligned}$$

are short exact. Since when we apply U we simply get the product of these two sequences, U is exact.

Similarly for Z since the two sequences we start with are assumed to be exact the resulting sequence will be exact by the characterization in Proposition 2.6. \square

Proposition 2.9. *[FGR75, Proposition 1.3] The pairs of functors (T, U) and (C, Z) form adjoint pairs.*

Proof. We want to establish an isomorphism

$$\mathrm{Hom}(T(A, B), (A', B', FA' \rightarrow B')) \cong \mathrm{Hom}((A, B), (A', B')).$$

A morphism in $\mathrm{Hom}(T(A, B), (A', B', FA' \rightarrow B'))$ is given by a commutative diagram

$$\begin{array}{ccc} FA & \xrightarrow{\begin{bmatrix} 0 \\ 1 \end{bmatrix}} & B \oplus FA \\ F\alpha \downarrow & & \downarrow \begin{bmatrix} \beta & \gamma \end{bmatrix} \\ FA' & \xrightarrow{f} & B' \end{array}$$

The isomorphism is then given by sending this to (α, β) . This is clearly surjective.

For injectivity assume $(\alpha, \beta) = 0$, then $\gamma = \begin{bmatrix} \beta & \gamma \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = fF\alpha = 0$. So the map is injective, and (T, U) is an adjoint pair.

Next we consider (C, Z) . We want an isomorphism

$$\mathrm{Hom}(C(A, B, f), (A', B')) = \mathrm{Hom}((A, \mathrm{Cok} f), (A', B')) \cong \mathrm{Hom}((A, B, f), (A', B', 0)).$$

A morphism in $\mathrm{Hom}((A, B, f), (A', B', 0))$ is a commutative diagram

$$\begin{array}{ccc} FA & \xrightarrow{f} & B \\ F\alpha \downarrow & & \downarrow \beta \\ FA' & \xrightarrow{0} & B' \end{array}$$

Since $\beta f = 0$, we have that β factors through the cokernel of f uniquely. Let the factorization be given by the map $\beta': \mathrm{Cok} f \rightarrow B'$. Then we send this diagram to (α, β') . Since the choice of β' was unique this is an isomorphism, so (C, Z) is an adjoint pair. \square

Corollary 2.9.1. *The functors T and C preserve projective objects.*

Proof. What we need to check is that for projective objects P and Q in $(\mathcal{A} \times \mathcal{B})$ and (F, \mathcal{B}) respectively we have that $\text{Hom}(TP, -)$ and $\text{Hom}(CQ, -)$ are exact. By adjointness these are equal to $\text{Hom}(P, U-)$ and $\text{Hom}(Q, Z-)$ respectively. Since U and Z are exact this holds, and so T and C preserve projective objects. \square

Proposition 2.10. *[FGR75, Corollary 1.6c] For a projective object P in (F, \mathcal{B}) we have that $T(C(P)) \cong P$, in particular all projectives are of the form $T(P')$ for a projective $P' \in \mathcal{A} \times \mathcal{B}$.*

Proof. Let P be given by $f: FA \rightarrow B$. Applying C we get $(A, \text{Cok } f)$. We have morphisms $P \rightarrow ZC(P)$ and $TC(P) \rightarrow ZC(P)$ given by the following diagram

$$\begin{array}{ccc} FA & \xrightarrow{f} & B \\ \parallel & & \downarrow \\ FA & \xrightarrow{0} & \text{Cok } f \\ \parallel & & \uparrow \\ FA & \hookrightarrow & \text{Cok } f \oplus FA \end{array}$$

By the projective property of P there is some morphism β factorizing the map $P \rightarrow ZC(P)$ giving us the diagram:

$$\begin{array}{ccc} FA & \xrightarrow{f} & B \\ \parallel & & \downarrow \beta \\ FA & \hookrightarrow & \text{Cok } f \oplus FA \\ \parallel & & \downarrow \\ FA & \xrightarrow{0} & \text{Cok } f \end{array}$$

Since $FA \hookrightarrow \text{Cok } f \oplus FA$ is split mono, f is split mono, and consequently β is an isomorphism. So we have $P \cong TC(P)$. \square

Proposition 2.11. *[FGR75, Lemma 4.16] Let $X = (A, B, f)$ be an object in the comma category. Then $\text{pd } X \geq \text{pd } A$, and if $A = 0$ then $\text{pd } X = \text{pd } B$.*

Proof. We first show that $\text{pd } X \geq \text{pd } A$. Note that $\text{pd } C(X) = \max\{\text{pd } A, \text{pd } \text{Cok } f\}$ so we always have $\text{pd } C(X) \geq \text{pd } A$. If $\text{pd } X = \infty$ then the statement holds so let us assume $\text{pd } X = n < \infty$. We proceed by induction on n . If $n = 0$ then $C(X)$ is projective so $\text{pd } X = \text{pd } C(X) = \text{pd } A = 0$. Next assume the statement holds for whenever the projective dimension is less than n . Let $P \rightarrow A$ and $P' \rightarrow \text{Cok } f$ be epimorphisms from projectives. Then we have an epimorphism $T(P, P') \rightarrow X$. If we let ΩA be the kernel of $P \rightarrow A$ and $X' = (\Omega A, K, \theta)$ be the kernel of $T(P, P') \rightarrow X$ as shown in the following diagram

$$\begin{array}{ccccccc} F\Omega A & \longrightarrow & FP & \longrightarrow & FA & \longrightarrow & 0 \\ \theta \downarrow & & \downarrow & & \downarrow f & & \\ 0 & \longrightarrow & K & \longrightarrow & P' \oplus FP & \longrightarrow & B \longrightarrow 0 \end{array}$$

Then we have $\text{pd } A \leq \text{pd } \Omega A + 1$ and $\text{pd } X = \text{pd } X' + 1$. By induction we have that $\text{pd } X' \geq \text{pd } \Omega A$ and so $\text{pd } X \geq \text{pd } \Omega A + 1 \geq \text{pd } A$.

If $A = 0$ then we can associate $C(X) = (0, B)$ with B . Any projective resolution P_B^\bullet of B gives a resolution of X by $T(0, P_B^\bullet)$, and any resolution P_X^\bullet of X gives a resolution of $(0, B)$ by $C(P_X^\bullet)$. \square

Theorem 2.12. [FGR75, Theorem 4.20] *The finitistic dimension of the comma category (F, \mathcal{B}) is bounded above by $\text{findim}(\mathcal{A}) + \text{findim}(\mathcal{B}) + 1$.*

Proof. Let $X = (A, B, f)$ be an element of the comma category with finite projective dimension. Let P_A^\bullet be a projective resolution of A shorter than $\text{findim}(\mathcal{A})$. Similar to what we did in Proposition 2.11 define P_X^0 to be $T(P_A^0, P(\text{Cok } f))$ where $P(\text{Cok } f)$ is a projective module with an epimorphism onto $\text{Cok } f$. Then we have that the kernel of $P_X^0 \rightarrow X$ is

$F\Omega A \xrightarrow{\theta^0} K^0$. We continue inductively defining P_X^n to be $T(P_A^n, \text{Cok } \theta^{n-1})$. Then $\Omega^{\text{findim}(\mathcal{A})+1} X = (0, K^{\text{findim}(\mathcal{A})}, 0)$. Then by Proposition 2.11 we know that $\text{pd } \Omega^{\text{findim}(\mathcal{A})+1} X = \text{pd } K^{\text{findim}(\mathcal{A})} \leq \text{findim}(\mathcal{B})$. So $\text{pd } X \leq \text{findim}(\mathcal{A}) + \text{findim}(\mathcal{B}) + 1$. \square

Example 2.13. If k is a field, $\mathcal{A} = \mathcal{B} = \text{mod } k$ and F is the identity, then the comma category (F, \mathcal{B}) is equivalent to the category of finite dimensional representations of A_2 over k . Then \mathcal{A} and \mathcal{B} both have finitistic dimension 0 while (F, \mathcal{B}) has finitistic dimension 1. So the bound shown above is tight.

Definition 2.14 (Triangular matrix ring). Let R and S be rings, and let M be an $S - R$ -bimodule. Then the *triangular matrix ring* $\begin{pmatrix} R & 0 \\ M & S \end{pmatrix}$ is the ring

of all matrices $\begin{bmatrix} r & 0 \\ m & s \end{bmatrix}$ with $r \in R$, $s \in S$, and $m \in M$. The multiplication is given by

$$\begin{bmatrix} r & 0 \\ m & s \end{bmatrix} \begin{bmatrix} r' & 0 \\ m' & s' \end{bmatrix} = \begin{bmatrix} rr' & 0 \\ mr' + sm' & ss' \end{bmatrix}.$$

Notice, if N is a module over the matrix ring $\begin{pmatrix} R & 0 \\ M & S \end{pmatrix}$, then as an abelian group N splits as a direct sum into

$$N = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} N \oplus \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} N.$$

By restriction of scalars we can think of $N_R := \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} N$ as an R -module and $N_S := \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} N$ as an S -module. Further multiplication by $\begin{bmatrix} 0 & 0 \\ m & 0 \end{bmatrix}$ is 0 on N_S and maps N_R into N_S . So N consists of an R -module N_R , an S -module N_S and a $S - R$ -linear map $M \rightarrow \text{Hom}_{\mathbb{Z}}(N_R, N_S)$, or equivalently a S -linear map $M \otimes_R N_R \rightarrow N_S$. This means that $\text{mod} \begin{pmatrix} R & 0 \\ M & S \end{pmatrix}$ is equivalent to the comma category $(\text{mod } R, \text{mod } S, M \otimes_R -)$. So we have that

$$\text{findim} \begin{pmatrix} R & 0 \\ M & S \end{pmatrix} \leq \text{findim}(R) + \text{findim}(S) + 1$$

Example

3 Contravariant finiteness

Definition 3.1 (Resolving). A full subcategory of an abelian category is called *resolving* if

- i) It is closed under extensions.
- ii) It contains the projectives.
- iii) It contains the kernels of its epimorphisms.

Note that the subcategory of modules with finite projective dimension is resolving.

Lemma 3.2. *Let \mathcal{X} be resolving. Then $\text{Ext}^1(\mathcal{X}, Y) = 0$ implies that $\text{Ext}^i(\mathcal{X}, Y) = 0$ for all $i \geq 1$.*

Proof. Since \mathcal{X} contains the projectives, ΩX is the kernel of an epimorphism in \mathcal{X} . Thus \mathcal{X} contains all syzygies. $\text{Ext}^i(X, Y) = \text{Ext}^1(\Omega^{i-1}X, Y) = 0$. \square

Proposition 3.3. *If \mathcal{X} is resolving, then $\mathcal{Y} := \ker \text{Ext}^{\geq 1}(\mathcal{X}, -) = \ker \text{Ext}^1(\mathcal{X}, -)$ is closed under extensions.*

Proof. Let $0 \rightarrow Y \rightarrow E \rightarrow Y' \rightarrow 0$ be an extension of objects in \mathcal{Y} , and let X be an object of \mathcal{X} . Then we get an exact sequence

$$0 = \text{Ext}^i(X, Y) \longrightarrow \text{Ext}^i(X, E) \longrightarrow \text{Ext}^i(X, Y') = 0$$

Thus $\text{Ext}^i(X, E) = 0$ for all $i \geq 1$ and E is in \mathcal{Y} . \square

Lemma 3.4. *Let \mathcal{X} be a contravariantly finite, resolving subcategory of $\text{mod } \Lambda$. Then for every object $C \in \text{mod } \Lambda$ there is a short exact sequence*

$$0 \rightarrow Y \rightarrow X \rightarrow C \rightarrow 0$$

with $X \rightarrow C$ minimal \mathcal{X} -approximation and $\text{Ext}^i(\mathcal{X}, Y) = 0$ for all $i \geq 1$.

Proof. Since \mathcal{X} is contravariantly finite, C has a minimal \mathcal{X} -approximation $X \rightarrow C$. Since \mathcal{X} contains the projective cover of C this approximation must be an epimorphism. So it is part of a short exact sequence

$$0 \rightarrow Y \rightarrow X \rightarrow C \rightarrow 0.$$

Let X' be an arbitrary object in \mathcal{X} . Taking the long exact sequence in $\text{Ext}(X', -)$ gives us

$$\begin{array}{ccccccc} \text{Hom}(X', Y) & \longrightarrow & \text{Hom}(X', X) & \longrightarrow & \text{Hom}(X', C) & & \\ & & & & & \searrow & \\ & & & & & & \text{Ext}^1(X', Y) \longrightarrow \text{Ext}^1(X', X) \longrightarrow \text{Ext}^1(X', C) \end{array}$$

Since $X \rightarrow C$ is an approximation, we know that $\text{Hom}(X', X) \rightarrow \text{Hom}(X', C)$ is epi. Thus if we can prove that $\text{Ext}^1(X', X) \rightarrow \text{Ext}^1(X', C)$ is mono we would have that $\text{Ext}^1(X', Y) = 0$. Assume we have an element of $\text{Ext}^1(X', X)$ that is mapped to 0, i.e. we have a commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & X & \longrightarrow & E & \longrightarrow & X' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & C & \longrightarrow & C \oplus X' & \longrightarrow & X' \longrightarrow 0
 \end{array}$$

Since \mathcal{X} is closed under extensions E is in \mathcal{X} . By composing with projection $C \oplus X' \rightarrow C$ we get a commutative triangle

$$\begin{array}{ccc}
 X & \longrightarrow & E \\
 \downarrow & \swarrow & \\
 C & &
 \end{array}$$

since $X \rightarrow C$ is an approximation we get that $E \rightarrow C$ factors through X . The endomorphism $X \rightarrow E \rightarrow X$ leaves the approximation unchanged, so by minimality it must be an isomorphism. Hence

$$0 \rightarrow X \rightarrow E \rightarrow X' \rightarrow 0$$

is split and $\text{Ext}(X', X) \rightarrow \text{Ext}(X', C)$ is injective. Thus $\text{Ext}(X', Y) = 0$. \square

Theorem 3.5. [AR91, 3.8] *Let \mathcal{X} be a contravariantly finite, resolving subcategory of $\text{mod } \Lambda$. Let X_i be the minimal approximation of S_i . Then any $X \in \mathcal{X}$ is a direct summand of an X_i -filtered module.*

Proof. The first part of the proof is to show by induction on length that any module C is in an exact sequence $0 \rightarrow Y \rightarrow X \rightarrow C \rightarrow 0$ with X X_i -filtered and $\text{Ext}^1(\mathcal{X}, Y) = 0$.

For the base case if $C = S_i$ is simple then by Lemma 3.4 we have an exact sequence $0 \rightarrow Y \rightarrow X_i \rightarrow C \rightarrow 0$ with the desired properties stated above.

For the induction step, assume it holds for all modules of length less than n , and let C be a module of length n . Then by Jordan-Hölder C is the extension of two modules of length less than n . Say

$$0 \longrightarrow C' \longrightarrow C \longrightarrow C'' \longrightarrow 0$$

Applying the induction hypothesis we get a diagram on the form

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & Y' & & Y'' & & \\
 & & \downarrow & & \downarrow & & \\
 & & X' & & X'' & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & C' & \longrightarrow & C & \longrightarrow & C'' \longrightarrow 0 \\
 & & \downarrow & & & & \downarrow \\
 & & 0 & & & & 0
 \end{array}$$

Taking the pullback of $X'' \rightarrow C''$ we get a diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C' & \longrightarrow & E & \longrightarrow & X'' \longrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C' & \longrightarrow & C & \longrightarrow & C'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Since Y' satisfies $\text{Ext}^1(\mathcal{X}, Y') = 0$ by Lemma 3.2 it also satisfies $\text{Ext}^2(\mathcal{X}, Y') = 0$. In particular from the long exact sequence

$$0 = \text{Ext}^1(X'', Y) \rightarrow \text{Ext}^1(X'', X') \rightarrow \text{Ext}^1(X'', C) \rightarrow \text{Ext}^2(X'', Y) = 0$$

we get that $X' \rightarrow C'$ induces an isomorphism $\text{Ext}^1(X'', X') \rightarrow \text{Ext}^1(X'', C)$. Thus the short exact sequence $0 \rightarrow C' \rightarrow E \rightarrow X'' \rightarrow 0$ must come from a sequence $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$. This gives us a diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & Y' & & Y'' & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & X' & \longrightarrow & X & \longrightarrow & X'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C' & \longrightarrow & C & \longrightarrow & C'' \longrightarrow 0 \\
 & & \downarrow & & & & \downarrow \\
 & & 0 & & & & 0
 \end{array}$$

Applying the Snake Lemma we can fill out the diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & Y' & \longrightarrow & Y & \longrightarrow & Y'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & X' & \longrightarrow & X & \longrightarrow & X'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & C' & \longrightarrow & C & \longrightarrow & C'' \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Since X is an extension of X_i -filtered modules, it is also X_i -filtered. Since Y is the extension of Y'' and Y' it follows from Proposition 3.3 that $\text{Ext}(\mathcal{X}, Y) = 0$.

Hence any C fits into a sequence $0 \rightarrow Y \rightarrow X \rightarrow C \rightarrow 0$ with X being X_i -filtered and $\text{Ext}^{\geq 1}(\mathcal{X}, Y) = 0$.


Now suppose that C is in \mathcal{X} , and let $0 \rightarrow Y \rightarrow X \rightarrow C \rightarrow 0$ be as before. Then we get that

$$\text{Hom}(C, X) \longrightarrow \text{Hom}(C, C) \longrightarrow \text{Ext}^1(C, Y) = 0$$

is exact, and thus C is a direct summand of X . So every object in \mathcal{X} is a direct summand of an X_i -filtered module. \square


Corollary 3.5.1. *If the subcategory of modules with finite projective dimension is contravariantly finite, then the finitistic dimension is the supremum of the projective dimension of X_i . In particular it is finite.*

Example 3.6. [IST90, Proposition 2.3] Let Λ be the path algebra of



with relations $\alpha\gamma$, $\beta\gamma$, and $\gamma\alpha$ over an algebraically closed field k . Then $\text{findim}(\Lambda) = 1$, but the subcategory of modules with finite projective dimension is not contravariantly finite.

Proof. The indecomposable projective Λ -modules are given by the following quivers



Note that both the indecomposable projectives have even dimension, so any projective module has even dimension. Then if X is a module with finite projective dimension, since $\dim X = \sum (-1)^i \dim P_X^i$ the dimension of X is also even. In particular the two simple modules have infinite projective dimension.

The radical of P_1 is P_2 and the radical of P_2 is S_1 , so the radical of an arbitrary projective looks like $P_2^n \oplus S_1^m$. Let $P \rightarrow X$ be the projective cover of a module with finite projective dimension. Then ΩX is a submodule of $JP = P_2^n \oplus S_1^m$. Let M be an indecomposable summand of ΩX , and consider the composition $M \rightarrow JP \rightarrow P_2$ for any possible projection to P_2 . If this is epi then we must have $M = P_2$. If none of these are epi then M is contained in $JP_2^n \oplus S_1^m = S_1^{m+n}$. This would mean $M = S_1$, but S_1 has infinite projective dimension. Thus we must have ΩX projective, and so $\text{pd } X \leq 1$.

Nextly we want to show that S_1 has no minimal approximation by modules with finite projective dimension. Assume for the sake of contradiction that


$X \rightarrow S_1$ is such a minimal approximation. Then we claim that P_2 is not a submodule of X . Since $\text{Hom}(P_2, S_1) = 0$ if this were the case then $X' = X/P_2$ would give an approximation of shorter length, because X' would also have finite projective dimension. Which can be seen in the diagram below.

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & P_X^1 & \longrightarrow & P_X^1 \oplus P_1 & \longrightarrow & P_1 \longrightarrow 0 \\
 & & \parallel & & \downarrow \lrcorner & & \downarrow \\
 0 & \longrightarrow & P_X^1 & \longrightarrow & P_X^0 & \longrightarrow & X \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & X' & = & X' \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

make a notation chapter setting notation for projective cover / resolution

This means that $\gamma X = 0$, because if there was an element $x \in X$ with $\gamma x \neq 0$, then $(e_2 x)$ would be a submodule of X isomorphic to P_2 . So X is a $\Lambda/(\gamma)$ module.

The algebra $\Lambda/(\gamma)$ is the path algebra of the 2-Kronecker quiver, whose representation theory is well understood. Specifically $\Lambda/(\gamma)$ can be associated with the subquiver highlighted below.



The indecomposable modules are as given in the table below.

$$\begin{array}{ccc}
 \begin{array}{c} \begin{array}{ccc} k^n & \xrightarrow{\begin{bmatrix} I_n \\ 0 \end{bmatrix}} & k^{n+1} \\ \text{preprojective} \end{array} \end{array} &
 \begin{array}{c} \begin{array}{ccc} k^n & \xrightarrow{\begin{array}{c} J(n,\lambda) \\ I_n \end{array}} & k^n \\ \text{regular} \end{array} \end{array} &
 \begin{array}{c} \begin{array}{ccc} k^{n+1} & \xrightarrow{\begin{bmatrix} I_n & 0 \\ 0 & I_n \end{bmatrix}} & k^n \\ \text{preinjective} \end{array} \end{array}
 \end{array}$$

We see that the preprojective and preinjective modules both have odd dimension, so they will have infinite projective dimension as Λ -modules. We

can easily verify that the $\Lambda/(\gamma)$ -modules $k \xrightleftharpoons[1]{\lambda} k$ all have finite projec-

tive dimension as Λ -modules and that they have a nonzero map onto S_1 . So each of these modules would need to have a nonzero map to X . But it is easy to verify that there is a nonzero homomorphism between the regular modules only if they have the same value of λ . So for it to be possible for X to factorize all these maps we would need X to have an infinite amount of direct summands. Since we are working with finitely generated modules this is impossible, hence S_1 has no approximation, and the subcategory is not contravariantly finite. \square

In the next example we look at the opposite algebra of Λ to show that there is not necessarily any link between the contravariant finiteness for Λ and for Λ^{op} .

Example 3.7. Let Γ be the opposite algebra of the one in Example 3.6. That is, Γ is the path algebra of

$$\begin{array}{ccc}
 & \hat{\alpha} & \\
 2 & \xrightarrow{\hat{\beta}} & 1 \\
 & \hat{\gamma} &
 \end{array}$$

with relations $\hat{\gamma}\hat{\alpha}$, $\hat{\gamma}\hat{\beta}$, and $\hat{\alpha}\hat{\gamma}$. Then the subcategory of modules with finite projective dimension is contravariantly finite. In other words the subcategory of Λ -modules with finite injective dimension is covariantly finite.

Proof. The indecomposable projective Γ -modules are given by the following quivers

$$\begin{array}{ccc} & & 1 \\ & & \downarrow \hat{\gamma} \\ & & 2 \\ & & \downarrow \hat{\beta} \\ & & 1 \end{array} \qquad \begin{array}{ccc} & 2 & \\ \hat{\alpha} \swarrow & & \searrow \hat{\beta} \\ 1 & & 1 \end{array}$$

Similar to before before, notice that the indecomposable projective modules are 3-dimensional and thus every module with finite projective dimension will have a k -dimension that is a multiple of 3. So in particular the simple modules have infinite projective dimension.

Let X be a module with finite projective dimension, and let P be its projective cover. We have that ΩX is a submodule of JP . Notice that $\hat{\alpha}J = \hat{\gamma}J = 0$, so ΩX is a $\Gamma/(\hat{\alpha}, \hat{\gamma})$ -module. But $\Gamma/(\hat{\alpha}, \hat{\gamma})$ is simply isomorphic to the path algebra of $2 \longrightarrow 1$, over which there are just 3 indecomposable modules. We already know that the simple modules cannot be summands of ΩX , because they have infinite projective dimension. The non-simple module $k \xrightarrow{1} k$ is 2-dimensional and thus also has infinite projective dimension over Γ . So we conclude that $\Omega X = 0$, so X is projective.

So the only modules with finite projective dimension are the projectives themselves. In particular there are only a finite number of indecomposable modules with finite projective dimension. So the subcategory is contravariantly finite. \square

4 Repdimension

Many results based on the survey [Opp09].

Definition 4.1 (dominated dimension). Let

$$\Lambda \longrightarrow I^0 \longrightarrow I^1 \longrightarrow \dots$$

be the minimal injective resolution of Λ . Then the *dominated dimension* of Λ is $\inf\{n \mid I^n \text{ is not projective}\}$.

Definition 4.2 (rep-dimesnion). Let A be defined by

$$A = \{\Gamma | \text{domdim}\Gamma \geq 2, \Lambda \text{ morita equivalent to } \text{End}_\Gamma(I_0(\Gamma))\}$$

where $I_0(\Gamma)$ is the injective envelope of Γ . Then the *representation dimension* of Λ is the minimal global dimension of $\Gamma \in A$.

Proposition 4.3. (all modules ar right modules) Repdim is the same as minimal global dimension of $\text{End}(M)$ for M being both a generator and cogenerator.

Proof. Consider $\Gamma \in A$. Since $\text{domdim}\Gamma \geq 1$, $I_0(\Gamma)$ is the sum of all projective-injective modules (some probably several times).

Let \mathcal{S} be the set of all Γ -modules with a copresentation

$$0 \longrightarrow X \longrightarrow I_0 \longrightarrow I_1$$

with I_i in $\text{add } I_0(\Gamma)$. In particular Γ is in \mathcal{S} , because $\text{domdim}\Gamma \geq 2$.

The Yoneda embedding gives an equivalence

$$\text{Hom}_\Gamma(-, I_0(\Gamma)): \text{add } I_0(\Gamma) \rightarrow \text{proj } \text{End}_\Gamma(I_0(\Gamma))^{\text{op}},$$

and thus we get an equivalence

$$D \text{Hom}_\Gamma(-, I_0(\Gamma)): \text{add } I_0(\Gamma) \rightarrow \text{inj } \text{End}_\Gamma(I_0(\Gamma))$$

Since $I_0(\Gamma)$ is injective $D \text{Hom}(-, I_0(\Gamma))$ is exact and preserves kernels, so extends to an equivalence

$$\text{Hom}_\Gamma(-, I_0(\Gamma)): \mathcal{S} \rightarrow \text{mod } \text{End}_\Gamma(I_0(\Gamma))$$

Since $\text{End}_\Gamma(I_0(\Gamma))$ is morita equivalent to Λ , \mathcal{S} is equivalent to $\text{mod } \Lambda$. $\Gamma \in \mathcal{S}$ is clearly a generator. To see that it is a cogenerator note that Γ contains all the projective-injective indecomposable objects as direct summands, so there is an injection $I_0(\Gamma) \rightarrow \Gamma^n$, and since $I_0(\Gamma)$ is a cogenerator in \mathcal{S} , Γ is aswell.

Thus by the equivalence $\mathcal{S} \rightarrow \text{mod } \Lambda$ there is a cogenerator-generator object M such that $\text{End}_\Lambda(M) = \text{End}_\Gamma(\Gamma) = \Gamma$.

The last step of the proof is showing that $\text{End}(M)$ is in A whenever M is a generator-cogenerator.

can probably reformulate this in terms of
I guess this is Auslander's original definition
importance to the distinction?

Let $0 \rightarrow M \rightarrow I_0(M) \rightarrow I_1(M)$ be a minimal injective copresentation of M . Since M is a cogenerator $I_i(M)$ is in $\text{add } M$, thus we get an exact sequence of projective $\text{End}(M)$ -modules

$$0 \rightarrow \text{End}(M) \rightarrow \text{Hom}(M, I_0(M)) \rightarrow \text{Hom}(M, I_1(M)). \quad (1)$$

Now we have the following isomorphisms of $\Lambda\text{-End}(M)$ -bimodules

$$\begin{aligned} \text{Hom}_\Lambda(M, D\Lambda) &= \text{Hom}_k(M \otimes \Lambda, k) \\ &= \text{Hom}_k(M, k) \\ &= DM \\ &= D \text{Hom}_\Lambda(\Lambda, M) \end{aligned}$$

Since Λ is in $\text{add } M$, $\text{Hom}(\Lambda, M)$ is projective, and thus $D \text{Hom}(\Lambda, M) = \text{Hom}(M, D\Lambda)$ is injective. This means that (1) is an injective copresentation, and thus $\text{domdim } \text{End}(M) \geq 2$.

Since $\text{Hom}(M, I_0(M))$ is the beginning of an injective resolution of $\text{End}(M)$, $I_0(\text{End}(M))$, must be a direct summand. Then $\text{Hom}(M, I_0(M))/I_0(\text{End}(M))$ would map injectively into $\text{Hom}(M, I_1(M))$, but that would mean theres a direct summand of $I_0(M)$ mapping injectively into $I_1(M)$, contradicting minimality. Thus $\text{Hom}(M, I_0(M)) = I_0(\text{End}(M))$.

Let $I = I_0(M)$ and $\Gamma = \text{End}_\Lambda(I)$, then $D \text{Hom}(-, I)$ is an exact equivalence from $\text{add } I$ to $\text{inj } \Gamma$. Since I is an injective cogenerator $\text{add } I = \text{inj } \Lambda$. Then because of exactness $D \text{Hom}(-, I)$ becomes an equivalence between $K^{+,b}(\text{inj } \Lambda)$ and $K^{+,b}(\text{inj } \Gamma)$. Considering only those complexes with homology in degree 0, we see that $\text{mod } \Lambda$ is equivalent to $\text{mod } \Gamma$. So Λ is morita equivalent to $\Gamma = \text{End}(I_0(M)) = \text{End}(I_0(\text{End}(M)))$. \square

Definition 4.4. Let X be an object of $\text{mod } \Lambda$ and M a contravariantly finite subcategory.

$$\begin{array}{ccccccc} \cdots & \longrightarrow & M_2 & \longrightarrow & M_1 & \longrightarrow & M_0 \\ & \searrow & \uparrow & \searrow & \uparrow & \searrow & \uparrow \\ & & \Omega_M^3 X & & \Omega_M^2 X & & \Omega_M X & \searrow & X \end{array}$$

If \rightarrow are minimal M -approximations (they need not be surjective), and \hookrightarrow are their kernels, then this is a minimal M -resolution of X . The M -res-dimension of X is the length of this sequence of (nonzero) M_i 's, and the M -res-dimension of Λ is the supremum of the dimension on its objects.

Proposition 4.5. *Repdim-2 is the minimum of M -res-dim(mod Λ) for M both generator and cogenerator (assuming repdim is at least 2).*

Proof. The functor $\text{Hom}(M, -)$ is an equivalence from $\text{add } M$ to $\text{proj End}(M)$, which maps minimal M -approximations to projective covers. Let X be any module in $\text{mod End}(M)$ with projective dimension at least 2. Then it has a projective presentation

$$\Omega^2 X \rightarrow (M, M_1) \rightarrow (M, M_0) \rightarrow X.$$

Because of the equivalence this is induced by a map $f: M_1 \rightarrow M_0$. Since Hom is left exact we have that $\Omega^2 X \cong \text{Hom}(M, \ker f)$, and so the projective dimension of X is 2 plus the M -res-dimension of $\ker f$.

Since M is a cogenerator any module Y in $\text{mod } \Lambda$ has a copresentation

$$0 \longrightarrow Y \longrightarrow M_0 \xrightarrow{f} M_1.$$

Applying $\text{Hom}(M, -) =: (M, -)$ we get

$$0 \longrightarrow (M, Y) \longrightarrow (M, M_0) \xrightarrow{(M, f)} (M, M_1) \longrightarrow \text{Cok}(M, f) \longrightarrow 0.$$

If the projective dimension of $\text{Cok}(M, f)$ is less than 2, then (M, Y) is a direct summand of (M, M_0) . This means that $(M, Y) \cong (M, M')$, so the minimal M -approximation of Y is M' , and $(M, \Omega_M Y) = 0$. Since M is a generator this means $\Omega_M Y = 0$ and thus the M -res-dimension of Y is 0.

So provided the projective dimension of $\text{Cok}(M, f)$ is larger than or equal to 2, it equals the M -res-dimension of Y plus 2. In particular the global dimension of $\text{End}(M)$ is 2 plus the M -res-dimension of $\text{mod } \Lambda$, provided it is at least 2. \square

Proposition 4.6. *The repdimension of an artin algebra is always finite. [Iya02]*

Theorem 4.7. *The repdimension of Λ is less than or equal to 2 if and only if Λ is representation finite.*

Proof. Assume Λ is representation finite and let M be the direct sum of all indecomposable modules (up to iso). Then M is a generator-cogenerator. Let X be an $\text{End}(M)^{\text{op}}$ -module with projective presentation

$$(M, M_1) \rightarrow (M, M_0) \rightarrow X \rightarrow 0.$$

Let M_2 be the kernel of $M_1 \rightarrow M_0$. Since M is the sum of all indecomposables M_2 is in $\text{add } M$, so

$$0 \rightarrow (M, M_2) \rightarrow (M, M_1) \rightarrow (M, M_0) \rightarrow X \rightarrow 0$$

is a projective resolution of X . So Λ has repdimension at most 2.

Assume Λ has repdimension at most 2, and let M be an auslander generator. We want to show that $\text{add } M = \text{mod } \Lambda$. Let X be any Λ -module, and let

$$0 \rightarrow X \rightarrow I_0 \rightarrow I_1$$

be a minimal injective presentation. If $I_0 \rightarrow I_1$ is split then X is injective and thus in $\text{add } M$. Let M_X be a minimal M -approximation of X , let $\Omega_M X$ be the kernel of the approximation, and let Y be the cokernel of $(M, I_0) \rightarrow (M, I_1)$. Then

$$(M, \Omega_M X) \rightarrow (M, M_X) \rightarrow (M, I_0) \rightarrow (M, I_1) \rightarrow Y \rightarrow 0$$

is a minimal exact sequence. Since the global dimension of $\text{End}(M)^{\text{op}}$ is at most 2 this means that $(M, \Omega_M X) = 0$. Consequently we have that $\Omega_M X = 0$ and that $X = M_X$, so X is in $\text{add } M$. Thus Λ is representation finite. \square

4.1 The Igusa-Todorov function

Let K be the free abelian group generated by isomorphism classes of modules, modulo the relations $[A \oplus B] = [A] + [B]$ and $[P] = 0$ when P is projective. Define the linear map $L: K \rightarrow K$ by $L[A] = [\Omega A]$. For any module X , $[\text{add } X]$ is a finitely generated subgroup of K . Fitting's lemma tells us that there is an integer η_X such that $L: L^m[\text{add } X] \rightarrow L^{m+1}[\text{add } X]$ is an isomorphism for every $m \geq \eta_X$. We define $\psi(X)$ to be $\eta_X + \sup\{\text{pd } Y \mid Y \in \text{add } \Omega^{\eta_X} X, \text{pd } Y < \infty\}$.

Lemma 4.8. *[IT05, Lemma 3]*

1. $\psi(M) = \text{pd } M$ when $\text{pd } M < \infty$.
2. $\psi(M^k) = \psi(M)$
3. $\psi(M) \leq \psi(M \oplus N)$
4. If Z is a direct summand of $\Omega^n(M)$ where $n \leq \eta_M$ and $\text{pd } Z < \infty$, then $\text{pd } Z + n \leq \psi(M)$.

Proof.

1. If $\text{pd } M < \infty$ then $L^m \neq 0$ for $m < \text{pd } M$, and $L^m = 0$ for $m \geq \text{pd } M$.
2. $\text{add } M^k = \text{add } M$, and ψ is only defined in terms of additive categories.
3. $\text{add } M \subseteq \text{add } M \oplus N$, so if L is injective when restricted to $L^m(\text{add } M \oplus N)$ then L is injective when restricted to $L^m(\text{add } M)$, so $\eta_M \leq \eta_{M \oplus N}$. Further $\Omega^{\eta_{M \oplus N} - \eta_M} \text{add } \Omega^{\eta_M} M \subset \text{add } \Omega^{\eta_{M \oplus N}} M \oplus N$, so $\psi(M) \leq \psi(M \oplus N)$.
4. Let $p = \text{pd } Z$ and $k = \eta_M - n$. Then $\Omega^k Z$ is in $\text{add } \Omega^{\eta_M} M$, so $\text{pd } \Omega^k Z + \eta_M \leq \psi(M)$. Thus

$$\text{pd } Z + n = p + n = (p - k) + \eta_M \leq \text{pd } \Omega^k Z + \eta_M \leq \psi(M).$$

□

Theorem 4.9. [IT05, Theorem 4] *Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be a short exact sequence of modules with $\text{pd } C < \infty$. Then $\text{pd } C \leq \psi(A \oplus B) + 1$.*

Proof. Let P_A^\bullet and P_C^\bullet be the minimal projective resolutions of A and C . Then we get a map of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & P_A^0 & \longrightarrow & P_A^0 \oplus P_C^0 & \longrightarrow & P_C^0 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \end{array}$$

Applying the snake lemma we get $0 \rightarrow \Omega A \rightarrow \Omega B \oplus P \rightarrow \Omega C \rightarrow 0$ for some projective P . Thus for some $n \leq \text{pd } C$ we have $L^n[A] = L^n[B]$, and let n be the minimal such number. Clearly $n \leq \eta_{A \oplus B}$. Let $X = \Omega^n A = \Omega^n B$, then our sequence of n -syzygies looks like

$$0 \longrightarrow X \longrightarrow X \oplus P \longrightarrow \Omega^n C \longrightarrow 0.$$

Let f be the composition $X \longrightarrow X \oplus P \xrightarrow{\pi_X} X$. Then by fittings lemma X breaks as a direct sum into two components $X = Z \oplus Y$ such that $f = f_Z \oplus f_Y$ with f_Y an isomorphism and f_Z nilpotent. In other words the sequence above can be written as

$$0 \longrightarrow Z \oplus Y \longrightarrow Z \oplus Y \oplus P \longrightarrow \Omega^n C \longrightarrow 0.$$

with the left map being

$$\begin{bmatrix} f_Z & 0 \\ 0 & f_Y \\ * & * \end{bmatrix} \sim \begin{bmatrix} f_Z & 0 \\ 0 & f_Y \\ * & 0 \end{bmatrix}$$

So we get another short exact sequence

$$0 \longrightarrow Z \longrightarrow Z \oplus P \longrightarrow \Omega^n C \longrightarrow 0.$$

Let $T = \Lambda/J$ and apply the long exact sequence in $\text{Ext}(-, T)$. Then we get an exact sequence

$$\text{Ext}^k(Z, T) \longrightarrow \text{Ext}^k(Z \oplus P, T) \longrightarrow \text{Ext}^{k+1}(\Omega^n C, T)$$

where the left map is induced by f_Z since $\text{Ext}^k(Z \oplus P, T) \cong \text{Ext}^k(Z, T)$. Since f_Z is nilpotent this map is surjective if and only if $\text{Ext}^k(Z, T) = 0$, and $\Omega^n C$ has finite projective dimension we have that Z has finite projective dimension. In particular $\text{pd } \Omega^n C - 1 \leq \text{pd } Z \leq \text{pd } \Omega^n C$.

Since Z is a direct summand of $\Omega^n A \oplus B$ by Lemma 4.8 we have that $\text{pd } Z + n \leq \psi(A \oplus B)$, and thus $\text{pd } \Omega^n C - 1 + n = \text{pd } C - 1 \leq \psi(A \oplus B)$. \square

Corollary 4.9.1. *Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be a short exact sequence of modules. If $\text{pd } A < \infty$ then $\text{pd } A \leq \psi(\Omega B \oplus \Omega C) + 1$, and if $\text{pd } B < \infty$ then $\text{pd } B \leq \psi(\Omega A \oplus \Omega^2 C) + 2$.*

Proof. Let $P_B \rightarrow B$ be a projective cover of B . Then we have a commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & 0 & \longrightarrow & P_B & \longrightarrow & P_B \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \end{array}$$

Applying the snake lemma we get a short exact sequence $0 \rightarrow \Omega B \rightarrow \Omega C \oplus P \rightarrow A \rightarrow 0$ for some projective P . Then using the theorem we have that $\text{pd } A \leq \psi(\Omega B \oplus \Omega C \oplus P) + 1 = \psi(\Omega B \oplus \Omega C) + 1$.

Applying the same reasoning to $0 \rightarrow \Omega B \rightarrow \Omega C \oplus P \rightarrow A \rightarrow 0$ gives us $\text{pd } \Omega B \leq \psi(\Omega A \oplus \Omega^2 C) + 1$. Hence $\text{pd } B \leq \psi(\Omega A \oplus \Omega^2 C) + 2$. \square

Theorem 4.10. [IT05, Corollary 8] *If $\Lambda = \text{End}_\Gamma(P)$ for an algebra Γ with global dimension at most 3, and P projective then $\text{findim}(\Lambda) < \infty$.*

Proof. Let X be any Λ -module with finite projective dimension. Then it has a projective presentation $(P, P_1) \rightarrow (P, P_0) \rightarrow X \rightarrow 0$ where $(P, P_i) = \text{Hom}_\Gamma(P, P_i)$ with $P_i \in \text{add } P$. Since $(P, -)$ is an equivalence from $\text{add } P$ to $\text{proj } \Lambda$ this corresponds to a map $P_1 \rightarrow P_0$ which we can extend to a projective resolution in Γ :

$$0 \longrightarrow P_3 \longrightarrow P_2 \longrightarrow P_1 \longrightarrow P_0.$$

Applying the exact functor $(P, -)$, we get an exact sequence

$$0 \longrightarrow (P, P_3) \longrightarrow (P, P_2) \longrightarrow (P, P_1) \longrightarrow (P, P_0) \longrightarrow X \longrightarrow 0.$$

Truncating this we get a short exact sequence

$$0 \longrightarrow (P, P_3) \longrightarrow (P, P_2) \longrightarrow \Omega^2 X \longrightarrow 0.$$

Then by Theorem 4.9 the projective dimension of $\Omega^2 X$ is bounded by $\psi((P, P_3) \oplus (P, P_2)) + 1$. Which means

$$\text{pd } X \leq \psi((P, P_3) \oplus (P, P_2)) + 3 \leq \psi((P, \Gamma)) + 3$$

Since this bound doesn't depend on X , Λ has finite finitistic dimension. \square

Corollary 4.10.1. *If $\text{repdim}(\Lambda) \leq 3$ then $\text{findim}(\Lambda) < \infty$.*

Proof. If Λ has rep-dimension less than or equal to 3 then by Proposition 4.3 there is a generator-cogenerator M in $\text{mod } \Lambda$ such that $\Gamma := \text{End}_\Lambda(M)$ has global dimension 3 or less. Then since M is a generator Λ is in $\text{add } M$ and so $\text{Hom}_\Lambda(M, \Lambda)$ is a projective Γ -module with $\text{End}_\Gamma(\text{Hom}_\Lambda(M, \Lambda)) = \text{End}_\Lambda(\Lambda) = \Lambda$. \square

4.2 Stably hereditary algebras

In this section we will show that the class of stably hereditary algebras has repdimension at most 3, and thus that they have finite finitistic dimension.

Definition 4.11 ((co)torsionfree). A module is called *torsionfree* if it is a submodule of a projective module. Dually, a module is called *cotorsionfree* if it is a factormodule of an injective.

Definition 4.12 (Stably hereditary algebra). An algebra is called *stably hereditary* if any indecomposable torsionfree module is projective or simple, and any indecomposable cotorsionfree module is injective or simple.

This generalizes the definition of hereditary algebra by also allowing simple modules to be (co)torsionfree.

Definition 4.13 (The stable category). For an algebra Λ , the *stable category* $\underline{\text{mod}}\Lambda$ has the same objects as $\text{mod}\Lambda$, but the sets of homomorphisms are given by

$$\text{Hom}_{\underline{\text{mod}}\Lambda}(M, N) = \text{Hom}_{\Lambda}(M, N) / \mathcal{P}(M, N)$$

where $\mathcal{P}(M, N)$ is the ideal of all morphisms factoring through a projective.

Proposition 4.14. *If for an algebra Λ there is a hereditary algebra H such that $\underline{\text{mod}}\Lambda \cong \underline{\text{mod}}H$ then Λ is stably hereditary.*

Proof. [AR91, Lemma 4.12] [AR73]

□

+ a bit
more...

The converse of the above proposition does not hold without more assumptions, but stably hereditary algebras generalize the idea of algebras stably equivalent to hereditary algebras.

Theorem 4.15. [Xi02, Theorem 3.5] *Stably hereditary algebras has repdimension at most 3.*

Proof. Let V be the direct sum of all the indecomposable projectives, all the indecomposable injectives, and all the simple modules. Then V is a generator-cogenerator. So by Proposition 4.3 if we can show that the global dimension of $\Gamma := \text{End}(V)^{op}$ is 3 or less, then we are done.

We will show that for any Λ -module M there is a short exact sequence $0 \rightarrow V_3 \rightarrow V_3 \rightarrow M \rightarrow 0$ with V_i in $\text{add } V$, and such that $0 \rightarrow (V, V_3) \rightarrow (V, V_2) \rightarrow (V, M) \rightarrow 0$ is exact. We will use this to construct short projective resolutions for $\text{mod } \Gamma$. To construct V_3 and V_2 let M' be the sum of the maximal injective summand of M and all simple submodules of M . Then let P be the projective cover of M/M' . Taking the pullback of $M \rightarrow M/M' \leftarrow P$ gives us the diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & K & \xlongequal{\quad} & K & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & M' & \longrightarrow & M' \oplus P & \longrightarrow & P \longrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M/M' \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

I claim that $0 \rightarrow K \rightarrow M' \oplus P \rightarrow M \rightarrow 0$ is the desired sequence. Firstly $M' \oplus P$ is clearly in $\text{add } V$ since it is the sum of an injective, a semisimple, and a projective module. Further K is a submodule of P , hence torsionfree. So since Λ is stably hereditary K is the sum of a projective and a semisimple module, so K is also in $\text{add } V$.

Next we need to show that $0 \rightarrow (V, K) \rightarrow (V, M' \oplus P) \rightarrow (V, M) \rightarrow 0$ is exact. The only thing needed to show here is that $(V, M' \oplus P) \rightarrow (V, M)$ is surjective. We do this by showing that $(W, M' \oplus P) \rightarrow (W, M)$ is surjective for any indecomposable summand of V . If W is projective this holds by definition. If W is simple then any map from W to M factors through the socle and hence through M' , so it's surjective. Lastly if W is injective then the image of W in M is a cotorsionfree module, so it is the sum of simple modules and an injective module. Hence the map from W to M factors through M' .

Now we use this to show that the global dimension of Γ is at most 3. Let N be any Γ -module. Then it has a projective presentation

$$(V, V_1) \xrightarrow{f \circ -} (V, V_0) \longrightarrow N \longrightarrow 0$$

If we let M denote the kernel of f and we choose V_3 and V_2 as above then we get a projective resolution of N by

$$0 \longrightarrow (V, V_3) \longrightarrow (V, V_2) \longrightarrow (V, V_1) \longrightarrow (V, V_0) \longrightarrow N \longrightarrow 0.$$

This shows that the projective dimension of N is at most 3, and since N was arbitrary the global dimension of Γ is at most 3. So the repdimension

of Λ is at most 3. \square

4.3 Special biserial algebras

[EHIS04]

5 Vanishing radical powers

Throughout this section Λ is a finite dimensional algebra, and J is its radical.

Theorem 5.1. *If $J^2 = 0$ then $\text{findim}(\Lambda) < \infty$.*

Proof. Let $d = \max\{\text{pd } S_i \mid \text{pd } S_i < \infty\}$ where S_i ranges over the simple Λ -modules. Let M be a module with $\text{pd } M < \infty$. Let $P \rightarrow M$ be a projective cover. Then ΩM is contained in JP and since $J^2P = 0$, ΩM is annihilated by J and is thus semisimple. This means $\text{pd } \Omega M \leq d$, and thus $\text{pd } M \leq d+1$. So $\text{findim}(\Lambda) \leq d+1 < \infty$. \square

Theorem 5.2. *[IT05, Corollary 6] If $J^3 = 0$ then $\text{findim}(\Lambda) < \infty$.*

Proof. Let M be a module with $\text{pd } M < \infty$, and let $P^0 \rightarrow M$ be its projective cover. Since $\Omega M \subseteq JP^0$ we have $J^2\Omega M = 0$. Let $P \rightarrow \Omega M$ be a projective cover. Since $J^2\Omega M = 0$ we can factorize this as $P \rightarrow P/J^2P \rightarrow \Omega M$, and we get a short exact sequence

$$0 \longrightarrow (\Omega^2 M + J^2 P)/J^2 P \longrightarrow P/J^2 P \longrightarrow \Omega M \longrightarrow 0$$

Let ψ be the Igusa-Todorov function as introduced in Section 4.1. Since $\Omega^2 M \subseteq JP$ we have that $(\Omega^2 M + J^2 P)/J^2 P$ is semisimple. Then by Lemma 4.8 $\psi((\Omega^2 M + J^2 P)/J^2 P) = \psi(\Lambda/J)$, and $\psi(P/J^2 P) = \psi(\Lambda/J^2)$.

Applying Theorem 4.9 to the short exact sequence above we thus get $\text{pd } \Omega M \leq \psi(\Lambda/J \oplus \Lambda/J^2) + 1$, and so $\text{pd } M \leq \psi(\Lambda/J \oplus \Lambda/J^2) + 2$, and $\text{findim}(\Lambda) < \infty$. \square

Theorem 5.3. *[Wan94] If $J^{2l+1} = 0$ and Λ/J^l is representation finite then $\text{findim}(\Lambda) < \infty$.*

Proof. Let M be a module with $\text{pd } M < \infty$. We have a short exact sequence

$$0 \longrightarrow J^l \Omega M \longrightarrow \Omega M \longrightarrow \Omega M / J^l \Omega M \longrightarrow 0.$$

Since $\Omega M \subseteq JP_M^0$ we have $J^{2l} \Omega M = 0$. This means that $J^l \Omega M$ and $\Omega M / J^l \Omega M$ are Λ / J^l -modules. We will use this, the fact that Λ / J^l is representation finite, and the Igusa-Todorov function to create a bound for $\text{pd } M$.

Applying Corollary 4.9.1 we have that:

$$\text{pd } \Omega M \leq \psi(\Omega(J^l \Omega M) \oplus \Omega^2(\Omega M / J^l \Omega M)) + 2.$$

Since Λ / J^l is representation finite there are only finitely many indecomposable Λ / J^l -modules, up to isomorphism. Let S be the sum of all of them. Then since $J^l \Omega M$ and $\Omega M / J^l \Omega M$ are in $\text{add } S$, using Lemma 4.8 we have that

$$\psi(\Omega(J^l \Omega M) \oplus \Omega^2(\Omega M / J^l \Omega M)) \leq \psi(\Omega S \oplus \Omega^2 S).$$

So $\text{pd } M \leq \psi(\Omega S \oplus \Omega^2 S) + 3$, and thus $\text{findim}(\Lambda) < \infty$. \square

6 Monomial algebras

[GKK91, IZ90]

In this section we will show a particularly nice way to construct a minimal projective resolution of the right module Λ / J for a monomial algebra Λ . We will use this to compute $\text{Tor}_i(\Lambda / J, M)$ and/or $\text{Ext}^i(M, D\Lambda / J)$ to get a bound on the projective dimension of all modules M .

Definition 6.1 (Monomial algebra). A *monomial algebra* is a path algebra with admissible relations that are generated by monomials. That is, we do not allow the generators for the relations to consist of nontrivial linear combinations of paths.

Definition 6.2 (m -chains). [GKK91] Let $\Lambda = k\Gamma / (\rho)$ be a monomial algebra, with ρ a minimal generating set of paths. As usual we define Γ_0 to be the vertices of Γ , and Γ_1 to be the arrows. Recursively define the set of $(m - 1)$ -chains, Γ_m , as the paths γ with the following criteria:

- i) $\gamma = \beta\delta\tau$ with $\beta \in \Gamma_{m-2}$, $\beta\delta \in \Gamma_{m-1}$, and τ a non-zero path of length at least 1.
- ii) $\delta\tau$ is 0 in Λ , i.e. it is in the ideal of relations.

We may assume ρ contains J^2

- iii) γ is left-minimal in the sense that if $\gamma = \gamma'\sigma$ such that γ' satisfies the above conditions, then $\gamma = \gamma'$.

The sets of m -chains will become the generating sets for the projectives in our projective resolution. But first we will prove some properties of them.

Lemma 6.3. *Any $\gamma \in \Gamma_m$ for $m \geq 1$ can be factored uniquely as $\gamma_1\gamma_0$ with $\gamma_1 \in \Gamma_{m-1}$, and γ_0 a non-zero path of length at least 1.*

Proof. When $m = 1$ this should be clear, since Γ_1 is the set of arrows, and Γ_0 is the set of vertices, so if $\gamma \in \Gamma_1$ is an arrow $i \rightarrow j$ then $\gamma = e_j\gamma$.

When $m > 1$ we know from the definition of Γ_m that γ can be written as $\gamma_1\gamma_0$. Assume there is another decomposition $\gamma = \gamma'_1\gamma'_0$. Then without loss of generality we may assume that γ'_1 is shorter than γ_1 . Then there is a σ such that $\gamma'_1\sigma = \gamma_1$. By minimality this means that $\gamma'_1 = \gamma_1$, and so the decomposition is unique. \square

From now on we will write R for the ring Λ/J , which we identify with the subring of Λ generated by the paths of length 0. Let $k\Gamma_m$ be the free vector space generated by Γ_m . Notice that $k\Gamma_m$ has a canonical structure as a $R - R$ -bimodule. This means we can get projective right Λ -modules $P^m := k\Gamma_m \otimes_R \Lambda$.

Proposition 6.4. *Define the map $\delta_m: P^m \rightarrow P^{m-1}$ by $\delta_m(\gamma \otimes \alpha) = \gamma_1 \otimes \gamma_0 \alpha$ where $\gamma_1\gamma_0$ is the unique decomposition of γ , and define $\delta_0: P^0 \rightarrow \Lambda/J$ by $\delta_0(e_i \otimes \alpha) = e_i \alpha + J$. Then we have a minimal projective resolution of the right Λ -module Λ/J by*

$$\begin{array}{ccccccc} \dots & \longrightarrow & P^3 & \xrightarrow{\delta_3} & P^2 & \xrightarrow{\delta_2} & P^1 & \xrightarrow{\delta_1} & P^0 & \longrightarrow & 0 \\ & & & & & & & & \downarrow \delta_0 & & \\ & & & & & & & & \Lambda/J & & \end{array}$$

Before proving this proposition we require a lemma.

Lemma 6.5. *[GKK91, Lemma 2.1] Let M be a Λ -module, and x an element in the kernel of $\delta_m \otimes M: P^m \otimes_R M \rightarrow P^{m-1} \otimes_R M$. Write x on the form*

$$x = \sum_j \sum_{k=0}^{n_j} \gamma_j \gamma_j^k \otimes m_j^k$$

with $\gamma_i \in \Gamma_{m-1}$ and $\gamma_i \neq \gamma_j$ and $\gamma_j^k \neq \gamma_j^l$ when $i \neq j$ and $k \neq l$. Then

$$\sum_{k=0}^{n_j} \gamma_j \gamma_j^k \otimes m_j^k$$

is also in the kernel for each j .

Proof. Let x be as given above. Applying $\delta_m \otimes M$ we get that

$$\sum_j \gamma_j \otimes \sum_{k=0}^{n_j} \gamma_j^k m_j^k = 0.$$

Since the γ_j 's are distinct we can deduce that

$$\sum_{k=0}^{n_j} \gamma_j^k m_j^k = 0.$$

Since Λ only has monomial relations, and by the minimality of the γ_j^k 's none of them divide each other, we have that $\gamma_j^k \alpha_j^k = 0$. Thus

$$\sum_{k=0}^{n_j} \gamma_j \gamma_j^k \otimes m_j^k$$

is also in the kernel of $\delta_m \otimes M$. □

Proof of Proposition 6.4. For all i the module P^i is projective as a right Λ -module and the image of δ_m is clearly contained in $P^{m-1}J$, so the only thing left to show is exactness. First we show that $\delta_m \delta_{m-1} = 0$. Let $\gamma \otimes \alpha$ be in P^m for $m \geq 2$. Then we can decompose γ uniquely as $\gamma_2 \gamma_1 \gamma_0$ and $\delta_m \delta_{m-1}(\gamma \otimes \alpha) = \gamma_2 \otimes \gamma_1 \gamma_0 \alpha$. By the way we defined Γ_m , $\gamma_1 \gamma_0$ is 0 in Λ , and so $\gamma_2 \otimes \gamma_1 \gamma_0 \alpha = 0$.

Next we want to show that $\text{Ker } \delta_{m-1} \subseteq \text{Im } \delta_m$. Let x be in the kernel of δ_{m-1} . By Lemma 6.5 it is sufficient to assume x is of the form

$$\sum_k \gamma \gamma_k \otimes \alpha_k.$$

Then $\sum_k \gamma \gamma_k \alpha_k = 0$. Because of this we have that $\gamma \gamma_k \alpha_k = \zeta_k \sigma_k$ for some m -chain ζ_k and some path σ_k (possibly of length 0). This gives us that x is the image of

$$\sum_k \zeta_k \otimes \sigma_k$$

by δ_m . Hence $\text{Ker } \delta_{m-1} \subseteq \text{Im } \delta_m$, and the sequence is exact. So this gives a minimal projective resolution of Λ/J as a right Λ -module. □

Definition 6.6. We call a path τ in Γ a *special segment* for $\Lambda = k\Gamma/(\rho)$ if there is a path γ such that $\gamma\tau$ is a minimal relation.

Note that when we decompose an m -chain γ in Lemma 6.3 into $\gamma_1\gamma_0$ then γ_0 is a special segment, and that the set of special segments is finite.

Lemma 6.7. [GKK91, Theorem 2.2] *Let d be the number of special segments for Λ . If $s \geq d + 3$ and γ is in Γ_s , then for any integer N there is an $n \geq N$ and a $\hat{\gamma} \in \Gamma_n$ such that for any path τ we have $\gamma\tau \in \Gamma_{s+r}$ if and only if $\hat{\gamma}\tau \in \Gamma_{n+r}$.*

Proof. Applying Lemma 6.3 recursively we get that γ can be written as $\gamma = \tau_0\tau_1 \cdots \tau_{s-1}$ where $\tau_0\tau_1 \cdots \tau_{i-1} \in \Gamma_i$. In particular each τ_i is a special segment.

Since $s \geq d + 3$ we must have that there exists i and j , $1 \leq i < j \leq s - 1$ such that $\tau_i = \tau_j$. Let $\beta = \tau_{i+1}\tau_{i+2} \cdots \tau_j$. Then

$$\gamma_k := \tau_0\tau_1 \cdots \tau_{j-1}\tau_j\beta^k\tau_{j+1} \cdots \tau_{s-1} \in \Gamma_{s+k(j-i)}$$

where β^k means β repeated k times. If we now choose k large enough such that $s + k(j - i) \geq N$ we can choose $n = s + k(j - i)$ and $\hat{\gamma} = \gamma_k$. Then we see that for any path τ , the composition $\gamma\tau$ is in Γ_{s+r} if and only if $\hat{\gamma}\tau$ is in Γ_{n+r} . \square

Theorem 6.8. [GKK91, Corollary 2.4] *Let $\Lambda = k\Gamma/(\rho)$ be a monomial relation algebra. Then $\text{findim}(\Lambda) \leq d + 3$ where d is the number of special segments for Λ .*

Proof. Let M be a module of finite projective dimension and let N be $\text{pd } M$. The projective dimension of M can be characterized as the largest integer c such that $\text{Tor}_c(\Lambda/J, M) \neq 0$. We will show that this is at most $d + 3$. Let $s \geq d + 3$ be an integer. Then we want to show that $\text{Tor}_{s+1}(\Lambda/J, M) = 0$. We compute this by taking the projective resolution of Λ/J found in Proposition 6.4 and tensoring with M .

$$\cdots \longrightarrow k\Gamma_{s+2} \otimes M \xrightarrow{\delta_{s+2} \otimes M} k\Gamma_{s+1} \otimes M \xrightarrow{\delta_{s+1} \otimes M} k\Gamma_s \otimes M \longrightarrow \cdots$$

Let x be in the kernel of $\delta_{s+1} \otimes M$. Then by Lemma 6.5 we may assume x is on the form

$$x = \sum_j \gamma_j \otimes m_j.$$

Now since γ is in Γ_s Lemma 6.7 gives us that there is an $n \geq N$ and a $\hat{\gamma} \in \Gamma_n$ such that $\gamma\tau$ is in Γ_{s+r} if and only if $\hat{\gamma}\tau$ is in Γ_{n+r} .

Then $\hat{x} = \sum \hat{\gamma}\gamma_j \otimes m_j$ is in the kernel of $\delta_{n+1} \otimes M$. Since $n+1 > N = \text{pd } M$ the complex is exact at $n+1$. This means that there are elements γ_j^k and m_j^k such that

$$\hat{x} = \delta_{n+2} \left(\sum_j \sum_{k=0}^{n_j} \hat{\gamma}\gamma_j \gamma_j^k \otimes m_j^k \right) = \sum_j \sum_{k=0}^{n_j} \hat{\gamma}\gamma_j \otimes \gamma_j^k m_j^k$$

Since $\hat{\gamma}\gamma_j \gamma_j^k$ is in Γ_{n+2} if and only if $\gamma_j \gamma_j^k$ is in Γ_{s+2} we have that

$$x = \delta_{s+2} \left(\sum_j \sum_{k=0}^{n_j} \gamma_j \gamma_j^k \otimes m_j^k \right)$$

and thus $\text{Tor}_{s+1}(\Lambda/J, M) = 0$ so $\text{pd } M \leq d+3$. \square

7 Unbounded derived category

If we go to the unbounded derived category we can get a sort of converse to Theorem 1.4.

Theorem 7.1. *[Ric19, Theorem 4.3] If the localizing category of $D\Lambda$ is the entire unbounded derived category then $\text{Findim}(\Lambda) < \infty$. (Note the capital F meaning the finitistic dimension of $\text{Mod } \Lambda$, which is bigger than or equal to that of $\text{mod } \Lambda$).*

Proof. Assume $\text{Findim}(\Lambda) = \infty$. Then there are modules M_i with projective dimension i for every $i \geq 0$. Let P_i be the minimal projective resolution of M_i , and consider $\bigoplus P_i[-i]$ and $\prod P_i[-i]$. Both of these have homology M_i in degree i , and are concentrated in non-negative degrees.

The inclusion from the sum to the product is clearly a quasi-isomorphism. We want to show that it is not a homotopy equivalence. Assume for the sake of contradiction that it was. Then tensoring with Λ/J would give us another homotopy equivalence. Since Λ/J is finitely presented tensoring preserves both products and coproducts. Because all the resolutions were minimal tensoring with Λ/J gives us 0 differentials. In degree 0 we get

$$\bigoplus \text{Tor}_i(\Lambda/J, M_i) \rightarrow \prod \text{Tor}_i(\Lambda/J, M_i).$$

Since $\text{Tor}_i(\Lambda/J, M_i)$ is nonzero for every M_i this map is not an isomorphism, and so we don't have a homotopy equivalence.

So the cone of the inclusion $\bigoplus P_i[-i] \rightarrow \prod P_i[-i]$, C , is 0 in the derived category, but non-zero in the homotopy category. Since Λ is artinian the product of projectives is projective [Cha60, Theorem 3.3], so $\prod P_i[-i]$ is a complex of projectives, which means that C is a complex of projectives.

In other words C is an acyclic lower bounded complex of projectives that is not contractible. Tensoring with $D\Lambda$ is an equivalence from projectives to injectives with inverse $\mathcal{D}(\Lambda)(D\Lambda, -)$, so $D\Lambda \otimes C$ is an lower bounded complex of injectives that is not contractible. Such a complex cannot be acyclic so $D\Lambda \otimes C$ has homology.

maybe I should prove this or find reference

The homology of C is 0, so $K(\Lambda)(\Lambda, C[i]) = 0$. Applying the equivalence $D\Lambda \otimes -$ we get

$$\mathcal{D}(\Lambda)(D\Lambda, D\Lambda \otimes C[i]) = K(\Lambda)(D\Lambda, D\Lambda \otimes C[i]) = 0.$$

This means that $D\Lambda \otimes C$ is not in the localizing category generated by $D\Lambda$, and so that can not be the entire derived category. \square

Theorem 7.2. [Ric19, Theorem 4.4] $\text{Findim}(\Lambda) < \infty$ if and only if $D\Lambda^\perp \cap \mathcal{D}^+(\Lambda) = 0$.

Proof. In the theorem above we proved that when the finitistic dimension is infinite then there is a non-zero complex in $\mathcal{D}^+(\Lambda)$ perpendicular to $D\Lambda$.

The proof of the converse is the same as for ???. If we have a non-zero object $X \in D\Lambda^\perp \cap \mathcal{D}^+(\Lambda)$, then $\mathcal{D}(\Lambda)(D\Lambda, X)$ is an acyclic minimal complex of projectives that continue arbitrarily to the right. So the cokernels have arbitrarily big projective dimension. \square

We see this by taking injective resolution of X

8 Summary

FDC holds for the following classes of algebras

- **Big FDC:**
- Representation finite algebras

Proof. The supremum over a finite set is finite so $\text{findim}(\Lambda) < \infty$ for a representation finite algebra. \square

- Monomial algebras

Proof. This was shown in Section 6. \square

- Gorenstein algebras

Proof. An algebra is said to be Gorenstein if all injectives have finite projective dimension and all projectives have finite injective dimension. In particular the Λ -module Λ is isomorphic to a finite injective resolution in the derived category. So Λ is in the localizing category generated by injectives. Then Theorem 7.1 gives us that $\text{Findim}(\Lambda) < \infty$, and therefor also $\text{findim}(\Lambda) < \infty$. \square

- Finite global dimension
- Self injective
- $J^2 = 0$
- Derived equivalent to the above
- Local algebras

Proof. Local algebras are local artinian rings. So if Λ is local then $\text{findim}(\Lambda) = 0$. \square

- **only small FDC is known?:**

- Stably hereditary algebras
- Special biserial algebras
- "half rep-finite" algebras, i.e. Λ/J^l rep-finite $J^{2l+1} = 0$.

Not sure where to put this, ill put it here for now

Theorem 8.1. *Local artinian rings have finitistic dimension zero.*

Proof. Assume there is a non-projective module with finite projective dimension. Then in particular we have one with projective dimension equal to 1. Since all finitely generated projectives are free this means we have a short exact sequence

$$0 \quad R^n \quad R^m \quad M \quad 0$$

with R^n contained in JR^m . Let k be the minimal integer such that $J^k = 0$. Let a be a generator in R^n and let r be a non-zero element of J^{k-1} . Then ra is non-zero, but is mapped to something in $J^{k-1}JR^m = 0$, thus the map is not injective which gives a contradiction. \square

9 Dual conjectures

Many of the cases are equivalent to their dual statements. Some are not.

- Given a recollement of the bounded derived category you get one for Λ^{op} .
- Just because the subcategory of modules with finite projective dimension is contravariantly finite does not mean the subcategory of modules with finite injective dimension has to be covariantly finite. See Example 3.6.
- repdim of Λ equals the repdim of Λ^{op} .

Look at examples of recollement to see how it translates.

Proof. If M is an auslander generator for Λ then DM is an auslander generator for Λ^{op} . \square

- If $J^{2l+1} = 0$ and Λ/J^l is repfinite then the same is true for Λ^{op} .
- If Λ is monomial then so is Λ^{op} .
- Injective generates implies the weaker property that projective cogenerate for the opposite algebra. This is also sufficient to prove the algebra satisfies FDC. [Ric19, Section 5]

Similarly for the weaker conjectures

- GSC says the injective dimension of Λ is finite if and only if the injective dimension of Λ^{op} is finite. This statement is symmetric with respect to Λ and Λ^{op} . So the dual is equivalent.
- NC: Certainly Λ is self injective if and only if Λ^{op} is.
- For all the others it seems just as difficult as solving the conjecture to connect it to its dual.

Can the dominant dimension of the opposite algebra be different? Arbitrary different?

10 Personal appendix

Theorem 10.1. *The global dimension of an artin algebra is the supremum of k with $\text{Ext}^k(T, T) \neq 0$ (T sum of simples). This is also the supremum of projective dimension and supremum of injective dimension.*

Proof. For a minimal projective resolution $\text{Hom}(-, T)$ makes the differentials 0, and similarly with $\text{Hom}(T, -)$ and injective resolutions. So $\text{Ext}^k(M, T)$ is only 0 exactly when $k > \text{pd } M$, similarly $\text{Ext}^k(T, M)$ is only 0 when k is bigger than the injective dimension. Since any module is built by extensions of simples you can prove by induction, and the long exact sequence in $\text{Ext}(-, T)$ you get that any module has projective dimension less than or equal to that of T . Similarly for injective dimension. \square

$\text{findim}(\Lambda)$ need not equal $\text{findim}(\Lambda^{\text{op}}) = \sup\{\text{injective dimension of } M \mid M \text{ has finite injective dimension}\}$.

Example 10.2. [hf] Let $\Lambda = k \left[\begin{array}{c} a \hookrightarrow 1 \xrightleftharpoons[b]{c} 2 \end{array} \right] / (a^2, ac, ba, cbc)$. Then $\text{findim}(\Lambda) \geq 1$, but $\text{findim}(\Lambda^{\text{op}}) = 0$.

Proof. The module $\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} = P_1/P_2$ (k^2 where a acts by $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, and b and c act trivially) has projective dimension 1, so $\text{findim}(\Lambda) \geq 1$. The projective/injective modules of Λ are:

$$P_1 = \begin{smallmatrix} & 1 \\ 1 & \\ & 2 \end{smallmatrix}, \quad P_2 = \begin{smallmatrix} & 2 \\ 1 & \\ & 2 \end{smallmatrix}, \quad I_1 = \begin{smallmatrix} & 1 \\ & 2 \\ 1 & \end{smallmatrix}, \quad I_2 = \begin{smallmatrix} & 1 \\ & 2 \\ 1 & \\ & 2 \end{smallmatrix}$$

If $\text{findim}(\Lambda^{\text{op}}) > 0$ there would be a module with finite non-zero injective resolution. In particular it would end with a non-split epimorphism between injectives. I claim this would mean there is a non-split epimorphism $I \rightarrow I_i$ from an injective to an indecomposable injective. Obviously we get epimorphisms by composing with the projections onto summands, so we want to show that they are not split. Assume that they are, that is the map looks like

$$\begin{array}{ccc}
 I_i \oplus I & \xrightarrow{\begin{bmatrix} 1 & 0 \\ f & g \end{bmatrix}} & I_i \oplus I' \\
 & \searrow \begin{bmatrix} 1 & 0 \end{bmatrix} & \downarrow \begin{bmatrix} 1 & 0 \end{bmatrix} \\
 & & I_i
 \end{array}$$

We see that by changing basis in the domain we get the matrix $\begin{bmatrix} 1 & 0 \\ 0 & g \end{bmatrix}$. Thus I_i is mapped isomorphically to itself, which doesn't happen in a minimal resolution.

The only thing left to show is that there are no non-split epimorphisms from injective modules to I_1 and I_2 . \square

Lemma 10.3. [CE99, Chapter I, theorem 3.2] *Let R be a noetherian ring. Then an R -module Q is injective if and only if it has the injective lifting property for inclusions of ideals into R .*

Proof. If Q is injective then Q has the lifting property for all monomorphisms, so one direction is clear. Assume we have a diagram

$$\begin{array}{ccc}
 & Q & \\
 f \uparrow & \nwarrow & \\
 M & \hookrightarrow & N
 \end{array}$$

We want to show that the dashed arrow exists. Let S be the partially ordered set $\{(M', f') : M \leq M', f'|_M = f\}$. By Zorn's lemma this has a maximal element (M', f') . Assume $M' \neq N$, then there is an element $x \in N - M'$. The set of r such that $rx \in M'$ forms an ideal I . Define the map $g : I \rightarrow Q$ by $I(r) = f'(rx)$. By hypothesis g lifts to a map $\tilde{g} : R \rightarrow Q$. Let q be $\tilde{g}(1)$. Then $\tilde{f} : M' + Rx \rightarrow Q$ defined by $\tilde{f}(m + rx) = f'(m) + rq$ gives us a bigger element of S , contradicting maximality. Thus $M' = N$ and Q is injective. \square

Theorem 10.4. *Let R be a noetherian ring. Then an arbitrary coproduct of injectives is injective.*

Proof. By the lemma above it is enough to show the lifting property on ideals of R . Let I be an ideal and $f : I \rightarrow \bigoplus_i Q_i$ be a map to a coproduct of injectives. Since R is noetherian I is finitely generated so f factors through

a finite sum $I \rightarrow \bigoplus_{i=0}^n Q_i \rightarrow \bigoplus Q_i$. Since finite coproducts of injectives are injective we are done.

$$\begin{array}{ccc}
 & \bigoplus Q_i & \\
 & \uparrow & \\
 & \bigoplus_{i=0}^n Q_i & \\
 & \uparrow & \nwarrow \text{---} \\
 I & \hookrightarrow & R
 \end{array}$$

□

Theorem 10.5. [CE99, Chapter I, Exercise 8] *Let R be a noetherian ring. Then direct limits of injectives is injective.*

Proof. By the lemma above it is enough to show the lifting property on ideals of R . Let I be an ideal and let $Q = \varinjlim Q_i$ be a direct limit of injectives.

Since R is noetherian I is finitely presented, say $R^n \rightarrow R^m \rightarrow I \rightarrow 0$. Applying $\text{Hom}(-, Q)$ we get an exact sequence

$$0 \longrightarrow \text{Hom}(I, Q) \longrightarrow \text{Hom}(R^m, Q) \longrightarrow \text{Hom}(R^n, Q)$$

Since direct limits are exact we also have an exact sequence

$$0 \longrightarrow \varinjlim \text{Hom}(I, Q_i) \longrightarrow \varinjlim \text{Hom}(R^m, Q_i) \longrightarrow \varinjlim \text{Hom}(R^n, Q_i)$$

We also have a natural map $\varinjlim \text{Hom}(-, Q_i) \rightarrow \text{Hom}(-, Q)$. $\text{Hom}(R^n, Q_i)$ just equals Q_i^n , so this map is an isomorphism at R^n . Then by the five lemma applied to the two sequences above we get that $\text{Hom}(I, Q) \cong \varinjlim \text{Hom}(I, Q_i)$ for all ideals I . So since

$$\varinjlim \text{Hom}(R, Q_i) \longrightarrow \varinjlim \text{Hom}(I, Q_i) \longrightarrow 0$$

is exact, we get that

$$\text{Hom}(R, Q) \longrightarrow \text{Hom}(I, Q) \longrightarrow 0$$

is exact. Hence Q is injective.

□

References

- [AR73] Maurice Auslander and Idun Reiten. Stable equivalence of Artin algebras. In *Proceedings of the Conference on Orders, Group Rings and Related Topics (Ohio State Univ., Columbus, Ohio, 1972)*, pages 8–71. Lecture Notes in Math., Vol. 353, 1973.
- [AR75] Maurice Auslander and Idun Reiten. On a generalized version of the Nakayama conjecture. *Proc. Amer. Math. Soc.*, 52:69–74, 1975.
- [AR91] Maurice Auslander and Idun Reiten. Applications of contravariantly finite subcategories. *Adv. Math.*, 86(1):111–152, 1991.
- [CE99] Henri Cartan and Samuel Eilenberg. *Homological algebra*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1999. With an appendix by David A. Buchsbaum, Reprint of the 1956 original.
- [Cha60] Stephen U. Chase. Direct products of modules. *Trans. Amer. Math. Soc.*, 97:457–473, 1960.
- [EHIS04] Karin Erdmann, Thorsten Holm, Osamu Iyama, and Jan Schröer. Radical embeddings and representation dimension. *Adv. Math.*, 185(1):159–177, 2004.
- [FGR75] Robert M. Fossum, Phillip A. Griffith, and Idun Reiten. *Trivial extensions of abelian categories*. Lecture Notes in Mathematics, Vol. 456. Springer-Verlag, Berlin-New York, 1975. Homological algebra of trivial extensions of abelian categories with applications to ring theory.
- [GKK91] Edward L. Green, Ellen Kirkman, and James Kuzmanovich. Finitistic dimensions of finite-dimensional monomial algebras. *J. Algebra*, 136(1):37–50, 1991.
- [Hap93] Dieter Happel. Reduction techniques for homological conjectures. *Tsukuba J. Math.*, 17(1):115–130, 1993.
- [hf] Jacob FG (<https://mathoverflow.net/users/157483/jacob fg>). Is there a finite dimensional algebra with left finitistic dimension different from its right finitistic dimension? MathOverflow. URL:<https://mathoverflow.net/q/376211> (version: 2020-11-11).
- [IST90] K. Igusa, S. O. Smalø, and G. Todorov. Finite projectivity and contravariant finiteness. *Proc. Amer. Math. Soc.*, 109(4):937–941, 1990.

- [IT05] Kiyoshi Igusa and Gordana Todorov. On the finitistic global dimension conjecture for Artin algebras. In *Representations of algebras and related topics*, volume 45 of *Fields Inst. Commun.*, pages 201–204. Amer. Math. Soc., Providence, RI, 2005.
- [Iya02] Osamu Iyama. Finiteness of representation dimension. In *Proceedings of the 34th Symposium on Ring Theory and Representation Theory (Maebashi, 2001)*, pages 21–25. Symp. Ring Theory Represent. Theory Organ. Comm., Tsukuba, 2002.
- [IZ90] Kiyoshi Igusa and Dan Zacharia. Syzygy pairs in a monomial algebra. *Proc. Amer. Math. Soc.*, 108(3):601–604, 1990.
- [Opp09] Steffen Oppermann. Representation dimension of artin algebras. https://folk.ntnu.no/opperman/repdim_survey.pdf, 2009.
- [Ric19] Jeremy Rickard. Unbounded derived categories and the finitistic dimension conjecture. *Adv. Math.*, 354:106735, 21, 2019.
- [Wan94] Yong Wang. A note on the finitistic dimension conjecture. *Comm. Algebra*, 22(7):2525–2528, 1994.
- [Xi02] Changchang Xi. Representation dimension and quasi-hereditary algebras. *Adv. Math.*, 168(2):193–212, 2002.