

1. Asian Citrus Psyllid

Names

English	Asian Citrus Psyllid
Hindi	एशियन साइट्रस सिल्ला
Marathi	लिंबू सिल्ला
Scientific Name	Diaphorina citri

Overview

English:

Asian Citrus Psyllid is one of the economically important citrus pests in India.

हिंदी:

एशियन साइट्रस सिल्ला भारत के प्रमुख साइट्रस कीटों में से एक है।

मराठी:

लिंबू सिल्ला हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Tiny brown insect (3–4 mm), wings held at 45°, yellow-orange nymphs on tender flush.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Sap-sucking insect and vector of Citrus Greening (HLB). Causes curling of tender leaves, stunted flush and reduced yield.

Life Cycle

Egg → Immature stage (nymph/larva) → Adult. Multiple generations may occur under warm conditions.

Monitoring

Inspect young flush and the underside of leaves every week. Record infestation level and capture photographs for AI datasets.

Integrated Pest Management

Regular monitoring, remove infected shoots, conserve *Tamarixia radiata*, neem oil, use registered systemic insecticides only when ETL is crossed.

AI Detection Features

- Images of eggs, immature stages and adults
- Healthy vs damaged leaves
- Different lighting, backgrounds and mobile cameras
- Bounding boxes and segmentation masks

JSON Metadata

```
{  
  "common_name": "Asian Citrus Psyllid",  
  "scientific_name": "Diaphorina citri",  
  "crop": "Citrus",  
  "country": "India"  
}
```

2. Citrus Leaf Miner

Names

English	Citrus Leaf Miner
Hindi	साइट्रस लीफ माइनर
Marathi	लिंबू पान खाणारी अळी
Scientific Name	Phyllocnistis citrella

Overview

English:

Citrus Leaf Miner is one of the economically important citrus pests in India.

हिंदी:

साइट्रस लीफ माइनर भारत के प्रमुख साइट्रस कीटों में से एक है।

मराठी:

लिंबू पान खाणारी अळी हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Silvery winding mines on young leaves, tiny moth, translucent larvae inside leaves.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Larvae create serpentine mines in young leaves causing curling and poor growth.

Life Cycle

Egg → Immature stage (nymph/larva) → Adult. Multiple generations may occur under warm conditions.

Monitoring

Inspect young flush and the underside of leaves every week. Record infestation level and capture photographs for AI datasets.

Integrated Pest Management

Prune damaged flush, avoid excess nitrogen, conserve parasitoids, neem products, registered insecticides if severe.

AI Detection Features

- Images of eggs, immature stages and adults
- Healthy vs damaged leaves

- Different lighting, backgrounds and mobile cameras
- Bounding boxes and segmentation masks

JSON Metadata

```
{  
  "common_name": "Citrus Leaf Miner",  
  "scientific_name": "Phyllocnistis citrella",  
  "crop": "Citrus",  
  "country": "India"  
}
```

3. Citrus Blackfly

Names

English	Citrus Blackfly
Hindi	साइट्रस ब्लैक फ्लाई
Marathi	लिंबू काळी पांढरी माशी
Scientific Name	Aleurocanthus woglumi

Overview

English:

Citrus Blackfly is one of the economically important citrus pests in India.

हिंदी:

साइट्रस ब्लैक फ्लाई भारत के प्रमुख साइट्रस कीटों में से एक है।

मराठी:

लिंबू काळी पांढरी माशी हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Black spiny immature stages on leaf underside, adults dark with white wax.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Sap sucking leads to honeydew and black sooty mold.

Life Cycle

Egg → Immature stage (nymph/larva) → Adult. Multiple generations may occur under warm conditions.

Monitoring

Inspect young flush and the underside of leaves every week. Record infestation level and capture photographs for AI datasets.

Integrated Pest Management

Monitor leaf undersides, encourage predators, horticultural oil and approved insecticides when required.

AI Detection Features

- Images of eggs, immature stages and adults
- Healthy vs damaged leaves

- Different lighting, backgrounds and mobile cameras
- Bounding boxes and segmentation masks

JSON Metadata

```
{  
  "common_name": "Citrus Blackfly",  
  "scientific_name": "Aleurocanthus woglumi",  
  "crop": "Citrus",  
  "country": "India"  
}
```

4. Citrus Whitefly

Names

English	Citrus Whitefly
Hindi	साइट्रस व्हाइटफ्लाई
Marathi	लिंबू पांढरी माशी
Scientific Name	Dialeurodes citri

Overview

English:

Citrus Whitefly is one of the economically important citrus pests in India.

हिंदी:

साइट्रस व्हाइटफ्लाई भारत के प्रमुख साइट्रस कीटों में से एक है।

मराठी:

लिंबू पांढरी माशी हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Tiny white powdery adults, flat nymphs beneath leaves.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Produces honeydew causing sooty mold and weakening trees.

Life Cycle

Egg → Immature stage (nymph/larva) → Adult. Multiple generations may occur under warm conditions.

Monitoring

Inspect young flush and the underside of leaves every week. Record infestation level and capture photographs for AI datasets.

Integrated Pest Management

Yellow sticky traps, neem oil, biological control, registered insecticides if infestation is high.

AI Detection Features

- Images of eggs, immature stages and adults
- Healthy vs damaged leaves

- Different lighting, backgrounds and mobile cameras
- Bounding boxes and segmentation masks

JSON Metadata

```
{  
  "common_name": "Citrus Whitefly",  
  "scientific_name": "Dialeurodes citri",  
  "crop": "Citrus",  
  "country": "India"  
}
```

5. Brown Citrus Aphid

Names

English	Brown Citrus Aphid
Hindi	भूरी साइट्स माहू
Marathi	तपकिरी लिंबू मावा
Scientific Name	Toxoptera citricida

Overview

English:

Brown Citrus Aphid is one of the economically important citrus pests in India.

हिंदी:

भूरी साइट्स माहू भारत के प्रमुख साइट्स कीटों में से एक है।

मराठी:

तपकिरी लिंबू मावा हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Glossy dark-brown aphids clustered on tender shoots.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Heavy sap loss and transmission of citrus tristeza virus.

Life Cycle

Egg → Immature stage (nymph/larva) → Adult. Multiple generations may occur under warm conditions.

Monitoring

Inspect young flush and the underside of leaves every week. Record infestation level and capture photographs for AI datasets.

Integrated Pest Management

Inspect new flush, conserve ladybird beetles, neem formulations and selective insecticides if necessary.

AI Detection Features

- Images of eggs, immature stages and adults
- Healthy vs damaged leaves

- Different lighting, backgrounds and mobile cameras
- Bounding boxes and segmentation masks

JSON Metadata

```
{  
  "common_name": "Brown Citrus Aphid",  
  "scientific_name": "Toxoptera citricida",  
  "crop": "Citrus",  
  "country": "India"  
}
```

6. Cotton Aphid

English	Cotton Aphid
Hindi	कॉटन एफिड
Marathi	कापूस मावा
Scientific Name	Aphis gossypii

Overview

Cotton Aphid is an important citrus pest found in India.

हिंदी: कॉटन एफिड साइट्रस की महत्वपूर्ण कीट है।

मराठी: कापूस मावा हा लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Small soft-bodied aphid that sucks sap from tender shoots and leaves.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Leaf curling, sticky honeydew, sooty mold.

Life Cycle

Egg → Immature stage → Adult. Multiple generations under suitable conditions.

Monitoring

Inspect young leaves, shoots and fruits weekly. Record infestation and capture images.

Integrated Pest Management

Inspect new flush, conserve ladybird beetles, use neem and selective insecticides.

AI Detection Features

Collect images of eggs, immature stages, adults, damage symptoms and healthy plant parts under different lighting.

JSON Metadata

```
{"common_name":"Cotton Aphid","scientific_name":"Aphis gossypii","crop":"Citrus"}
```

7. Citrus Mealybug

English	Citrus Mealybug
Hindi	साइट्रस मिलीबग
Marathi	लिंबू मिलीबग
Scientific Name	Planococcus citri

Overview

Citrus Mealybug is an important citrus pest found in India.

हिंदी: साइट्रस मिलीबग साइट्रस की महत्वपूर्ण कीट है।

मराठी: लिंबू मिलीबग हा लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

White wax-covered sap-sucking insect found on leaves, shoots and fruits.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Honeydew, sooty mold, poor fruit quality.

Life Cycle

Egg → Immature stage → Adult. Multiple generations under suitable conditions.

Monitoring

Inspect young leaves, shoots and fruits weekly. Record infestation and capture images.

Integrated Pest Management

Prune infested twigs, encourage predators, neem oil, registered insecticides if needed.

AI Detection Features

Collect images of eggs, immature stages, adults, damage symptoms and healthy plant parts under different lighting.

JSON Metadata

```
{"common_name":"Citrus Mealybug","scientific_name":"Planococcus citri","crop":"Citrus"}
```

8. California Red Scale

English	California Red Scale
Hindi	कैलिफोर्निया रेड स्केल
Marathi	कॅलिफोर्निया रेड स्केल
Scientific Name	Aonidiella aurantii

Overview

California Red Scale is an important citrus pest found in India.

हिंदी: कैलिफोर्निया रेड स्केल साइटस की महत्वपूर्ण कीट है।

मराठी: कॅलिफोर्निया रेड स्केल हा लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Armored scale insect attacking fruits, leaves and branches.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Yellow spots, fruit blemishes, twig weakening.

Life Cycle

Egg → Immature stage → Adult. Multiple generations under suitable conditions.

Monitoring

Inspect young leaves, shoots and fruits weekly. Record infestation and capture images.

Integrated Pest Management

Monitor regularly, horticultural oil, conserve parasitoids.

AI Detection Features

Collect images of eggs, immature stages, adults, damage symptoms and healthy plant parts under different lighting.

JSON Metadata

```
{"common_name":"California Red Scale","scientific_name":"Aonidiella aurantii","crop":"Citrus"}
```

9. Purple Scale

English	Purple Scale
Hindi	पर्पल स्केल
Marathi	जांभळी स्केल
Scientific Name	Lepidosaphes beckii

Overview

Purple Scale is an important citrus pest found in India.

हिंदी: पर्पल स्केल साइट्रस की महत्वपूर्ण कीट है।

मराठी: जांभळी स्केल हा लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Scale insect commonly found on leaves and fruits.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Yellowing leaves, reduced vigor and fruit quality.

Life Cycle

Egg → Immature stage → Adult. Multiple generations under suitable conditions.

Monitoring

Inspect young leaves, shoots and fruits weekly. Record infestation and capture images.

Integrated Pest Management

Pruning, orchard sanitation, horticultural oils and biological control.

AI Detection Features

Collect images of eggs, immature stages, adults, damage symptoms and healthy plant parts under different lighting.

JSON Metadata

```
{"common_name":"Purple Scale","scientific_name":"Lepidosaphes beckii","crop":"Citrus"}
```

10. Green Shield Scale

English	Green Shield Scale
Hindi	ग्रीन शील्ड स्केल
Marathi	हिरवी स्केल
Scientific Name	Coccus viridis

Overview

Green Shield Scale is an important citrus pest found in India.

हिंदी: ग्रीन शील्ड स्केल साइट्रस की महत्वपूर्ण कीट है।

मराठी: हिरवी स्केल हा लिंबूवर्गीय पिकांवरील महत्वाचा किड आहे.

Identification

Soft scale producing abundant honeydew.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Sticky leaves, black sooty mold and reduced photosynthesis.

Life Cycle

Egg → Immature stage → Adult. Multiple generations under suitable conditions.

Monitoring

Inspect young leaves, shoots and fruits weekly. Record infestation and capture images.

Integrated Pest Management

Encourage natural enemies, prune heavily infested branches, neem-based sprays.

AI Detection Features

Collect images of eggs, immature stages, adults, damage symptoms and healthy plant parts under different lighting.

JSON Metadata

```
{"common_name":"Green Shield Scale","scientific_name":"Coccus viridis","crop":"Citrus"}
```

11. Citrus Red Mite

English	Citrus Red Mite
Hindi	साइट्रस रेड माइट
Marathi	लाल कोळी

Scientific Name	Panonychus citri
-----------------	------------------

Overview

English:

Citrus Red Mite is an economically important citrus pest found in India.

हिंदी:

साइट्रस रेड माइट भारत में पाया जाने वाला एक महत्वपूर्ण साइट्रस कीट/माइट है।

मराठी:

लाल कोळी हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड/कोळी आहे.

Identification

Tiny red mite feeding on leaf sap.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Leaf bronzing, yellowing and reduced photosynthesis.

Life Cycle

Egg → Immature stage → Adult. Several generations may occur each year under favorable weather.

Monitoring

Inspect leaves and fruits weekly. Record infestation level and capture field images for AI training.

Integrated Pest Management

Monitor leaf undersides, conserve predatory mites, use miticides only when required.

AI Detection Features

Capture images of all life stages, healthy and damaged leaves, fruits, multiple angles, lighting conditions and devices.

JSON Metadata

```
{"common_name":"Citrus Red Mite","scientific_name":"Panonychus citri","crop":"Citrus","country":"India"}
```

12. Oriental Spider Mite

English	Oriental Spider Mite
Hindi	ओरिएंटल स्पाइडर माइट
Marathi	पूर्वेकडील कोळी
Scientific Name	Eutetranychus orientalis

Overview

English:

Oriental Spider Mite is an economically important citrus pest found in India.

हिंदी:

ओरिएंटल स्पाइडर माइट भारत में पाया जाने वाला एक महत्वपूर्ण साइट्रस कीट/माइट है।

मराठी:

पूर्वेकडील कोळी हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड/कोळी आहे.

Identification

Spider mite causing chlorotic patches on citrus leaves.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Leaf discoloration, premature leaf fall.

Life Cycle

Egg → Immature stage → Adult. Several generations may occur each year under favorable weather.

Monitoring

Inspect leaves and fruits weekly. Record infestation level and capture field images for AI training.

Integrated Pest Management

Avoid dusty conditions, encourage natural enemies, apply registered miticides if needed.

AI Detection Features

Capture images of all life stages, healthy and damaged leaves, fruits, multiple angles, lighting conditions and devices.

JSON Metadata

```
{"common_name":"Oriental Spider Mite","scientific_name":"Eutetranychus orientalis","crop":"Citrus","country":"India"}
```


13. Citrus Rust Mite

English	Citrus Rust Mite
Hindi	साइट्रस रस्ट माइट
Marathi	लिंबू रस्ट माइट
Scientific Name	Phyllocoptruta oleivora

Overview

English:

Citrus Rust Mite is an economically important citrus pest found in India.

हिंदी:

साइट्रस रस्ट माइट भारत में पाया जाने वाला एक महत्वपूर्ण साइट्रस कीट/माइट है।

मराठी:

लिंबू रस्ट माइट हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड/कोळी आहे.

Identification

Microscopic mite attacking leaves and fruits.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Fruit russetting and dull fruit appearance.

Life Cycle

Egg → Immature stage → Adult. Several generations may occur each year under favorable weather.

Monitoring

Inspect leaves and fruits weekly. Record infestation level and capture field images for AI training.

Integrated Pest Management

Monitor fruits regularly, maintain orchard health, use appropriate miticides when ETL is exceeded.

AI Detection Features

Capture images of all life stages, healthy and damaged leaves, fruits, multiple angles, lighting conditions and devices.

JSON Metadata

```
{"common_name": "Citrus Rust Mite", "scientific_name": "Phyllocoptruta  
oleivora", "crop": "Citrus", "country": "India"}
```

14. Yellow Citrus Thrips

English	Yellow Citrus Thrips
Hindi	येलो साइट्रस थ्रिप्स
Marathi	पिवळी फुलकिडी
Scientific Name	Scirtothrips dorsalis

Overview

English:

Yellow Citrus Thrips is an economically important citrus pest found in India.

हिंदी:

येलो साइट्रस थ्रिप्स भारत में पाया जाने वाला एक महत्वपूर्ण साइट्रस कीट/माइट है।

मराठी:

पिवळी फुलकिडी हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड/कोळी आहे.

Identification

Tiny yellow thrips feeding on young leaves and fruits.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Leaf curling and fruit scarring.

Life Cycle

Egg → Immature stage → Adult. Several generations may occur each year under favorable weather.

Monitoring

Inspect leaves and fruits weekly. Record infestation level and capture field images for AI training.

Integrated Pest Management

Blue/yellow sticky traps, weed management, conserve predators, selective insecticides.

AI Detection Features

Capture images of all life stages, healthy and damaged leaves, fruits, multiple angles, lighting conditions and devices.

JSON Metadata

```
{"common_name":"Yellow Citrus Thrips","scientific_name":"Scirtothrips dorsalis","crop":"Citrus","country":"India"}
```


15. Citrus Thrips

English	Citrus Thrips
Hindi	साइट्रस थ्रिप्स
Marathi	लिंबू थ्रिप्स
Scientific Name	Scirtothrips citri

Overview

English:

Citrus Thrips is an economically important citrus pest found in India.

हिंदी:

साइट्रस थ्रिप्स भारत में पाया जाने वाला एक महत्वपूर्ण साइट्रस कीट/माइट है।

मराठी:

लिंबू थ्रिप्स हा भारतातील लिंबूवर्गीय पिकांवरील महत्वाचा किड/कोळी आहे.

Identification

Thrips damaging tender foliage and young fruits.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Silvering, scarring and distorted fruits.

Life Cycle

Egg → Immature stage → Adult. Several generations may occur each year under favorable weather.

Monitoring

Inspect leaves and fruits weekly. Record infestation level and capture field images for AI training.

Integrated Pest Management

Regular monitoring, orchard sanitation, biological control and selective insecticides.

AI Detection Features

Capture images of all life stages, healthy and damaged leaves, fruits, multiple angles, lighting conditions and devices.

JSON Metadata

```
{"common_name":"Citrus Thrips","scientific_name":"Scirtothrips citri","crop":"Citrus","country":"India"}
```

16. Fruit Sucking Moth

English	Fruit Sucking Moth
Hindi	फल चूसने वाला पतंगा
Marathi	फळरस शोषण करणारा पतंग
Scientific Name	Eudocima materna

Overview

English:

Fruit Sucking Moth is an economically important citrus pest affecting yield and fruit quality.

हिंदी:

फल चूसने वाला पतंगा साइट्रस फसलों का एक महत्वपूर्ण कीट है जो उत्पादन और गुणवत्ता को प्रभावित करता है।

मराठी:

फळरस शोषण करणारा पतंग हा लिंबूवर्गीय पिकांवरील महत्वाचा किड असून उत्पादन व गुणवत्तेवर परिणाम करतो.

Identification

Large nocturnal moth that pierces ripening citrus fruits to suck juice.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Pierced fruits, juice leakage, rotting and secondary fungal infection.

Life Cycle

Egg → Larva/Caterpillar or Maggot → Pupa → Adult. Development varies with weather conditions.

Monitoring

Inspect fruits, stems and branches weekly. Record pest incidence and capture field images for AI datasets.

Integrated Pest Management

Collect and destroy damaged fruits, use fruit bagging where practical, light traps and integrated pest management.

AI Detection Features

Collect images of eggs, larvae, pupae, adults, entry holes, damaged fruits, bark symptoms, healthy plants, and different environmental conditions.

JSON Metadata

```
{"common_name":"Fruit Sucking Moth","scientific_name":"Eudocima  
materna","crop":"Citrus","country":"India"}
```

17. Lemon Butterfly

English	Lemon Butterfly
Hindi	नींबू तितली
Marathi	लिंबू फुलपाखरू
Scientific Name	Papilio demoleus

Overview

English:

Lemon Butterfly is an economically important citrus pest affecting yield and fruit quality.

हिंदी:

नींबू तितली साइट्रस फसलों का एक महत्वपूर्ण कीट है जो उत्पादन और गुणवत्ता को प्रभावित करता है।

मराठी:

लिंबू फुलपाखरू हा लिंबूवर्गीय पिकांवरील महत्वाचा किड असून उत्पादन व गुणवत्तेवर परिणाम करतो.

Identification

Caterpillars feed aggressively on citrus leaves, especially in young orchards.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Severe defoliation, reduced plant growth and delayed establishment.

Life Cycle

Egg → Larva/Caterpillar or Maggot → Pupa → Adult. Development varies with weather conditions.

Monitoring

Inspect fruits, stems and branches weekly. Record pest incidence and capture field images for AI datasets.

Integrated Pest Management

Hand-pick larvae, conserve natural enemies, use biological control and approved insecticides only if needed.

AI Detection Features

Collect images of eggs, larvae, pupae, adults, entry holes, damaged fruits, bark symptoms, healthy plants, and different environmental conditions.

JSON Metadata

```
{"common_name": "Lemon Butterfly", "scientific_name": "Papilio  
demoleus", "crop": "Citrus", "country": "India"}
```

18. Bark Eating Caterpillar

English	Bark Eating Caterpillar
Hindi	छाल खाने वाली इल्ली
Marathi	साल खाणारी अळी
Scientific Name	Indarbela quadrinotata

Overview

English:

Bark Eating Caterpillar is an economically important citrus pest affecting yield and fruit quality.

हिंदी:

छाल खाने वाली इल्ली साइट्रस फसलों का एक महत्वपूर्ण कीट है जो उत्पादन और गुणवत्ता को प्रभावित करता है।

मराठी:

साल खाणारी अळी हा लिंबूवर्गीय पिकांवरील महत्वाचा किड असून उत्पादन व गुणवत्तेवर परिणाम करतो.

Identification

Caterpillar bores into bark and branches while feeding internally.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Holes in bark, webbing with frass, branch weakening and dieback.

Life Cycle

Egg → Larva/Caterpillar or Maggot → Pupa → Adult. Development varies with weather conditions.

Monitoring

Inspect fruits, stems and branches weekly. Record pest incidence and capture field images for AI datasets.

Integrated Pest Management

Remove webbing, insert wire into tunnels, prune badly affected branches and maintain orchard sanitation.

AI Detection Features

Collect images of eggs, larvae, pupae, adults, entry holes, damaged fruits, bark symptoms, healthy plants, and different environmental conditions.

JSON Metadata

```
{"common_name": "Bark Eating Caterpillar", "scientific_name": "Indarbela  
quadrinotata", "crop": "Citrus", "country": "India"}
```

19. Citrus Trunk Borer

English	Citrus Trunk Borer
Hindi	साइट्रस तना छेदक
Marathi	लिंबू खोड पोखरणारा भुंगा
Scientific Name	Anoplophora versteegi

Overview

English:

Citrus Trunk Borer is an economically important citrus pest affecting yield and fruit quality.

हिंदी:

साइट्रस तना छेदक साइट्रस फसलों का एक महत्वपूर्ण कीट है जो उत्पादन और गुणवत्ता को प्रभावित करता है।

मराठी:

लिंबू खोड पोखरणारा भुंगा हा लिंबूवर्गीय पिकांवरील महत्वाचा किड असून उत्पादन व गुणवत्तेवर परिणाम करतो.

Identification

Wood-boring beetle attacking trunk and major branches.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Round exit holes, frass, branch drying and reduced tree vigor.

Life Cycle

Egg → Larva/Caterpillar or Maggot → Pupa → Adult. Development varies with weather conditions.

Monitoring

Inspect fruits, stems and branches weekly. Record pest incidence and capture field images for AI datasets.

Integrated Pest Management

Regular inspection, destroy infested wood, protect trunk and use recommended management practices.

AI Detection Features

Collect images of eggs, larvae, pupae, adults, entry holes, damaged fruits, bark symptoms, healthy plants, and different environmental conditions.

JSON Metadata

```
{"common_name": "Citrus Trunk Borer", "scientific_name": "Anoplophora  
versteegi", "crop": "Citrus", "country": "India"}
```

20. Oriental Fruit Fly

English	Oriental Fruit Fly
Hindi	ओरिएंटल फ्रूट फ्लाई
Marathi	फळ माशी
Scientific Name	Bactrocera dorsalis

Overview

English:

Oriental Fruit Fly is an economically important citrus pest affecting yield and fruit quality.

हिंदी:

ओरिएंटल फ्रूट फ्लाई साइट्रस फसलों का एक महत्वपूर्ण कीट है जो उत्पादन और गुणवत्ता को प्रभावित करता है।

मराठी:

फळ माशी हा लिंबूवर्गीय पिकांवरील महत्वाचा किड असून उत्पादन व गुणवत्तेवर परिणाम करतो.

Identification

Adult flies lay eggs inside fruits; maggots feed internally.

Host Plants

Orange, Lemon, Lime, Mandarin, Sweet Lime and Grapefruit.

Damage Symptoms

Fruit punctures, rotting, premature fruit drop and market losses.

Life Cycle

Egg → Larva/Caterpillar or Maggot → Pupa → Adult. Development varies with weather conditions.

Monitoring

Inspect fruits, stems and branches weekly. Record pest incidence and capture field images for AI datasets.

Integrated Pest Management

Collect fallen fruits, methyl eugenol traps, orchard sanitation, bait sprays and integrated management.

AI Detection Features

Collect images of eggs, larvae, pupae, adults, entry holes, damaged fruits, bark symptoms, healthy plants, and different environmental conditions.

JSON Metadata

```
{"common_name":"Oriental Fruit Fly","scientific_name":"Bactrocera dorsalis","crop":"Citrus","country":"India"}
```