

Guideline for the Management of Chronic Condition A

Strength of recommendation: Strong. Certainty of evidence: Moderate (GRADE). Benefits are judged to outweigh harms for most patients in this population.

1. Scope and Purpose

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

2. Screening Recommendations

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

3. Pharmacologic Management

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

4. Monitoring and Follow-up

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.