

Abstract

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts. This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts. This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts. This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.

This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts. This recommendation applies to adult patients presenting in the ambulatory setting. Clinicians should assess baseline risk before initiating therapy and reassess at each follow-up visit. Shared decision making is advised when the balance of benefits and harms is uncertain. The strength of this recommendation reflects moderate-certainty evidence derived from randomized controlled trials and well-conducted observational cohorts.