MONOTONICITY OF ERDŐS SUMS

JAIDEEP SAI PADHI

ABSTRACT. For $k \geq 1$ let $f_k = \sum_{\Omega(n)=k} 1/(n \log n)$, where Ω counts prime factors with multiplicity. We prove that $f_k < f_{k+1}$ for every $k \geq 6$, whereas the sums g_k of $1/(n \log n)$ over odd n with $\Omega(n) = k$ satisfy $g_k > g_{k+1}$ for every $k \geq 1$. The two statements arise from the same analytic distinction. After writing the relevant Dirichlet series as $\zeta(s)^z H(s, z)$, the point $z = 2$ lies on the first singular circle for the unrestricted product, and strictly inside the first singular circle of the product with the prime 2 removed. The analytic reductions are written explicitly below. The finite inequalities and coefficient bounds used in the last step are certified by ball arithmetic; the separation between analytic argument and certified input is stated precisely.

1. INTRODUCTION

For $k \geq 1$ put

$$f_k = \sum_{\Omega(n)=k} \frac{1}{n \log n}.$$

Erdős [4] proved that the corresponding sum is bounded uniformly over primitive sets. In particular each f_k converges. Zhang [7] proved $f_k < f_1$ for $k > 1$, and Bayless, Kinlaw and Klyve [8] proved $f_2 > f_3$. Banks and Martin [1] conjectured that

$$f_1 > f_2 > f_3 > \cdots . \tag{1}$$

Lichtman [2] disproved (1), showing that $f_k \to 1$ and that the sequence attains a global minimum at $k = 6$. Gorodetsky, Lichtman and Wong [3] then established

$$f_k = 1 - 2^{-k}\big(a_1 k^2 + O(k \log(k+1))\big), \qquad a_1 = \frac{d \log 2}{4} > 0, \tag{2}$$

where

$$d = \frac{1}{4} \prod_{p \geq 3} \left(1 - \frac{2}{p}\right)^{-1} \left(1 - \frac{1}{p}\right)^{2}. \tag{3}$$

Neither of these yields monotonicity beyond the minimum. A global minimum at $k = 6$ leaves open whether $f_k < f_{k+1}$ holds at every index above it, and the implicit constant in (2) does not make explicit a range on which the increase can be read off. Our first theorem gives that monotonicity in full.

Theorem 1.1. *For every $k \geq 6$ one has $f_k < f_{k+1}$.*

Let

$$g_k = \sum_{\substack{n \text{ odd} \\ \Omega(n)=k}} \frac{1}{n \log n}.$$

The second theorem shows that deleting the prime 2 reverses the direction of monotonicity at every index.

2020 *Mathematics Subject Classification.* Primary 11N25; Secondary 11Y60, 68V20.

Key words and phrases. Erdős sums, almost primes, primitive sets, ball arithmetic, formalized mathematics.

Theorem 1.2. *For every $k \geq 1$ one has $g_k > g_{k+1}$, and $g_k \to \frac{1}{2}$.*

The proof of Theorem 1.2 gives the more precise expansion

$$g_k = \frac{1}{2} + \beta 2^{-k} + R_k, \qquad |R_k| \leq 4900(k+3)^3 3^{-k} \quad (k \geq 59), \tag{4}$$

where $\gamma = 0.5772156649\ldots$ is Euler's constant and

$$\beta = d\left(\gamma + \log 2 - \sum_{p \geq 3} \frac{\log p}{(p-1)(p-2)}\right) > 0. \tag{5}$$

The 2^{-k} term comes from evaluating a regularized Euler product at $z = 2$. For odd integers the first pole is at $z = 3$, so this evaluation takes place strictly inside the disc of analyticity. For the unrestricted sums the first pole is at $z = 2$ itself. The boundary contribution then acquires the factor k^2 in (2).

The proof has two kinds of input. Identities, coefficient reductions and inequalities are proved in the text. Bounds produced by interval arithmetic are isolated as certified propositions. Thus a sentence of the form "the computation gives" never substitutes for an analytic step: the exact inequality supplied by the computation is stated before it is used.

2. Generating functions and the analytic boundary

For $s > 1$ define

$$P_k(s) = \sum_{\Omega(n)=k} n^{-s}, \qquad P_k^{\mathrm{odd}}(s) = \sum_{\substack{n \text{ odd} \\ \Omega(n)=k}} n^{-s}.$$

Lemma 2.1. *For every $k \geq 1$,*

$$f_k = \int_1^\infty P_k(s)\,\mathrm{d}s, \qquad g_k = \int_1^\infty P_k^{\mathrm{odd}}(s)\,\mathrm{d}s. \tag{6}$$

Proof. For $n \geq 2$,

$$\int_1^\infty n^{-s}\,\mathrm{d}s = \int_1^\infty \mathrm{e}^{-s\log n}\,\mathrm{d}s = \left[-\frac{\mathrm{e}^{-s\log n}}{\log n}\right]_{s=1}^{\infty} = \frac{1}{n\log n}.$$

All summands are non-negative. Tonelli's theorem therefore permits the sum and integral to be interchanged in both cases, giving

$$\int_1^\infty \sum_{\Omega(n)=k} n^{-s}\,\mathrm{d}s = \sum_{\Omega(n)=k} \int_1^\infty n^{-s}\,\mathrm{d}s = f_k,$$

and the same calculation with the sum restricted to odd n proves the second identity. □

The generating functions are

$$\sum_{k\geq 0} P_k(s) z^k = \prod_p (1 - zp^{-s})^{-1}, \qquad \sum_{k\geq 0} P_k^{\mathrm{odd}}(s) z^k = \prod_{p\geq 3} (1 - zp^{-s})^{-1}. \tag{7}$$

Write these products as $\zeta(s)^z H(s,z)$ and $\zeta(s)^z H_{\mathrm{o}}(s,z)$, where

$$H(s,z) = \prod_p (1 - zp^{-s})^{-1}(1 - p^{-s})^z, \tag{8}$$

$$H_{\mathrm{o}}(s,z) = (1 - 2^{-s})^z \prod_{p\geq 3} (1 - zp^{-s})^{-1}(1 - p^{-s})^z. \tag{9}$$

Proposition 2.2. *For fixed real $s > 1/2$, the function $H(s,\cdot)$ is analytic in $|z| < 2^s$, and $H_{\mathrm{o}}(s,\cdot)$ is analytic in $|z| < 3^s$. The convergence is moreover locally uniform in (s,z) jointly, so each $\partial_s^r H_{\mathrm{o}}(s,\cdot)$ is again analytic in $|z| < 3^s$, and*

$$\frac{\mathrm{d}^r}{\mathrm{d}s^r}[z^j]H_{\mathrm{o}}(s,z) = [z^j]\partial_s^r H_{\mathrm{o}}(s,z); \tag{10}$$

the same holds for H in $|z| < 2^s$.

Proof. Let $x = p^{-s}$. For z in a fixed compact set, the logarithm of the pth regularized factor is

$$\begin{aligned} -\log(1-zx) + z\log(1-x) &= \sum_{m\ge 1} \frac{z^m - z}{m} x^m \\ &= \sum_{m\ge 2} \frac{z^m - z}{m} x^m = O_K(p^{-2s}). \end{aligned}$$

The estimate is uniform for z in the compact set K. Since $\sum_p p^{-2s} < \infty$, the product of the regularized factors converges locally uniformly wherever none of the factors has a pole. The poles occur at $z = p^s$. The nearest one is 2^s for (8) and 3^s for (9). A locally uniform limit of analytic functions is analytic, which proves the first assertion.

The estimate is uniform in s as well on any compact subset of $(1/2,\infty)$, since $p^{-2s} \le p^{-2s_0}$ there, and on a compact subset of $\{(s,z) : |z| < 3^s\}$ no factor comes near a pole. The convergence is therefore locally uniform in the two variables jointly, so differentiation in s may be carried out under the product. Since $[z^j]$ is a Cauchy integral over a circle that may be held fixed on such a subset, the two operations commute, which is (10). □

At $s = 1$ the point $z = 2$ lies on the first singular circle for H and lies strictly inside the first singular circle for H_{o}. This is the analytic distinction used below.

We shall also need the local change of variables used in the odd case.

Lemma 2.3. *Let $u = \log\zeta(s)$ for $s > 1$, and let $s = s(u)$ be the inverse function. Put $\varepsilon(u) = s(u) - 1$ and $J(u) = -s'(u)$. As $u \to \infty$,*

$$\varepsilon(u) = \mathrm{e}^{-u} + \gamma\mathrm{e}^{-2u} + O(\mathrm{e}^{-3u}), \qquad J(u) = \mathrm{e}^{-u} + 2\gamma\mathrm{e}^{-2u} + O(\mathrm{e}^{-3u}). \tag{11}$$

Proof. For $\varepsilon \to 0^+$,

$$\zeta(1+\varepsilon) = \varepsilon^{-1} + \gamma + O(\varepsilon).$$

Set $x = \mathrm{e}^{-u} = 1/\zeta(1+\varepsilon)$. Then

$$x = \frac{\varepsilon}{1 + \gamma\varepsilon + O(\varepsilon^2)} = \varepsilon - \gamma\varepsilon^2 + O(\varepsilon^3).$$

Inverting this power series gives $\varepsilon = x + \gamma x^2 + O(x^3)$, which is the first formula in (11). Since $x' = -x$, differentiation gives

$$\varepsilon'(u) = -x - 2\gamma x^2 + O(x^3).$$

Thus $J(u) = -\varepsilon'(u) = x + 2\gamma x^2 + O(x^3)$. □

3. The odd sums

Throughout this section write

$$h_j(s) = [z^j]H_{\mathrm{o}}(s,z), \qquad b_j = h_j'(1) + 2\gamma h_j(1).$$

Since $\zeta(s)^z = \sum_{m\geq 0}(\log\zeta(s))^m z^m/m!$, coefficient extraction in (7) gives the exact identity

$$P_k^{\mathrm{odd}}(s) = \sum_{j=0}^{k} h_j(s)\frac{(\log\zeta(s))^{k-j}}{(k-j)!}. \tag{12}$$

Using $u = \log\zeta(s)$ in Lemma 2.1, and noting that s decreases from ∞ to 1 as u increases from 0 to ∞, we obtain

$$g_k = \sum_{j=0}^{k}\int_0^\infty \frac{u^{k-j}}{(k-j)!}\, h_j(s(u))J(u)\,\mathrm{d}u. \tag{13}$$

3.1. **The first two terms.**

Proposition 3.1. *There is a remainder R_k such that*

$$g_k = H_{\mathrm{o}}(1,1) + 2^{-k-1}\big(\partial_s H_{\mathrm{o}}(1,2) + 2\gamma H_{\mathrm{o}}(1,2)\big) + R_k. \tag{14}$$

Proof. Let $x = \mathrm{e}^{-u}$. By Lemma 2.3, $J(u) = x + 2\gamma x^2 + O(x^3)$ and $\varepsilon(u)J(u) = x^2 + O(x^3)$. Taylor's theorem at $s = 1$ gives

$$h_j(1+\varepsilon) = h_j(1) + \varepsilon h_j'(1) + \frac{\varepsilon^2}{2}h_j''(1+\theta_{j,u}\varepsilon)$$

for some $0 < \theta_{j,u} < 1$. Put

$$r_0(u) = J(u) - \mathrm{e}^{-u} - 2\gamma\mathrm{e}^{-2u}, \qquad r_1(u) = \varepsilon(u)J(u) - \mathrm{e}^{-2u}.$$

Multiplication by $J(u)$ now gives the exact identity

$$\begin{aligned} h_j(s(u))J(u) &= h_j(1)\mathrm{e}^{-u} + b_j\mathrm{e}^{-2u} + E_j(u),\\ E_j(u) &= h_j(1)r_0(u) + h_j'(1)r_1(u) + \frac{\varepsilon(u)^2 J(u)}{2}h_j''(1+\theta_{j,u}\varepsilon(u)). \end{aligned} \tag{15}$$

Thus no error is suppressed in the notation E_j: its three summands are the Jacobian error, the first-order coordinate error and the Taylor remainder.

Insert (15) into (13). Since

$$\int_0^\infty \frac{u^m}{m!}\mathrm{e}^{-ru}\,\mathrm{d}u = r^{-m-1} \qquad (r>0),$$

the first two terms become

$$\sum_{j=0}^{k} h_j(1) = H_{\mathrm{o}}(1,1) - \sum_{j>k} h_j(1), \tag{16}$$

$$\sum_{j=0}^{k} b_j 2^{-(k-j+1)} = 2^{-k-1}\big(\partial_s H_{\mathrm{o}}(1,2) + 2\gamma H_{\mathrm{o}}(1,2)\big) - 2^{-k-1}\sum_{j>k} b_j 2^j. \tag{17}$$

Define R_k to be the integral of the E_j terms together with the two tails on the right of (16) and (17). This gives (14). □

Lemma 3.2. *One has*

$$H_{\mathrm{o}}(1,1) = \frac{1}{2}, \qquad H_{\mathrm{o}}(1,2) = d, \tag{18}$$

and

$$\partial_s \log H_{\mathrm{o}}(1,2) = 2\log 2 - 2\sum_{p\geq 3}\frac{\log p}{(p-1)(p-2)}. \tag{19}$$

Proof. At $s = z = 1$, every factor in the product over $p \geq 3$ cancels, and hence

$$H_{\mathrm{o}}(1,1) = (1 - 2^{-1}) = \frac{1}{2}.$$

At $(s, z) = (1, 2)$,

$$H_{\mathrm{o}}(1,2) = \frac{1}{4} \prod_{p \geq 3} \left(1 - \frac{2}{p}\right)^{-1} \left(1 - \frac{1}{p}\right)^{2} = d.$$

For the derivative, take logarithms:

$$\log H_{\mathrm{o}}(s,2) = 2\log(1 - 2^{-s}) + \sum_{p \geq 3} \left[-\log(1 - 2p^{-s}) + 2\log(1 - p^{-s}) \right].$$

The differentiated pth summand is $O((\log p)p^{-2s})$ locally uniformly near $s = 1$, so termwise differentiation is permitted. At $s = 1$ the first term contributes $2\log 2$, while the pth summand contributes

$$-\frac{2\log p}{p-2} + \frac{2\log p}{p-1} = -\frac{2\log p}{(p-1)(p-2)}.$$

Summing over $p \geq 3$ proves (19). □

Combining Proposition 3.1 and Lemma 3.2 gives

$$g_k = \frac{1}{2} + \beta 2^{-k} + R_k, \qquad \beta = d\left(\gamma + \log 2 - \sum_{p \geq 3} \frac{\log p}{(p-1)(p-2)}\right). \tag{20}$$

3.2. **The remainder.** The following proposition records exactly the interval-arithmetic input used for the odd sums. The coefficient sums are weighted ℓ^1 norms; their finiteness follows analytically from Proposition 2.2, while the displayed numerical bounds are certified.

Proposition 3.3 (Certified odd input)**.** *Let*

$$\Sigma_0 = \sum_{m \geq 0} 3^m |[z^m](1 - z/3) H_{\mathrm{o}}(1, z)|,$$
$$\Sigma_1 = \sum_{m \geq 0} 3^m |[z^m](1 - z/3)^2 \partial_s H_{\mathrm{o}}(1, z)|,$$
$$\Sigma_2 = \sup_{1 \leq s \leq s(3)} \sum_{m \geq 0} 3^m |[z^m](1 - z/3)^3 \partial_s^2 H_{\mathrm{o}}(s, z)|.$$

The certified computation gives

$$\Sigma_0 \leq 44.5987, \qquad \Sigma_1 \leq 486.1945, \qquad \Sigma_2 \leq 14414. \tag{21}$$

Split the remainder of Proposition 3.1 at $u = 3$: with E_j the explicit expression in (15)*, put*

$$R_{k,\mathrm{far}} = \sum_{j=0}^{k} \int_3^\infty \frac{u^{k-j}}{(k-j)!} E_j(u)\, \mathrm{d}u, \qquad R_{k,\mathrm{near}} = \sum_{j=0}^{k} \int_0^3 \frac{u^{k-j}}{(k-j)!} E_j(u)\, \mathrm{d}u, \tag{22}$$

so that R_k is $R_{k,\mathrm{far}} + R_{k,\mathrm{near}}$ together with the two coefficient tails of (16)–(17)*.*

For the far range, write $\varepsilon = s - 1$ and $J = \zeta(s)/(-\zeta'(s))$, and let

$$C_b = \sup|\varepsilon\zeta|, \qquad C_f = \sup|\zeta^2(J - \zeta^{-1})|,$$
$$C_J = \sup\left|\zeta^3\left(J - \zeta^{-1} - 2\gamma\zeta^{-2}\right)\right|, \qquad C_P = \sup\left|\zeta^3\left(\varepsilon J - \zeta^{-2}\right)\right|, \tag{23}$$
$$C_E = C_J + C_P + C_b^2\,(C_f + 1),$$

every supremum being over $1 < s \le s(3)$*. Each of the four is a ratio in which the pole of* ζ *at* $s = 1$ *cancels, so none is evaluated on boxes of* s *– arb treats each occurrence of* ε *independently and the cancellation would be lost; they are computed instead from a Taylor model about* $1 + \varepsilon(3)/2$*, which makes the cancellation symbolic and covers the whole range at once. The computation certifies the aggregate inequality*

$$\sum_{j=0}^{k} \int_3^\infty \frac{u^{k-j}}{(k-j)!} |E_j(u)| \, \mathrm{d}u \le \frac{C_E \Sigma_2}{18} (k+3)^3 3^{-k}, \qquad C_E \le 5.48910. \tag{24}$$

For the near range the two groups of terms in (15) *are certified separately. With* $s(3)$ *the value of* s *at* $u = 3$*,*

$$\sum_{j=0}^{k} \int_0^3 \frac{u^{k-j}}{(k-j)!} |h_j(s(u))| J(u) \, \mathrm{d}u \le C_N 3^{-k} \quad (k \ge 59), \qquad C_N \le 367.8435, \tag{25}$$

and the two subtracted main terms satisfy

$$\sum_{j=0}^{k} \int_0^3 \frac{u^{k-j}}{(k-j)!} \big(|h_j(1)| \mathrm{e}^{-u} + |b_j| \mathrm{e}^{-2u} \big) \, \mathrm{d}u \le N_U (k+3)^3 3^{-k}, \qquad N_U \le 26.0962. \tag{26}$$

The restriction to $k \ge 59$ *in* (25) *is the range in which* (29) *is applied, and it is needed: the left-hand side exceeds* $C_N 3^{-k}$ *for* $k \le 23$*, peaking near* $k = 12$ *at some* $15329 \cdot 3^{-k}$*. That costs nothing, because monotonicity below the analytic threshold comes from the certified sweep and never appeals to* (29)*. On* $k \ge 59$ *the certified value is* $34.14 \cdot 3^{-k}$*; we carry the weaker* $C_N \le 367.8435$*, which is the bound the first group of terms satisfies for every* $k \ge 1$*, so that one constant serves both and the ledger below is unchanged.*

Finally, $\beta > 0$*, the inequality*

$$4900(62)^3 3^{-59} < \frac{\beta}{3} 2^{-59} \tag{27}$$

is certified, and direct interval evaluation gives $g_k > g_{k+1}$ *for* $1 \le k \le 75$*.*

Lemma 3.4. *Let* $A(z) = \sum_{j \ge 0} a_j z^j$*.*
(i) If $\sum_m 3^m |[z^m](1 - z/3)A(z)| \le S$*, then* $|a_j| \le S 3^{-j}$*.*
(ii) If $\sum_m 3^m |[z^m](1 - z/3)^2 A(z)| \le S$*, then* $|a_j| \le S(j+1)3^{-j}$*.*
(iii) If $\sum_m 3^m |[z^m](1 - z/3)^3 A(z)| \le S$*, then*

$$\sum_{j=0}^{k} 3^j |a_j| \le \binom{k+3}{3} S. \tag{28}$$

Proof. Write $(1 - z/3)^r A(z) = \sum_m w_m z^m$. Then

$$A(z) = \left(\sum_{q \ge 0} \binom{q + r - 1}{r - 1} 3^{-q} z^q \right) \left(\sum_{m \ge 0} w_m z^m \right).$$

For $r=1$ this gives $|a_j| \le 3^{-j}\sum_{m\le j} 3^m|w_m| \le S3^{-j}$. For $r=2$ it gives $|a_j| \le (j+1)3^{-j}S$. For $r=3$,

$$\begin{aligned}\sum_{j=0}^{k} 3^j|a_j| &\le \sum_{m=0}^{k} 3^m|w_m| \sum_{q=0}^{k-m} \binom{q+2}{2} \\ &= \sum_{m=0}^{k} 3^m|w_m| \binom{k-m+3}{3} \le \binom{k+3}{3} S.\end{aligned}$$

□

Proposition 3.5. *For every $k \ge 59$,*

$$|R_k| \le 4900(k+3)^3 3^{-k}. \tag{29}$$

The threshold is not an artefact of the method but of the near range: the coefficients h_j alternate in sign, so the near-range term has to be estimated through the majorant (32), and the resulting bound exceeds (29) for small k. Nothing is lost, since monotonicity for $k \le 75$ is certified directly and does not use (29); see the proof of Theorem 1.2.

Proof. Write R_k as the sum of the far contribution, the near contribution, and the two coefficient tails of (16)–(17). The far term and the near-range main term are cubic in k; the other two are bounded by constant multiples of 3^{-k}, and since $(k+3)^3 \ge 1$ the cubic factor may be absorbed once, at the end.

Far range $u \ge 3$. The exact error is the sum of the three terms of (15)—no asymptotic series is summed—so the certified Taylor model of (24) applies to the integrand. Its k-dependence is supplied not by that computation, which certifies finitely many numbers, but by Lemma 3.4(iii): applied to the second s-derivative of H_{o} with $S=\Sigma_2$, (28) gives $\sum_{j\le k} 3^j|\partial_s^2 h_j| \le \binom{k+3}{3}\Sigma_2 \le (k+3)^3\Sigma_2/6$, and the remaining factor 3 comes from the width of the u-range in (24). Hence $|R_{k,\mathrm{far}}| \le \frac{C_E\Sigma_2}{18}(k+3)^3 3^{-k}$, the $18 = 6\cdot 3$ recording those two steps.

First tail. By Lemma 3.4(i), $\left|\sum_{j>k} h_j(1)\right| \le \Sigma_0 \sum_{j>k} 3^{-j} = \frac{\Sigma_0}{2} 3^{-k}$.

Second tail. Put $B(z) = \partial_s H_{\mathrm{o}}(1,z) + 2\gamma H_{\mathrm{o}}(1,z)$. Multiply by $(1-z/3)^2$ and treat the two summands separately. The first contributes Σ_1 by its definition in Proposition 3.3. For the second, write $(1-z/3)^2 H_{\mathrm{o}} = (1-z/3)\cdot\big[(1-z/3)H_{\mathrm{o}}\big]$; the weighted ℓ^1 norm $\sum_m 3^m|[z^m]\cdot|$ is submultiplicative under this extra factor $(1-z/3)$, whose own norm is 2, so the second summand contributes at most $2\gamma\cdot 2\Sigma_0 = 4\gamma\Sigma_0$ — this is where the 4 comes from, rather than the 2 of the prefactor. Since $\gamma < 0.578$,

$$\sum_{m\ge 0} 3^m|[z^m](1-z/3)^2 B(z)| \le \Sigma_1 + 4\gamma\Sigma_0 < 590,$$

so $|b_j| \le 590(j+1)3^{-j}$ by Lemma 3.4(ii). Summing the resulting geometric series,

$$2^{-k-1}\sum_{j>k}|b_j|2^j \le 590\cdot 2^{-k-1}\sum_{j>k}(j+1)(2/3)^j = 590(k+4)3^{-k},$$

and $(k+4) \le \frac{5}{64}(k+3)^3$ for $k \ge 1$, with equality at $k=1$.

Near range $0 < u \le 3$. Here $|E_j|$ is at most the true integrand plus the two subtracted main terms, and the latter contribute at most $N_U(k+3)^3 3^{-k}$ by (26). For the former, (12) and the substitution $s = s(u)$, under which $u = \log\zeta(s)$ and $J(u)\,\mathrm{d}u = \mathrm{d}s$, give

$$\sum_{j=0}^{k}\int_0^3 \frac{u^{k-j}}{(k-j)!}|h_j(s(u))|J(u)\,\mathrm{d}u = \int_{s(3)}^{\infty} A_k(s)\,\mathrm{d}s, \qquad A_k(s) = \sum_{j=0}^{k}\frac{(\log\zeta(s))^{k-j}}{(k-j)!}|h_j(s)|. \tag{30}$$

It is A_k, and not the signed sum, that has to be estimated: writing $\Pi(t) = \sum_{p\geq 3} p^{-t}$, the z-linear coefficient of $\log H_{\mathrm{o}}$ in (9) is

$$h_1(s) = \Pi(s) - \log\zeta(s) = \log(1-2^{-s}) - \sum_{m\geq 2}\frac{\Pi(ms)}{m} < 0, \tag{31}$$

both terms being negative, so the h_j are not of one sign and the identity $\sum_j u^{k-j}h_j(s)/(k-j)! = P_k^{\mathrm{odd}}(s)$ bounds A_k from below rather than above.

We majorize instead. By (31) every coefficient of $\log H_{\mathrm{o}}(s,z) = h_1(s)z + \sum_{m\geq 2}\Pi(ms)z^m/m$ beyond the first is positive, so replacing h_1 by $|h_1|$ produces

$$H_{\mathrm{o}}^{+}(s,z) = H_{\mathrm{o}}(s,z)\mathrm{e}^{-2h_1(s)z},$$

a power series with non-negative coefficients dominating $|h_j|$ termwise. Hence, with $a(s) = 2|h_1(s)|$,

$$A_k(s) \leq [z^k]\big(\zeta(s)^z H_{\mathrm{o}}^{+}(s,z)\big) = \sum_{i=0}^{k} P_{k-i}^{\mathrm{odd}}(s)\frac{a(s)^i}{i!} =: P_k^{+}(s). \tag{32}$$

Two features of (32) make it effective. First, every P_j^{odd} is a Dirichlet series with positive terms and, by (31), a is a positive combination of such series; so P_k^{+} is decreasing in s, and a finite left-endpoint Riemann sum is an upper bound for $\int P_k^{+}$ outright, with no quadrature error to estimate. Second, the integral converges at a certified rate: for $s \geq s_0$ every odd n with $\Omega(n) = k-i$ satisfies $n \geq 3^{k-i}$, so $P_{k-i}^{\mathrm{odd}}(s) \leq 3^{-(k-i)(s-s_0)}P_{k-i}^{\mathrm{odd}}(s_0)$, while $a(s) \leq 2^{-(s-s_0)}a(s_0)$ by (31); since $3^{-x} \leq 2^{-x}$, each term of P_k^{+} then decays by at least $2^{-k(s-s_0)}$ and

$$\int_{s_0}^{\infty} P_k^{+}(s)\,\mathrm{d}s \leq \frac{P_k^{+}(s_0)}{k\log 2}. \tag{33}$$

Together with (25) this gives $\int_{s(3)}^{\infty} A_k(s)\,\mathrm{d}s \leq C_N 3^{-k}$ for $k \geq 59$, which is the range in which (29) is used.

Substituting the certified constants, the four contributions are

$$|R_k| \leq \Big(\underbrace{\frac{5.48910\cdot 14414}{18}}_{4395.55} + \underbrace{\frac{44.5987}{2} + \frac{5\cdot 590}{64} + C_N + N_U}_{462.33}\Big)(k+3)^3 3^{-k} < 4900(k+3)^3 3^{-k},$$

with $C_N = 367.8435$ and $N_U = 26.0962$. □

Proof of Theorem 1.2. The limit follows at once from (20) and Proposition 3.5. For monotonicity,

$$g_k - g_{k+1} = \beta 2^{-k-1} + R_k - R_{k+1}. \tag{34}$$

It is enough to prove $|R_m| < \beta 2^{-m}/3$ for $m = k, k+1$, because then

$$|R_k| + |R_{k+1}| < \frac{2\beta}{3}2^{-k-1} + \frac{\beta}{3}2^{-k-1} = \beta 2^{-k-1}.$$

Define

$$q(k) = \frac{3\cdot 4900}{\beta}(k+3)^3(2/3)^k.$$

The certified inequality (27) says $q(59) < 1$. Moreover

$$\frac{q(k+1)}{q(k)} = \frac{2}{3}\left(\frac{k+4}{k+3}\right)^3 < 1 \qquad (k \geq 4).$$

Thus $q(k) < 1$ for every $k \geq 59$, and Proposition 3.5 gives $|R_k| < \beta 2^{-k}/3$ throughout that range. Equation (34) proves monotonicity for $k \geq 59$. The certified direct computation covers $1 \leq k \leq 75$, so the two ranges cover every $k \geq 1$. They overlap on the seventeen indices $59 \leq k \leq 75$. □

4. The unrestricted sums: an exact criterion

Write

$$f_k = 1 - 2^{-k}\big(a_1 k^2 + E(k)\big), \qquad a_1 = \frac{d \log 2}{4}. \tag{35}$$

Then

$$f_{k+1} - f_k = 2^{-k-1}\left(a_1(k^2 - 2k - 1) + 2E(k) - E(k+1)\right). \tag{36}$$

Thus $f_k < f_{k+1}$ if and only if

$$a_1(k^2 - 2k - 1) + 2E(k) - E(k+1) > 0. \tag{37}$$

We now derive the decomposition of $E(k)$ used to bound the second term.

Set

$$F(s,z) = \prod_p (1 - zp^{-s})^{-1}, \qquad G_2(s,z) = (s-1)^z (1 - z2^{-s}) F(s,z),$$

and write $c_i = [z^i] G_2(1,z)$. Since $G_2(1,z) = H_{\mathrm{o}}(1,z)$, the power series has radius 3 and

$$\sum_{i \geq 0} c_i = \frac{1}{2}, \qquad \sum_{i \geq 0} 2^i c_i = d. \tag{38}$$

The coefficients are those of Section 3:

$$c_i = h_i(1). \tag{39}$$

Define the absolutely convergent signed moments

$$U = \sum_{i \geq 0} i 2^i c_i, \qquad V = \sum_{i \geq 0} i^2 2^i c_i. \tag{40}$$

Following [3, §3], put

$$I_k = \int_1^2 [z^k] F(s,z)\, \mathrm{d}s, \tag{41}$$

$$I_k' = \sum_{i+j+\ell = k} c_i \int_1^2 \frac{(-\log(s-1))^\ell}{\ell!} 2^{-js}\, \mathrm{d}s. \tag{42}$$

For $m \geq 0$ define

$$A_m = 2^m \sum_{j+\ell = m} \int_1^2 \frac{(-\log(s-1))^\ell}{\ell!} 2^{-js}\, \mathrm{d}s \tag{43}$$

and

$$\delta_m = A_m - 2^{m+1} + \frac{\log 2}{4} m^2, \qquad \Delta(k) = \sum_{i=0}^{k} 2^i c_i \delta_{k-i}. \tag{44}$$

Proposition 4.1. *Let*

$$E_{S_1}(k) = -2^k(I_k - I'_k), \tag{45}$$

$$L_k = 2\sum_{i>k} c_i, \qquad Q_k = \sum_{i>k} 2^i c_i (k-i)^2, \tag{46}$$

$$T_k = f_k - I_k, \tag{47}$$

$$E_{\exp}(k) = 2^k L_k - \frac{\log 2}{4} Q_k - 2^k T_k. \tag{48}$$

Then the identity

$$E(k) = E_{S_1}(k) + \frac{\log 2}{4}(V - 2kU) - \Delta(k) + E_{\exp}(k) \tag{49}$$

holds for every $k \geq 1$.

Proof. From (42) and (43),

$$I'_k = 2^{-k} \sum_{i=0}^{k} 2^i c_i A_{k-i}.$$

Insert the definition of δ_{k-i}:

$$I'_k = 2\sum_{i=0}^{k} c_i - \frac{\log 2}{4} 2^{-k} \sum_{i=0}^{k} 2^i c_i (k-i)^2 + 2^{-k}\Delta(k). \tag{50}$$

By (38) and (40),

$$2\sum_{i=0}^{k} c_i = 1 - L_k,$$

$$\begin{aligned} \sum_{i=0}^{k} 2^i c_i (k-i)^2 &= \sum_{i\geq 0} 2^i c_i (k-i)^2 - Q_k \\ &= dk^2 - 2kU + V - Q_k. \end{aligned}$$

Since $a_1 = \frac{\log 2}{4} d$, substitution into (50) gives

$$I'_k = 1 - 2^{-k}\left(a_1 k^2 + \frac{\log 2}{4}(V - 2kU) - \Delta(k) + 2^k L_k - \frac{\log 2}{4} Q_k \right).$$

Finally $f_k = I'_k + (I_k - I'_k) + T_k$. Comparing with (35) yields (49). □

5. Bounds for the four source terms

5.1. **The S_1 term.** Set

$$\alpha_i = \sup_{1\leq t\leq 2} |[z^i]\partial_s G_2(t,z)|, \qquad \alpha_+ = \frac{1}{2}\sum_{i\geq 0} \alpha_i 2^i.$$

The series converges, and a pole location at one value of t is not enough to see it: what is needed is a bound on the coefficients uniform in t. The first z-pole of $\partial_s G_2(t,\cdot)$ is at $z = 3^t$, so for $t \in [1,2]$ all of these functions are analytic in $|z| < 3$. Fix r with $2 < r < 3$. Then $M = \sup_{1\leq t\leq 2} \sup_{|z|=r} |\partial_s G_2(t,z)|$ is finite, the set $[1,2] \times \{|z| = r\}$ being compact and $\partial_s G_2$ continuous on it, and Cauchy's estimate on the fixed circle $|z| = r$ gives $\alpha_i \leq M r^{-i}$ uniformly in t. Hence $\sum_{i\geq 0} \alpha_i 2^i \leq M(1 - 2/r)^{-1} < \infty$.

Lemma 5.1. *For every $k \geq 1$,*

$$|E_{S_1}(k)| \leq \alpha_+(k+1), \tag{51}$$

and hence

$$|2E_{S_1}(k) - E_{S_1}(k+1)| \leq \alpha_+(3k+4). \tag{52}$$

Proof. The mean value theorem gives

$$|[z^i]G_2(s,z) - c_i| \leq (s-1)\alpha_i.$$

Subtracting (42) from the coefficient expansion for I_k and then setting $s = 1 + \mathrm{e}^{-t}$ gives

$$\begin{aligned} |I_k - I_k'| &\leq \sum_{i+j+\ell=k} \alpha_i \int_0^\infty \frac{t^\ell}{\ell!} 2^{-j(1+\mathrm{e}^{-t})} \mathrm{e}^{-2t}\,\mathrm{d}t \\ &\leq \sum_{i+j+\ell=k} \alpha_i 2^{-j} \int_0^\infty \frac{t^\ell}{\ell!} \mathrm{e}^{-2t}\,\mathrm{d}t \\ &= \sum_{i+j+\ell=k} \alpha_i 2^{-j-\ell-1}. \end{aligned}$$

Multiplying by 2^k and using $k = i + j + \ell$,

$$|E_{S_1}(k)| \leq \frac{1}{2}\sum_{i=0}^{k} \alpha_i 2^i (k-i+1) \leq \alpha_+(k+1).$$

The triangle inequality now gives

$$|2E_{S_1}(k) - E_{S_1}(k+1)| \leq 2\alpha_+(k+1) + \alpha_+(k+2) = \alpha_+(3k+4).$$

□

5.2. **The signed moment term.**

Lemma 5.2. *If $|U| \leq U_{\max}$ and $|V| \leq V_{\max}$, then for $k \geq 2$,*

$$\frac{\log 2}{4}\big(V + (2-2k)U\big) \geq -\frac{\log 2}{4}\big(V_{\max} + (2k-2)U_{\max}\big). \tag{53}$$

Proof. The coefficient of V is 1, while the coefficient $2-2k$ of U is negative. The minimum over the rectangle $[-U_{\max}, U_{\max}] \times [-V_{\max}, V_{\max}]$ is therefore attained at $U = U_{\max}$ and $V = -V_{\max}$, giving (53). □

Indeed,

$$2(V - 2kU) - \big(V - 2(k+1)U\big) = V + (2-2k)U. \tag{54}$$

Thus Lemma 5.2 is an exact optimization, not a termwise absolute-value estimate.

5.3. **The δ term.** The next lemma supplies the self-contained bound used for $\Delta(k)$.

Lemma 5.3. *For $j \geq 0$ and $u \geq 0$ put*

$$B_j(u) = 2^{-j\mathrm{e}^{-u}} - 1 + j\log 2\,\mathrm{e}^{-u}$$

and

$$\eta_i(j) = \frac{1}{i!}\int_0^\infty u^i \mathrm{e}^{-u} B_j(u)\,\mathrm{d}u.$$

Then $B_j(u) \geq 0$, $\eta_i(j) \geq 0$, and

$$\delta_m = -1 - \frac{\log 2}{4} m + \sum_{i=0}^{m} 2^i \eta_i(m-i). \tag{55}$$

Proof. With $s = 1 + \mathrm{e}^{-u}$ in (43),

$$A_m = \sum_{i=0}^{m} \frac{2^i}{i!} \int_0^\infty u^i \mathrm{e}^{-u} 2^{-(m-i)\mathrm{e}^{-u}} \, \mathrm{d}u.$$

Use $2^{-j\mathrm{e}^{-u}} = 1 - j \log 2 \, \mathrm{e}^{-u} + B_j(u)$. The constant term contributes

$$\sum_{i=0}^{m} 2^i = 2^{m+1} - 1.$$

The linear term contributes

$$\begin{aligned} -\log 2 \sum_{i=0}^{m} \frac{2^i (m-i)}{i!} \int_0^\infty u^i \mathrm{e}^{-2u} \, \mathrm{d}u &= -\frac{\log 2}{2} \sum_{i=0}^{m} (m-i) \\ &= -\frac{\log 2}{4} m(m+1). \end{aligned}$$

The remaining term is $\sum_{i=0}^{m} 2^i \eta_i(m-i)$. Subtracting 2^{m+1} and adding $(\log 2) m^2/4$ gives (55). Finally, for $x \geq 0$, $\mathrm{e}^{-x} \geq 1 - x$; applying this with $x = j \log 2 \, \mathrm{e}^{-u}$ proves $B_j(u) \geq 0$. □

Lemma 5.4. *For every $m \geq 0$,*

$$|\delta_m| \leq D(m), \qquad D(m) = (\log 2) m \log(m+1) + 1. \tag{56}$$

Proof. For the lower bound, (55) gives $\delta_m \geq -1 - \frac{\log 2}{4} m \geq -D(m)$ for $m \geq 1$, because $\log(m+1) \geq \log 2 > 1/4$; and $\delta_0 = -1 = -D(0)$.

For the upper bound put $\Lambda_m = \sum_{i=0}^{m} 2^i \eta_i(m-i)$. Since $\mathrm{e}^{-x} - 1 + x \leq \min(x, x^2/2)$ for $x \geq 0$, taking $x = (m-i) \log 2 \, \mathrm{e}^{-u}$ gives

$$\Lambda_m \leq \int_0^\infty \mathrm{e}^{-u} \sum_{i=0}^{m} \frac{(2u)^i}{i!} \min\left((m-i) \log 2 \, \mathrm{e}^{-u}, \frac{(m-i)^2 (\log 2)^2 \mathrm{e}^{-2u}}{2} \right) \mathrm{d}u. \tag{57}$$

Let $v_m = \max\left(0, \log \frac{m+1}{3}\right)$. On $0 \leq u \leq v_m$ use the linear bound together with $\sum_{i=0}^{m} (m-i)(2u)^i/i! \leq m\mathrm{e}^{2u}$, and on $u \geq v_m$ the quadratic bound together with $\sum_{i=0}^{m} (m-i)^2 (2u)^i/i! \leq m^2 \mathrm{e}^{2u}$. Then

$$\Lambda_m \leq m \log 2 \, v_m + \frac{m^2 (\log 2)^2}{2} \mathrm{e}^{-v_m}. \tag{58}$$

For $m \geq 2$ we have $v_m = \log \frac{m+1}{3}$ and $\mathrm{e}^{-v_m} = \frac{3}{m+1}$, so (58) and $\frac{3m^2}{2(m+1)} \leq \frac{3}{2} m$ give

$$\begin{aligned} \Lambda_m &\leq m \log 2 \log \frac{m+1}{3} + \frac{3m^2 (\log 2)^2}{2(m+1)} \\ &\leq m \log 2 \left(\log(m+1) - \log 3 + \tfrac{3}{2} \log 2 \right) \leq m \log 2 \log(m+1), \end{aligned}$$

the last step because $3 > 2^{3/2}$. For $m \leq 1$ we have $v_m = 0$, and (58) reads $\Lambda_m \leq m^2 (\log 2)^2/2$, which is at most $m \log 2 \log(m+1)$ since $\frac{m \log 2}{2} \leq \log(m+1)$ there. In both cases

$$\delta_m = -1 - \frac{\log 2}{4} m + \Lambda_m \leq \Lambda_m \leq (\log 2) m \log(m+1) < D(m),$$

which with the lower bound proves (56). □

Lemma 5.5. *If $S^* = \sum_{i\ge 0} 2^i|c_i|$, then, with D as in Lemma 5.4,*

$$|2\Delta(k) - \Delta(k+1)| \le S^*\big(2D(k) + D(k+1)\big). \tag{59}$$

Proof. Since D is increasing,

$$|\Delta(k)| \le \sum_{i=0}^{k} 2^i|c_i|D(k-i) \le S^*D(k).$$

Apply the triangle inequality to $2\Delta(k) - \Delta(k+1)$. □

5.4. **The exponentially small term.** The three terms in (48) are respectively a coefficient tail at $z = 1$, a weighted coefficient tail at $z = 2$, and the integral over $s \ge 2$. The following is the exact numerical input used for them.

Proposition 5.6 (Certified unrestricted input)**.** *The interval computations give*

$$\begin{aligned} d \ge 0.3786949, \qquad S^* \le 6.9764, \qquad \alpha_+ \le 15.425661, \\ |U| \le U_{\max} = 0.08436, \qquad |V| \le V_{\max} = 1.907462. \end{aligned} \tag{60}$$

For every $k \ge 900$,

$$|E_{\exp}(k)| \le \frac{1.45}{k} + 10^{-3}. \tag{61}$$

The direct certificates establish $f_k < f_{k+1}$ on

$$6 \le k \le 199, \qquad 78 \le k \le 899, \qquad 900 \le k \le 2439, \tag{62}$$

three separate runs whose union is the single range $6 \le k \le 2439$. The quadrature behind these values, and the six error sources entering each enclosure, are described in Appendix A.

The coefficient tails in (61) are obtained only after the pole at $z = 3$ has been removed. Put

$$\widetilde{G}_2(z) = (1 - z/3)G_2(1, z) = \sum_{m\ge 0} w_m z^m.$$

The certified contour bound is $\sup_{|z|=4.2}|\widetilde{G}_2(z)| \le 101.245$. Hence $|w_m| \le 101.245\, 4.2^{-m}$, and from $G_2(1, z) = (1 - z/3)^{-1}\widetilde{G}_2(z)$ we obtain

$$|c_i| \le 101.245\, 3^{-i} \sum_{m=0}^{i} (3/4.2)^m < 354.36\, 3^{-i}. \tag{63}$$

By (39) the c_i are the coefficients $h_i(1)$, so Lemma 3.4(i) sharpens (63) to $|c_i| \le \Sigma_0 3^{-i}$, dispensing with the contour bound; the weaker form suffices below. With $C = 354.36$,

$$\begin{aligned} 2^k|L_k| &\le C(2/3)^k, \\ \frac{\log 2}{4}|Q_k| &\le \frac{\log 2}{4} C(2/3)^k \sum_{r\ge 1} r^2(2/3)^r = \frac{30C\log 2}{4}(2/3)^k. \end{aligned}$$

Both terms are less than 10^{-3} in total for $k \ge 900$. The remaining certified tail inequality is $2^k|T_k| \le 1.45/k$. Substitution into (48) proves (61); no asymptotic O-term enters the final criterion. The first two terms are in fact below 10^{-3} from $k \ge 37$ on, so the threshold 900 comes from T_k alone.

6. Assembly of the unrestricted theorem

For $k \geq 900$ define

$$R(k) = \frac{\log 2}{4}\big((2k-2)U_{\max} + 2V_{\max} + 4U_{\max}\big) + S^*\big(2D(k) + D(k+1)\big) + \alpha_+(3k+4) + 3\left(\frac{1.45}{k} + 10^{-3}\right), \tag{64}$$

and

$$\varphi(k) = a_1(k^2 - 2k - 1) - R(k). \tag{65}$$

Proposition 6.1. *For $k \geq 900$,*

$$2E(k) - E(k+1) \geq -R(k). \tag{66}$$

Consequently $\varphi(k) > 0$ implies $f_k < f_{k+1}$.

Proof. Apply (49) at k and $k+1$. The S_1 contribution is bounded by (52). The signed moment contribution is

$$\frac{\log 2}{4}\big(V + (2-2k)U\big),$$

which is bounded below by (53). The first line of (64) is slightly larger than the resulting quantity, since

$$V_{\max} + (2k-2)U_{\max} \leq (2k-2)U_{\max} + 2V_{\max} + 4U_{\max}.$$

The Δ contribution is bounded by (59). Finally, (61) gives

$$|2E_{\exp}(k) - E_{\exp}(k+1)| \leq 2\left(\frac{1.45}{k} + 10^{-3}\right) + \left(\frac{1.45}{k+1} + 10^{-3}\right) \leq 3\left(\frac{1.45}{k} + 10^{-3}\right).$$

Adding the four lower bounds proves (66). Equation (36) then gives

$$f_{k+1} - f_k \geq 2^{-k-1}\varphi(k),$$

which proves the last assertion. □

Lemma 6.2. *With the certified constants in Proposition 5.6, $\varphi(2440) > 1155$.*

Proof. Substitution in (65) gives the following certified one-sided bounds:

$$\begin{aligned} a_1(2440^2 - 2\cdot 2440 - 1) &> 390371.6, \\ \frac{\log 2}{4}\big(4878U_{\max} + 2V_{\max} + 4U_{\max}\big) &< 72.1, \\ S^*\big(2D(2440) + D(2441)\big) &< 276166.8, \\ \alpha_+(3\cdot 2440 + 4) &< 112977.6, \\ 3\left(\frac{1.45}{2440} + 10^{-3}\right) &< 0.005. \end{aligned}$$

Therefore

$$\varphi(2440) > 390371.6 - 72.1 - 276166.8 - 112977.6 - 0.005 > 1155.$$

The same substitution gives $\varphi(2432) < 0 < \varphi(2433)$, so the criterion gives $\varphi(2433) > 135$, so the criterion is already in force at $k = 2433$. Direct evaluation puts the sign change just below that, at $\varphi(2432) < 0 < \varphi(2433)$; we do not rely on it, since certifying $\varphi(2432) < 0$ would need a *lower* bound on $R(2432)$, whereas every constant in (60) is one-sided the other way. Nothing below requires the first index: we carry the threshold at 2440, which is one past the end of (62), so that the sweep and the criterion meet without a gap. □

Lemma 6.3. *The function $k \mapsto \varphi(k)$ is strictly increasing for integers $k \geq 2440$.*

Proof. First,

$$a_1\big((k+1)^2 - 2(k+1) - 1 - (k^2 - 2k - 1)\big) = a_1(2k-1).$$

Moreover

$$\begin{aligned} D(m+1) - D(m) &= \log 2 \left(\log(m+2) + m \log\left(1 + \frac{1}{m+1}\right)\right) \\ &\leq \log 2\big(\log(m+2) + 1\big). \end{aligned}$$

The last term in (64) decreases with k. Hence

$$\begin{aligned} R(k+1) - R(k) \leq \frac{\log 2}{2} U_{\max} + 3\alpha_+ \\ + S^* \log 2\big(2\log(k+2) + \log(k+3) + 3\big). \end{aligned} \tag{67}$$

Define, for real $x \geq 2440$,

$$\begin{aligned} M(x) = a_1(2x-1) - \frac{\log 2}{2} U_{\max} - 3\alpha_+ \\ - S^* \log 2\big(2\log(x+2) + \log(x+3) + 3\big). \end{aligned}$$

Then

$$M'(x) = 2a_1 - S^* \log 2 \left(\frac{2}{x+2} + \frac{1}{x+3} \right) > 0,$$

because the certified lower bound for d gives $2a_1 > 0.1312$, whereas the subtracted term is < 0.006 for $x \geq 2440$. Direct substitution gives $M(2440) > 146$. Thus $M(k) > 0$ for every $k \geq 2440$. Together with (67), this yields $\varphi(k+1) - \varphi(k) > 0$. □

Proof of Theorem 1.1. By Lemmas 6.2 and 6.3, $\varphi(k) > 0$ for every $k \geq 2440$. Proposition 6.1 therefore gives $f_k < f_{k+1}$ on that range. The certified ranges in (62) cover every integer from 6 through 2439. Together the analytic and certified ranges cover all $k \geq 6$. □

7. A general principle

Conjecture 7.1. Let $\mathcal{P}$ be a set of primes with least element q and thin complement,

$$\sum_{p \notin \mathcal{P}} \frac{\log p}{p} < \infty, \tag{68}$$

and let $f_k^{\mathcal{P}}$ be the sum of $1/(n \log n)$ over integers n with $\Omega(n) = k$ and all prime factors in $\mathcal{P}$. The sequence $f_k^{\mathcal{P}}$ is eventually increasing if and only if $q = 2$.

Some hypothesis of this kind is needed. For $\mathcal{P} = \{2\}$ one has $f_k^{\mathcal{P}} = 1/(2^k k \log 2)$, and for $\mathcal{P} = \{2, 3\}$ the sequence again decreases to 0; in both cases $q = 2$ and the conclusion fails. What goes wrong is that a sparse $\mathcal{P}$ leaves $\sum_{p \in \mathcal{P}} p^{-s}$ bounded as $s \to 1^+$, so the generating function has no $\zeta(s)^z$ factor and $f_k^{\mathcal{P}}$ decays geometrically to 0 whatever q is.

Under (68) that cannot happen, and the singularity structure is the one the two theorems above exploit:

$$\prod_{p \in \mathcal{P}} (1 - zp^{-s})^{-1} = \prod_{p} (1 - zp^{-s})^{-1} \prod_{p \notin \mathcal{P}} (1 - zp^{-s}),$$

where the second product converges locally uniformly by (68) and is therefore entire in z. Its zeros cancel exactly the poles at $z = p^s$ with $p \notin \mathcal{P}$, leaving the first z-singularity at $z = q^s$. At $s = 1$ the point $z = 2$ lies on that singular circle exactly when $q = 2$. If $q \geq 3$ the coefficient

extraction at $z = 2$ is interior and gives a constant multiple of 2^{-k}, as in Section 3; if $q = 2$ it takes place at the boundary and may produce the factor k^2.

Three points remain to turn this mechanism into a theorem for arbitrary $\mathcal{P}$ satisfying (68). First, the regularized product must provide enough uniform control to identify the limiting constant and the coefficient at $z = 2$. Second, in the boundary case one must rule out cancellation of the k^2 coefficient. Third, in the interior case one must show that the constant multiplying 2^{-k} is positive, which is what makes the sequence decrease rather than increase; for $\mathcal{P} = \{p : p \geq 3\}$ this is the inequality $\beta > 0$ of (5). The present paper proves the two extremal cases $\mathcal{P} = \{\text{all primes}\}$ and $\mathcal{P} = \{p : p \geq 3\}$.

8. Certification and formalization

What the formalization establishes is an *implication*. The certified quantities enter as fields of five hypothesis structures — `NumericalFacts`, `NumericalFactsFull`, `OddNumericalFacts`, `OddCoeffFacts`, `OddDomFacts` — and no instance of any of them is constructed. Lean therefore checks that those hypotheses imply the theorems, and nothing in the kernel checks that the hypotheses are true, or even that they are mutually consistent: an unsatisfiable field would make the theorems assuming it vacuous rather than false, and would not be detected. That the imported intervals agree with what Arb computed is verified outside Lean, by the script `verify.py` against the certificate. The reader should take the Lean development as discharging the reductions below and nothing else; the trusted base is the structure fields, and it is listed in Table 1.

With that said, the division is as follows. Formalized in Lean 4 without `sorry` are Lemmas 2.1 and 2.3, the coefficient identities (13) and (49), the assembly of the ranges in both theorems, and the lemmas named in the rest of this section.

Two items a reader would expect to find among them are not there, and we state the position precisely. First, Proposition 2.2: a radius-specific consequence of it is proved, on the ball $|z| < 3$ for $s \geq 1$, but its operative content (10) — that $\mathrm{d}/\mathrm{d}s$ commutes with $[z^j]$ — is not, and the s-differentiability of the coefficient functions is itself a hypothesis, as field `g_deriv` of `NumericalFactsFull` in the full case and as `deriv_exists`, `deriv2_exists` of `OddCoeffFacts` in the odd case. No interval computation can establish a differentiability claim, and none is attempted. Second, Proposition 3.5: its statement is imported as field `remainder` of `OddNumericalFacts` rather than derived from Proposition 3.3; the derivation is the proof given in Section 3 above, which has not been formalized. The Lean development contains the intended route to it, but that route still rests on assumed coefficient and dominating-sequence data of its own, so it does not discharge the hypothesis.

Likewise the source estimates of Section 5 are assumed, not proved, by the theorem named below: `NumericalFacts` carries $|2E(k) - E(k+1)| \leq R(k)$ as the single field `source_bound`, which is the aggregate of that whole section. Only the second formal version discussed below derives it, from the four source terms separately.

The numerical propositions are Proposition 3.3 and Proposition 5.6. Their statements are finite systems of interval inequalities, established by Arb/FLINT through its Python bindings. The Lean formalization imports the resulting intervals as hypotheses — together with Proposition 3.5 and the differentiability noted above, neither of which is an interval inequality — and proves that those hypotheses imply Theorem 1.1 and both assertions of Theorem 1.2; the three formal statements are `erdos_sums_increase_from_six`, `fOdd_decreasing` and `fOdd_tendsto_half`. The differentiability field belongs to `NumericalFactsFull` and so is a hypothesis of the second, primitive version of the full theorem rather than of `erdos_sums_increase_from_six`, which does not mention it. The limit $g_k \to \frac{1}{2}$ was formerly left unformalized, the monotonicity

argument making no use of it; it is now proved there as well, and from the same hypothesis, since Proposition 3.5 already bounds $|g_k - \frac{1}{2} - \beta 2^{-k}|$ by a multiple of $(k+3)^3 3^{-k}$, and both that and 2^{-k} vanish.

For Theorem 1.1 the hypotheses are not only the interval bounds of Proposition 5.6. Proposition 6.1 enters as a hypothesis in one of the two formal versions, and is proved in the other, from the primitive certified quantities rather than from the aggregate bound (66). (The bound (61) is not what the primitive route replaces; it remains a hypothesis, `NumericalFactsFull.Eexp_bound`, in both versions.) What the formalization proves from these is the passage to $f_k < f_{k+1}$ for $k \geq 2440$, together with the assembly of the four ranges.

Lemmas 6.2 and 6.3 were formerly hypotheses of that development as well. They are now proved in it, as `phiBound_pos_at_2440` and `phiBound_mono_of_bounds`, from upper bounds on the parameters alone: φ is an explicit expression in k, so neither lemma is a statement about primes, contours or limits, and what the formalization had been assuming was arithmetic. Proving them needed one quantity the certified hypotheses did not carry, the bound $|U| \leq U_{\max}$ of (60), which is now among them. The sharpness required is not uniform: the base case turns on $\log 2441$ to a tenth of a percent, where $\log 2441 \leq 12 \log 2$ is already too weak, while the monotonicity step needs only $\log 2441 \leq 12 \log 2$ itself.

For the odd theorem the lower bound on γ was likewise a hypothesis and is now proved, as `gamma_lower_bound`, from $\mathrm{harmonic}(n) - \log(n+1) < \gamma$ at $n = 8191$, chosen so that $n+1 = 2^{13}$ and the logarithm is $13 \log 2$ exactly.

The constants d and ρ were hypotheses of both developments and are now proved in them, as `dConst_lower_bound`, `dConst_lower_bound_sharp` and `rhoConst_upper_bound`. What makes this possible is that the Euler factors are rational and exceed 1,

$$\Big(1 - \tfrac{2}{p}\Big)^{-1}\Big(1 - \tfrac{1}{p}\Big)^{2} = \frac{(p-1)^2}{p(p-2)} = 1 + \frac{1}{p(p-2)},$$
$$\Big(1 - \tfrac{3}{p}\Big)^{-1}\Big(1 - \tfrac{1}{p}\Big)^{3} = \frac{(p-1)^3}{p^2(p-3)} = 1 + \frac{3p-1}{p^2(p-3)},$$

the second because $(p-1)^3 - p^2(p-3) = 3p-1$. The constant regularized by the second display is

$$\rho = -\frac{1}{9} \prod_{p \geq 5} \Big(1 - \frac{3}{p}\Big)^{-1}\Big(1 - \frac{1}{p}\Big)^{3}, \tag{69}$$

which is recorded here only so that every constant proved in the formalization has a counterpart in the text; it is used nowhere in the proofs above. A product of factors each at least 1 is bounded below by the product over any finite subset of them, with no tail estimate entering, so a one-sided bound on such a constant is a finite computation in the kernel even where the two-sided enclosure is not. Only one-sided bounds are used: d enters through $a_1 = d \log 2/4$ and through $\beta = d(\log 2 - S + \gamma)$, in both cases from below, and ρ is not used at all. The bounds proved are weaker than those of Proposition 5.6, which are unchanged. The 94 primes $p \leq 499$ give $d \geq 0.3785913$, which suffices for $a_1 \geq 0.0656$; the 668 primes $p \leq 4999$ give $d \geq 0.3786823$, which is what $\beta \geq 0.1556$ requires; and the 93 primes $5 \leq p \leq 499$ give $\rho \leq -0.1747885$. In each case the requirement is computed against the bound for $\log 2$ available in the formalization rather than against $\log 2$ itself, the two differing by enough to change how many primes are needed.

The same device does not reach S. There the bound required is an upper one on a sum of positive terms, so the tail cannot be discarded, and the summands carry $\log p$ rather than being rational. The budget is narrow: with d at the bound just proved, $\beta \geq 0.1556$ permits

$S \le 0.8593986$, while the partial sum over $p < 2\cdot 10^6$ already reaches 0.8593917, so the admissible tail is below $7 \cdot 10^{-6}$ and primes beyond about $1.5 \cdot 10^5$ are needed. Accordingly S remains a certified hypothesis, as do the sweeps and the Cauchy estimates of Propositions 3.3 and 5.6.

Two constants are stated above more conservatively than the computation requires, so that the text and the formalization carry the same values: the threshold 2440 of Lemma 6.2 in place of 2433, and (63) in place of the sharper bound that Lemma 3.4(i) supplies. Neither changes the ranges covered. Two further discrepancies should be recorded rather than left for the reader to find. The certified odd sweep is stated above for $1 \le k \le 75$, whereas the field `OddNumericalFacts.sweep_low` assumes only $1 \le k \le 58$; since the analytic branch begins at $k = 59$ that is all the assembly needs, and assuming less is the safe direction, but the two ranges are not the same. And the field `NumericalFacts.Gamma_bound`, carrying $\Gamma \le 104.52$, has no counterpart above and no consumer in the development either; it is retained only so that the certified package and the text can be compared field by field.

TABLE 1. What the Lean development proves and what it assumes. The assumed rows are the trusted base: no instance of any structure is constructed, so these are not checked in the kernel.

Statement	Status	Lean name
Theorem 1.1	proved	`erdos_sums_increase_from_six`
Theorem 1.2, monotonicity	proved	`fOdd_decreasing`
Theorem 1.2, limit	proved	`fOdd_tendsto_half`
Lemma 6.2	proved	`phiBound_pos_at_2440`
Lemma 6.3	proved	`phiBound_mono_of_bounds`
γ from below	proved	`gamma_lower_bound`
d from below	proved	`dConst_lower_bound`, `_sharp`
ρ from above	proved	`rhoConst_upper_bound`
Proposition 3.5	assumed	`OddNumericalFacts.remainder`
Section 5 aggregate	assumed	`NumericalFacts.source_bound`
(61)	assumed	`NumericalFactsFull.Eexp_bound`
g_i differentiable on $(1,2)$	assumed	`NumericalFactsFull.g_deriv`
∂_s, ∂_s^2 of coefficients exist	assumed	`OddCoeffFacts.deriv_exists`, `deriv2_exists`
S from above	assumed	`NumericalFacts.Sstar_bound`
Sweeps	assumed	`OddNumericalFacts.sweep_low` and companions
Σ_i, C_E, C_N, N_U, α_+, U, V	assumed	fields of the five structures

For the odd theorem, the coefficient bounds in (21) are obtained by cancelling the pole at $z = 3$ before applying coefficient recursion or Cauchy bounds. The range $0 < u \le 3$ is evaluated directly, while the range $u \ge 3$ uses the Taylor model made explicit in Proposition 3.1.

The constant C_E is certified on the whole of $(1, s(3)]$. The four quantities entering it are ratios in which the pole of ζ at $s = 1$ cancels. Interval evaluation on boxes of s loses that cancellation, since each occurrence of $s - 1$ is widened independently, and the enclosures diverge however finely $(1, s(3)]$ is partitioned. Expanding ζ as a power series in $\sigma = s - s_0$ about $s_0 = 1 + E/2$, where $E = s(3) - 1$, makes the cancellation symbolic. The resulting series are evaluated on $|\sigma| \le E/2$, which is the whole range; their coefficients fall off geometrically, and truncating after nine terms contributes less than 10^{-20}.

The constant C_N requires $\Pi(t) = \sum_{p\ge3} p^{-t}$ at $t = s, 2s, \dots, 200s$ for $s = s(3)$. At $s \approx 1.05$ these converge too slowly to truncate: the Rosser–Schoenfeld bound applied beyond 10^8 still overestimates the tail twofold, and that excess compounds through the coefficient recursion. Instead Π is evaluated by the Möbius series $\Pi(t) = \sum_{n\ge1} \mu(n)n^{-1} \log\zeta(nt) - 2^{-t}$, which requires no prime enumeration. The recursion itself is run at 600 bits, since cancellation costs about one digit every 2.3 steps over the 200 terms used; at 1200 bits the head agrees to every printed digit.

For the unrestricted theorem, the s-uniform coefficient bounds defining α_+ are obtained by carrying s as an interval over a partition of $[1, 2]$. The signed moments U and V are evaluated before absolute values are taken. This order is essential: replacing them by $\sum_i 2^i|c_i|$ loses the cancellation used in (54).

Appendix A. Certified quadrature

Since $1/(n\log n) = \int_1^\infty n^{-s}\,\mathrm{d}s$, we have $f_k = \int_1^\infty P_k(s)\,\mathrm{d}s$, and the substitution $s = 1 + \mathrm{e}^{-u}$ turns this into

$$f_k = \int_{-\infty}^{\infty} F(u)\,\mathrm{d}u, \qquad F(u) = P_k(1+\mathrm{e}^{-u})\mathrm{e}^{-u}, \tag{70}$$

the integrand being analytic on the strip $|\mathrm{Im}\,u| < \pi/2$. The integral is split into a truncated range $[u_-, u_+]$ with $u_- = -5.10$ and $u_+ = 3.5k_{\max}$, taken once for the largest k of the run and so uniform in k, and the two tails beyond it. The truncated range is divided into panels of width $H = 0.8$, each mapped affinely to $[-1, 1]$ and integrated by Gauss–Legendre quadrature with $N = 52$ nodes at 1400 bits of precision; the number of panels therefore grows linearly with $k_{\max}$.

Four of the six error sources in the certificate are arithmetic: ball arithmetic throughout, the explicit truncation balls in the Möbius series for the prime zeta function, the discretization bound derived below, and interval-Newton enclosures for the Gauss nodes and weights. The remaining two are the tails of (70), which are bounded as follows and are not negligible by inspection.

For the left tail, $u \le u_-$ corresponds to $s \ge s_- = 1 + \mathrm{e}^{5.10} = 165.02$, where $P(s) \le 2.78 \cdot 2^{-s}$ and hence $P_k(s) \le 2.78^k 2^{-ks}$, so

$$\int_{-\infty}^{u_-} |F|\,\mathrm{d}u = \int_{s_-}^{\infty} P_k(s)\,\mathrm{d}s \le \frac{2.78^k 2^{-ks_-}}{k\log 2}. \tag{71}$$

For the right tail, $P_k(s) \le (\log\zeta(s))^k/k!$ for $s > 1$ [2, Lemma 5.6], so with $u = \log\zeta(s)$ the integral is dominated by $\int_{u_+}^\infty u^k\mathrm{e}^{-u}\,\mathrm{d}u/k!$; the resulting incomplete-Gamma integral obeys $\Gamma(k+1, u_+)/k! \le 2u_+^k\mathrm{e}^{-u_+}/k!$ provided $u_+ \ge 2k+2$, so

$$\int_{u_+}^{\infty} |F|\,\mathrm{d}u \le \frac{2u_+^k\mathrm{e}^{-u_+}}{k!}. \tag{72}$$

The hypothesis $u_+ \ge 2k+2$ holds throughout, since $u_+ = 3.5k_{\max}$ and $k \le k_{\max}$. Both bounds are added to every reported enclosure. They are largest at $k = k_{\max}$, and even there they are far below the differences $f_{k+1} - f_k \asymp a_1 k^2 2^{-k}$ that the sweep has to resolve: for $k_{\max} = 199$ the right-hand side of (72) is below 10^{-100} while the smallest difference in range exceeds 10^{-57}.

For f analytic inside the Bernstein ellipse E_ρ and bounded there by M, the N-node Gauss rule satisfies

$$\left|\int_{-1}^{1} f(x)\,\mathrm{d}x - \widehat{I}_N\right| \le \frac{64}{15}\,\frac{M\rho^{-2N}}{1-\rho^{-2}} \tag{73}$$

[5, Thm. 4.5], where the rule is indexed so that I_n uses $n+1$ nodes and the stated exponent $\rho_B^{-(2n+2)}$ is the ρ_B^{-2N} above; the bound appears as [6, Thm. 19.3]. Here ρ_B is the Bernstein-ellipse parameter, unrelated to the arithmetic constant ρ of Section 8. The implementation evaluates the right side of (73) with the interval-scaling factor $H/2$ and a further factor 10 of margin. The ellipse is taken with semi-minor axis $\delta = 1.45 < \pi/2$ in the u variable, giving $\rho_B \approx 7.385$, and M is bounded using $|P_k(s)| \le P_k(\operatorname{Re} s)$ together with the monotonicity of P_k on the reals.

The per-panel bound is not monotone in the panel's position, and it would be wrong to locate its maximum by inspection: on the right-hand side $F(u) \approx u^k \mathrm{e}^{-u}/k!$ by (70), which increases until $u \approx k$ and only then decays, so M rises across the bulk of the range before falling. What is used is therefore the maximum over all panels and all $k \le k_{\max}$, which the certificate records; it is 1.1×10^{-87}, and so is negligible beside the widths actually reported. It is also, at $k_{\max} = 199$, some twenty orders of magnitude larger than either tail (71)–(72), so the discretization bound and not the truncation is what limits the accuracy of the f_k.

The pinned coefficient sweeps use primes up to 10^5 and 300 bits of precision, with 1600 boxes in s for α_+ and 800 boxes for the second-derivative sweep. The certificate records these parameters together with a SHA-256 hash of the numerical payload.

Acknowledgements

I thank Jared Duker Lichtman for his guidance and for many helpful discussions throughout this work.

Statement on the use of AI

PLACEHOLDER — see the TODO above.

References

[1] W. D. Banks and G. Martin, *Optimal primitive sets with restricted primes*, Integers **13** (2013), #A69, 10 pp.; arXiv:1301.0948.
[2] J. D. Lichtman, *Almost primes and the Banks–Martin conjecture*, J. Number Theory **211** (2020), 513–529; arXiv:1909.00804.
[3] O. Gorodetsky, J. D. Lichtman and M. W. Wong, *On Erdős sums of almost primes*, C. R. Math. Acad. Sci. Paris **362** (2024), no. G12, 1571–1596; arXiv:2303.08277.
[4] P. Erdős, *Note on sequences of integers no one of which is divisible by any other*, J. London Math. Soc. **10** (1935), no. 2, 126–128.
[5] L. N. Trefethen, *Is Gauss quadrature better than Clenshaw–Curtis?*, SIAM Rev. **50** (2008), 67–87.
[6] L. N. Trefethen, *Approximation Theory and Approximation Practice*, SIAM, Philadelphia, 2013.
[7] Z. Zhang, *On a problem of Erdős concerning primitive sequences*, Math. Comp. **60** (1993), no. 202, 827–834.
[8] J. Bayless, P. Kinlaw and D. Klyve, *Sums over primitive sets with a fixed number of prime factors*, Math. Comp. **88** (2019), 3063–3077.

Department of Computer Science, Purdue University, West Lafayette, IN 47907, USA
Email address: jpadhi@purdue.edu