

Hyperbolicity in Complex Analysis

Jaikrishnan Janardhanan
Indian Institute of Technology Madras

`jaikrishnan@iitm.ac.in`

Talk at IIT Madras, August 23, 2018

Some famous results

- 1 **Liouville's Theorem (1847):** Any bounded entire function is constant.

Some famous results

- ① **Liouville's Theorem (1847):** Any bounded entire function is constant.
- ② **The Schwarz Lemma (Schwarz, 1869 and Carathéodory, 1912):** If $f : \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic with $f(0) = 0$ then $|f(z)| \leq |z|$ and $|f'(0)| \leq 1$.

Some famous results

- ① **Liouville's Theorem (1847):** Any bounded entire function is constant.
- ② **The Schwarz Lemma (Schwarz, 1869 and Carathéodory, 1912):** If $f : \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic with $f(0) = 0$ then $|f(z)| \leq |z|$ and $|f'(0)| \leq 1$.
- ③ **The big Picard Theorem (1879):** If $f : \mathbb{D}^* \rightarrow \mathbb{C} \setminus \{0, 1\}$ is holomorphic then 0 *cannot* be an essential singularity for f .

Some famous results

- ➊ **Liouville's Theorem (1847):** Any bounded entire function is constant.
- ➋ **The Schwarz Lemma (Schwarz, 1869 and Carathéodory, 1912):** If $f : \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic with $f(0) = 0$ then $|f(z)| \leq |z|$ and $|f'(0)| \leq 1$.
- ➌ **The big Picard Theorem (1879):** If $f : \mathbb{D}^* \rightarrow \mathbb{C} \setminus \{0, 1\}$ is holomorphic then 0 *cannot* be an essential singularity for f .
- ➍ **Montel's Theorem (1910):** The collection of all uniformly bounded holomorphic mappings from a domain in \mathbb{C} is a normal family.

Some famous results

- ➊ **Liouville's Theorem (1847):** Any bounded entire function is constant.
- ➋ **The Schwarz Lemma (Schwarz, 1869 and Carathéodory, 1912):** If $f : \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic with $f(0) = 0$ then $|f(z)| \leq |z|$ and $|f'(0)| \leq 1$.
- ➌ **The big Picard Theorem (1879):** If $f : \mathbb{D}^* \rightarrow \mathbb{C} \setminus \{0, 1\}$ is holomorphic then 0 *cannot* be an essential singularity for f .
- ➍ **Montel's Theorem (1910):** The collection of all uniformly bounded holomorphic mappings from a domain in \mathbb{C} is a normal family.
- ➎ **Montel's Theorem 2nd version (1910):** The collection of all holomorphic functions from a domain into $\mathbb{C} \setminus \{0, 1\}$ is a normal family.

Some famous results

- ➊ **Liouville's Theorem (1847):** Any bounded entire function is constant.
- ➋ **The Schwarz Lemma (Schwarz, 1869 and Carathéodory, 1912):** If $f : \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic with $f(0) = 0$ then $|f(z)| \leq |z|$ and $|f'(0)| \leq 1$.
- ➌ **The big Picard Theorem (1879):** If $f : \mathbb{D}^* \rightarrow \mathbb{C} \setminus \{0, 1\}$ is holomorphic then 0 *cannot* be an essential singularity for f .
- ➍ **Montel's Theorem (1910):** The collection of all uniformly bounded holomorphic mappings from a domain in \mathbb{C} is a normal family.
- ➎ **Montel's Theorem 2nd version (1910):** The collection of all holomorphic functions from a domain into $\mathbb{C} \setminus \{0, 1\}$ is a normal family.
- ➏ **The de Franchis Theorem (1913):** Let R and S be compact Riemann surfaces of genus higher than 1. Then there are only finitely many non-constant holomorphic mappings from R into S .

Conformal Metrics

Conformal Metric

Let $D \subset \mathbb{C}$ be a domain. A *conformal metric* on D is a C^2 -smooth function $\rho : D \rightarrow \mathbb{R}^+ \cup \{0\}$. We also allow ρ to have only a discrete set of zeroes at which points ρ is not required to be differentiable. If $z \in D$ and $v \in \mathbb{C}$ is a vector, we define the *length of v at z* to be

$$\|v\|_{\rho,z} := \rho(z) \cdot |v|.$$

Conformal Metrics

Conformal Metric

Let $D \subset \mathbb{C}$ be a domain. A *conformal metric* on D is a C^2 -smooth function $\rho : D \rightarrow \mathbb{R}^+ \cup \{0\}$. We also allow ρ to have only a discrete set of zeroes at which points ρ is not required to be differentiable. If $z \in D$ and $v \in \mathbb{C}$ is a vector, we define the *length of v at z* to be

$$\|v\|_{\rho,z} := \rho(z) \cdot |v|.$$

Recall that one can identify the tangent space of D at any point with \mathbb{C} itself. A conformal metric provides for us an inner product at each point of D in the following way:

Conformal Metrics

Conformal Metric

Let $D \subset \mathbb{C}$ be a domain. A *conformal metric* on D is a C^2 -smooth function $\rho : D \rightarrow \mathbb{R}^+ \cup \{0\}$. We also allow ρ to have only a discrete set of zeroes at which points ρ is not required to be differentiable. If $z \in D$ and $v \in \mathbb{C}$ is a vector, we define the *length of v at z* to be

$$\|v\|_{\rho,z} := \rho(z) \cdot |v|.$$

Recall that one can identify the tangent space of D at any point with \mathbb{C} itself. A conformal metric provides for us an inner product at each point of D in the following way: if $v, w \in T_z \mathbb{D}$, then

$$\langle v, w \rangle_z := \rho^2(z) v \cdot \overline{w}.$$

Conformal Metrics

Conformal Metric

Let $D \subset \mathbb{C}$ be a domain. A *conformal metric* on D is a C^2 -smooth function $\rho : D \rightarrow \mathbb{R}^+ \cup \{0\}$. We also allow ρ to have only a discrete set of zeroes at which points ρ is not required to be differentiable. If $z \in D$ and $v \in \mathbb{C}$ is a vector, we define the *length of v at z* to be

$$\|v\|_{\rho,z} := \rho(z) \cdot |v|.$$

Recall that one can identify the tangent space of D at any point with \mathbb{C} itself. A conformal metric provides for us an inner product at each point of D in the following way: if $v, w \in T_z\mathbb{D}$, then

$$\langle v, w \rangle_z := \rho^2(z) v \cdot \overline{w}.$$

Angles can now be defined using the inner-product and they agree with the Euclidean angles at those points where the metric is non-singular.

Length of curves

If $\gamma : [a, b] \rightarrow D$ is a piecewise smooth curve then we define the length of γ with respect to ρ as

$$\ell(\gamma) := \int_a^b \rho(\gamma(t)) |\gamma'(t)| dt.$$

The distance associated to a conformal metric

If $D \subset \mathbb{C}$ is a domain and ρ is a conformal metric, we define

$$d_\rho(z, w) := \inf \{ \ell(\gamma) : \gamma : [0, 1] \rightarrow D \text{ is a curve with } \gamma(0) = a \text{ and } \gamma(1) = b \}.$$

It is straightforward to show that d_ρ defines a distance (i.e., a metric in the usual sense) on D .

Pullback of a conformal metric

Definition

Let $D_1, D_2 \subset \mathbb{C}$ be domains and let $f : D_1 \rightarrow D_2$ be a thrice continuously differentiable function with isolated zeroes. We define the *pullback* of the conformal metric ρ of D_2 under f to be the conformal metric on D_1 given by

$$f^*\rho(z) = \rho(f(z)) \cdot |\partial_z f|.$$

If ρ_1 is another conformal metric such that for some holomorphic mapping $g : D_1 \rightarrow D_2$, $g^*\rho \equiv \rho_1$ then we say g is an *isometry* of the pair (D_1, ρ_1) with (D_2, ρ) .

Pullback of a conformal metric

Definition

Let $D_1, D_2 \subset \mathbb{C}$ be domains and let $f : D_1 \rightarrow D_2$ be a thrice continuously differentiable function with isolated zeroes. We define the *pullback* of the conformal metric ρ of D_2 under f to be the conformal metric on D_1 given by

$$f^*\rho(z) = \rho(f(z)) \cdot |\partial_z f|.$$

If ρ_1 is another conformal metric such that for some holomorphic mapping $g : D_1 \rightarrow D_2$, $g^*\rho \equiv \rho_1$ then we say g is an *isometry* of the pair (D_1, ρ_1) with (D_2, ρ) .

Straightforward computations reveal that holomorphic isometries preserve lengths of curves and therefore distances as well.

The Schwarz–Pick Lemma

A nice exercise in a first course in complex analysis is to deduce the following result known as the Schwarz–Lemma using the standard Schwarz Lemma.

Schwarz–Pick Lemma, 1916

Let $f : \mathbb{D} \rightarrow \mathbb{D}$ be holomorphic. Then

$$\frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \forall z \in \mathbb{D}.$$

The Schwarz–Pick Lemma

A nice exercise in a first course in complex analysis is to deduce the following result known as the Schwarz–Lemma using the standard Schwarz Lemma.

Schwarz–Pick Lemma, 1916

Let $f : \mathbb{D} \rightarrow \mathbb{D}$ be holomorphic. Then

①

$$\frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \forall z \in \mathbb{D}.$$

②

$$\frac{|f(z_1) - f(z_2)|}{|1 - f(z_1)\overline{f(z_2)}|} \leq \frac{|z_1 - z_2|}{|1 - z_1\overline{z_2}|} \quad \forall z_1, z_2 \in \mathbb{D}.$$

The Poincaré metric, 1881-84

- ❶ Recall that holomorphic mappings are conformal (i.e., angle preserving) at those points where they are non-singular. However, non-singular holomorphic mappings are not length preserving under the standard Euclidean metric.

The Poincaré metric, 1881-84

- 1 Recall that holomorphic mappings are conformal (i.e., angle preserving) at those points where they are non-singular. However, non-singular holomorphic mappings are not length preserving under the standard Euclidean metric.
- 2 Consider the unit disk. It is easy to construct a conformal metric ρ on \mathbb{D} under which each automorphism is an isometry. In fact, it is immediate from the Schwarz–Pick lemma that

$$\rho(z) = \frac{1}{1 - |z|^2},$$

is such a metric.

The Poincaré metric, 1881-84


- ❶ Recall that holomorphic mappings are conformal (i.e., angle preserving) at those points where they are non-singular. However, non-singular holomorphic mappings are not length preserving under the standard Euclidean metric.
- ❷ Consider the unit disk. It is easy to construct a conformal metric ρ on \mathbb{D} under which each automorphism is an isometry. In fact, it is immediate from the Schwarz–Pick lemma that

$$\rho(z) = \frac{1}{1 - |z|^2},$$


is such a metric.

- ❸ The above metric is known as the *Poincaré metric*. It is straightforward to show that the only conformal metrics on the disk that preserve the length of curves under automorphisms are constant multiples of the Poincaré metric.

Artistic visualization



Geodesics



The distance decreasing property

- ① Automorphisms of the disk are isometries of the Poincaré distance.

The distance decreasing property

- ① Automorphisms of the disk are isometries of the Poincaré distance.
- ② Observe that the Schwarz–Pick lemma implies the more general fact that holomorphic self-maps are contracting under the Poincaré distance.

The distance decreasing property

- ① Automorphisms of the disk are isometries of the Poincaré distance.
- ② Observe that the Schwarz–Pick lemma implies the more general fact that holomorphic self-maps are contracting under the Poincaré distance.
- ③ This geometric interpretation of the Schwarz lemma led Ahlfors to formulate a far-reaching generalization of the Schwarz lemma. This work kick-started the study of invariant metrics which is now a prominent field of research in complex analysis.

Curvature

Definition

Let $D \subset \mathbb{C}$ be a domain equipped with a conformal metric ρ . The *curvature of ρ* is defined to be

$$\kappa_{D,\rho}(z) := -\frac{1}{\rho^2(z)} \nabla^2 \log \rho(z).$$

Curvature

Definition

Let $D \subset \mathbb{C}$ be a domain equipped with a conformal metric ρ . The *curvature of ρ* is defined to be

$$\kappa_{D,\rho}(z) := -\frac{1}{\rho^2(z)} \nabla^2 \log \rho(z).$$

- 1 The above formula for the curvature comes from Riemannian geometry.

Curvature

Definition

Let $D \subset \mathbb{C}$ be a domain equipped with a conformal metric ρ . The *curvature of ρ* is defined to be

$$\kappa_{D,\rho}(z) := -\frac{1}{\rho^2(z)} \nabla^2 \log \rho(z).$$

- 1 The above formula for the curvature comes from Riemannian geometry.
- 2 The curvature of the Euclidean metric is clearly 0 and a calculation reveals that the curvature of the Poincaré metric is constant -4 .

Curvature

Definition

Let $D \subset \mathbb{C}$ be a domain equipped with a conformal metric ρ . The *curvature of ρ* is defined to be

$$\kappa_{D,\rho}(z) := -\frac{1}{\rho^2(z)} \nabla^2 \log \rho(z).$$

- 1 The above formula for the curvature comes from Riemannian geometry.
- 2 The curvature of the Euclidean metric is clearly 0 and a calculation reveals that the curvature of the Poincaré metric is constant -4 .
- 3 An easy computation shows that if D_1 is a domain and $f : D_1 \rightarrow D$ is a holomorphic map with $f'(z) \neq 0$ for some $z \in D_1$ then

$$\kappa_{D_1, f^*\rho}(z) = \kappa_{D,\rho}(f(z)).$$

Curvature

Definition

Let $D \subset \mathbb{C}$ be a domain equipped with a conformal metric ρ . The *curvature of ρ* is defined to be

$$\kappa_{D,\rho}(z) := -\frac{1}{\rho^2(z)} \nabla^2 \log \rho(z).$$

- ❶ The above formula for the curvature comes from Riemannian geometry.
- ❷ The curvature of the Euclidean metric is clearly 0 and a calculation reveals that the curvature of the Poincaré metric is constant -4 .
- ❸ An easy computation shows that if D_1 is a domain and $f : D_1 \rightarrow D$ is a holomorphic map with $f'(z) \neq 0$ for some $z \in D_1$ then

$$\kappa_{D_1, f^*\rho}(z) = \kappa_{D,\rho}(f(z)).$$

- ❹ This means that the notion of curvature is invariant under biholomorphism.

The Ahlfors–Schwarz lemma

Theorem (Ahlfors, 1938)

Let ρ be the Poincaré metric on \mathbb{D} and let U be a domain equipped with a conformal metric σ such that $\kappa_{U,\sigma}(w) \leq -4 \forall w \in U$. Then for any holomorphic map $f : \mathbb{D} \rightarrow U$, we have

$$f^* \sigma(z) \leq \rho(z) \quad \forall z \in \mathbb{D}.$$

The Ahlfors–Schwarz lemma

Theorem (Ahlfors, 1938)

Let ρ be the Poincaré metric on \mathbb{D} and let U be a domain equipped with a conformal metric σ such that $\kappa_{U,\sigma}(w) \leq -4 \forall w \in U$. Then for any holomorphic map $f : \mathbb{D} \rightarrow U$, we have

$$f^* \sigma(z) \leq \rho(z) \quad \forall z \in \mathbb{D}.$$

The proof of the lemma is not hard. What is more interesting is that several classical results including Montel's theorem and Picard's theorems can be given easy proofs using this theorem.

The Ahlfors–Schwarz lemma

Theorem (Ahlfors, 1938)

Let ρ be the Poincaré metric on \mathbb{D} and let U be a domain equipped with a conformal metric σ such that $\kappa_{U,\sigma}(w) \leq -4 \forall w \in U$. Then for any holomorphic map $f : \mathbb{D} \rightarrow U$, we have

$$f^* \sigma(z) \leq \rho(z) \quad \forall z \in \mathbb{D}.$$

The proof of the lemma is not hard. What is more interesting is that several classical results including Montel's theorem and Picard's theorems can be given easy proofs using this theorem.

The only non-trivial step in the proofs is the construction of a conformal metric on $\mathbb{C} \setminus \{0, 1\}$ whose curvature is bounded above by -4 . There are several such constructions available.

First definition of hyperbolicity

- ① We say a domain $D \subset \mathbb{C}$ is *hyperbolic* if it admits a conformal metric whose curvature is bounded above by -4 .

First definition of hyperbolicity

- ① We say a domain $D \subset \mathbb{C}$ is *hyperbolic* if it admits a conformal metric whose curvature is bounded above by -4 .
- ② Using the famous uniformization theorem for Riemann surfaces, one can prove that the hyperbolic domains are precisely those whose universal cover is the unit disk.

First definition of hyperbolicity

- 1 We say a domain $D \subset \mathbb{C}$ is *hyperbolic* if it admits a conformal metric whose curvature is bounded above by -4 .
- 2 Using the famous uniformization theorem for Riemann surfaces, one can prove that the hyperbolic domains are precisely those whose universal cover is the unit disk.
- 3 It also turns out that the distance generated on a hyperbolic domain makes it into a complete metric space.

On to higher dimensions: The Carathéodory pseudodistance

Let M be a complex manifold. The Carathéodory pseudodistance $C_M(z, w)$ is defined as follows:

$$C_M(z, w) := \sup\{\rho_{\mathbb{D}}(f(z), f(w)) : f \in O(M, \mathbb{D})\}.$$

On to higher dimensions: The Carathéodory pseudodistance

Let M be a complex manifold. The Carathéodory pseudodistance $C_M(z, w)$ is defined as follows:

$$C_M(z, w) := \sup\{\rho_{\mathbb{D}}(f(z), f(w)) : f \in O(M, \mathbb{D})\}.$$

Here $\rho_{\mathbb{D}}$ is the Poincaré distance on the disk. Observe that the Carathéodory pseudodistance of \mathbb{D} is the Poincaré distance.

On to higher dimensions: The Carathéodory pseudodistance

Let M be a complex manifold. The Carathéodory pseudodistance $C_M(z, w)$ is defined as follows:

$$C_M(z, w) := \sup\{\rho_{\mathbb{D}}(f(z), f(w)) : f \in O(M, \mathbb{D})\}.$$

Here $\rho_{\mathbb{D}}$ is the Poincaré distance on the disk. Observe that the Carathéodory pseudodistance of \mathbb{D} is the Poincaré distance.

It is a triviality that holomorphic mappings are contractions when both the domain and codomain are equipped with the respective Carathéodory pseudodistances.

On to higher dimensions: The Carathéodory pseudodistance

Let M be a complex manifold. The Carathéodory pseudodistance $C_M(z, w)$ is defined as follows:

$$C_M(z, w) := \sup\{\rho_{\mathbb{D}}(f(z), f(w)) : f \in O(M, \mathbb{D})\}.$$

Here $\rho_{\mathbb{D}}$ is the Poincaré distance on the disk. Observe that the Carathéodory pseudodistance of \mathbb{D} is the Poincaré distance.

It is a triviality that holomorphic mappings are contractions when both the domain and codomain are equipped with the respective Carathéodory pseudodistances.

Observe also that the Carathéodory pseudodistance of a compact complex manifold is trivial.

The second definition of hyperbolicity

Definition (Carathéodory, 1926)

We say a complex manifold M is *C-hyperbolic* if the Carathéodory pseudodistance of M is a distance. If the distance is complete, we say that M is *complete C-hyperbolic*

The second definition of hyperbolicity

Definition (Carathéodory, 1926)

We say a complex manifold M is *C-hyperbolic* if the Carathéodory pseudodistance of M is a distance. If the distance is complete, we say that M is *complete C-hyperbolic*

The unit disk is (complete) *C-hyperbolic* whereas \mathbb{C} and $\mathbb{C} \setminus \{0, 1\}$ and compact complex manifolds are not.

Invariant metrics

- ❶ The Carathéodory pseudodistance, though useful, is not ideal. Even on a C -hyperbolic complex manifold, it is not true that the topology generated by the Carathéodory distance coincides with the underlying topology.

Invariant metrics

- ① The Carathéodory pseudodistance, though useful, is not ideal. Even on a C -hyperbolic complex manifold, it is not true that the topology generated by the Carathéodory distance coincides with the underlying topology.
- ② This prompted the study of the more general idea of a *contractible family of invariant metrics*. A system of invariant metrics is an association of a pseudodistance to each complex manifold such that holomorphic mappings become distance decreasing. We also stipulate that the pseudodistance associated to \mathbb{D} is the Poincaré distance.

Invariant metrics

- ❶ The Carathéodory pseudodistance, though useful, is not ideal. Even on a C -hyperbolic complex manifold, it is not true that the topology generated by the Carathéodory distance coincides with the underlying topology.
- ❷ This prompted the study of the more general idea of a *contractible family of invariant metrics*. A system of invariant metrics is an association of a pseudodistance to each complex manifold such that holomorphic mappings become distance decreasing. We also stipulate that the pseudodistance associated to \mathbb{D} is the Poincaré distance.
- ❸ In other words, it is a way of putting a pseudodistance on complex manifolds so that the analogue of the Schwarz lemma is built-in.

Invariant metrics

- ❶ The Carathéodory pseudodistance, though useful, is not ideal. Even on a C -hyperbolic complex manifold, it is not true that the topology generated by the Carathéodory distance coincides with the underlying topology.
- ❷ This prompted the study of the more general idea of a *contractible family of invariant metrics*. A system of invariant metrics is an association of a pseudodistance to each complex manifold such that holomorphic mappings become distance decreasing. We also stipulate that the pseudodistance associated to \mathbb{D} is the Poincaré distance.
- ❸ In other words, it is a way of putting a pseudodistance on complex manifolds so that the analogue of the Schwarz lemma is built-in.
- ❹ The Carathéodory pseudodistance provides one way of creating such a system.

The Kobayashi pseudodistance

- 1 If ρ_M is a pseudodistance on the complex manifold M that is distance decreasing for holomorphic maps $f : M \rightarrow \mathbb{D}$ then

$$\rho_{\mathbb{D}}(f(z), f(w)) \leq \rho_M(z, w) \quad \forall z, w \in M.$$

The Kobayashi pseudodistance

- 1 If ρ_M is a pseudodistance on the complex manifold M that is distance decreasing for holomorphic maps $f : M \rightarrow \mathbb{D}$ then

$$\rho_{\mathbb{D}}(f(z), f(w)) \leq \rho_M(z, w) \quad \forall z, w \in M.$$

- 2 Taking supremum on the LHS reveals that Carathéodory system of assigning pseudodistances gives the “smallest” possible pseudodistance on each complex manifold.

The Kobayashi pseudodistance

- 1 If ρ_M is a pseudodistance on the complex manifold M that is distance decreasing for holomorphic maps $f : M \rightarrow \mathbb{D}$ then

$$\rho_{\mathbb{D}}(f(z), f(w)) \leq \rho_M(z, w) \quad \forall z, w \in M.$$

- 2 Taking supremum on the LHS reveals that Carathéodory system of assigning pseudodistances gives the “smallest” possible pseudodistance on each complex manifold.
- 3 The Kobayashi pseudodistance is the dual of the Carathéodory distances and gives the “largest” possible distance on each complex manifold.

The Kobayashi pseudodistance, 1967

If $z, w \in M$ define the *Lempert function* $\delta_M(z, w)$ to be

$$\delta_M(z, w) := \inf \rho(x, y)$$

where the infimum ranges over all holomorphic mappings $f : \mathbb{D} \rightarrow M$ with $f(x) = z$ and $f(y) = w$.

The Kobayashi pseudodistance, 1967

If $z, w \in M$ define the *Lempert function* $\delta_M(z, w)$ to be

$$\delta_M(z, w) := \inf \rho(x, y)$$

where the infimum ranges over all holomorphic mappings $f : \mathbb{D} \rightarrow M$ with $f(x) = z$ and $f(y) = w$.

The Lempert function need not satisfy the triangle inequality so we use the Kobayashi trick to create a pseudodistance as follows: we define the *Kobayashi pseudodistance* $K(z, w)$ to be

$$K(z, w) := \inf \left\{ \sum_{i=1}^n \delta_M(p_i, p_{i+1}) \right\},$$

where the sum runs through all finite chains of points $z = p_1, \dots, p_n = w$.

Basic properties of the Kobayashi pseudodistance

- 1 It is clear that the Kobayashi pseudodistance is the “largest” possible invariant metric and also that the Kobayashi pseudodistance of the disk is the Poincaré distance.

Basic properties of the Kobayashi pseudodistance

- ① It is clear that the Kobayashi pseudodistance is the “largest” possible invariant metric and also that the Kobayashi pseudodistance of the disk is the Poincaré distance.
- ② Again, it is trivial the holomorphic functions are contractions under the Kobayashi pseudodistance.

Basic properties of the Kobayashi pseudodistance

- ① It is clear that the Kobayashi pseudodistance is the “largest” possible invariant metric and also that the Kobayashi pseudodistance of the disk is the Poincaré distance.
- ② Again, it is trivial the holomorphic functions are contractions under the Kobayashi pseudodistance.
- ③ Analogous definitions of K -hyperbolicity and completeness can be made. If a complex manifold is K -hyperbolic then the topology generated by the Kobayashi distance coincides with the manifold topology.

Basic properties of the Kobayashi pseudodistance

- 1 It is clear that the Kobayashi pseudodistance is the “largest” possible invariant metric and also that the Kobayashi pseudodistance of the disk is the Poincaré distance.
- 2 Again, it is trivial the holomorphic functions are contractions under the Kobayashi pseudodistance.
- 3 Analogous definitions of K -hyperbolicity and completeness can be made. If a complex manifold is K -hyperbolic then the topology generated by the Kobayashi distance coincides with the manifold topology.
- 4 An extremely useful property of the Kobayashi pseudodistance is its good behaviour under coverings: If a complex manifold M' is a covering manifold of the complex manifold M then M is (complete) K -hyperbolic iff M' is.

A central fact

The following fact is one of the central facts that make invariant metrics so powerful in geometric function theory:

Equicontinuity of the space of holomorphic mappings

If M and N are complex manifolds with N being K -hyperbolic then the space of holomorphic mappings $O(M, N)$ is an equicontinuous family.

Wu's version of Montel's theorem

H. Wu initiated the study of normal families in a seminal paper in 1967. Using the classical Arzela–Ascoli theorem and the central fact from the previous slide the following theorem becomes straightforward.

Theorem (Wu)

Let M and N be complex manifolds and let N be complete K -hyperbolic. Then the family $\mathcal{O}(M, N)$ is a normal family.

A version of Montel's theorem for Riemann surfaces

Theorem (J., 2014)

Let X be a connected complex manifold and let R be a hyperbolic open connected subset of a compact Riemann surface S . Then, given any sequence $\{f_v\} \subset O(X, R)$, there exists a subsequence $\{f_{v_k}\}$ and a holomorphic map $f_0 : X \rightarrow \overline{R}$ (the closure taken in S whenever R is non-compact) such that $f_{v_k} \rightarrow f_0$ uniformly on compact subsets of X .

But where is the geometry?

So far, all the geometry in this talk has been differential and the familiar classical geometry has disappeared. However, there is one deep and shocking result that is poorly understood where Euclidean geometry crops up in an unexpected manner.

Theorem (Lempert, 1984)

Let $D \subset \mathbb{C}^n$, $n > 1$ be a bounded and convex domain. Then D is K -hyperbolic and the Lempert function, the Kobayashi distance and the Carathéodory distance all coincide.

Structure theorem for products of Riemann surfaces

Theorem (J., 2014)

Let R_j and S_j , $j = 1, \dots, n$, be compact Riemann surfaces, and let X_j (resp. Y_j) be a connected, hyperbolic open subset of R_j (resp. S_j) for each $j = 1, \dots, n$. Let $F = (F_1, \dots, F_n) : X_1 \times \dots \times X_n \rightarrow Y_1 \times \dots \times Y_n$ be a finite proper holomorphic map. Then, each F_i is of the form $F_i(z_{\pi(i)})$, where π is a permutation of $\{1, \dots, n\}$.

Finiteness of the space of holomorphic mappings

Theorem (Divakaran and J., 2017)

Let $X := X_1 \times \cdots \times X_n$ be a product of hyperbolic Riemann surfaces of finite type and let $Y = \Omega/\Gamma$ be an m -dimensional complex manifold where $\Omega \subset \mathbb{C}^m$ is a bounded domain and Γ is fixed-point-free discrete subgroup of $\text{Aut}(\Omega)$.

- ❶ *If N is a tautly embedded complex submanifold of Y then $O_{\text{dom}}(X, N)$ is a finite set.*
- ❷ *If Y is geometrically finite and Ω is complete hyperbolic then $O_{\text{dom}}(X, Y)$ is a finite set.*
- ❸ *If in addition to the conditions in (2), the essential boundary dimension of Ω is zero, then $O_*(X, Y)$ is a finite set.*

Structure theorem for the symmetric product of a Riemann surface

Theorem (Biswas, Bharali, Divakaran and J., 2018)

Let $X = X_1 \times \cdots \times X_n$ be a complex manifold where each X_j is a connected non-compact Riemann surface obtained by excising a non-empty indiscrete set from a compact Riemann surface R_j . Let Y be a connected bordered Riemann surface with C^2 -smooth boundary. Let $F : X \rightarrow \text{Sym}^n(Y)$ be a proper holomorphic map. Then, there exist proper holomorphic maps $F_j : X_j \rightarrow Y$, $j = 1, \dots, n$, such that

$$F(x_1, \dots, x_n) = \pi_{\text{Sym}} \circ (F_1(x_1), \dots, F_n(x_n)) \quad \forall (x_1, \dots, x_n) \in X.$$

THANK YOU