

YET ANOTHER TABLE OF INTEGRALS

RICHARD J. MATHAR

ABSTRACT. This collection of sums and integrals has been harvested from the mathematical and physical literature in unstructured ways. Its main use is backtracking the original sources whenever an integral of the reader's application resembles one of the items in the collection.

INTRODUCTION

Dealing with the analysis of real numbers in the physical sciences shows a strange attraction towards integrals. Closed-form integration beats numerical integration, and often adaptive series expansion helps to crumble cumbersome integral kernels to digestable pieces.

The current table started as a incoherent list of bookmarks pointing to “interesting” formulas that complement or correct the Gradstein-Rhyshik tables [91], see <http://www.mathtable.com/gr/>. As such it does not replicate the original sources in full but is to be merely regarded as an aid to find places at which certain forms and classes of integrals or sums have been targeted.

The notation is generally not harmonized. Stirling numbers appear in bracketed and indexed notations, and at least two different meanings of harmonic numbers H with lower and upper indices are met.

There is only one hint of use: The list of references appears *prior* to each formula.

0.1. Finite Series. [181, 176]

$$(0.1) \quad \sum_{j=1}^n j^k = \frac{1}{n} \left(\rho(n, i) + \sum_{i=1}^n \sigma_{k+1}(i) \right) = \frac{B_{k+1}(n+1) - B_{k+1}}{k+1}$$

where $\rho(n, k) \equiv \sum_{d=1}^n d^k (n \bmod d)$ is a sum over $n \bmod d$ multiplied, then summed, over d^k , and $\sigma_k(n) = \sum_{d|n} d^k$.

[55, 67]

$$(0.2) \quad \sum_{k=1}^{n-1} k^r = \sum_{k=0}^r \frac{B_k}{k!} \frac{r!}{(r-k+1)!} n^{r-k+1}.$$

[182]

$$(0.3) \quad \sum_{k=0}^n k^m = \sum_{j=0}^m \left\{ \begin{matrix} m \\ j \end{matrix} \right\} \binom{n+1}{j+1} j!.$$

Date: August 30, 2013.

2010 Mathematics Subject Classification. Primary 33-00, 40-00; Secondary 44-00.

Key words and phrases. sums, integrals, tables.

By telescoping the last term in this formula becomes

$$(0.4) \quad n^m = \sum_{j=0}^m \left\{ \begin{matrix} m \\ j \end{matrix} \right\} \binom{n}{j} j!.$$

[186]

$$(0.5) \quad \sum_{0 \leq j \leq n/m} \binom{n-jm}{k} = P(n+m, k, m) - P(r, k, m)$$

for $n \geq 0$, $k \geq 1$, $m \geq 1$ where

$$(0.6) \quad P(x, k, m) \equiv \frac{1}{m} \sum_{j=1}^{k+1} \binom{x}{j} A(m, k+1-j)$$

with g.f.

$$(0.7) \quad \frac{mx}{(1+x)^m - 1} = \sum_{j=0}^{\infty} A(m, j) x^j.$$

[166]

$$(0.8) \quad \sum_{k=0}^n \binom{r+k}{k} = \binom{r+n+1}{n}, \quad n = 0, 1, 2, \dots$$

[182]

$$(0.9) \quad \sum_k \binom{n}{2k} k = n2^{n-3}, \quad n \geq 2.$$

[182]

$$(0.10) \quad \sum_k \binom{n}{2k+1} k = (n-2)2^{n-3}, \quad n \geq 2.$$

[182]

$$(0.11) \quad \sum_k \binom{n}{2k} k^{\underline{m}} = n(n-m-1)^{\underline{m-1}} 2^{n-2m-1}, \quad n \geq m+1,$$

[182]

$$(0.12) \quad \sum_k \binom{n}{2k+1} k^{\underline{m}} = n(n-m-1)^{\underline{m}} 2^{n-2m-1}, \quad n \geq m+1,$$

where $k^{\underline{m}}$ is the falling factorial $k(k-1)(k-2) \cdots (k-m+1)$.

[182]

$$(0.13) \quad \sum_{k=0}^n \binom{n}{k} k^m = \sum_{j=0}^m \left\{ \begin{matrix} m \\ j \end{matrix} \right\} \binom{n}{j} j! 2^{n-j}.$$

[182]

$$(0.14) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k k^m = \left\{ \begin{matrix} m \\ n \end{matrix} \right\} (-1)^n n!.$$

[182]

$$(0.15) \quad \sum_{k=0}^n \binom{n}{2k} k^m = n \sum_{j=1}^{\min(m, n-1)} \left\{ \begin{matrix} m \\ j \end{matrix} \right\} \binom{n-j-1}{j-1} (j-1)! 2^{n-2j-1}.$$

[182]

$$(0.16) \quad \sum_{k=0}^n \binom{n}{2k+1} k^m = \sum_{j=1}^{\min(m, n-1)} \left\{ \begin{matrix} m \\ j \end{matrix} \right\} \binom{n-j-1}{j-1} j! 2^{n-2j-1}.$$

[107]

$$(0.17) \quad \sum_{0 \leq k \leq N, k \neq K} \binom{N}{k} (-1)^k \frac{1}{(k-K)^m} = \binom{N}{K} (-1)^{K+1} \frac{1}{m!} Y_m(\dots, (i-1)! [H_{N-K}^{(i)} + (-1)^i H_K^{(i)}], \dots)$$

where $0 \leq K \leq N$, $Y_m(\dots, x_i, \dots)$ are the Bell polynomials and $H_r^{(i)} = \sum_{j=1}^r j^{-i}$ Harmonic numbers of the i -th order.

[107]

$$(0.18) \quad \sum_{0 \leq k \leq N, k \neq K} \binom{N}{k} (-1)^k \frac{1}{k-K} = \binom{N}{K} (-1)^{K+1} (H_{N-K} - H_K)$$

where $0 \leq K \leq N$, $H_r^{(i)} = \sum_{j=1}^r j^{-i}$ Harmonic numbers of the i -th order.

[107]

$$(0.19) \quad \sum_{0 \leq k \leq N} \binom{N}{k} (-1)^k \frac{1}{k^m} = -\frac{1}{m!} Y_m(\dots, (i-1)! H_N^{(i)}, \dots)$$

where $H_r^{(i)} = \sum_{j=1}^r j^{-i}$ Harmonic numbers of the i -th order.

[107]

$$(0.20) \quad \sum_{0 \leq k \leq N} \binom{N}{k} (-1)^k \frac{1}{(k-\xi)^m} = \Gamma(-\xi) \frac{\Gamma(N+1)}{\Gamma(N+1-\xi)} \frac{1}{(m-1)!} Y_{m-1}(\dots, (i-1)! \zeta_N(i, -\xi), \dots),$$

where $\zeta_N(i, -\xi) = \sum_{j=0}^N (j-\xi)^{-j}$.

[161, §4.3]

$$(0.21) \quad a_n = \sum_{k=0}^n \binom{p+k}{k} b_{n-k} \Leftrightarrow b_n = \sum_{k=0}^n (-1)^k \binom{p+k}{k} a_{n-k}.$$

[161, §4.3]

$$(0.22) \quad a_n = \sum_{k=0}^n \binom{p+k}{k} b_{n-qk} \Leftrightarrow b_n = \sum_{k=0}^n (-1)^k \binom{p+k}{k} a_{n-qk}.$$

[161, §4.3]

$$(0.23) \quad a_n = \sum_{k=0}^n \binom{2k}{k} b_{n-k} \Leftrightarrow b_n = \sum_{k=0}^n \frac{1}{1-2k} \binom{2k}{k} a_{n-k}.$$

[161, §4.3]

$$(0.24) \quad a_n = \sum_{k=0}^n \frac{1}{k+1} \binom{2k}{k} b_{n-k} \Leftrightarrow b_n = a_n - \sum_{k=1}^n \frac{1}{k} \binom{2k}{k} a_{n-k}.$$

[156]

$$(0.25) \quad x_n = a_n + 2^{1-n} \sum_{k=0}^n \binom{n}{k} x_k \Leftrightarrow x_n = x_0 + \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{k\bar{x}_1 + \hat{a}_k - \bar{x}_0}{1 - 2^{1-k}}$$

where $\bar{x}_0 \equiv a_0 + x_0$, $\bar{x}_1 = a_1 + x_0$ and

$$(0.26) \quad \hat{a}_n = \sum_{k=0}^n \binom{n}{k} (-1)^k a_k.$$

[156]

$$(0.27) \quad x_{n+1} = a_{n+1} + 2^{1-n} \sum_{k=0}^n \binom{n}{k} x_k \Leftrightarrow x_n = - \sum_{k=0}^n (-1)^k \binom{n}{k} \hat{x}_{k-2}$$

where $\bar{x}_0 = 0$,

$$(0.28) \quad \hat{x}_n = Q_n \sum_{i=1}^{n+1} (\hat{a}_i - \hat{a}_{i+1} - a_1) / Q_{i-1}, \quad Q_n \equiv \prod_{k=1}^n (1 - 2^{-k}).$$

[106]

$$(0.29) \quad u_n = Au_{n-1} - Bu_{n-2}, \quad v_n = Av_{n-1} - Bv_{n-2},$$

$$(0.30)$$

$$\rightsquigarrow u_{n+1} = \prod_{j=1}^n \left(A - 2i\sqrt{-B} \cos \frac{\pi j}{n+1} \right) = (i\sqrt{-B})^n \frac{\sin \left((n+1) \cos^{-1} \frac{-iA}{2\sqrt{-B}} \right)}{\sin \left(\cos^{-1} \frac{-iA}{2\sqrt{-B}} \right)},$$

$$(0.31)$$

$$\rightsquigarrow v_n = \prod_{k=1}^n \left(A - 2i\sqrt{-B} \cos \frac{\pi(k-1/2)}{n} \right) = 2(i\sqrt{-B})^n \cos \left(n \cos^{-1} \frac{-iA}{2\sqrt{-B}} \right),$$

where $u_0 = 0$, $u_1 = 1$, $v_0 = 2$, $v_1 = A$

[201]

$$(0.32) \quad \sum_{x=1}^{p-1} F(x) + \sum_{x=1}^{p-1} \left(\frac{x}{p} \right) F(x) = \sum_{x=1}^{p-1} F(x^2),$$

if p is an odd prime and $F(x) = F(x+p)$, where $(\frac{x}{p})$ is the Legendre Symbol.

[201]

$$(0.33) \quad \sum_{x=1}^{p-1} \left(\frac{x}{p} \right) e(k(x+\bar{x})) = \left(\frac{k}{p} \right) i^{(p-1)^2/4} p^{1/2} (e(2k) + e(-2k)).$$

where \bar{x} is the unique solution to $x\bar{x} \equiv 1 \pmod{p}$, and $e(t) = e^{2\pi it/p}$.

[48]

$$(0.34) \quad \sum_{k=0}^m \binom{m}{k} \frac{(n+k)!}{(n+k+s)!} a_{n+k+s} = \sum_{k=0}^n \binom{n}{k} \frac{(-1)^{n-k} (m+k)!}{(m+k+s)!} b_{m+k+s} \\ + \sum_{j=0}^{s-1} \sum_{i=0}^{s-1-j} \binom{s-1-j}{i} \binom{s-1}{j} \frac{(-1)^{n+1+i} a_j}{(s-1)!(m+n+1+i) \binom{m+n+i}{n}},$$

where $b_n \equiv \sum_{k=0}^n \binom{n}{k} a_k$.

[48]

$$(0.35) \quad \sum_{k=0}^m \binom{m}{k} \binom{n+k}{s} a_{n+k+s} = \sum_{k=0}^n \binom{n}{k} \binom{m+k}{s} (-1)^{n-k} b_{m+k+s},$$

where $b_n \equiv \sum_{k=0}^n \binom{n}{k} a_k$.

[48]

$$(0.36) \quad \sum_{k=0}^m \frac{\binom{m}{k}}{\binom{n+k+s}{s}} x^{m-k} A_{n+k+s}(y) = \sum_{k=0}^n \frac{\binom{n}{k}}{\binom{m+k+s}{s}} (-1)^{n+m+s} x^{n-k} A_{m+k+s}^*(z) \\ + \sum_{j=0}^{s-1} \sum_{i=0}^{s-1-j} \binom{s-1-j}{i} \binom{s-1}{j} \frac{(-1)^{n+1+i} x^{m+n+s-j} s A_j(y)}{(m+n+1+i) \binom{m+n+i}{n}},$$

and

$$(0.37) \quad \sum_{k=0}^m \binom{m}{k} \binom{n+k}{s} x^{m-k} A_{n+k+s}(y) = \sum_{k=0}^n \binom{n}{k} \binom{m+k}{s} (-1)^{n+m+s} x^{n-k} A_{m+k+s}^*(z)$$

where $A_n(x) \equiv \sum_{k=0}^n \binom{n}{k} (-1)^k a_k x^{n-k}$ and $A_n^*(x) \equiv \sum_{k=0}^n \binom{n}{k} (-1)^k a_k^* x^{n-k}$ and $a_n^* \equiv \sum_{k=0}^n \binom{n}{k} (-1)^k a_k$ and $x+y+z=1$.

[48]

$$(0.38) \quad a_{n,m} = \sum_{k=0}^n \binom{n}{k} \alpha^{n-k} \beta^k a_{0,m+k},$$

and

$$(0.39) \quad \sum_{k=0}^n \binom{n}{k} \alpha^{n-k} \beta^k a_{0,m+k} = \sum_{k=0}^m \binom{m}{k} (-\alpha)^{m-k} \beta^{-m} a_{n+k,0},$$

where $a_{n,m} \equiv \alpha a_{n-1,m} + \beta a_{n-1,m+1}$ for $n \geq 1, m \geq 0$.

[92]

$$(0.40) \quad x_{nm} = \sum_{k=0}^m x_{mk} \binom{n+k}{2m} \rightsquigarrow x_{nm} = \sum_{k=0}^m \frac{2k+1}{m+k+1} \binom{n+k}{m+k} \binom{n-1-k}{m-k} x_{kk},$$

for $m < n, n \geq 0, 0 \leq m \leq n$.

[33]

$$(0.41) \quad F(x) \equiv \sum_{k \geq 1} f(x/k); \quad G(x) \equiv \sum_{k \geq 1} (-)^k f(x/k) \rightsquigarrow F(x) = 2^{-n} F(2^n x) + \sum_{k \geq 1} 2^{-k} G(2^k x).$$

[33]
(0.42)

$$F(x) = r_1 F(m_1 x) + r_2 G(m_2 x) \rightsquigarrow r_2 \sum_{k=1}^n r_1^{k-1} G(m_1^{k-1} m_2 x) = F(x) - r_1^n F(m_1^n x).$$

[182]

(0.43)
$$\sum_{k=0}^n \binom{n}{k} H_k = 2^n \left(H_n - \sum_{k=1}^n \frac{1}{k 2^k} \right).$$

where H_k are the harmonic numbers.

[182]

(0.44)
$$\sum_k \binom{n}{2k} \frac{1}{k+1} = \frac{n 2^{n+1} + 2}{(n+1)(n+2)}.$$

[182]

(0.45)
$$\sum_k \binom{n}{2k+1} \frac{1}{k+1} = \frac{2^{n+1} - 2}{n+1}.$$

[164]

(0.46)
$$\sum_{j=0}^n \binom{m-a+b}{j} \binom{n+a-b}{n-j} \binom{a+j}{m+n} = \binom{a}{m} \binom{b}{n},$$

where n, m are integer and a, b real.

[164]

(0.47)
$$\sum_{m=0}^P (-)^m \binom{P}{m} \binom{a-m}{M} = \binom{a-P}{M-P},$$

where P, M are integer and a, b real.

[166]

(0.48)
$$\sum_k \binom{r}{k} \binom{s}{n+k} = \binom{r+s}{r+n}, \quad r = 0, 1, 2, \dots, \quad n = \dots - 2, -1, 0, 1, 2, 3, \dots$$

[164, 166]

(0.49)
$$\sum_{m=0}^n (-)^m \binom{n}{m} \binom{a+m}{p} = (-)^n \binom{a}{p-n},$$

where n, p are integer and a is real.

[166]

(0.50)
$$\sum_{k \geq 0} \binom{n+k}{m+2k} \binom{2k}{k} \frac{(-1)^k}{k+1} = \binom{n-1}{m-1}.$$

[166]

(0.51)
$$\sum_{\nu=0}^s (-)^{\nu} \binom{\beta}{\nu} \binom{\beta+s-\nu}{\beta} \frac{\alpha}{\alpha+s-\nu} = \frac{(\alpha-\beta)_s}{(\alpha+1)_s}.$$

[166]
(0.52)

$$\sum_{k=-l}^l (-)^k \binom{2l}{l+k} \binom{2m}{m+k} \binom{2n}{n+k} = \frac{(l+m+n)!(2l)!(2m)!(2n)!}{(l+m)!(m+n)!(n+l)!l!m!n!}, \quad l = \min(l, m, n).$$

[199]
(0.53)

$$\sum_k \sum_j \binom{n}{k} \binom{n+k}{k} \binom{k}{j}^3 = \sum_k \binom{n}{k}^2 \binom{n+k}{k}^2.$$

[199]
(0.54)

$$\sum_{i=0}^n \sum_{j=0}^n \binom{i+j}{i}^2 \binom{4n-2i-2j}{2n-2i} = (2n+1) \binom{2n}{n}.$$

[199]
(0.55)

$$\sum_{k_1} \sum_{k_2 \leq k_1} \sum_{k_3 \leq k_2} (k_1 - k_2)(k_1 - k_3)(k_2 - k_3) \binom{n}{k_1} \binom{n}{k_2} \binom{n}{k_3} = n^2(n-1)8^{n-2} \frac{(3/2)_{n-2}}{(3)_{n-2}}.$$

[199]
(0.56)

$$\sum_i \sum_j \binom{i+j}{i} \binom{n-i}{j} \binom{n-j}{n-i-j} = \sum_{k=0}^n \binom{2k}{k}.$$

[199]
(0.57)

$$\sum_{i=0}^n \sum_{j=0}^m \binom{i+j}{j}^2 \binom{m+m-i-j}{n-j}^2 = \frac{1}{2} \binom{2m+2n+2}{2n+1}.$$

[199]
(0.58)

$$\sum_{s=0}^k \sum_{b \geq 0} (-)^b \binom{s}{b} \binom{k-s}{2v-b} \binom{k-2v}{s-b} = \binom{k-v}{k-2v} 2^{k-2v}, \quad k \geq 2v.$$

[199]
(0.59)

$$\sum_{j,k} (-)^{j+k} \binom{j+k}{k+l} \binom{r}{j} \binom{n}{k} \binom{s+n-j-k}{m-j} = (-)^l \binom{n+r}{n+l} \binom{s-r}{m-n-l}.$$

[49]
(0.60)

$$\sum_{i=0}^n \sum_{j=0}^n \binom{i+j}{i} \binom{m-i+j}{j} \binom{n-j+i}{i} \binom{m+n-i-j}{m-i} = \frac{(m+n+1)!}{m!n!} \sum_k \frac{1}{2k+1} \binom{m}{k} \binom{n}{k}.$$

[49]
(0.61)

$$\sum_{i=0}^n \sum_{j=0}^n \binom{i+j}{i}^2 \binom{4n-2i-2j}{2n-2i} = (2n+1) \binom{2n}{n}^2.$$

$$\begin{aligned} & \text{[49]} \\ (0.62) \quad & \sum_{i=0}^{\lfloor m/2 \rfloor} \sum_{j=0}^{\lfloor n/2 \rfloor} \binom{i+j}{i}^2 \binom{m+n-2i-2j}{n-2i} = \frac{\lfloor (m+n+1)/2 \rfloor! \lfloor (m+n+2)/2 \rfloor!}{\lfloor m/2 \rfloor! \lfloor (m+1)/2 \rfloor! \lfloor (n+1)/2 \rfloor!}. \end{aligned}$$

$$\begin{aligned} & \text{[49]} \\ (0.63) \quad & \sum_i \sum_j \binom{n}{j} \binom{n+j}{j} \binom{j}{i}^2 \binom{2i}{i} \binom{2i}{j-i} = \sum_k \binom{n}{k}^3 \binom{n+k}{k}^3. \end{aligned}$$

$$\begin{aligned} & \text{[49]} \\ (0.64) \quad & \sum_j \sum_k (-1)^{j+k} \binom{j+k}{k+l} \binom{r}{j} \binom{n}{k} \binom{s+n-j-k}{m-j} = (-1)^l \binom{n+r}{n+l} \binom{s-r}{m-n-l}. \end{aligned}$$

$$\begin{aligned} & \text{[49, 199]} \\ (0.65) \quad & \sum_r \sum_s (-1)^{n+r+s} \binom{n}{r} \binom{n}{s} \binom{n+s}{s} \binom{n+r}{r} \binom{2n-r-s}{n} = \sum_k \binom{n}{k}^4. \end{aligned}$$

$$\begin{aligned} & \text{[172]} \\ (0.66) \quad & \binom{n}{i}^k = \binom{n+ik-i}{ik} + A_2 \binom{n+ik-i-1}{ik} + \cdots + A_{ik-i} \binom{n+1}{ik} + \binom{n}{ik}, \end{aligned}$$

$$\begin{aligned} S_i^k(n, 0) &\equiv \binom{n}{i}^k; \\ S_i^k(n, p) &\equiv S_i^k(1, p-1) + S_i^k(2, p-1) + \cdots + S_i^k(n, k-1) \\ &= \binom{n+ik-i+p}{ik+p} + A_2 \binom{n+ik-i+p-1}{ik+p} + \cdots + A_{ik-i} \binom{n+p+1}{ik+p} + \binom{n+p}{ik+p} \end{aligned}$$

where $A_j = A_{ik-i-j+2}$ for $j = 2, 3, \dots, ik-i$,

$$(0.67) \quad A_j = \begin{vmatrix} 1 & 0 & 0 & \cdots & 0 & \binom{i}{i}^k \\ \binom{ik+1}{ik} & 1 & 0 & \cdots & 0 & \binom{i+1}{i}^k \\ \binom{ik+2}{ik} & \binom{ik+1}{ik} & 1 & \cdots & 0 & \binom{i+2}{i}^k \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \binom{ik+j-1}{ik} & \binom{ik+j-2}{ik} & \binom{ik+j-3}{ik} & \cdots & \binom{ik+1}{ik} & \binom{i+j-1}{i}^k \end{vmatrix},$$

[189]

$$(0.68) \quad \sum_{k=0}^{2n} (-)^k \frac{\binom{2n}{k}}{\binom{4n}{2k}} = \frac{4n+1}{2n+1}.$$

[189]

$$(0.69) \quad \sum_{k=0}^{2n} (-)^k \frac{\binom{4n}{2k}}{\binom{2n}{k}} = -\frac{1}{2n-1}.$$

[189]

$$(0.70) \quad \sum_{k=0}^m \frac{\binom{m}{k}}{\binom{n+m}{p+k}} = \frac{n+m+1}{n+1} \binom{n}{p}^{-1},$$

with m, n, p nonnegative integers and $p \leq n$.

[189]

$$(0.71) \quad \sum_{k=0}^n (-)^k \frac{1}{\binom{n+m}{m+k}} = \frac{n+m+1}{m+n+2} \left(\binom{m+n+1}{m}^{-1} + (-)^n \right),$$

with m and n nonnegative integers.

[182]

$$(0.72) \quad \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right] H_k = n! \sum_{k=1}^n \frac{c_{k-1}}{k!},$$

where H_k are the harmonic numbers and $[\dots]$ unsigned Stirling numbers of the first kind.

[182]

$$(0.73) \quad \sum_{k=0}^n s(n, k) = s(n-1, 0) + s(n-1, -1),$$

where $s(.,.)$ are the Stirling numbers of the first kind.

[182]

$$(0.74) \quad \sum_{k=0}^n s(n, k) k = s(n-1, 1) + s(n-1, 0).$$

[182]

$$(0.75) \quad n! \sum_{k=0}^n s(k, m) (-1)^{n-k} / k! = s(n+1, m+1).$$

[182]

$$(0.76) \quad \sum_{k=0}^n s(n, k) k^{\underline{m}} = m! [s(n-1, m) + s(n-1, m-1)],$$

where $k^{\underline{m}} = k(k-1)(k-2) \cdots (k-m+1)$.

[182]

$$(0.77) \quad \sum_{k=0}^n s(n, k) k^m = \sum_{j=0}^m \left\{ \begin{matrix} m \\ j \end{matrix} \right\} (s(n-1, j) + s(n-1, j-1)) j!.$$

[182]

$$(0.78) \quad \sum_{k=0}^n \frac{s(n, k)}{k+1} = b_n,$$

where $b_n = \int_0^1 x^n dx$ are the Cauchy numbers of the first type [175, A006232].

[182]

$$(0.79) \quad n! \sum_{k=0}^n \left[\begin{matrix} k \\ m \end{matrix} \right] \frac{1}{k!} = \left[\begin{matrix} n+1 \\ m+1 \end{matrix} \right].$$

where $[\cdots]$ are the unsigned Stirling numbers of the first kind.
[\[182\]](#)

$$(0.80) \quad \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right] k = \left[\begin{matrix} n+1 \\ 2 \end{matrix} \right].$$

[\[182\]](#)

$$(0.81) \quad \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right] k^{\underline{m}} = \left[\begin{matrix} n+1 \\ m+1 \end{matrix} \right] m!,$$

where $k^{\underline{m}} = k(k-1)(k-2)\cdots(k-m+1)$.
[\[182\]](#)

$$(0.82) \quad \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right] k^m = \sum_{j=0}^m \left[\begin{matrix} n+1 \\ j+1 \end{matrix} \right] \left\{ \begin{matrix} m \\ j \end{matrix} \right\} j!.$$

[\[182\]](#)

$$(0.83) \quad \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right] \frac{1}{k+1} = c_n,$$

where $c_n = \int_0^1 (x)_n dx$ are the Cauchy numbers of the second type [\[175, A002657\]](#),
 using Pochhammer's symbol.
[\[48\]](#)

$$(0.84) \quad \sum_{k=0}^m \binom{m}{k} (n+k)! \left\{ \begin{matrix} n+k+s \\ q \end{matrix} \right\} = \sum_{k=0}^n \binom{n}{k} (m+k)! (-1)^{n-k} \frac{(m+k+s)^q}{(m+k+s)!} \\ + \sum_{j=0}^{s-1} \sum_{i=0}^{s-1-j} \binom{s-1-j}{i} \frac{(-1)^{n+1+i} \left\{ \begin{matrix} j \\ q \end{matrix} \right\}}{(s-1-j)!(m+n+1+i) \binom{m+n+i}{n}}$$

and

$$(0.85) \quad \sum_{k=0}^m (n+k)! \left\{ \begin{matrix} n+k-s \\ q \end{matrix} \right\} = \sum_{k=0}^n \binom{n}{k} (m+k)! (-1)^{n-k} \frac{(m+k-s)^q}{(m+k-s)!}.$$

[\[138\]](#)

$$(0.86) \quad \left\{ \begin{matrix} n \\ m \end{matrix} \right\}_r = \sum_{k=2}^n \binom{n}{k} \sum_{l=1}^{k-1} (-1)^{l-1} \binom{l+r-2}{l-1} \left\{ \begin{matrix} k-l \\ m-1 \end{matrix} \right\}_{r-1}$$

where $\{\}_r$ are r -Stirling numbers of the second kind, namely

$$(0.87) \quad \left\{ \begin{matrix} n \\ m \end{matrix} \right\}_r = \begin{cases} 0, & n < r \\ \delta_{mr}, & n = r \\ m \left\{ \begin{matrix} n-1 \\ m \end{matrix} \right\}_r + \left\{ \begin{matrix} n-1 \\ m-1 \end{matrix} \right\}_r, & n > r. \end{cases}$$

0.2. **Numerical Series.** [56, 52][175, A152649]

$$(0.88) \quad \sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^3} = \frac{1}{2}\zeta^2(2),$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.
[52]

$$(0.89) \quad \sum_{n=1}^{\infty} \frac{H_n^{(1)^2}}{n^4} = \frac{97}{24}\zeta(6) - 2\zeta^2(3) \approx 1.22187994531988.$$

[52]

$$(0.90) \quad \sum_{n=1}^{\infty} \frac{H_n^{(1)^3}}{n^3} = \zeta^2(3) - \frac{1}{3}\zeta(6) \approx 1.1058264444388.$$

[56]

$$(0.91) \quad \sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^{2p+1}} = \frac{1}{2} \sum_{j=2}^{2p} (-1)^j \zeta(j) \zeta(2p-j+2),$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.
[35, 137, 179]

$$(0.92) \quad \sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^m} = \frac{m+2}{2} \zeta(m+1) - \frac{1}{2} \sum_{k=1}^{m-2} \zeta(m-k) \zeta(k+1),$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.
[179]

$$(0.93) \quad \sum_{n=1}^{\infty} \frac{H_n}{n(n+\alpha)} = \frac{1}{2\alpha} [3\zeta(2) + \psi^2(\alpha) + 2\gamma\psi(\alpha) + \gamma^2 - \psi'(x)],$$

and

$$(0.94) \quad \sum_{n=1}^{\infty} \frac{H_n}{(n+\alpha)^2} = \gamma\psi'(\alpha) + \psi(\alpha)\psi'(\alpha) - \frac{1}{2}\psi''(\alpha),$$

and

$$(0.95) \quad \sum_{n=1}^{\infty} \frac{H_n}{(n+x)^q} = \frac{(-)^q}{(q-1)!} \left[(\psi(x) + \gamma) \psi^{(q-1)}(x) - \frac{1}{2} \psi^{(q)}(x) + \sum_{m=1}^{q-2} \binom{q-2}{m} \psi^{(m)} \psi^{(q-m-1)}(x) \right].$$

The paper also demonstrates a finite expansion of $\sum_{n \geq 1} H_n / [n^q \binom{an+k}{k}^t]$ in terms of ζ and ψ functions for $t = 1$ and 2.

[137]

$$(0.96) \quad S(r, m) = S(1, m) + \sum_{k=1}^{r-1} \frac{S(k, m-1) - B(k, m)}{k}.$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$, where $S(r, m) \equiv \sum_{n=1}^{\infty} \frac{H_n^{(r)}}{n^m}$, where $B(k, m) \equiv {}_{m+1}F_m(1, 1, \dots, 1, k+1; 2, 2, \dots, 2; 1)$.

[137]

$$(0.97) \quad S(2, 3) = \frac{\pi^4}{72} - \frac{\pi^2}{6} + 2\zeta(3) \approx 2.1120837816098848.$$

[137]

$$(0.98) \quad S(2, 4) = \frac{\pi^4}{72} + 3\zeta(5) - \zeta(3) \left(1 + \frac{\pi^2}{6}\right).$$

[21, 52]

$$(0.99) \quad \sum_{k=1}^{\infty} \frac{H_k^{(1)2}}{k^2} = \frac{17}{4}\zeta(4).$$

[21]

$$(0.100) \quad \sum_{k=1}^{\infty} \frac{H_k^{(1)2}}{(k+1)^n} = \frac{1}{3}n(n+1)\zeta(n+2) + \zeta(2)\zeta(n) - \frac{1}{n} \sum_{k=0}^{n-2} \zeta(n-k)\zeta(k+2) \\ + \frac{1}{3} \sum_{k=2}^{n-2} \zeta(n-k) \sum_{j=1}^{k-1} \zeta(j+1)\zeta(k+1-j) + \sum_{k=1}^{\infty} \frac{H_k^{(2)}}{(k+1)^n}.$$

[21]

$$(0.101) \quad \sum_{k=1}^{\infty} \frac{H_k^{(2)}}{(k+1)^2} = \frac{1}{2}\zeta^2(2) - \frac{1}{2}\zeta(4) = \frac{1}{120}\pi^4.$$

[167, 21][175, A214508]

$$(0.102) \quad A_4 \equiv \sum_{k=1}^{\infty} (-)^{k+1} \frac{1}{(k+1)^2} H_k^{(2)} = -4 \operatorname{Li}_4(1/2) + \frac{13\pi^4}{288} + \log(2) \left[-\frac{7}{2}\zeta(3) + \frac{\pi^2}{6} \log 2 - \frac{\log^3 2}{6} \right].$$

[21]

$$(0.103) \quad \sum_{k=1}^{\infty} \frac{H_k^{(2)}}{(k+1)^4} = -6\zeta(6) + \frac{8}{3}\zeta(2)\zeta(4) + \zeta^2(3) = \zeta^2(3) - \frac{4}{2835}\pi^6.$$

[21]

$$(0.104) \quad \sum_{k=1}^{\infty} \frac{H_k^{(1)}}{(k+1)^n} = \frac{1}{2}n\zeta(n+1) - \frac{1}{2} \sum_{k=1}^{n-2} \zeta(n-k)\zeta(k+1).$$

[21]

$$(0.105) \quad \sum_{k=1}^{\infty} \frac{H_k^{(2)}}{(k+1)^{2n-1}} = -\frac{1}{2}(2n^2+n+1)\zeta(2n+1) + \zeta(2)\zeta(2n-1) + \sum_{k=1}^{n-1} 2k\zeta(2k+1)\zeta(2n-2k).$$

[21]

(0.106)

$$\sum_{k=1}^{\infty} \frac{H_k^{(1)2}}{(k+1)^{2n-1}} = \frac{1}{6}(2n^2-7n-3)\zeta(2n+1) + \zeta(2)\zeta(2n-1) - \frac{1}{2} \sum_{k=1}^{n-2} (2k-1)\zeta(2n-1-2k)\zeta(2k+2) \\ + \frac{1}{3} \sum_{k=1}^{n-2} \zeta(2k+1) \sum_{j=1}^{n-2-k} \zeta(2j+1)\zeta(2n-1-2k-2j).$$

[138]

(0.107)

$$\sum_{n=1}^{\infty} \frac{H_n^{(2)}}{n!} = e \left[1 + \frac{1}{4} {}_2F_2 \left(\begin{matrix} 1 & 1 \\ 3 & 3 \end{matrix} \middle| -1 \right) \right],$$

(0.108)

$$\sum_{n=1}^{\infty} \frac{H_n^{(4)}}{2^n n!} = \sqrt{e} \left[\frac{H_1^{(3)}}{2^1 1!} + \frac{H_2^{(2)}}{2^2 2!} + \frac{H_3^{(1)}}{2^3 3!} + \frac{3!}{2^4 (4!)^2} {}_2F_2 \left(\begin{matrix} 1 & 1 \\ 5 & 5 \end{matrix} \middle| -\frac{1}{2} \right) \right],$$

where $H_n^{(r)}$ are hyperharmonic numbers (1.77).

[74]

(0.109)

$$\sum_{-\infty}^{\infty} \frac{1}{n^2 + q^2} = \frac{\pi \coth \pi q}{q}.$$

[74]

(0.110)

$$\sum_{-\infty}^{\infty} \frac{1}{(n^2 + q_1^2)((N-n)^2 + q_2^2)} = \frac{2\pi}{2q_1 2q_2} \left[\frac{1 + n_b(q_1) + n_b(q_2)}{Ni + q_1 + q_2} + \frac{n_b(q_1) - n_b(q_2)}{Ni - q_1 + q_2} \right. \\ \left. - \frac{n_b(q_1) - n_b(q_2)}{Ni + q_1 - q_2} - \frac{1 + n_b(q_1) + n_b(q_2)}{Ni - q_1 - q_2} \right]$$

where $n_b(z) \equiv 1/(e^{2\pi z} - 1)$.

[74] Define the Matsubara sum associated to the Graph G (loopless multigraph such that the degree of each vertex is at least 2, with I lines) by

(0.111)

$$S_G \equiv \sum_{n_1=-\infty}^{\infty} \sum_{n_2=-\infty}^{\infty} \cdots \sum_{n_I=-\infty}^{\infty} \frac{\delta_g(n_1, n_2, \dots, n_I; \{N_v\})}{(n_1^2 + q_1^2)(n_2^2 + q_2^2) \cdots (n_I^2 + q_I^2)},$$

where

(0.112)

$$\delta_g(n_1, \dots; \{N_v\}) = \prod_{v=1}^V \delta_{T_v, N_v}$$

imposes a series of constraints over the vertices v , and $T_v \equiv \sum_i s_i^v n_i$ is an algebraic sum at vertex v , with s_i^v having values ± 1 or 0 depending on the orientation of the line i with respect to the vertex v . The q_i are weights associated with the lines i . Then the integral

(0.113)

$$I(N, q_1, q_2, \dots) \int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \cdots \frac{1}{(x_1^2 + q_1^2)(x_2^2 + q_2^2) \cdots ((N - x_1 - x_2 \cdots)^2 + q_I^2)}$$

is related to the sum via

$$(0.114) \quad S_G = \hat{O}_G I_G$$

where the operator $\hat{O}_G = \prod_{i=1}^I [1 + n_{b_i} (1 - \hat{R}_i)]$ is composed of the functions n_b of the previous formula and the reflection operator \hat{R}_i (which switches the sign of the variable q_i).

[175, A152416][126]

$$(0.115) \quad \sum_{n=2}^{\infty} \frac{1}{n^s(n-1)} = s - \sum_{l=2}^s \zeta(l).$$

[36]

$$(0.116) \quad \sum_{n \geq 1} \frac{1}{n(n^2+1)} = \gamma + \Re \psi(1+i) \approx 0.67186598552400983.$$

[36]

$$(0.117) \quad \sum_{n \geq 1} \frac{1}{n^2(n^2+1)} = \frac{\pi^2}{6} - \frac{\pi \coth \pi - 1}{2} \approx 0.56826001937964526.$$

[132]

$$(0.118) \quad \sum_{k=1}^{\infty} \frac{1}{(2k)^{2s}(2k+1)^{2s}} = \sum_{t=1}^{2s} \binom{4s-t-1}{2s-1} \left\{ \left[1 - \frac{1-(-)^t}{2^t} \right] \zeta(t) - 1 \right\}.$$

[132]

$$(0.119) \quad \sum_{k=1}^{\infty} \frac{1}{(2k)^2(2k+1)^2} = -3 + \frac{\pi^2}{6} + 2 \log 2.$$

[132]

$$(0.120) \quad \sum_{k=1}^{\infty} \frac{1}{(2k)^4(2k+1)^4} = -35 + \frac{\pi^4}{90} + 3\zeta(3) + \frac{5\pi^2}{3} + 20 \log 2.$$

[132]

$$(0.121) \quad \sum_{n=1}^{\infty} \frac{1}{n^{2s}(n+1)^{2s}} = \sum_{t=1}^{2s} \binom{4s-t-1}{2s-1} \{ [1 + (-)^t] \zeta(t) - 1 \}.$$

[132]

$$(0.122) \quad \sum_{n=1}^{\infty} \frac{1}{n^2(n+1)^2} = \frac{\pi^2}{3} - 3.$$

[132]

$$(0.123) \quad \sum_{n=1}^{\infty} \frac{1}{n^4(n+1)^4} = -35 + \frac{10\pi^2}{3} + \frac{\pi^4}{45}.$$

[119, (1)]

$$(0.124) \quad \sum_{n=0}^{\infty} \frac{\binom{2n}{n}}{8^n} = \sqrt{2}.$$

[119, (1)]

$$(0.125) \quad \sum_{n=0}^{\infty} \frac{\binom{2n}{n}}{10^n} = \sqrt{5/3}.$$

[119, (1)]

$$(0.126) \quad \sum_{n=0}^{\infty} \frac{(-1)^n \binom{2n}{n}}{8^n} = \sqrt{2/3}.$$

[119][175, A145439]

$$(0.127) \quad \sum_{n=0}^{\infty} \frac{\binom{4n}{2n}}{64^n} = \frac{3\sqrt{2} + \sqrt{6}}{6}$$

[119]

$$(0.128) \quad 4 \sum_{n=0}^{\infty} \frac{\binom{8n}{4n}}{8^{4n}} = \frac{3\sqrt{2} + \sqrt{6}}{3} + \frac{2\sqrt{2 + \sqrt{5}}}{\sqrt{5}}.$$

[119, (6)]

$$(0.129) \quad \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n4^n} = \log 4.$$

[119, (6)][175, A157699]

$$(0.130) \quad \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \binom{2n}{n}}{n4^n} = 2 \log \frac{1 + \sqrt{2}}{2},$$

The formula above corrects a factor 2 in [119], see [127].

[175, A091648]

$$(0.131) \quad \sum_{n=1,3,5,7,\dots}^{\infty} \frac{\binom{2n}{n}}{n4^n} = \log(1 + \sqrt{2}).$$

[119, (7)]

$$(0.132) \quad \sum_{n=1}^{\infty} \frac{\binom{2n}{n}}{n(n+1)4^n} = \log 4 - 1.$$

[119, (8)]

$$(0.133) \quad \sum_{n=1}^{\infty} \frac{n \binom{2n}{n}}{8^n} = 1/\sqrt{2}.$$

[119]

$$(0.134) \quad \sum_{n=1}^{\infty} \frac{n^2 \binom{2n}{n}}{8^n} = \frac{5\sqrt{2}}{4}.$$

[119][175, A019670]

$$(0.135) \quad \sum_{n=0}^{\infty} \frac{\binom{2n}{n}}{(2n+1)16^n} = \frac{\pi}{3}.$$

[175, A086466]

$$(0.136) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{m \binom{2m}{m}} = \frac{2}{\sqrt{5}} \log \frac{1+\sqrt{5}}{2}.$$

This corrects a factor 2 in [119] and two typos in [17, 4.1.42], see [127].
[119, (15)] [66, 189][175, A073016]

$$(0.137) \quad \sum_{m=1}^{\infty} \frac{1}{\binom{2m}{m}} = \frac{9+2\pi\sqrt{3}}{27}.$$

[189]

$$(0.138) \quad \sum_{k=0}^{\infty} \frac{1}{\binom{mk}{nk}} = \int_0^1 \frac{1+(m-1)t^n(1-t)^{m-n}}{(1-t^n(1-t)^{m-n})^2} dt,$$

where m and n are positive integers with $m > n$.

[189]

$$(0.139) \quad \sum_{k=0}^{\infty} \frac{1}{\binom{4k}{2k}} = \frac{16}{15} + \frac{\pi\sqrt{3}}{27} - \frac{2\sqrt{5}}{25} \ln \frac{1+\sqrt{5}}{2}.$$

[119][175, A086465]

$$(0.140) \quad \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\binom{2n}{n}} = \frac{1}{5} + \frac{4\sqrt{5}}{25} \log \frac{1+\sqrt{5}}{2}.$$

[119, 66][175, A145429]

$$(0.141) \quad \sum_{m=1}^{\infty} \frac{m}{\binom{2m}{m}} = \frac{2}{27}(\pi\sqrt{3}+9).$$

[39] And by linear combination with (0.137):

$$(0.142) \quad \sum_{n \geq 1} \frac{18-9n}{\binom{2n}{n}} = 2 \frac{\pi}{\sqrt{3}}.$$

[119, 66]

$$(0.143) \quad \sum_{m=1}^{\infty} \frac{m^2}{\binom{2m}{m}} = \frac{2}{81}(5\pi\sqrt{3}+54).$$

[119, 66]

$$(0.144) \quad \sum_{m=1}^{\infty} \frac{m^3}{\binom{2m}{m}} = \frac{2}{243}(37\pi\sqrt{3}+405).$$

[119][175, A145433]

$$(0.145) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1}m}{\binom{2m}{m}} = \frac{2}{125}(2\sigma+15), \quad \sigma \equiv \sqrt{5} \log \frac{1+\sqrt{5}}{2}.$$

[119]

$$(0.146) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1}m^2}{\binom{2m}{m}} = \frac{2}{125}(5-\sigma).$$

Erratum to [17, 4.1.40]:

$$(0.147) \quad \sum_{n=0}^{\infty} \frac{(-1)^{n-1} n^2}{\binom{2n}{n}} = \frac{4}{125} \left[5 - \sqrt{5} \ln \left(\frac{1 + \sqrt{5}}{2} \right) \right].$$

$$(0.148) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1} m^3}{\binom{2m}{m}} = \frac{2}{625} (14\sigma - 5),$$

which corrects a factor 2 in [119], see [127].

[119, 66]

$$(0.149) \quad \sum_{m=1}^{\infty} \frac{2^m}{m^2 \binom{2m}{m}} = \frac{\pi^2}{8},$$

a special case of (1.17).

[119]

$$(0.150) \quad \sum_{m=1}^{\infty} \frac{2^m}{m \binom{2m}{m}} = \frac{\pi}{2}.$$

[119]

$$(0.151) \quad \sum_{m=1}^{\infty} \frac{2^m}{\binom{2m}{m}} = \frac{\pi}{2} + 1.$$

[119]

$$(0.152) \quad \sum_{m=1}^{\infty} \frac{m 2^m}{\binom{2m}{m}} = \pi + 3.$$

[119, 66]

$$(0.153) \quad \sum_{m=1}^{\infty} \frac{m^2 2^m}{\binom{2m}{m}} = \frac{1}{5} \sum_{m=1}^{\infty} \frac{m^3 2^m}{\binom{2m}{m}} = \frac{7\pi}{2} + 11.$$

[119, 66]

$$(0.154) \quad \sum_{m=1}^{\infty} \frac{m^4 2^m}{\binom{2m}{m}} = 113\pi + 355.$$

[119, 66]

$$(0.155) \quad \sum_{m=1}^{\infty} \frac{m^{10} 2^m}{\binom{2m}{m}} = 229093376\pi + 719718067.$$

[119]

$$(0.156) \quad \sum_{m=1}^{\infty} \frac{3^m}{m^2 \binom{2m}{m}} = \frac{2\pi^2}{9},$$

a special case of (1.17).

[119][175, A186706]

$$(0.157) \quad \sum_{m=1}^{\infty} \frac{3^m}{m \binom{2m}{m}} = \frac{2\pi}{\sqrt{3}} \equiv \nu.$$

[119]

$$(0.158) \quad \sum_{m=1}^{\infty} \frac{3^m}{\binom{2m}{m}} = 2\nu + 3.$$

[119]

$$(0.159) \quad \sum_{m=1}^{\infty} \frac{m3^m}{\binom{2m}{m}} = 10\nu + 18.$$

[119]

$$(0.160) \quad \sum_{m=1}^{\infty} \frac{m^2 3^m}{\binom{2m}{m}} = 2(43\nu + 78).$$

[119]

$$(0.161) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1} 2^m}{m \binom{2m}{m}} = \rho/3, \quad \rho \equiv \sqrt{3} \log(2 + \sqrt{3}).$$

[12]

$$(0.162) \quad \sum_{m=1}^{\infty} \frac{(2t)^{2m+2k}}{m(2m+2k) \binom{2m}{m}} = \arcsin^2(t) \binom{2k}{k} + \sum_{j=1}^k \binom{2k}{k-j} \frac{(-)^{j+1}}{j^2} \\ + \sum_{j=1}^k (-)^j \binom{2k}{k-j} \left(\frac{2 \arcsin t \sin(2j \arcsin t)}{j} + \frac{\cos(2j \arcsin t)}{j^2} \right),$$

and a similar logarithmic result for an alternating sign sum on the left hand side.

[12]

(0.163)

$$\sum_{m=1}^{\infty} \frac{(2t)^{2m+2k}}{m^2(2m+2k) \binom{2m}{m}} = -\frac{1}{k} \arcsin^2(t) + \frac{(2t)^{2k}}{k} \arcsin^2 t - \sum_{j=1}^k \binom{2k}{k-j} \frac{(-)^{j+1}}{kj^2} \\ - \sum_{j=1}^k (-)^j \binom{2k}{k-j} \left(\frac{2 \arcsin t \sin(2j \arcsin t)}{kj} + \frac{\cos(2j \arcsin t)}{kj^2} \right),$$

and a similar logarithmic result for an alternating sign sum on the left hand side.

[119]

$$(0.164) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1} 2^m}{\binom{2m}{m}} = \frac{\rho + 3}{9}.$$

[119]

$$(0.165) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1} m 2^m}{\binom{2m}{m}} = \frac{1}{3}.$$

[119]

$$(0.166) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1} m^2 2^m}{\binom{2m}{m}} = \frac{1}{27}(3 - \rho).$$

[119]

$$(0.167) \quad \sum_{m=1}^{\infty} \frac{(-1)^m m^3 2^m}{\binom{2m}{m}} = \frac{1}{81}(\rho + 15).$$

[119]

$$(0.168) \quad \sum_{m=1}^{\infty} \frac{(2 - \sqrt{2})^m}{\binom{2m}{m}} = \frac{3 - 2\sqrt{2}}{4}(\pi\sqrt{2} + 4).$$

[119]

$$(0.169) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1} 3^{2m}}{4^m \binom{2m}{m}} = \frac{48}{125}(\log 2 + \frac{15}{16}).$$

[119][175, A152422]

$$(0.170) \quad \sum_{m=1}^{\infty} \frac{2^m (2 - \sqrt{3})^m}{m^2 \binom{2m}{m}} = 2(\arcsin \tau)^2, \quad \tau = \frac{\sqrt{3} - 1}{2} = \sqrt{2} \sin \frac{\pi}{12} = \sin \frac{\pi}{3} - \sin \frac{\pi}{6},$$

where both right hand sides in [119] are erroneous, see [127].

[39]

$$(0.171) \quad \sum_{n \geq 0} \frac{50n - 6}{\binom{3n}{n} 2^n} = \pi.$$

[39]

$$(0.172) \quad \sum_{n \geq 1} \frac{1}{\binom{3n}{n} 2^n} = \frac{2}{25} - \frac{6}{125} \ln 2 + \frac{11}{250} \pi.$$

[39]

$$(0.173) \quad \sum_{n \geq 1} \frac{n}{\binom{3n}{n} 2^n} = \frac{81}{625} - \frac{18}{3125} \ln 2 + \frac{79}{3125} \pi.$$

[39]

$$(0.174) \quad \sum_{n \geq 1} \frac{n^2}{\binom{3n}{n} 2^n} = \frac{561}{3125} + \frac{42}{15625} \ln 2 + \frac{673}{31250} \pi.$$

[39]

$$(0.175) \quad \sum_{n \geq 1} \frac{-150n^2 + 230n - 36}{\binom{3n}{n} 2^n} = \pi.$$

[39]

$$(0.176) \quad \sum_{n \geq 1} \frac{575n^2 - 965n + 273}{\binom{3n}{n} 2^n} = 6 \log 2.$$

[39]

$$(0.177) \quad \sum_{n \geq 1} \frac{(-1)^n}{\binom{3n}{n} 4^n} = -\frac{1}{28} - \frac{3}{32} \ln 2 + \frac{13}{112} \frac{\arctan(\sqrt{7/5})}{\sqrt{7}}.$$

[39]

$$(0.178) \quad \sum_{n \geq 1} \frac{(-1)^n n}{\binom{3n}{n} 4^n} = -\frac{81}{1568} - \frac{9}{256} \ln 2 + \frac{17}{6272} \frac{\arctan(\sqrt{7/5})}{\sqrt{7}}.$$

[39]

$$(0.179) \quad \sum_{n \geq 1} \frac{1}{n \binom{3n}{n} 2^n} = \frac{1}{10} \pi - \frac{1}{5} \ln 2.$$

[39]

$$(0.180) \quad \sum_{n \geq 1} \frac{1}{n^2 \binom{3n}{n} 2^n} = \frac{1}{24} \pi^2 - \frac{1}{2} \ln^2 2.$$

[34]

$$(0.181) \quad \sum_{j=1}^{\infty} \frac{(-1)^j}{j!} (j/2)! = 1 - \frac{1}{2} e^{1/4} \sqrt{\pi} (1 - \operatorname{erf}(1/2)).$$

Erratum to [17, 4.1.47][175, A145438]: Using twice 9.14.+13, then [91, 9.121.6] and [91, 9.121.26], then substituting $tt'/4 = z^2$, then $y = \sqrt{t'/4}$, then $\arcsin y = v$

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n^3 \binom{2n}{n}} &= \frac{1}{2} {}_4F_3(1, 1, 1, 1; 2, 2, 3/2; 1/4) = \frac{1}{2} \int_0^1 dt {}_3F_2(1, 1, 1; 2, 3/2; t/4) \\ &= \frac{1}{4} \int_0^1 dt \int_0^1 dt' {}_2F_1(1, 1; 3/2; tt'/4) = \frac{1}{4} \int_0^1 dt \int_0^1 dt' \frac{1}{\sqrt{1-tt'/4}} {}_2F_1(1/2, 1/2; 3/2; tt'/4) \\ &= \frac{1}{4} \int_0^1 dt \int_0^1 dt' \frac{1}{\sqrt{1-tt'/4}} \frac{\arcsin(\sqrt{tt'/4})}{\sqrt{tt'/4}} = \int_0^1 \frac{dt'}{t'} \int_0^{\sqrt{t'/4}} dz \frac{\arcsin z}{\sqrt{1-z^2}} = \frac{1}{2} \int_0^1 \frac{dt'}{t'} \arcsin^2(\sqrt{t'/4}) \\ &= \int_0^{\sqrt{1/2}} \frac{dy}{y} \arcsin^2 y = \int_0^{\sin \sqrt{1/2}} v^2 \cot v dv. \end{aligned}$$

This becomes [39, (35)][119][38, Theorem 3.3]

(0.182)

$$\frac{2\pi}{3} \Im \operatorname{Li}_2(e^{i\pi/4}) - \frac{4}{3} \zeta(3) = \frac{2\pi}{3} \operatorname{Cl}_2(\pi/3) - \frac{4}{3} \zeta(3) = \frac{\pi\sqrt{3}}{18} \{\psi'(1/3) - \psi'(2/3)\} - \frac{4\zeta(3)}{3}.$$

So the r.h.s. of [119] is missing a factor 4, and [15, 4.1.47] is in addition misleading to imply that the digamma (instead of the trigamma) functions are in effect [127].

[178]

$$(0.183) \quad \sum_{n=1}^{\infty} \frac{t^n}{n^{k+1} \binom{an+j+1}{j+1}} = \begin{cases} \frac{(j+1)t(-)^k}{k!} \int_0^1 \int_0^1 \frac{(1-x)^j x^a (\ln y)^k}{1-tx^a y} dx dy, & k \geq 1 \\ at \int_0^1 \frac{(1-x)^{j+1} x^{a-1}}{1-tx^a} dx, & k = 1 \end{cases}$$

$$= T_{0 \ a+k+1} F_{a+k} \left(\begin{matrix} 1, 1, \dots, 1; (a+1)/a, \dots, (2a-1)/a \\ 2, 2, \dots, 2; (a+j+2)/a, \dots, (a+j+a+1)/a \end{matrix} \middle| t \right)$$

where $T_0 = t(j+1)B(j+1, a+1)$.

[67]

(0.184)

$$\frac{1}{2^m} \sum_{i=1}^{\infty} \binom{i+1}{2}^{-m} = (-)^{m-1} \binom{2m-1}{m} + (-)^m 2 \sum_{i=1}^{\lfloor m/2 \rfloor} \binom{2m-2i-1}{m-1} \zeta(2i).$$

[39, 119][175, A073010]

$$(0.185) \quad \sum_{n \geq 1} \frac{1}{n \binom{2n}{n}} = \frac{\pi}{3\sqrt{3}}.$$

$$(0.186) \quad \sum_{n \geq 1} \frac{1}{n^2 \binom{2n}{n}} = \frac{1}{3} \zeta(2).$$

a special case of (1.17).

[94]

$$(0.187) \quad \sum_{n \geq 1} \frac{1}{n^4 \binom{2n}{n}} = \frac{17}{36} \zeta(4).$$

[94]

$$(0.188) \quad \sum_{n \geq 1} \frac{1}{n^5 \binom{2n}{n}} = 2\pi \operatorname{Cl}_4(\pi/3) - \frac{19}{3} \zeta(5) + \frac{2}{3} \zeta(3) \zeta(2).$$

[94]

$$(0.189) \quad \sum_{n \geq 1} \frac{1}{n^6 \binom{2n}{n}} = -\frac{4\pi}{3} \Im L_{4,1}(e^{i\pi/3}) + \frac{3341}{1296} \zeta(6) - \frac{4}{3} \zeta^2(3).$$

[94]

$$(0.190) \quad \sum_{n \geq 1} \frac{(-1)^n}{n \binom{2n}{n}} = -2 \frac{\operatorname{arctanh}(1/\sqrt{5})}{\sqrt{5}}.$$

Correction of a sign error in [119], see [127]:

$$(0.191) \quad \sum_{n \geq 1} \frac{(-1)^n}{n^2 \binom{2n}{n}} = -2 \left(\ln \frac{\sqrt{5}-1}{2} \right)^2 = -2 \left(\ln \frac{\sqrt{5}+1}{2} \right)^2.$$

$$(0.192) \quad \sum_{n \geq 1} \frac{(-1)^n}{n^3 \binom{2n}{n}} = -\frac{2}{5} \zeta(3).$$

(0.193)

$$\sum_{n \geq 1} \frac{(-1)^n}{n^4 \binom{2n}{n}} = -4K_4(\rho) + 4K_4(-\rho) + \frac{1}{2} \ln^4 \rho + 7\zeta(4), \quad K_k(x) \equiv \sum_{r=0}^{k-1} \frac{(-\ln|x|)^r}{r!} L_{k-r}(x).$$

$$(0.194) \quad \sum_{n \geq 1} \frac{1}{n^3 \binom{3n}{n} 2^n} = -\frac{33}{16} \zeta(3) + \frac{1}{6} \ln^3 2 - \frac{1}{24} \pi^2 \ln 2 + \pi \Im L_2(i).$$

(0.195)

$$\sum_{n \geq 1} \frac{1}{n^4 \binom{3n}{n} 2^n} = -\frac{143}{16} \zeta(3) \ln 2 + \frac{91}{640} \pi^4 - \frac{3}{8} \ln^4 2 + \frac{3}{8} \pi^2 \ln^2 2 - 8L_4(1/2) - 8\Re L_{3,1}\left(\frac{1+i}{2}\right) - 8\Re L_4\left(\frac{1+i}{2}\right).$$

$$\begin{aligned} \sum_{n \geq 1} \frac{(-1)^n}{8^n n \binom{6n}{2n}} &= \left(-\frac{1}{3} + \frac{2}{57} \sqrt{114\sqrt{57}-342} \right) \ln 2 - \frac{1}{114} \sqrt{114\sqrt{57}-342} \ln \left(13 + \sqrt{57} + \sqrt{-30+26\sqrt{57}} \right) \\ &\quad + \frac{1}{57} \sqrt{114\sqrt{57}+342} \arctan \frac{\sqrt{2\sqrt{57}+9}}{7}. \end{aligned}$$

$$(0.196) \quad \sum_{n \geq 1} \frac{(-1)^n}{2^n n \binom{4n}{n}} = \sum_{13+12r+2r^3=0} \frac{\ln(r+2)}{r+3} - \frac{6}{7} \ln 3 + \frac{1}{7} \ln 2.$$

$$(0.197) \quad \sum_{n \geq 1} \frac{(-1)^n}{n \binom{3n}{n}} = \sum_{8+4r+r^3=0} \frac{\ln(r+2)}{r+3} - \ln 2.$$

[206] Let $A_k \equiv \sum_{n=0}^{\infty} \binom{n+k}{n} / \binom{2n}{n}$, then

$$(0.198) \quad A_{k+1} = -\frac{2}{3(k+1)} + \frac{2(2k+3)}{3(k+1)} A_k.$$

[206] Let $B_k \equiv \sum_{n=0}^{\infty} (-1)^n \binom{n+k}{n} / \binom{2n}{n}$, then

$$(0.199) \quad B_{k+1} = -\frac{2}{5(k+1)} + \frac{2(2k+3)}{5(k+1)} B_k.$$

and similar recurrences with an additional factor n or n^2 in the denominator of the sums.

[33]

$$(0.200) \quad \sum_{k=1}^{\infty} \arctan \frac{2}{k^2} = \frac{3\pi}{4}.$$

[33][175, A091007]

$$(0.201) \quad \sum_{k=1}^{\infty} \arctan \frac{1}{k^2} = \arctan \frac{\tan(\pi/\sqrt{2}) - \tanh(\pi/\sqrt{2})}{\tan(\pi/\sqrt{2}) + \tanh(\pi/\sqrt{2})}.$$

[33]

$$(0.202) \quad \sum_{k=1}^{\infty} \arctan \frac{a}{a^2 k^2 + a(a+2b)k + 1 + ab + b^2} = \frac{\pi}{2} - \arctan(a+b).$$

[33]

$$(0.203) \quad \sum_{k=1}^{\infty} \arctan \frac{a^2 k^2 + a^2 k - 1 - b^2}{a^4 k^4 + 2a^3(a+2b)k^3 + a^2(2+a^2+6ab+6b^2)k^2 + 2a(a+2b)(1+ab+b^2) + (1+b^2)(1+[a+b]^2)} = \frac{1}{1+(a+b)^2}.$$

[33]

$$(0.204) \quad \sum_{k=1}^{\infty} \arctan \frac{2ak + a + b}{a_4 k^4 + a_3 k^3 + a_2 k^2 + a_1 k + a_0} = \frac{\pi}{2} - \arctan(a+b+c).$$

[33]

$$(0.205) \quad \sum_{k=1}^n \arctan \frac{f(k+1) - f(k-1)}{1 + f(k+1)f(k-1)} = \arctan f(n+1) - \arctan f(1) + \arctan f(n) - \arctan f(0).$$

[33]

$$(0.206) \quad \sum_{k=1}^{\infty} \arctan \frac{8k}{k^4 - 2k^2 + 5} = \pi - \arctan \frac{1}{2}.$$

[33]

$$(0.207) \quad \sum_{k=1}^{\infty} \arctan \frac{4ak}{k^4 + a^2 + 4} = \arctan \frac{a}{2} + \arctan a.$$

[33]

$$(0.208) \quad \sum_{k=1}^{\infty} \arctan \frac{2xy}{k^2 - x^2 + y^2} = \arctan \frac{y}{x} - \arctan \frac{\tanh \pi y}{\tan \pi x}.$$

[33]

$$(0.209) \quad \sum_{k=1}^{\infty} \arctan \frac{1}{2k^2} = \pi/4.$$

[33]

$$(0.210) \quad \sum_{k=1}^{\infty} \frac{k^2}{k^4 + 4x^4} = \frac{\pi}{4x} \frac{\sin 2\pi x - \sinh 2\pi x}{\cos 2\pi x - \cosh 2\pi x}.$$

[33]

$$(0.211) \quad \sum_{k=1}^{\infty} \frac{k^2}{k^4 + 4} = \frac{\pi}{4} \coth \pi.$$

[33]

$$(0.212) \quad \sum_{k=1}^{\infty} \frac{k^2}{k^4 + x^2 k^2 + x^4} = \frac{\pi}{2x\sqrt{3}} \frac{\sinh \pi x \sqrt{3} - \sqrt{3} \sin \pi x}{\cosh \pi x \sqrt{3} - \cos \pi x}.$$

[153]

$$(0.213) \quad \sum_{k=1}^{\infty} \frac{A_0 + B_0 k + C_0 k^2}{(k^2 - a^2)(k^2 - b^2)} = \sum_{n=1}^{\infty} \frac{d_n}{\prod_{m=1}^n (m^2 - a^2)(m^2 - b^2)}$$

for $|a| < 1$, $|b| < 1$, with d_n defined via a recurrence in the reference.

$$(0.214) \quad \sum_{n=0}^{\infty} n x^n = \frac{x}{(1-x)^2}.$$

$$(0.215) \quad \sum_{n=0}^{\infty} n^2 x^n = \frac{x(1+x)}{(1-x)^3}.$$

[175, A000578]

$$(0.216) \quad \sum_{n=0}^{\infty} n^3 x^n = \frac{x(1+4x+x^2)}{(1-x)^4}.$$

[175, A000583]

$$(0.217) \quad \sum_{n=0}^{\infty} n^4 x^n = \frac{x(1+x)(1+10x+x^2)}{(1-x)^5}.$$

[175, A000584]

$$(0.218) \quad \sum_{n=0}^{\infty} n^5 x^n = \frac{x(1 + 26x + 66x^2 + 26x^3 + x^4)}{(1-x)^6}.$$

[175, A001014]

$$(0.219) \quad \sum_{n=0}^{\infty} n^6 x^n = \frac{x(x+1)(1 + 56x + 246x^2 + 56x^3 + x^4)}{(1-x)^7}.$$

[175, A001015]

$$(0.220) \quad \sum_{n=0}^{\infty} n^7 x^n = \frac{x(1 + 120x + 1191x^2 + 2416x^3 + 1191x^4 + 120x^5 + x^6)}{(1-x)^8}.$$

See [175, A008292] for coefficients in the numerator polynomial if exponents of n are higher than 7. The pole at $x = 1$ in the generating functions obviously delimits the radius of convergence.

[98] Let

$$(0.221) \quad S_{a,p}(n) \equiv \sum_{k=0}^n a^k k^p;$$

then

$$(0.222) \quad S_{a,p}(n) = \frac{a^{n+1}}{a-1} \sum_{r=0}^{p-1} \binom{p}{r} f_r(a) (n+1)^{p-r} + f_p(a) \frac{a^{n+1} - 1}{a-1}$$

with

$$(0.223) \quad f_0(a) = 1, \quad a \sum_{j=0}^r \binom{r}{j} f_j(a) - f_r(a) = 0, \quad r = 1, 2, \dots$$

[6]

$$(0.224) \quad \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2} = G.$$

[6, 42]

$$(0.225) \quad 2 \sum_{k=0}^{\infty} \frac{1}{(4k+1)^2} = G + \frac{\pi^2}{8}.$$

[6, 42]

$$(0.226) \quad -2 \sum_{k=0}^{\infty} \frac{1}{(4k+3)^2} = G - \frac{\pi^2}{8}.$$

[42]

$$(0.227) \quad \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \sum_{k=0}^{\infty} \binom{n}{k} \frac{(-1)^k}{(2k+1)^2} = G.$$

[42]

$$(0.228) \quad 1 - \sum_{n=1}^{\infty} \frac{n\zeta(2n+1)}{16^n} = G.$$

[42]

$$(0.229) \quad \frac{1}{4} \sum_{n=1}^{\infty} n 16^{-n} (3^{2n} - 1) \zeta(2n+1) = G - \frac{1}{6}$$

(and similar ζ -sums).

[42]

$$(0.230) \quad \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{n+1/2}{n}^{-2} = {}_4F_3(1, 1, 1, 1; 2, \frac{3}{2}, \frac{3}{2}; 1) = 2\pi G - \frac{7}{2}\zeta(3).$$

[42]

$$(0.231) \quad 2 \sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{n^2} \sum_{k=0}^{n-1} \frac{1}{2k+1} = 2\pi G - \frac{7}{2}\zeta(3).$$

[42]

$$(0.232) \quad \sum_{n=0}^{\infty} \frac{2^n}{(2n+1) \binom{2n}{n}} \sum_{k=0}^n \frac{1}{2k+1} = 2G.$$

[42]

$$(0.233) \quad \sum_{n=0}^{\infty} \frac{4^n}{(2n+1)^2 \binom{2n}{n}} = 2G.$$

[42]

$$(0.234) \quad \frac{3}{8} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2 \binom{2n}{n}} = G - \frac{1}{8}\pi \log(2 + \sqrt{3}).$$

[42]

$$(0.235) \quad \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \sum_{k=1}^n \frac{1}{k} = G - \frac{1}{2}\pi \log 2.$$

[42]

$$(0.236) \quad -2 \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \sum_{k=0}^{n-1} \frac{1}{2k+1} = G - \frac{1}{4}\pi \log 2.$$

[42]

$$(0.237) \quad -\frac{1}{32}\pi \sum_{n=0}^{\infty} \frac{(2n+1)^2}{(n+1)^3 16^n} \binom{2n}{n}^2 = G - \frac{1}{2}\pi \log 2.$$

[42]

$$(0.238) \quad \sum_{n=0}^{\infty} \frac{\sqrt{2}}{(2n+1)^2 8^n} \binom{2n}{n} = G + \frac{1}{4}\pi \log 2.$$

[42]

$$(0.239) \quad \sum_{n=1}^{\infty} \frac{\sin(n\pi/4)}{n^2 2^{n/2}} = G - \frac{1}{8}\pi \log 2.$$

[42]

$$(0.240) \quad \sum_{n=0}^{\infty} \frac{2^{n+1}(n!)^2}{(2n+1)!(n+1)^2} = 2\pi G - \frac{35}{8}\zeta(3) + \frac{1}{4}\pi^2 \log 2.$$

[42]

$$(0.241) \quad \frac{1}{4}\pi {}_3F_2(1/2, 1/2, n+1/2; 1, n+3/2; 1) - \frac{1}{2} \sum_{k=0}^{n-1} \frac{(k!)^2}{(\frac{3}{2})_k^2} = G.$$

[76, 77] Let T denote Tornheim double sums

$$(0.242) \quad T(a, b, c) \equiv \sum_{r=1}^{\infty} \sum_{s=1}^{\infty} \frac{1}{r^a s^b (r+s)^c}$$

with $a+c > 1$, $b+c > 1$ and $a+b+c > 2$. Then

(0.243)

$$T(m, k, n) = \sum_{i=1}^m \binom{m+k-i-1}{m-i} T(i, 0, N-i) + \sum_{i=1}^k \binom{m+k-i-1}{k-i} T(i, 0, N-i).$$

$$\begin{aligned} T(m, 0, n) &= (-1)^m \sum_{j=0}^{\lfloor (n-1)/2 \rfloor} \binom{m+n-2j-1}{m-1} \zeta(2j) \zeta(m+n-2j) \\ &\quad + (-1)^m \sum_{j=0}^{\lfloor m/2 \rfloor} \binom{m+n-2j-1}{n-1} \zeta(2j) \zeta(m+n-2j) - \frac{1}{2} \zeta(m+n), \end{aligned}$$

if $N \equiv m+k+n$ is odd.

$$(0.244) \quad T(1, 0, 5) = -\frac{1}{2}\zeta(3)^2 + \frac{3}{4}\zeta(6).$$

$$(0.245) \quad T(2, 0, 4) = \zeta^2(3) - \frac{4}{3}\zeta(6).$$

$$(0.246) \quad T(3, 0, 3) = \frac{1}{2}\zeta(3)^2 - \frac{1}{2}\zeta(6).$$

$$(0.247) \quad T(4, 0, 2) = -\zeta^2(3) + \frac{25}{12}\zeta(6).$$

$$(0.248) \quad T(0, 0, N) = \zeta(N-1) - \zeta(N), \quad N \geq 3; \quad T(0, 0, 4) = \zeta(3) - \frac{\pi^4}{90}.$$

$$(0.249) \quad T(n, 0, n) = \frac{1}{2}\zeta^2(n) - \frac{1}{2}\zeta(2n). \quad T(2, 0, 2) = \frac{\pi^4}{120}.$$

$$(0.250) \quad T(0, 0, 8) = \zeta(7) - \zeta(8).$$

$$(0.251) \quad T(1, 0, 7) = \frac{5}{4}\zeta(8) - \zeta(3)\zeta(5).$$

$$(0.252) \quad T(4, 0, 4) = \frac{1}{12}\zeta(8).$$

$$(0.253) \quad T(0, 0, 14) = \zeta(13) - \zeta(14).$$

$$(0.254) \quad T(1, 0, 13) = \frac{11}{4}\zeta(14) - \zeta(3)\zeta(11) - \zeta(5)\zeta(9) - \frac{1}{2}\zeta(7)^2.$$

$$(0.255) \quad T(1, 0, 2) = \zeta(3).$$

$$(0.256) \quad T(1, 1, 1) = 2\zeta(3).$$

$$(0.257) \quad T(1, 0, 3) = \frac{\pi^4}{360}.$$

$$(0.258) \quad T(1, 1, 2) = \frac{\pi^4}{180}.$$

$$(0.259) \quad T(2, 1, 1) = \frac{\pi^4}{72}.$$

$$(0.260) \quad T(2, 2, 0) = \frac{\pi^4}{36}.$$

$$(0.261) \quad T(1, 0, 5) = \frac{3}{4}\zeta(6) - \frac{1}{2}\zeta^2(3).$$

$$(0.262) \quad T(1, 1, 4) = \frac{3}{2}\zeta(6) - \zeta^2(3).$$

$$(0.263) \quad T(3, 3, 0) = \zeta^2(3).$$

$$(0.264) \quad T(4, 1, 1) = \frac{7}{6}\zeta(6) - \frac{1}{2}\zeta^2(3).$$

$$(0.265) \quad T(4, 2, 0) = \frac{7}{4}\zeta(6).$$

$$(0.266) \quad T(1, 1, 4) = \frac{3}{2}\zeta(6) - \zeta^2(3).$$

$$(0.267) \quad \sum_{n=0}^{\infty} \frac{1}{(n+a)(n+b)} = \frac{\psi(a) - \psi(b)}{a-b}.$$

$$(0.268) \quad \sum_{n=0}^{\infty} \frac{(-)^n}{(n+a)(n+b)} = -\frac{1}{2} \frac{\psi(1/2+a/2) - \psi(a/2) - \psi(1/2+b/2) + \psi(b/2)}{a-b}.$$

[89]

$$(0.269) \quad \log 2 = \frac{1}{2} - \sum_{k \geq 1} \frac{(-1)^k}{k(4k^4 + 1)}.$$

After inserting $x = 1$ in (1.65), [17, 4.1.13]

$$(0.270) \quad 2 \ln 2 = 1 + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k(k+1)} = 1 + 2 \sum_{k=1,3,5,\dots}^{\infty} \frac{1}{k(k+1)(k+2)} \\ = 1 + 4 \sum_{k=1,5,9,13,\dots}^{\infty} \frac{6 + 4k + k^2}{k(k+1)(k+2)(k+3)(k+4)}.$$

[17, 4.1.20]

$$2 \ln 2 = \frac{5}{4} + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k(k+1)(k+2)} = \frac{5}{4} + 3 \sum_{k=1,3,5,\dots}^{\infty} \frac{1}{k(k+1)(k+2)(k+3)} \\ = \frac{17}{12} - 3 \sum_{k=2,4,6,\dots}^{\infty} \frac{1}{k(k+1)(k+2)(k+3)}.$$

$$(0.271) \quad \frac{4}{3} \ln 2 = \frac{8}{9} + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k(k+1)(k+2)(k+3)} = \frac{8}{9} + 4 \sum_{k=1,3,5,\dots}^{\infty} \frac{1}{k(k+1)(k+2)(k+3)(k+4)}.$$

More irregular denominators follow from hybridization. For example we can multiply the penultimate formula by 4, the previous formula by 3, and subtract

$$(0.272) \quad 4 \ln 2 = \frac{7}{3} + 12 \sum_{k=1,3,5,\dots}^{\infty} \frac{1}{k(k+1)(k+2)(k+4)}.$$

$$\frac{2}{3} \ln 2 = \frac{131}{288} + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k(k+1)(k+2)(k+3)(k+4)} \\ = \frac{131}{288} + 5 \sum_{k=1,3,5,\dots}^{\infty} \frac{1}{k(k+1)(k+2)(k+3)(k+4)(k+5)}.$$

Inserting $x = -1/2$ in (1.65) [89, 22] yields

$$(0.273) \quad \ln 2 = 1 - \sum_{k=1}^{\infty} \frac{1}{2^k k(k+1)}.$$

$$(0.274) \quad \ln 2 = \frac{1}{2} + \sum_{k=1}^{\infty} \frac{1}{2^{k-1} k(k+1)(k+2)}.$$

$$(0.275) \quad \ln 2 = \frac{5}{6} - 3 \sum_{k=1}^{\infty} \frac{1}{2^{k-1} k(k+1)(k+2)(k+3)}.$$

More irregular denominators follow from hybridization. For example we can multiply the penultimate formula by 3 and add to the previous formula,

$$(0.276) \quad 4 \ln 2 = \frac{7}{3} + 3 \sum_{k=1}^{\infty} \frac{1}{2^{k-1} k(k+1)(k+3)}.$$

$$(0.277) \quad \ln 2 = \frac{7}{12} + 3 \sum_{k=1}^{\infty} \frac{1}{2^{k-3} k(k+1)(k+2)(k+3)(k+4)}.$$

[89]

$$(0.278) \quad \log 2 = \frac{1327}{1920} + \frac{45}{4} \sum_{k \geq 4} \frac{(-1)^k}{k(k^2-1)(k^2-4)(k^2-9)}.$$

[89]

$$(0.279) \quad \log 2 = \sum_{k \geq 1} \left(\frac{1}{3^k} + \frac{1}{4^k} \right) \frac{1}{k}.$$

$$(0.280) \quad \log 2 = \sum_{k \geq 0} \left(\frac{1}{8k+8} + \frac{1}{4k+2} \right) \frac{1}{4^k}.$$

[89]

$$(0.281) \quad \log 2 = \frac{2}{3} + \sum_{k \geq 1} \left(\frac{1}{2k} + \frac{1}{4k+1} + \frac{1}{8k+4} + \frac{1}{16k+12} \right) \frac{1}{16^k}.$$

[89]

$$(0.282) \quad \log 2 = \frac{2}{3} \sum_{k \geq 0} \frac{1}{(2k+1)9^k}.$$

$$(0.283) \quad \log 2 = \frac{3}{4} \sum_{k \geq 0} \frac{(-1)^k k!^2}{2^k (2k+1)!}.$$

[89]

$$(0.284) \quad \log 2 = \frac{3}{4} + \frac{1}{4} \sum_{k \geq 1} \frac{(-1)^k (5k+1)}{k(2k+1)16^k} \binom{2k}{k}.$$

[175, A154920][23]

$$(0.285) \quad \log 3 = \sum_{k \geq 1} \left[\frac{9}{2k-1} + \frac{1}{2k} \right] \frac{1}{9^k}.$$

[175, A164985][23]

$$(0.286) \quad 27 \log 5 = 4 \sum_{k \geq 0} \left[\frac{9}{4k+1} + \frac{3}{4k+2} + \frac{1}{4k+3} \right] \frac{1}{81^k}.$$

$$(0.287) \quad \log 5 = 2 \log 3 - \log 2 + \sum_{k=1}^{\infty} \frac{1}{k10^k}.$$

$$(0.288) \quad 4 \log 7 = 5 \log 2 + \log 3 + 2 \log 5 + \sum_{k=1}^{\infty} \frac{(-1)^k}{k2400^k}.$$

TABLE 1. Formulas of the type $s \log p = t \log q + \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} r^k$. [128]

s	p	t	q	r
1	3	1	2	$1/2 \approx 0.50000000$
1	3	2	2	$-1/4 \approx -0.25000000$
2	3	3	2	$1/8 \approx 0.12500000$
5	3	8	2	$-13/256 \approx -0.05078125$
12	3	19	2	$7153/524288 \approx 0.01364326$
1	5	2	2	$1/4 \approx 0.25000000$
3	5	7	2	$-3/128 \approx -0.02343750$
1	7	2	2	$3/4 \approx 0.75000000$
1	7	3	2	$-1/8 \approx -0.12500000$
5	7	14	2	$423/16384 \approx 0.02581787$
1	11	3	2	$3/8 \approx 0.37500000$
2	11	7	2	$-7/128 \approx -0.05468750$
11	11	38	2	$10433763667/274877906944 \approx 0.03795781$
1	13	3	2	$5/8 \approx 0.62500000$
1	13	4	2	$-3/16 \approx -0.18750000$
3	13	11	2	$149/2048 \approx 0.07275391$
7	13	26	2	$-4360347/67108864 \approx -0.06497423$
10	13	37	2	$419538377/137438953472 \approx 0.00305254$
1	17	4	2	$1/16 \approx 0.06250000$
1	19	4	2	$3/16 \approx 0.18750000$
4	19	17	2	$-751/131072 \approx -0.00572968$
1	23	4	2	$7/16 \approx 0.43750000$
1	23	5	2	$-9/32 \approx -0.28125000$
2	23	9	2	$17/512 \approx 0.03320312$
1	29	4	2	$13/16 \approx 0.81250000$
1	29	5	2	$-3/32 \approx -0.09375000$
7	29	34	2	$70007125/17179869184 \approx 0.00407495$
1	31	4	2	$15/16 \approx 0.93750000$
1	31	5	2	$-1/32 \approx -0.03125000$
1	37	5	2	$5/32 \approx 0.15625000$
4	37	21	2	$-222991/2097152 \approx -0.10633039$
5	37	26	2	$2235093/67108864 \approx 0.03330548$
1	41	5	2	$9/32 \approx 0.28125000$
2	41	11	2	$-367/2048 \approx -0.17919922$
3	41	16	2	$3385/65536 \approx 0.05165100$
1	43	5	2	$11/32 \approx 0.34375000$
2	43	11	2	$-199/2048 \approx -0.09716797$
5	43	27	2	$12790715/134217728 \approx 0.09529825$
7	43	38	2	$-3059295837/274877906944 \approx -0.01112965$
1	47	5	2	$15/32 \approx 0.46875000$
1	47	6	2	$-17/64 \approx -0.26562500$
2	47	11	2	$161/2048 \approx 0.07861328$
1	53	5	2	$21/32 \approx 0.65625000$
1	53	6	2	$-11/64 \approx -0.17187500$
3	53	17	2	$17805/131072 \approx 0.13584137$
4	53	23	2	$-498127/8388608 \approx -0.05938137$
1	59	5	2	$27/32 \approx 0.84375000$
1	59	6	2	$-5/64 \approx -0.07812500$
1	61	5	2	$29/32 \approx 0.90625000$
1	61	6	2	$-3/64 \approx -0.04687500$
1	67	6	2	$3/64 \approx 0.04687500$

[23]

$$(0.289) \quad 3^5 \log 7 = \sum_{k \geq 0} \left[\frac{405}{6k+1} + \frac{81}{6k+2} + \frac{72}{6k+3} + \frac{9}{6k+4} + \frac{5}{6k+5} \right] \frac{1}{3^{6k}}.$$

[23]

$$(0.290) \quad 2 \times 3^9 \log 11 = \sum_{k \geq 0} \left[\frac{85293}{10k+1} + \frac{10935}{10k+2} + \frac{9477}{10k+3} + \frac{1215}{10k+4} + \frac{648}{10k+5} + \frac{135}{10k+6} \right. \\ \left. + \frac{117}{10k+7} + \frac{15}{10k+8} + \frac{13}{10k+9} \right] \frac{1}{3^{10k}}.$$

$$(0.291) \quad \log 11 = \log 2 + \log 5 + \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k10^k}.$$

$$(0.292) \quad \log 11 = 2 \log 2 + \log 3 - \sum_{k=1}^{\infty} \frac{1}{k12^k}.$$

$$(0.293) \quad \log 11 = 2(\log 2 + \log 5 - \log 3) - \sum_{k=1}^{\infty} \frac{1}{k100^k}.$$

$$(0.294) \quad \log 13 = 2 \log 2 + \log 3 + \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k12^k}.$$

$$(0.295) \quad 3 \log 17 = 3 \log 2 + 4 \log 5 - \sum_{k=1}^{\infty} \frac{1}{k} \left(\frac{87}{5000} \right)^k.$$

$$(0.296) \quad \log 19 = 2 \log 2 + \log 5 - \sum_{k=1}^{\infty} \frac{1}{k20^k}.$$

[115]

$$(0.297) \quad \log 11 = \sum_{k \geq 1} \frac{1}{k} \left(\frac{3}{2^k} + \frac{1}{2^{2k}} + \frac{1}{2^{5k}} - \frac{1}{2^{10k}} \right) = \frac{1}{2} \sum_{k \geq 1} \frac{1}{k} \left(\frac{13}{3^k} - \frac{4}{3^{2k}} - \frac{1}{3^{5k}} \right).$$

[115]

$$(0.298) \quad \log 13 = \sum_{k \geq 1} \frac{1}{k} \left(\frac{3}{2^k} + \frac{1}{2^{2k}} + \frac{1}{2^{3k}} + \frac{1}{2^{4k}} - \frac{1}{2^{12k}} \right) = \sum_{k \geq 1} \frac{1}{k} \left(\frac{7}{3^k} - \frac{2}{3^{2k}} - \frac{1}{3^{3k}} \right).$$

[115]

$$(0.299) \quad \log 17 = \sum_{k \geq 1} \frac{1}{k} \left(\frac{4}{2^k} + \frac{1}{2^{4k}} - \frac{1}{2^{8k}} \right).$$

[115]

$$(0.300) \quad \log 19 = \sum_{k \geq 1} \frac{1}{k} \left(\frac{3}{2^k} + \frac{3}{2^{2k}} + \frac{1}{2^{9k}} - \frac{1}{2^{18k}} \right).$$

Similar formulas as the four above are obtained by inserting

(0.301)

$$\log[(2^s - 1)^\tau] = s\tau \log 2 - \sum_{k \geq 1} \frac{\tau}{k 2^{ks}} = -s\tau \log \frac{1}{2} - \sum_{k \geq 1} \frac{\tau}{k 2^{ks}} = \sum_{k \geq 1} \frac{\tau}{k} \left(\frac{s}{2^k} - \frac{1}{2^{ks}} \right),$$

—immediate consequence of putting $x = 1/2$ in (1.64)—into the right hand sides of [46]:

$$(0.302) \quad \log 41 = \log(2^{20} - 1) - \log(2^{10} - 1) + \log[(2^2 - 1)^2] - \log[(2^4 - 1)^2].$$

$$(0.303) \quad \log 43 = \log(2^{14} - 1) - \log(2^2 - 1) - \log(2^7 - 1).$$

$$(0.304) \quad \log 73 = \log(2^9 - 1) - \log(2^3 - 1).$$

$$(0.305) \quad \log 151 = \log(2^{15} - 1) - \log(2^3 - 1) - \log(2^5 - 1).$$

$$(0.306) \quad \log 241 = \log(2^{24} - 1) - \log(2^{12} - 1) - \log(2^8 - 1) + \log(2^4 - 1).$$

$$(0.307) \quad \log 257 = \log(2^{16} - 1) - \log(2^8 - 1).$$

$$(0.308) \quad \log 331 = \log(2^{30} - 1) - \log(2^{15} - 1) - \log(2^{10} - 1) + \log(2^5 - 1) - \log(2^2 - 1).$$

$$(0.309) \quad \log 337 = \log(2^{21} - 1) - \log(2^7 - 1) - \log[(2^6 - 1)^2] + \log[(2^2 - 1)^4].$$

$$(0.310) \quad \log 683 = \log(2^{22} - 1) - \log(2^{11} - 1) - \log(2^2 - 1).$$

$$(0.311) \quad \log 2731 = \log(2^{26} - 1) - \log(2^{13} - 1) - \log(2^2 - 1).$$

$$(0.312) \quad \log 5419 = \log(2^{42} - 1) - \log(2^{21} - 1) - \log(2^{14} - 1) + \log(2^7 - 1) - \log(2^2 - 1).$$

$$(0.313) \quad \log 43691 = \log(2^{34} - 1) - \log(2^{17} - 1) - \log(2^2 - 1).$$

$$(0.314) \quad \log 61681 = \log(2^{40} - 1) - \log(2^{20} - 1) - \log(2^8 - 1) + \log(2^4 - 1).$$

$$(0.315) \quad \log 174763 = \log(2^{38} - 1) - \log(2^{18} - 1) - \log(2^2 - 1).$$

$$(0.316) \quad \log 262657 = \log(2^{27} - 1) - \log(2^9 - 1).$$

$$(0.317) \quad \log 599479 = \log(2^{33} - 1) - \log(2^{11} - 1) - \log(2^3 - 1).$$

$$(0.318) \quad \sum_{k \geq 0} \frac{\log(2k+1)}{(2k+1)^l} = -\frac{\log 2}{2^l} \zeta(l) - (1 - 2^{-l}) \zeta'(l).$$

$$(0.319) \quad \sum_{k \geq 0} \frac{\log(ak+b)}{(ak+b)^l} = \frac{\log a}{a^l} \zeta(l, b/a) - \frac{1}{a^l} \zeta'(l, b/a).$$

[34]

$$(0.320) \quad \sum_{k=1}^{\infty} \frac{kk!}{(2k)!} = \frac{1}{8} \left(2 + 3e^{1/4} \sqrt{\pi} \operatorname{erf}(1/2) \right).$$

$$(0.321) \quad \sum_{k=1}^{\infty} \frac{(-1)^k \alpha^k x^k}{kk!} = -\gamma - \Gamma(0, \alpha x) - \ln \alpha - \ln x.$$

[6]

$$\begin{aligned} & \frac{3}{2} \sum_{k=0}^{\infty} \frac{1}{16^k} \left(\frac{1}{(8k+1)^2} - \frac{1}{(8k+2)^2} + \frac{1}{2(8k+3)^2} - \frac{1}{4(8k+5)^2} + \frac{1}{4(8k+6)^2} - \frac{1}{8(8k+7)^2} \right) \\ & - \frac{1}{4} \sum_{k=0}^{\infty} \frac{1}{4096^k} \left(\frac{1}{(8k+1)^2} + \frac{1}{2(8k+2)^2} + \frac{1}{8(8k+3)^2} - \frac{1}{64(8k+5)^2} - \frac{1}{128(8k+6)^2} - \frac{1}{512(8k+7)^2} \right) = G, \end{aligned}$$

which is Catalan's constant [175, A006752].

[44]

$$(0.322) \quad \sum_{k \geq 0} \frac{1}{16^k} \left(\frac{4}{8k+1} - \frac{2}{8k+4} - \frac{1}{8k+5} - \frac{1}{8k+6} \right) = \pi.$$

[90]

$$(0.323) \quad \frac{1}{16807} \sum_{n=0}^{\infty} \frac{1}{2^n \binom{7n}{2n}} \left(\frac{59296}{7n+1} - \frac{10326}{7n+2} - \frac{3200}{7n+3} - \frac{1352}{7n+4} - \frac{792}{7n+5} + \frac{552}{7n+6} \right) = \pi.$$

[6]

$$(0.324) \quad \frac{1}{2} \sum_{k=0}^{\infty} \frac{4^k k!^2}{(2k)!(2k+1)^2} = G.$$

[6]

$$(0.325) \quad -\frac{\pi}{32} \sum_{k=0}^{\infty} \frac{(2k+1)!^2}{16^k k!^4 (k+1)^3} = G - \frac{\pi}{2} \log 2.$$

[6]

$$(0.326) \quad \sqrt{2} \sum_{k=0}^{\infty} \frac{(2k)!}{8^k k!^2 (2k+1)^2} = G + \frac{\pi}{4} \log 2.$$

[6, 41]

$$(0.327) \quad \frac{3}{8} \sum_{k=0}^{\infty} \frac{k!^2}{(2k)!(2k+1)^2} = G + \frac{\pi}{8} \log(2 + \sqrt{3}).$$

[41]

$$(0.328) \quad \frac{5}{8} \sum_{k=0}^{\infty} \frac{L(2k+1)}{(2k+1)^2 \binom{2k}{k}} = G - \frac{\pi}{8} \log\left(\frac{10 + \sqrt{50 - 22\sqrt{5}}}{10 - \sqrt{50 - 22\sqrt{5}}}\right).$$

where L are the Lucas numbers.

[155]

$$(0.329) \quad \sqrt{e} = \frac{16}{31} \left[1 + \sum_{n=1}^{\infty} \frac{1 + n^3/2 + n/2}{2^n n!} \right],$$

which is a special case of $\frac{1}{2}P_3(1/2) + \frac{1}{2}P_1(1/2) + P_0(1/2)$ of [100]

$$(0.330) \quad \sum_{n=0}^{\infty} n^j \frac{t^n}{n!} = P_j(t) \exp(t)$$

where

$$(0.331) \quad P_0(t) \equiv 1; \quad P_j(t) \equiv t [P'_{j-1}(t) + P_{j-1}(t)], j > 0.$$

[4]

$$(0.332) \quad \sum_{k \geq 1} \left[\begin{matrix} k \\ p \end{matrix} \right] \frac{1}{k!k} = \zeta(p+1).$$

[4]

$$(0.333) \quad \sum_{k \geq 2} \left[\begin{matrix} k \\ 2 \end{matrix} \right] \frac{1}{k!k^q} = \frac{q+1}{2} \zeta(q+2) - \frac{1}{q} \sum_{k=1}^{q-1} k \zeta(k+1) \zeta(q+1-k).$$

[4]

$$(0.334) \quad \sum_{k \geq 1} \left[\begin{matrix} k \\ p \end{matrix} \right] \frac{z^k}{k!k} = \zeta(p+1) + \sum_{k=1}^p \left(\frac{(-)^{k-1}}{k!} \text{Li}_{p+1-k}(1-z) \log^k(1-z) \right).$$

[4]

$$(0.335) \quad \sum_{k \geq p} \left[\begin{matrix} k \\ p \end{matrix} \right] \frac{1}{k!k^q} = \sum_{k \geq q} \left[\begin{matrix} k \\ q \end{matrix} \right] \frac{1}{k!k^p} = \frac{(-)^{q-1}}{(q-1)!p!} \lim_{\beta \rightarrow 0} \lim_{\alpha \rightarrow 0} \frac{d^{q+p-1}}{d\alpha^q d\beta^{q-1}} \frac{\Gamma(1-\alpha)\Gamma(1+\beta)}{\Gamma(\beta)\Gamma(1-\alpha+\beta)},$$

which can always be represented in finite terms of zeta functions.

0.3. Formulae from Differential Calculus. [112]

$$(0.336) \quad \left(\frac{d}{dx} \right)^m e^{g(x)} = e^{g(x)} Y_m \left(g'(x), g''(x), \dots, g^{(m)}(x) \right)$$

where Y_m is the m th exponential complete Bell polynomial,

$$(0.337) \quad Y_n(x_1, \dots, x_n) = \sum_{\pi(n)} \frac{n!}{k_1! \dots k_n!} \left(\frac{x_1}{1!} \right)^{k_1} \dots \left(\frac{x_n}{n!} \right)^{k_n}.$$

$$(0.338) \quad \frac{d}{dx} \left[f(x)^{g(x)} \right] = f(x)^{g(x)} \left[(\log f) g' + g \frac{f'}{f} \right]$$

[62]

$$(0.339) \quad \sum_{n \geq 0} \frac{g^{(n)}(0)}{n!} f(n) x^n = \sum_{n \geq 0} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k g^{(k)}(x).$$

where the braces are Stirling numbers of the second kind.

1. ELEMENTARY FUNCTIONS

1.1. Powers of Binomials. [30]

$$(1.1) \quad \left(\frac{1 + \sqrt{1+4x}}{2} \right)^n = (1+x)^{n/2} \sum_{k \geq 0} \frac{(-)^{k+1} \Gamma(\frac{3k-2}{2})}{\Gamma(\frac{3k-n}{2} - k + 1) k!} \left(\frac{x}{(1+x)^{3/2}} \right)^k.$$

[119]

$$(1.2) \quad \sum_{n=0}^{\infty} \binom{2n}{n} x^n = \frac{1}{\sqrt{1-4x}}.$$

[119][175, A000108]

$$(1.3) \quad \sum_{n=0}^{\infty} \binom{2n}{n} \frac{x^n}{n+1} = \frac{1 - \sqrt{1-4x}}{2x}.$$

[119]

$$(1.4) \quad \sum_{n=1}^{\infty} \frac{1}{n} \binom{2n}{n} x^n = 2 \log \frac{1 - \sqrt{1-4x}}{2x}.$$

[119]

$$(1.5) \quad x \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \binom{2n}{n} x^n = 2x \log \frac{1 - \sqrt{1-4x}}{x} + \frac{\sqrt{1-4x}}{2} - x(\log 4 - 1) - \frac{1}{2},$$

which corrects a sign error in [119], see [127].

[40]

$$(1.6) \quad \sum_{n=0}^{\infty} \frac{1}{n+1} \binom{2n}{n} H_n x^{n+1} = \sqrt{1-4x} \log(2\sqrt{1-4x}) - (1+\sqrt{1-4x}) \log(1+\sqrt{1-4x}) + \log 2,$$

$$(1.7) \quad \sum_{n=0}^{\infty} \binom{2n}{n} H_n x^n = \frac{2}{\sqrt{1-4x}} \log\left(\frac{1+\sqrt{1-4x}}{2\sqrt{1-4x}}\right),$$

$$(1.8) \quad \sum_{n=0}^{\infty} \binom{2n}{n} H_n (-1)^{n-1} x^n = \frac{2}{\sqrt{1+4x}} \log\left(\frac{2\sqrt{1+4x}}{1+\sqrt{1+4x}}\right),$$

where $H_n = \sum_{k=1}^n 1/k$.

[40]

$$(1.9) \quad \sum_{n=0}^{\infty} \binom{2n}{n} P_q(n) x^{n+1} = \frac{1}{\sqrt{1-4x}} \sum_{k=0}^q \binom{2k}{k} k! \left(\frac{x}{1-4x}\right)^k \sum_{m=k}^q a_m S(m, k)$$

where $P_q(z) = a_q z^q + a_{q-1} z^{q-1} + \dots + q_0$ is a polynomial and S are the Stirling Numbers of the Second Kind.

[175, A005430]

$$(1.10) \quad \sum_{n=1}^{\infty} n \binom{2n}{n} x^n = 2x(1-4x)^{-3/2}.$$

[119][175, A002736]

$$(1.11) \quad \sum_{n=1}^{\infty} n^2 \binom{2n}{n} x^n = \frac{2x(2x+1)}{(1-4x)^{5/2}}.$$

[119]

$$(1.12) \quad \sum_{n=0}^{\infty} \frac{1}{2n+1} \binom{2n}{n} x^n = \frac{1}{2x} \arcsin(2x).$$

[40]

$$(1.13) \quad \sum_{n=0}^{\infty} \frac{1}{n+m+1} \binom{2n}{n} x^n = \frac{1}{2^{2m+1} x^{m+1}} \sum_{k=0}^m \binom{m}{k} \frac{(-1)^k}{2k+1} [1 - (1-4x)^{k+1/2}].$$

[40]

$$(1.14) \quad \sum_{n=0}^{\infty} \frac{1}{n^2} \binom{2n}{n} x^n = 2 \operatorname{Li}_2 \left(\frac{1 - \sqrt{1 - 4x}}{2} \right) \\ - \log^2(1 + \sqrt{1 - 4x}) - 2 \log 2 \log \frac{1 - \sqrt{1 - 4x}}{x} + 3 \log^2 2.$$

[119]

$$(1.15) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1}}{\binom{2m}{m}} (2x/y)^{2m} = \frac{xy^2}{h^3} \left[\log \frac{x+h}{y} + \frac{xh}{y^2} \right]$$

where $h \equiv \sqrt{x^2 + y^2}$.

[119]

$$(1.16) \quad \sum_{m=1}^{\infty} \frac{(2x)^{2m}}{m \binom{2m}{m}} = \frac{2x \arcsin x}{\sqrt{1 - x^2}}.$$

[119]

$$(1.17) \quad \sum_{m=1}^{\infty} \frac{(2x)^{2m}}{m^2 \binom{2m}{m}} = 2(\arcsin x)^2.$$

[119]

$$(1.18) \quad \sum_{m=1}^{\infty} \frac{(2x)^{2m}}{\binom{2m}{m}} = \frac{x^2}{1 - x^2} + \frac{x \arcsin x}{(1 - x^2)^{3/2}}.$$

(1.19)

$$\frac{1}{(\sigma^2 + \sigma_L^2)^{\gamma/2}} = \frac{1}{\sigma_L^{\gamma}} \exp \left(-\frac{\gamma}{2} \frac{\sigma^2}{\sigma_L^2} \right) \left(1 + \frac{\gamma}{4} \left(\frac{\sigma}{\sigma_L} \right)^4 - \frac{\gamma}{6} \left(\frac{\sigma}{\sigma_L} \right)^6 + \frac{\gamma(\gamma+4)}{32} \left(\frac{\sigma}{\sigma_L} \right)^8 \right. \\ \left. - \frac{\gamma(5\gamma+12)}{120} \left(\frac{\sigma}{\sigma_L} \right)^{10} + \frac{\gamma(3\gamma^2+52\gamma+96)}{1152} \left(\frac{\sigma}{\sigma_L} \right)^{12} - \frac{\gamma(35\gamma^2+308\gamma+480)}{6720} \left(\frac{\sigma}{\sigma_L} \right)^{14} + \dots \right)$$

[162, 3.41(a)]

$$(1.20) \quad \frac{1}{a^2 - x^2} = \frac{2}{a\sqrt{a^2 - 1}} \sum_{j=0}^{\infty \prime} (a - \sqrt{a^2 - 1})^{2j} T_{2j}(x)$$

where the prime at the sum symbols means taking only half of the value at index zero.

[31]

$$(1.21) \quad (1 + x + x^{-1})^n = \sum_{j=-n}^n \binom{n}{j}_2 x^j, \quad \binom{n}{m}_2 \equiv \sum_{j \geq 0} \frac{n!}{j!(m+j)!(n-2j-m)!}.$$

[153]

$$(1.22) \quad \sum_{s=0}^{\infty} \zeta(4s+3) x^{4s} = \sum_{k=1}^{\infty} \frac{k}{k^4 - x^4} = \frac{5}{2} \sum_{k=1}^{\infty} \frac{(-)^{k-1}}{\binom{2k}{k}} \frac{k}{k^4 - x^4} \prod_{m=1}^{k-1} \frac{m^4 + 4x^4}{m^4 - x^4}.$$

[28] Let $B_0(x)$ be a period function of period 1. Assume $B_0(x)$ has a continuous derivative in the open interval $(0,1)$. Let $a_0 \equiv \int_0^1 B_0(x)dx$ and define

$$(1.23) \quad B_n(x) = \int_0^2 B_{n-1}(y)dy + \int_0^1 (y-1)B_{n-1}(y)dy.$$

Then

$$(1.24) \quad \sum_{k=0}^{\infty} B_k(x)t^k = \frac{te^{xt}}{e^t - 1} \left[a_0 - \int_0^1 B'_0(1-y) \frac{e^{ty} - 1}{t} dy \right] + \int_0^x e^{t(x-y)} B'_0(y) dy,$$

and

$$(1.25) \quad B_n(x) = \Re \left[2 \sum_{j=1}^{\infty} \frac{e^{2\pi i j x}}{(2\pi i j)^n} (a_j - a_0 - i b_j) \right]$$

where a_j and b_j are the Fourier coefficients of $B_0(x)$. For example $B_0(x) = \cos(\pi x)$ yields $B_2(x) = (1 - 2x - \cos(\pi x))/\pi^2$ and

$$(1.26) \quad \frac{\pi^3}{2} B_2(x) = - \sum_{j=1}^{\infty} \frac{\sin(2\pi j x)}{j(4j^2 - 1)}.$$

Consider also a periodic sequence $\lambda_{j+T} = \lambda_j$, the Dirichlet series

$$(1.27) \quad f(s) = \sum_{j=1}^{\infty} \frac{a_j - a_0}{j^s} \lambda_j, \quad g(s) = \sum_{j=1}^{\infty} \frac{b_j}{j^s} \lambda_j,$$

and the Fourier coefficients

$$(1.28) \quad \alpha_j \equiv \sum_{k=1}^T \lambda_k \cos(2\pi j k / T), \quad \beta_j \equiv \sum_{k=1}^T \lambda_k \sin(2\pi j k / T).$$

If $\lambda_{T-j} = \lambda_j$ for $1 \leq j < T$, then

$$(1.29) \quad f(2n) = \frac{(-)^n}{2T} (2\pi)^{2n} \sum_{j=1}^T \alpha_j B_{2n}(j/T),$$

$$(1.30) \quad g(2n+1) = \frac{(-)^{n+1}}{2T} (2\pi)^{2n+1} \sum_{j=1}^T \alpha_j B_{2n+1}(j/T),$$

and similar expressions for other even-odd symmetries of the λ .

[28]

$$(1.31) \quad 8 \log 2 - 4 = \zeta(3) + \sum_{j=1}^{\infty} \frac{1}{j^2(4j^2 - 1)} = \sum_{k=0}^{\infty} \frac{\zeta(3+2k)}{4^k}.$$

[28]

$$(1.32) \quad \sum_{j=1}^{\infty} \frac{\cos(2\pi j/3)}{j^{2k}(4\pi^2 j^2 + 1)} = \frac{(-)^k}{4} (2\pi)^{2k} \alpha_{2k},$$

with g.f.

$$(1.33) \quad \sum_{k=0}^{\infty} \alpha_k t^k = \frac{e + e^{4/3} - e^{1+t} - e^{t+4/3} + t^3 [e^{(2t+5)/3} - e^{(2+2t)/3} - e^{(2+t)/3} + e^{(5+t)/3}]}{(e-1)e^{2/3}(e^t-1)(t^2-1)} - \frac{2te^{t/3}}{e^t-2}.$$

1.2. The Exponential Function. [84], [175, A001469]

$$(1.34) \quad \frac{2t}{e^t + 1} = \sum_{N=0}^{\infty} \frac{G_N}{N!} t^N;$$

with $(G+1)^N + G_N = 1$ for $N > 1$, $G_1 = 1$, $G_2 = -1$, $G_4 = 1$, $G_6 = -3$, $G_8 = 17$, $G_{10} = -155$, $G_{12} = 2073$, Genocchi numbers.

[31]

$$(1.35) \quad \exp(xt + yt^2) = \sum_{n=0}^{\infty} \frac{t^n}{n!} H_n(x, y), \quad H_n(x, y) \equiv n! \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{x^{n-2k} y^k}{k!(n-2k)!}.$$

1.3. Fourier Series. [61, B2a]

$$(1.36) \quad \cos(m\theta) = \sum_{k=0}^{\lfloor m/2 \rfloor} (-1)^k \binom{m}{2k} \cos^{m-2k} \theta \sin^{2k} \theta.$$

$$(1.37) \quad \sin^{2k} \alpha + \cos^{2k} \alpha = 1 - \sum_{l=1}^{k-1} \binom{k}{l} \sin^{2l} \alpha \cos^{2k-2l} \alpha.$$

[60]

$$(1.38) \quad S_{\nu}(\alpha) \equiv \sum_{k=0}^{\infty} \frac{\sin(2k+1)\alpha}{(2k+1)^{\nu}};$$

$$(1.39) \quad C_{\nu}(\alpha) \equiv \sum_{k=0}^{\infty} \frac{\cos(2k+1)\alpha}{(2k+1)^{\nu}};$$

$$(1.40) \quad S_{2n+1}(\alpha) = \frac{(-1)^n}{4(2n)!} \pi^{2n+1} E_{2n} \left(\frac{\alpha}{\pi} \right);$$

$$(1.41) \quad C_{2n}(\alpha) = \frac{(-1)^n}{4(2n-1)!} \pi^{2n} E_{2n-1} \left(\frac{\alpha}{\pi} \right);$$

$$(1.42) \quad S_{\nu}(2\pi p/q) = \frac{1}{q^{\nu}} \sum_{s=1}^{q-1} \zeta(\nu, s/q) \left[\sin(s2\pi p/q) - \frac{\sin(s4\pi p/q)}{2^{\nu}} \right];$$

$$(1.43) \quad C_{\nu}(2\pi p/q) = \frac{1}{q^{\nu}} \left\{ \zeta(\nu) \left(1 - \frac{1}{2^{\nu}} \right) + \sum_{s=1}^{q-1} \zeta(\nu, s/q) \left[\cos(s2\pi p/q) - \frac{\cos(s4\pi p/q)}{2^{\nu}} \right] \right\}.$$

[91, 1.448.1]

$$(1.44) \quad \sum_{k=1}^{\infty} \frac{p^k \sin(kx)}{k} = \arctan \frac{p \sin x}{1 - p \cos x}, \quad 0 < x < 2\pi, p^2 \leq 1,$$

with special case

$$(1.45) \quad \sum_{k=1}^{\infty} \frac{\sin(kx)}{2^k k} = \arctan \frac{\sin x}{2 - \cos x}, \quad 0 < x < 2\pi.$$

[91, 1.448.2]

$$(1.46) \quad \sum_{k=1}^{\infty} \frac{p^k \cos(kx)}{k} = \ln \frac{1}{\sqrt{1 - 2p \cos x + p^2}}, \quad 0 < x < 2\pi,$$

with special case

$$(1.47) \quad \sum_{k=1}^{\infty} \frac{\cos(kx)}{2^k k} = -\ln \sqrt{1 - \cos x + 1/4}, \quad 0 < x < 2\pi.$$

[99, (1.10)]

$$(1.48) \quad -\sqrt{2 - 2 \cos \theta} = -\frac{4}{\pi} + \frac{2}{\pi} \sum_{k=1}^{\infty} \frac{\cos k\theta}{(k - 1/2)(k + 1/2)}$$

[184]

$$(1.49) \quad \text{Cl}_{2n+1}(\pi/2) = -2^{-2n-1}(1 - 2^{-2n})\zeta(2n + 1), n \in \mathbb{N}.$$

[184]

$$(1.50) \quad \text{Cl}_{2n+1}(\pi/3) = \frac{1}{2}(1 - 2^{-2n})(1 - 3^{-2n})\zeta(2n + 1), n \in \mathbb{N}.$$

[184]

$$(1.51) \quad \text{Cl}_{2n+1}(2\pi/3) = -\frac{1}{2}(1 - 3^{-2n})\zeta(2n + 1), n \in \mathbb{N}.$$

[184]

$$(1.52) \quad \sum_{k \geq 1} \frac{\cos(k\pi/2)}{k^s} = -2^{-s}(1 - 2^{1-s})\zeta(s), \Re s > 1.$$

[119]

$$(1.53) \quad \sum_{m=1}^{\infty} \frac{m^{k-2}(2x)^{2m}}{\binom{2m}{m}} = \frac{x}{2^{k-2}(1-x^2)^{k-1/2}} \left[\arcsin x V_x(x^2) + x \sqrt{1-x^2} W_k(x^2) \right], \quad k \geq 0,$$

[119]

$$(1.54) \quad \sum_{m=1}^{\infty} \frac{m^{k-2} 4^m (\sin \theta)^{2m}}{\binom{2m}{m}} = \frac{\sin 2\theta}{(2 \cos^2 \theta)^k} [2\theta V_k(\sin^2 \theta) + \sin 2\theta W_k(\sin^2 \theta)],$$

[119]

$$(1.55) \quad \sum_{m=1}^{\infty} \frac{(-1)^{m-1} m^{k-2} 4^m (\sinh z)^{2m}}{\binom{2m}{m}} \\ = \frac{\sinh 2z}{(2 \cosh^2 z)^k} [2 \log\{\sinh z + \cosh z\} V_k(-\sinh^2 z) + \sinh 2z W_k(-\sinh^2 z)],$$

where $V_1(t) = 1$, (see [175, A156919]) $W_1(t) = 0$,

$$V_{k+1}(t) = \{(2k-2)t + 1\} V_k(t) + 2(1-t) \delta V_k(t)$$

$$W_{k+1}(t) = \{(2k-4)t + 2\} W_k(t) + 2(1-t) \delta W_k(t) + V_k(t)$$

and δ is the operator $x \frac{d}{dx}$.

1.4. Expansions of Hyperbolic Functions. [30]

$$(1.56) \quad \cos x \cosh x = \sum_{k \geq 0} \frac{(-)^k (2x^2)^{2k}}{(4k)!}.$$

[30]

$$(1.57) \quad \sin x \sinh x = \sum_{k \geq 0} \frac{(-)^k (2x^2)^{2k+1}}{(4k+2)!}.$$

[30]

$$(1.58) \quad \coth(2x) \tanh x = 1 - \sum_{k \geq 1} \frac{2^{2k-1} (2^{2k} - 1) (2k-1) B_{2k} x^{2k-2}}{(2k)!}.$$

[33]

$$(1.59) \quad \sum_{j=0}^{\infty} \frac{x}{\sinh 2^{-j} x} - 2^j = 1 - \frac{x}{\tanh x}.$$

[33]

$$(1.60) \quad \sum_{j=0}^{\infty} \frac{2^j - \coth 2^{-j}}{2^j \sinh 2^{-j}} = \frac{1 + 4e^2 - e^4}{1 - 2e^2 + e^4}.$$

[86]

$$(1.61) \quad \sum_{m,n=-\infty}^{\infty} \frac{F(|2m+2n+1|)}{\cosh(2m+1)u \cosh 2nu} = 2 \sum_{n=0}^{\infty} \frac{(2n+1)F(2n+1)}{\sinh(2n+1)u},$$

for any summable function $F(x)$.

[86]

$$(1.62) \quad \sum_{m,n=-\infty}^{\infty} \frac{F(m+n+1) + F(m-n)}{\sinh(2m+1)u \cosh(2n+1)u} = 8 \sum_{n=1}^{\infty} \frac{nF(n)}{\cosh(2nu)},$$

and

$$(1.63) \quad \sum_{k,m,n=-\infty}^{\infty} \frac{F(k+m+n+1) + F(k-m+n)}{\cosh(2ku) \cosh(2m+1)u \sinh(2n+1)u} = 8 \sum_{n=1}^{\infty} \frac{n^2 F(n)}{\sinh(2nu)},$$

where F is any sine transform (and hence odd).

1.5. The Logarithmic Function.

$$(1.64) \quad \ln(1-x) = - \sum_{k=1}^{\infty} \frac{x^k}{k}.$$

(1.65)

$$(1+x) \ln(1+x) = x + \frac{1}{1 \cdot 2} x^2 - \frac{1}{2 \cdot 3} x^3 + \frac{1}{3 \cdot 4} x^4 - \frac{1}{4 \cdot 5} x^5 + \dots = x + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^{k+1}}{k(k+1)}.$$

By integration of this w.r.t. x [91, 2.729], [135]

$$(1.66) \quad \frac{1}{2} (1+x)^2 \ln(1+x) = \frac{x}{2} + \frac{3x^2}{4} + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^{k+2}}{k(k+1)(k+2)}.$$

$$(1.67) \quad \frac{1}{6}(1+x)^3 \ln(1+x) = \frac{x}{6} + \frac{5x^2}{12} + \frac{11x^3}{36} + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^{k+3}}{k(k+1)(k+2)(k+3)}.$$

$$(1.68) \quad \frac{1}{24}(1+x)^4 \ln(1+x) = \frac{x}{24} + \frac{7x^2}{48} + \frac{13x^3}{72} + \frac{25x^4}{288} + \sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^{k+4}}{k(k+1)(k+2)(k+3)(k+4)}.$$

$$(1.69) \quad x + (1-x)^l \ln(1-x) = \sum_{i=2}^l \tau_{i,l} x^i - (-1)^l l! \sum_{k=1}^{\infty} \frac{x^{k+l}}{k(k+1)(k+2) \cdots (k+l)},$$

where

$$(1.70) \quad \tau_{2,l} = l - 1/2, \quad l \geq 2; \quad i\tau_{i,l} = (-1)^i \binom{l-1}{i-1} - l\tau_{i-1,l-1}, \quad i \geq 3.$$

[116]

$$(1.71) \quad \log n = - \sum_{s \geq 1} \frac{\alpha(s, n)}{s}$$

with

$$(1.72) \quad \alpha(s, n) \equiv \begin{cases} n-1, & n \mid s; \\ -1 & n \nmid s. \end{cases}$$

Superposition of [91, 1.513.1] and [91, 1.511], see [175, A165998]:

$$(1.73) \quad \frac{1}{3x} \ln \frac{1+x}{(1-x)^2} = 1 + \frac{x}{6} + \frac{x^2}{3} + \frac{x^3}{12} + \cdots + \frac{x^{2j}}{2j+1} + \frac{x^{2j+1}}{3(2j+1)} + \cdots$$

[11]

$$(1.74) \quad \sum_{j=k}^{\infty} \frac{|S_j^{(k)}| b^j}{j!} = \frac{\ln^k(1+b)}{k!}.$$

[101]

$$(1.75) \quad \sum_{k=0}^{\infty} \frac{x^k}{\binom{k+L}{L}} = {}_2F_1(1, 1; L+1; x) = L \sum_{j=0}^{L-2} \frac{(x-1)^j}{(L-j-1)x^{j+1}} - L(x-1)^{L-1} \frac{\ln(1-x)}{x^L}; \quad L \geq 1.$$

[43]

$$(1.76) \quad \frac{1}{2x} \left\{ 1 - \ln(1+x) - \frac{1-x}{\sqrt{x}} \arctan \sqrt{x} \right\} = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} x^{k-1}}{(2k-1)2k(2k+1)};$$

$0 < x \leq 1$.

[138]

$$(1.77) \quad -\frac{\ln(1-t)}{(1-t)^r} = \sum_{n=0}^{\infty} H_n^{(r)} t^n,$$

where the Hyperharmonic numbers are defined by $H_n = \sum_{k=1}^n (1/k)$ and $H_n^{(r)} = \sum_{k=1}^n H_k^{(r-1)} = \binom{n+r-1}{r-1} (H_{n+r-1} - H_{r-1})$.

1.6. The Inverse Trigonometric and Hyperbolic Function. [33]

$$(1.78) \quad \sum_{k=1}^{\infty} 2^{-k} \tan \frac{x}{2^k} = \frac{1}{2^n} \cot \frac{x}{2^n} - \cot x.$$

[33]

$$(1.79) \quad \sum_{k=1}^{\infty} \csc \frac{x}{2^{k-1}} = \cot \frac{x}{2^n} - \cot x.$$

[33]

$$(1.80) \quad \sum_{k=1}^{\infty} \arctan \frac{2x^2}{k^2} = \frac{\pi}{4} - \arctan \frac{\tanh \pi x}{\tan \pi x}.$$

[33]

$$(1.81) \quad \sum_{k=1}^{\infty} (-1)^{k-1} \arctan \frac{2x^2}{k^2} = -\frac{\pi}{4} - \arctan \frac{\sinh \pi x}{\sin \pi x}.$$

2. INDEFINITE INTEGRALS OF ELEMENTARY FUNCTIONS

2.1. Rational Functions. Aids to partial fraction decompositions: [205]

$$(2.1) \quad \frac{1}{s^n(s^2 + as + b)} = \frac{-\alpha_{n-1}s + \alpha_n}{b^n(s^2 + as + b)} + \sum_{k=0}^{n-1} \frac{\alpha_k}{b^{k+1}s^{n-k}};$$

where $b > 0$, $a^2 - 4b < 0$, $\alpha_0 \equiv 1$,

$$(2.2) \quad \alpha_m \equiv (-1)^m \sqrt{b^m} U_m \left(\frac{a}{2\sqrt{b}} \right) = \sum_{k=0}^{\lfloor m/2 \rfloor} (-1)^{k+m} \binom{m-k}{k} a^{m-2k} b^k.$$

[205]

$$(2.3) \quad \frac{\alpha s + \beta}{(s^2 + as + b)(s^2 + kas + kb)} = \frac{1}{(k-1)b^2} \left(\frac{L(s+a) + \beta b}{s^2 + as + b} - \frac{L(s+ka) + \beta b}{s^2 + kas + kb} \right),$$

$$k \neq 1, ab \neq 0, L \equiv \begin{vmatrix} \alpha & \beta \\ a & c \end{vmatrix}.$$

[205]

$$(2.4) \quad \frac{1}{(pq + ap + b)(pq + cp + b)} = \begin{vmatrix} a & c \\ b & b \end{vmatrix}^{-1} \left(\frac{q+a}{pq + ap + b} - \frac{q+c}{pq + cp + b} \right),$$

$a \neq c, b \neq 0$.

[205]

$$(2.5) \quad \frac{\alpha p + \beta}{(pq + ap + b)(pq + cp + b)} = \begin{vmatrix} a & c \\ b & b \end{vmatrix}^{-1} \left(\frac{\begin{vmatrix} \alpha & \beta \\ q+c & b \end{vmatrix}}{pq + cp + b} - \frac{\begin{vmatrix} \alpha & \beta \\ q+a & b \end{vmatrix}}{pq + ap + b} \right),$$

$a \neq c, b \neq 0$.

[205]
(2.6)

$$\frac{as+b}{(s^2+\alpha s+\beta)(s^2+\gamma s+\delta)} = \frac{1}{k} \left(\frac{\left(b - a\frac{\delta-\beta}{\gamma-\alpha}\right)s + b\alpha - a\beta - b\frac{\delta-\beta}{\gamma-\alpha}}{s^2+\alpha s+\beta} - \frac{\left(b - a\frac{\delta-\beta}{\gamma-\alpha}\right)s + b\gamma - a\delta - b\frac{\delta-\beta}{\gamma-\alpha}}{s^2+\gamma s+\delta} \right),$$

$\alpha \neq \gamma$.
[132]

$$(2.7) \quad \frac{1}{n^{2s}(n+1)^{2s}} = \sum_{t=1}^{2s} \binom{4s-t-1}{2s-1} \left[\frac{(-1)^t}{n^t} + \frac{1}{(n+1)^t} \right].$$

[65, 170.]

$$(2.8) \quad \int \frac{dx}{a^4+x^4} = \frac{1}{4a^3\sqrt{2}} \log \frac{x^2+ax\sqrt{2}+a^2}{x^2-ax\sqrt{2}+a^2} + \frac{1}{2a^3\sqrt{2}} \arctan \frac{ax\sqrt{2}}{a^2-x^2}.$$

[65, 171.]

$$(2.9) \quad \int \frac{dx}{a^4-x^4} = \frac{1}{4a^3} \log \left| \frac{a+x}{a-x} \right| + \frac{1}{2a^3} \arctan \frac{x}{a}.$$

[65, 170.1]

$$(2.10) \quad \int \frac{xdx}{a^4+x^4} = \frac{1}{2a^2} \arctan \frac{x^2}{a^2}.$$

[65, 170.2]

$$(2.11) \quad \int \frac{x^2 dx}{a^4+x^4} = -\frac{1}{4a\sqrt{2}} \log \frac{x^2+ax\sqrt{2}+a^2}{x^2-ax\sqrt{2}+a^2} + \frac{1}{2a\sqrt{2}} \arctan \frac{ax\sqrt{2}}{a^2-x^2}.$$

[65, 170.3]

$$(2.12) \quad \int \frac{x^3 dx}{a^4+x^4} = \frac{1}{4} \log(a^4+x^4).$$

[65, 171.1]

$$(2.13) \quad \int \frac{xdx}{a^4-x^4} = \frac{1}{4a^2} \log \left| \frac{a^2+x^2}{a^2-x^2} \right|.$$

[65, 171.2]

$$(2.14) \quad \int \frac{x^2 dx}{a^4-x^4} = \frac{1}{4a} \log \left| \frac{a+x}{a-x} \right| - \frac{1}{2a} \arctan \frac{x}{a}.$$

[65, 171.3]

$$(2.15) \quad \int \frac{x^3 dx}{a^4-x^4} = -\frac{1}{4} \log |a^4-x^4|.$$

[65, 173]

$$(2.16) \quad \int \frac{dx}{x(a+bx^m)} = \frac{1}{am} \log \left| \frac{x^m}{a+bx^m} \right|.$$

2.2. **Algebraic Functions.** [65, 186.11,188.11]

$$(2.17) \quad \int \frac{dx}{(a^2 + b^2x)x^{1/2}} = \frac{2}{ab} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.18) \quad \int \frac{dx}{(a^2 - b^2x)x^{1/2}} = \frac{1}{2ab} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 185.11,187.11]

$$(2.19) \quad \int \frac{x^{1/2}dx}{a^2 + b^2x} = \frac{2x^{1/2}}{b^2} - \frac{2a}{b^3} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.20) \quad \int \frac{x^{1/2}dx}{a^2 - b^2x} = -\frac{2x^{1/2}}{b^2} + \frac{a}{b^3} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 185.13,187.13]

$$(2.21) \quad \int \frac{x^{3/2}dx}{a^2 + b^2x} = \frac{2}{3} \frac{x^{3/2}}{b^2} - \frac{2a^2x^{1/2}}{b^4} + \frac{2a^3}{b^5} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.22) \quad \int \frac{x^{3/2}dx}{a^2 - b^2x} = -\frac{2}{3} \frac{x^{3/2}}{b^2} - \frac{2a^2x^{1/2}}{b^4} + \frac{a^3}{b^5} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 186.21,188.21]

$$(2.23) \quad \int \frac{dx}{(a^2 + b^2x)^2x^{1/2}} = \frac{x^{1/2}}{a^2(a^2 + b^2x)} + \frac{1}{a^3b} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.24) \quad \int \frac{dx}{(a^2 - b^2x)^2x^{1/2}} = \frac{x^{1/2}}{a^2(a^2 - b^2x)} + \frac{1}{2a^3b} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 185.21,187.21]

$$(2.25) \quad \int \frac{x^{1/2}dx}{(a^2 + b^2x)^2} = -\frac{x^{1/2}}{b^2(a^2 + b^2x)} + \frac{1}{ab^3} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.26) \quad \int \frac{x^{1/2}dx}{(a^2 - b^2x)^2} = \frac{x^{1/2}}{b^2(a^2 - b^2x)} - \frac{1}{2ab^3} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 185.23,187.23]

$$(2.27) \quad \int \frac{x^{3/2}dx}{(a^2 + b^2x)^2} = \frac{2x^{3/2}}{b^2(a^2 + b^2x)} + \frac{3a^2x^{1/2}}{b^4(a^2 + b^2x)} - \frac{3a}{b^5} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.28) \quad \int \frac{x^{3/2}dx}{(a^2 - b^2x)^2} = \frac{3a^2x^{1/2} - 2b^2x^{3/2}}{b^4(a^2 - b^2x)} - \frac{3a}{2b^5} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 186.13,188.13]

$$(2.29) \quad \int \frac{dx}{(a^2 + b^2x)x^{3/2}} = -\frac{2}{a^2x^{1/2}} - \frac{2b}{a^3} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.30) \quad \int \frac{dx}{(a^2 - b^2x)x^{3/2}} = -\frac{2}{a^2x^{1/2}} + \frac{b}{a^3} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 186.23,188.23]

$$(2.31) \quad \int \frac{dx}{(a^2 + b^2x)^2 x^{3/2}} = -\frac{2}{a^2(a^2 + b^2x)x^{1/2}} - \frac{3b^2x^{1/2}}{a^4(a^2 + b^2x)} - \frac{3b}{a^5} \arctan \frac{bx^{1/2}}{a}.$$

$$(2.32) \quad \int \frac{dx}{(a^2 - b^2x)^2 x^{3/2}} = -\frac{2}{a^2(a^2 - b^2x)x^{1/2}} + \frac{3b^2x^{1/2}}{a^4(a^2 - b^2x)} + \frac{3b}{2a^5} \log \left| \frac{a + bx^{1/2}}{a - bx^{1/2}} \right|.$$

[65, 189.2,189.4]

$$(2.33) \quad \int \frac{dx}{(a^4 + x^2)x^{1/2}} = \frac{1}{2a^3\sqrt{2}} \log \frac{x + a\sqrt{2x} + a^2}{x - a\sqrt{2x} + a^2} + \frac{1}{a^3\sqrt{2}} \arctan \frac{a\sqrt{2x}}{a^2 - x}.$$

$$(2.34) \quad \int \frac{dx}{(a^4 - x^2)x^{1/2}} = \frac{1}{2a^3} \log \left| \frac{a + x^{1/2}}{a - x^{1/2}} \right| + \frac{1}{a^3} \arctan \frac{x^{1/2}}{a}.$$

[65, 188.23,189.3]

$$(2.35) \quad \int \frac{x^{1/2}dx}{a^4 + x^2} = -\frac{1}{2a\sqrt{2}} \log \frac{x + a\sqrt{2x} + a^2}{x - a\sqrt{2x} + a^2} + \frac{1}{a\sqrt{2}} \arctan \frac{a\sqrt{2x}}{a^2 - x}.$$

$$(2.36) \quad \int \frac{x^{1/2}dx}{a^4 - x^2} = \frac{1}{2a} \log \left| \frac{a + x^{1/2}}{a - x^{1/2}} \right| - \frac{1}{a} \arctan \frac{x^{1/2}}{a}.$$

$$(2.37) \quad \int \frac{x^n}{\sqrt{a + bx}} dx = \frac{2\sqrt{a + bx}}{b^{n+1}} (-a)^n \sum_{k=0}^n \binom{n}{k} \frac{(-z/a)^k}{2k+1}.$$

This corrects two sign errors in [91] at the b :

$$(2.38) \quad \int \frac{z^m dx}{t^n \sqrt{z}} = -z^m \sqrt{z} \left\{ \frac{1}{(n-1)\Delta} \frac{1}{t^{n-1}} \right. \\ \left. + \sum_{k=2}^{n-1} \frac{(2n-2m-3)(2n-2m-5) \cdots (2n-2m-2k+1)(-b)^{k-1}}{2^{k-1}(n-1)(n-2) \cdots (n-k)\Delta^k} \frac{1}{t^{n-k}} \right\} \\ - \frac{(2n-3m-3)(2n-3m-5) \cdots (-2m+3)(-2m+1)(-b)^{n-1}}{2^{n-1}(n-1)!\Delta^n} \int \frac{z^m dx}{t\sqrt{z}}$$

where $z = a + bx$ and $t = \alpha + \beta x$ and $\Delta \equiv a\beta - \alpha b$.

[191]

$$(2.39) \quad \int \frac{f_m(z)}{\sqrt{D_m(z)}} dz = \log(x_m(z) + y_m(z)\sqrt{D_m(z)}),$$

if for example

$$(2.40) \quad f = 4z+2; \quad D = z^4+8(z+1); \quad x = z^4-2z^3+2z^2+4z-4; \quad y = z^2-2z+2.$$

or

$$(2.41) \quad f = 5z+1; \quad D = (z^2+1)^2+4z; \quad x = z^5-z^4+3z^3+z^2+2; \quad y = z^3-z^2+2z.$$

or

$$(2.42) \quad f = 6z+2; \quad D = (z^2+2)^2+8z; \quad x = z^6-2z^5+8z^4-4z^3+8z^2+8z; \quad y = z^4-2z^3+6z^2-4z+4.$$

or

(2.43)

$$f = 3z - s; \quad D = (z^2 - s^2)^2 + t(z - s); \quad x = 1 + 2(z + s)(z^2 - s^2)/t; \quad y = 2(z + s)/t.$$

2.3. The Exponential Function.

$$(2.44) \quad \int x^{m+2} e^{-ax^2} dx = \frac{m+1}{2a} \left[-x^{m+1} e^{-ax^2} + \int x^m e^{-ax^2} dx \right];$$

$$m \neq -1.$$

2.4. Hyperbolic Functions.

2.5. Trigonometric Functions.

$$\begin{aligned} \int \sin x \frac{(a + \sin x)^2}{(a + \sin x)^2 + (b + \cos x)^2} dx &= \frac{1}{4} \left[\frac{2b^2}{a^2 + b^2} + \frac{a^2 - b^2}{(a^2 + b^2)^2} - 3 \right] \cos(x) \\ &\quad - \frac{ab}{2} \left[\frac{1}{a^2 + b^2} - \frac{1}{(a^2 + b^2)^2} \right] \sin(x) \\ &+ \frac{1}{8} \frac{b \cos(2x) - a \sin(2x)}{a^2 + b^2} + \frac{b}{4} \left[1 + \frac{2(a^2 - b^2)}{(a^2 + b^2)^2} + \frac{-3a^2 + b^2}{(a^2 + b^2)^3} \right] \ln \sqrt{(\cos x + b)^2 + (\sin x + a)^2} \\ &+ \frac{a}{4} \left[1 - \frac{4b^2}{(a^2 + b^2)^2} - \frac{a^2 - 3b^2}{(a^2 + b^2)^3} \right] \arctan \frac{\sin x + a}{\cos x + b} + \frac{a}{4} \left[\frac{4b^2}{(a^2 + b^2)^2} + \frac{-3b^2 + a^2}{(a^2 + b^2)^3} \right] x. \end{aligned}$$

2.6. Rational Functions of Trigonometric Functions.

2.7. The Logarithm. [137]

$$(2.45) \quad \int \frac{\log z}{(1-z)z} dz = \text{Li}_2(1-z) + \frac{1}{2} \log^2 z.$$

[17, 3.1.6.]

$$(2.46) \quad \int \frac{\ln x}{1+ax} dx = \frac{1}{a} [\ln x \ln(1+ax) + \text{Li}_2(-ax)].$$

Correcting a sign error in [17, 3.1.7]:

$$(2.47) \quad \int \frac{\ln(a+bx)}{c+hx} dx = \frac{1}{h} \left[\ln \left(\frac{ah-bc}{h} \right) \ln(c+hx) - \text{Li}_2 \left(\frac{bc+bhx}{bc-ah} \right) \right].$$

(2.48)

$$\begin{aligned} \int z \ln(c+z+1/z) dz &= \frac{z^2 - c^2/2 + 1}{2} \ln(z^2 + cz + 1) - \frac{c\sqrt{c^2/4 - 1}}{2} \ln \frac{z + c/2 + \sqrt{c^2/4 - 1}}{z + c/2 - \sqrt{c^2/4 - 1}} \\ &\quad - \frac{(z-c)^2}{4} - \frac{1}{2} z^2 \ln z. \end{aligned}$$

3. DEFINITE INTEGRALS OF ELEMENTARY FUNCTIONS I.

3.1. General formulae. [8]

$$(3.1) \quad \int_0^\infty f([ax - b/x]^2) dx = \frac{1}{a} \int_0^\infty f(y^2) dy, \quad a, b > 0.$$

[34] Let

$$\varphi(x) = \sum_{k=0}^{\infty} A_k x^k,$$

then

$$(3.2) \quad \int_0^\infty x^{\beta-1} e^{-x} \varphi(x) dx = \sum_{k=0}^{\infty} A_k \Gamma(k + \beta), \quad \beta > 0.$$

Let

$$\varphi(x) = \sum_{k=0}^{\infty} A_k x^{k/p},$$

then

$$(3.3) \quad \int_0^\infty x^{\beta-1} e^{-x} \varphi(x) dx = \sum_{k=0}^{\infty} A_k \Gamma(k/p + \beta), \quad \beta > 0.$$

Let

$$\varphi(x) = \sum_{k=0}^{\infty} A_k x^k,$$

then

$$(3.4) \quad p \int_0^\infty x^{\beta-1} e^{-x^p} \varphi(x) dx = \sum_{k=0}^{\infty} A_k \Gamma\left(\frac{\beta + k}{p}\right), \quad \beta > 0.$$

[117]

$$(3.5) \quad \int_a^u \frac{x dx}{\sqrt{(x-a)(x-b)(x-c)}} = \frac{2}{\sqrt{a-c}} [(a-b)\Pi(\mu, 1, q) + bF(\mu, q)], \quad u > a > b > c.$$

[117]

$$(3.6) \quad \int_u^c \frac{dx}{(r-x)\sqrt{(a-x)(b-x)(c-x)}} = \frac{2(c-b)}{(r-b)(r-c)\sqrt{a-c}} \\ \times \Pi\left(\beta, \frac{r-b}{r-c}, p\right) + \frac{2}{(r-b)\sqrt{a-c}} F(\beta, p), \quad a > b > c > u, r \neq c.$$

[117]

$$(3.7) \quad \int_a^u \frac{dx}{(x-r)\sqrt{(x-a)(x-b)(x-c)}} = \frac{2}{(b-r)(a-r)\sqrt{a-c}} \\ \times \left[(b-a)\Pi\left(\mu, \frac{b-r}{a-b}, q\right) + (a-r)F(\mu, q) \right], \quad u > a > b > c, r \neq a.$$

[117]

$$(3.8) \quad \int_u^b \sqrt{\frac{(x-c)(b-x)}{a-x}} dx = \frac{2}{3} \sqrt{a-c} [2(b-a)F(\delta, q) \\ + (2a-b-c)E(\delta, q)] + \frac{2}{3} (b+c-a-u) \sqrt{\frac{(b-u)(u-c)}{a-u}}, \quad a > b > u \geq c.$$

[117]

$$(3.9) \quad \int_a^u \sqrt{\frac{(x-b)(x-c)}{x-a}} dx = \frac{2}{3} \sqrt{a-c} [2(a-b)F(\mu, q) \\ + (b+c-2a)E(\mu, q)] + \frac{2}{3} (u+2a-2b-c) \sqrt{\frac{(u-a)(u-c)}{u-b}}, \quad u > a > b > c.$$

[117]

$$(3.10) \quad \int_a^u \sqrt{\frac{(x-a)(x-c)}{x-b}} dx = \frac{2}{3} \sqrt{a-c} [(a+c-2b)E(\mu, q) \\ - (a-b)F(\mu, q)] + \frac{2}{3} (u+b-a-c) \sqrt{\frac{(u-a)(u-c)}{u-b}}, \quad u > a > b > c.$$

[58]

$$(3.11) \quad \int_0^\infty \frac{u^\alpha du}{(au+1)^\beta (bu+1)^\gamma} = A_0 \log a + B_0 \log b + \sum_{r=1}^{\beta-1} \frac{A_r}{r} + \sum_{s=1}^{\gamma-1} \frac{B_s}{s};$$

where

(3.12)

$$A_r = (-1)^{\alpha+\beta+r+1} a^{\gamma-\alpha-1} \sum_{j=0}^{\beta-\gamma-1} \binom{\alpha}{j} \binom{\beta+\gamma-r-j-2}{\gamma-1} \frac{b^{\beta-r-j-1}}{(a-b)^{\beta+\gamma-r-j-1}};$$

$$(3.13) \quad B_s = (-1)^\beta b^{\beta-\alpha-1} \sum_{j=0}^{\gamma-s-1} \binom{\alpha}{j} \binom{\beta+\gamma-s-j-2}{\beta-1} \frac{a^{\gamma-s-j-1}}{(a-b)^{\beta+\gamma-s-j-1}}.$$

(3.14)

$$r = 0, 1, \dots, \beta-1; \quad s = 0, 1, \dots, \gamma-1.$$

3.2. Powers of x , of binomials of the form $\alpha + \beta x^p$, and of polynomials in

 x . [64]

(3.15)

$${}_3F_2(-n, b/2, (b+1)/2; c/2, (c+1)/2; x) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-xt^2)^n dt, \quad \Re c > \Re b > 0.$$

[32]

$$(3.16) \quad \int_0^\infty \left[\frac{x^2}{x^4 + 2ax^2 + 1} \right]^r \cdot \frac{x^2 + 1}{x^b + 1} \frac{dx}{x^2} \\ = \int_0^\infty \left[\frac{x^2}{x^4 + 2ax^2 + 1} \right]^r \frac{dx}{x^2} = \int_0^\infty \left[\frac{x^2}{x^4 + 2ax^2 + 1} \right]^r dx \\ = \int_0^\infty \left[\frac{x^2}{x^4 + 2ax^2 + 1} \right]^r \frac{x^2 + 1}{x^2} dx = 2^{-1/2-r} (1+a)^{1/2-r} B(r-1/2, 1/2),$$

with B Euler's beta function, $a > 1, r > 1/2$, any b .

From this by specialization [32]

$$(3.17) \quad \int_0^\infty \frac{x^4}{(x^4 + x^2 + 1)^3} dx = \frac{\pi}{48\sqrt{3}}.$$

[32]

$$(3.18) \quad \int_0^\infty \frac{x^3}{(x^4 + 7x^2 + 1)^{5/2}} dx = \frac{2}{243}.$$

[32]

$$(3.19) \quad \int_0^\infty \frac{\sqrt{x}}{(x^4 + 14x^2 + 1)^{5/4}} dx = \frac{\Gamma^2(3/4)}{4\sqrt{2\pi}}.$$

[32]

$$(3.20) \quad \int_0^\infty \left[\frac{x^2}{bx^4 + 2ax^2 + c} \right]^r dx = \frac{B(r - 1/2, 1/2)}{2^{r+1/2}\sqrt{b}[a + \sqrt{bc}]^{r-1/2}}$$

with $b > 0, c \geq 0, a > -\sqrt{bc}$ and $r > 1/2$.

[32]

$$(3.21) \quad \int_0^\infty \left[\frac{x^2}{x^4 - x^2 + 1} \right]^r \frac{dx}{x^2} = \frac{\sqrt{\pi}}{2} \frac{\Gamma(r - 1/2)}{\Gamma(r)}$$

[10, 143]

$$(3.22) \quad \int_0^\infty \frac{dx}{(x^4 + 2ax^2 + 1)^{m+1}} = \frac{\pi}{2} \frac{\sum_{l=0}^m d_l(m) a^l}{[2(a+1)]^{m+1/2}}$$

where

$$(3.23) \quad d_l(m) \equiv 2^{-2m} \sum_{k=l}^m 2^k \binom{2m-2k}{m-k} \binom{m+k}{m} \binom{k}{l}.$$

[143]

$$(3.24) \quad \int_0^\infty \frac{dx}{bx^4 + 2ax^2 + 1} = \frac{\pi}{2\sqrt{2}} \frac{1}{\sqrt{a + \sqrt{b}}}.$$

[121]

$$(3.25) \quad \frac{(-)^n}{4^{n-1}} \int_0^1 \frac{x^{4n}(1-x)^{4n}}{1+x^2} dx = \pi - \sum_{k=0}^{n-1} (-)^k \frac{2^{4-2k}(4k)!(4k+3)!}{(8k+7)!} (820k^3 + 1533k^2 + 902k + 165).$$

[121]

$$(3.26) \quad \int_0^1 \frac{x^m(1-x)^n}{1+x^2} dx = \frac{\sqrt{\pi}\Gamma(m+1)\Gamma(n+1)}{2^{m+n+1}} {}_3F_2 \left(\begin{matrix} 1, (m+1)/2, (m+2)/2 \\ (m+n+2)/2, (m+n+3)/2 \end{matrix} \mid -1 \right).$$

[19]

$$(3.27) \quad \int_0^1 \frac{x^m(1-x)^n}{1+x^2} dx = R_{m,n} + A_{m,n}\pi + B_{m,n} \ln \sqrt{2}$$

induced by the partial fraction decomposition

$$(3.28) \quad \frac{x^m(1-x)^n}{1+x^2} = Q_{m,n}(x) + \frac{A_{m,n}}{1+x^2} + \frac{B_{m,n}x}{1+x^2}.$$

[170][175, A093341]

$$(3.29) \quad \int_0^\infty \frac{dx}{\sqrt{1+x^4}} = \mathbf{K}\left(\frac{1}{\sqrt{2}}\right).$$

[170]

$$(3.30) \quad \int_0^\infty \frac{dx}{\sqrt{x+x^4}} = \mathbf{K}\left(\frac{\sqrt{6}-\sqrt{2}}{4}\right).$$

[170]

$$(3.31) \quad \int_0^1 \frac{dx}{\sqrt[4]{1-x^2}} = \sqrt{3} \left[2\mathbf{E}\left(\frac{1}{\sqrt{2}}\right) - \mathbf{K}\left(\frac{1}{\sqrt{2}}\right) \right].$$

[170][175, A062539]

$$(3.32) \quad \int_0^1 \frac{dx}{(1-x^2)^{3/4}} = \sqrt{2}\mathbf{K}\left(\frac{1}{\sqrt{2}}\right).$$

[170]

$$(3.33) \quad \int_0^1 \frac{x^2}{(1-x^2)^{3/4}} dx = \frac{2^{3/2}}{3} \mathbf{K}\left(\frac{1}{\sqrt{2}}\right).$$

[170]

$$(3.34) \quad \int_0^1 \frac{1}{\sqrt{1-x^6}} dx = \frac{1}{\sqrt[4]{3}} \mathbf{K}\left(\frac{\sqrt{6}-\sqrt{2}}{4}\right).$$

[170]

$$(3.35) \quad \int_0^\infty \frac{1}{\sqrt{1+x^6}} dx = \frac{2}{\sqrt[4]{27}} \mathbf{K}\left(\frac{\sqrt{6}-\sqrt{2}}{4}\right).$$

[170]

$$(3.36) \quad \int_0^1 \frac{1}{\sqrt{1-x^8}} dx = \frac{1}{\sqrt{2}} \mathbf{K}(\sqrt{2}-1).$$

[170]

$$(3.37) \quad \int_0^1 \frac{x^2}{\sqrt{1-x^8}} dx = \left(1 - \frac{1}{\sqrt{2}}\right) \mathbf{K}(\sqrt{2}-1).$$

[170]

$$(3.38) \quad \int_0^1 \frac{x^{a-1}}{\sqrt{1-x^n}} dx = \cos(a\pi/n) \int_0^\infty \frac{z^{a-1}}{\sqrt{1+z^n}} dz, \quad 2a < n.$$

[170]

$$(3.39) \quad \int_0^\infty \frac{z^{n-a-1}}{\sqrt{1+z^n}} dz \cdot \int_0^1 \frac{x^{a-1}}{\sqrt{1-x^n}} dx = \frac{2\pi}{n(2a-n)\sin(\pi a/n)}, \quad n/2 < a < n.$$

[170]

$$(3.40) \quad \int_0^\infty \frac{1}{\sqrt{1+x^8}} dx = \int_0^\infty \frac{x^2}{\sqrt{1+x^8}} dx = \sqrt{2-\sqrt{2}} \mathbf{K}(\sqrt{2}-1).$$

[170]

$$(3.41) \quad \int_0^1 \frac{x^4}{\sqrt{1-x^8}} dx = \frac{\pi}{8} \frac{\sqrt{2}}{\mathbf{K}(\sqrt{2}-1)}.$$

[170]

$$(3.42) \quad \int_0^1 \frac{x^6}{\sqrt{1-x^8}} dx = \frac{\pi}{24} \frac{2+\sqrt{2}}{\mathbf{K}(\sqrt{2}-1)}.$$

[170]

$$(3.43) \quad \int_0^\infty \frac{1}{\sqrt[3]{1+x^6}} dx = \frac{\sqrt[3]{4}}{\sqrt[4]{3}} \mathbf{K}\left(\frac{\sqrt{6}-\sqrt{2}}{4}\right).$$

[170]

$$(3.44) \quad \int_0^1 \frac{1}{\sqrt[3]{1-x^6}} dx = \frac{\sqrt[3]{4}}{\sqrt[4]{27}} \mathbf{K}\left(\frac{\sqrt{6}-\sqrt{2}}{4}\right).$$

[170]

$$(3.45) \quad \int_1^\infty \frac{1}{\sqrt[m]{(x^n+a)(x^n+b)}} dx = \frac{m}{2n-m} F_1\left(\begin{matrix} \frac{2n-m}{m}; 1/m, 1/n \\ \frac{2n-m+mn}{mn} \end{matrix} \mid -a, -b\right), \quad 2n-m > 0, \quad a, b > 0.$$

[170]

$$(3.46) \quad \int_0^1 \frac{1}{\sqrt[m]{(x^n+a)(x^n+b)}} dx = \frac{1}{\sqrt[m]{ab}} F_1\left(\begin{matrix} 1/n; 1/m, 1/m \\ 1+1/n \end{matrix} \mid -1/a, -1/b\right), \quad a, b > 0.$$

3.3. The Exponential function. [11]

$$(3.47) \quad -\int_0^2 \frac{x^p - x^q}{1-x} dx = \psi(p+1) - \psi(q+1)$$

[110]

$$(3.48) \quad \int_u^\infty \exp\left(-\frac{x^2}{4\beta} - \gamma x\right) dx = \sqrt{\pi\beta} e^{\beta\gamma^2} \left[1 - \Phi\left(\gamma\sqrt{\beta} + \frac{u}{2\sqrt{\beta}}\right)\right], \quad \Re\beta > 0, u \geq 0.$$

[110]

$$(3.49) \quad \int_{-\infty}^\infty \exp(-p^2 x^2 \pm qx) dx = \exp\left(\frac{q^2}{4p^2}\right) \frac{\sqrt{\pi}}{|p|}.$$

[110]

$$(3.50) \quad \int_0^\infty \frac{x^n e^{-\mu x}}{x+\beta} dx = (-1)^{n-1} \beta^n e^{\beta\mu} \operatorname{Ei}(-\beta\mu) + \sum_{k=1}^n (k-1)! (-\beta)^{n-k} \mu^{-k}, \quad |\arg \beta| < \pi, \Re\mu > 0, n \geq 0.$$

3.4. Rational functions of powers and exponentials. [110]

$$(3.51) \quad \int_0^\infty \frac{x^{\nu-1} e^{-\mu x}}{1-\beta e^{-x}} dx = \Gamma(\nu) \sum_{n=0}^\infty (\mu+n)^{-\nu} \beta^n.$$

[110]

$$(3.52) \quad \int_0^\infty \frac{(1+ix)^{2n-1} - (1-ix)^{2n-1}}{i} \frac{dx}{e^{\pi x} + 1} = \frac{1}{2n} [1 - 2^{2n} B_{2n}].$$

[110]

$$(3.53) \quad \int_0^\infty \frac{x^q e^{-px}}{(1-ae^{-px})^2} dx = \frac{\Gamma(q+1)}{ap^{q+1}} \sum_{k=1}^\infty \frac{a^k}{k^q}, \quad -1 \leq a < 1, q > -1, p > 0.$$

[110]

$$(3.54) \quad \int_0^\infty \frac{(1+a)e^x + a}{(1+e^x)^2} e^{-ax} x^n dx = -n! \sum_{k=1}^\infty \frac{(-1)^k}{(a+k)^n}, \quad a > -1, \quad n = 1, 2, \dots$$

[34]

$$(3.55) \quad p \int_0^\infty x^{\beta-1} e^{ax-x^p} dx = \sum_{k=0}^\infty \frac{a^k}{k!} \Gamma\left(\frac{k+\beta}{p}\right).$$

$$(3.56) \quad p^2 \int_0^\infty t^{p-1} e^{at-t^p} dt = \sum_{k=1}^\infty \frac{a^k}{(k-1)!} \Gamma(k/p).$$

$$(3.57) \quad \int_0^\infty x^{\beta-1} e^{-x-x^p} dx = \sum_{j=0}^\infty \frac{(-1)^j}{j!} \Gamma(\beta + jp).$$

[171][175, A002161]

$$(3.58) \quad \int_0^\infty (1 - e^{-1/x^2}) dx = \sqrt{\pi}.$$

[110][175, A155739]

$$(3.59) \quad \int_0^\infty \left\{ e^{-x^2} - \frac{1}{1+x^{2n}} \right\} \frac{dx}{x} = -\frac{1}{2}C \approx -0.288607.$$

[192]

$$(3.60) \quad \int_0^\infty \exp(nx - \beta \operatorname{sh} x) dx = \frac{1}{2} [S_n(\beta) - \pi \mathbf{E}_n(\beta) - \pi N_n(\beta)].$$

[192]

$$(3.61) \quad \int_{-\infty}^\infty \frac{\exp(\nu \operatorname{Arsh} x - iax)}{\sqrt{1+x^2}} = \begin{cases} 2 \exp\left(-\frac{i\nu\pi}{2}\right) K_\nu(a) & \text{if } a > 0, \\ 2 \exp\left(\frac{i\nu\pi}{2}\right) K_\nu(-a) & \text{if } a < 0. \end{cases}$$

[$|\Re \nu| < 1$]

3.5. Hyperbolic Functions. [6, 42]

$$(3.62) \quad \frac{1}{2} \int_0^\infty \frac{x}{\cosh x} dx = G.$$

[14]

$$(3.63) \quad \int_0^\infty \frac{du}{(b^2 + u^2) \sinh(au)} = \frac{1}{2b} \left[\psi\left(\frac{ab}{2\pi} + \frac{3}{4}\right) - \psi\left(\frac{ab}{2\pi} + \frac{1}{4}\right) \right]$$

for $\Re a > 0$, $\Re b > \max(-\Re a, -\Re 3a)$.

[14]

$$(3.64) \quad \int_0^\infty \frac{u du}{(b^2 + u^2) \sinh(au)} = \frac{1}{2} \left[\psi\left(\frac{ab}{2\pi} + \frac{1}{2}\right) - \psi\left(\frac{ab}{2\pi}\right) \right] - \frac{\pi^2}{4a^2b}$$

for $\Re a > 0$, $\Re b > 0$.

[34]

$$(3.65) \quad \int_0^\infty x^{\beta-1} e^{-x} \frac{\sinh b\sqrt{x}}{b\sqrt{x}} dx = \sum_{j=0}^\infty \frac{\Gamma(j+\beta)}{(2j+1)!} b^{2j}.$$

$$(3.66) \quad \int_0^\infty t^{2\beta-2} e^{-t^2/b^2} \sinh t dt = \frac{1}{2} \sum_{j=0}^\infty \frac{\Gamma(j+\beta)}{(2j+1)!} b^{2(j+\beta)}.$$

$$(3.67) \quad \int_0^\infty x^{\beta-1} e^{-x} \sinh \sqrt{x} dx = \sum_{k=1}^\infty \frac{\Gamma(\beta+k)}{\Gamma(2k)}.$$

[42]

$$(3.68) \quad \int_0^{\pi/2} \sinh^{-1}(\sin x) dx = \int_0^{\pi/2} \sinh^{-1}(\cos x) dx = G.$$

[42]

$$(3.69) \quad \int_0^{\pi/2} \operatorname{csch}^{-1}(\csc x) dx = \int_0^{\pi/2} \operatorname{csch}^{-1}(\sec x) dx = G.$$

3.6. Rational Functions of Sines and Cosines. [110]

$$(3.70) \quad \int_0^{\pi/2} \frac{\sin 2nx \cos^{2m+1} x}{\sin x} dx = \frac{\pi}{2}, \quad n > m \geq 0.$$

[61, B2b]

$$(3.71) \quad \int_0^{2\pi} \sin^m \theta \cos^n \theta d\theta = 2\pi \epsilon_m \epsilon_n \frac{(m-1)!!(n-1)!!}{(m+n)!!}$$

where $\epsilon_j = 1$ if j is even and $\epsilon_j = 0$ otherwise.

[26]

$$(3.72) \quad \int_0^z \sin^\mu t \sin^\nu(z-t) dt = \frac{\sqrt{\pi} \Gamma(\mu+1) \Gamma(\nu+1)}{2^{(\mu+\nu+1)/2} \Gamma(\mu/2 + \nu/2 + 1)} \sin^{(\mu+\nu+1)/2} z P_{(\mu-\nu-1)/2}^{-(\mu+\nu+1)/2}(\cos z), \quad \Re \mu > -1, \Re \nu > -1.$$

[26]

$$(3.73) \quad \frac{2^m \Gamma(m+1/2)}{\sqrt{\pi} \Gamma(m+n+1) \Gamma(m-n)} \int_0^z \sin^{m+n} t \sin^{m-n-1}(z-t) dt = \sin^m z P_n^{-m}(\cos z).$$

[26]

$$(3.74) \quad \int_0^z \left(\frac{\sin t}{\sin(z-t)} \right)^\mu dt = \frac{\pi \sin \mu z}{\sin \mu \pi}, \quad -1 < \Re \mu < 1.$$

[124]

$$(3.75) \quad \int_0^{\arcsin q} \left(\cos \phi \pm \sqrt{q^2 - \sin^2 \phi} \right)^{n+2} T_m(\cos \phi) d\phi = \dots$$

[6]

$$(3.76) \quad \int_0^{\pi/2} \sinh^{-1}(\sin x) dx = G.$$

[6]

$$(3.77) \quad \int_0^{\pi/2} \sinh^{-1}(\cos x) dx = G.$$

[6]

$$(3.78) \quad \int_0^{\pi/2} \operatorname{csch}^{-1}(\csc x) dx = G.$$

[6]

$$(3.79) \quad \int_0^{\pi/2} \operatorname{csch}^{-1}(\sec x) dx = G.$$

[122, p40]

$$(3.80) \quad \int_0^\infty \cos(T_n(t, -x)) dt = \frac{\pi\sqrt{x}}{2n \sin \frac{\pi}{2n}} \left[J_{1/n}(2x^{n/2}) - J_{-1/n}(2x^{n/2}) \right],$$

(3.81)

$$\int_0^\infty \cos(T_{2m}(t, x)) dt = \frac{\pi\sqrt{x}}{4m \sin \frac{\pi}{4m}} \left[J_{-1/(2m)}(2x^m) - J_{1/(2m)}(2x^m) \right], \quad m = 1, 2, 3 \dots$$

(3.82)

$$\int_0^\infty \cos(T_{2m+1}(t, x)) dt = \frac{2\sqrt{x} \cos \frac{\pi}{4m+2}}{2m+1} K_{1/(2m+1)}(2x^{m+1/2}), \quad m = 1, 2, 3 \dots$$

where x real positive, where

$$(3.83) \quad T_n(t, x) \equiv t^n {}_2F_1\left(-\frac{n}{2}, \frac{1-n}{2}; 1-n; -\frac{4x}{t^2}\right), \quad n = 2, 3, 4, \dots$$

for example $T_2 = t^2 + 2x$, $T_3 = t^3 + 3tx$.

3.7. Trigonometric and Rational Functions. [6, 42]

$$(3.84) \quad \frac{1}{2} \int_0^{\pi/2} \frac{x}{\sin x} = G.$$

[6]

$$(3.85) \quad \frac{3}{4} \int_0^{\pi/6} \frac{x}{\sin x} = G - \frac{\pi}{8} \log(1 + \sqrt{3}).$$

[6, 42]

$$(3.86) \quad -\frac{\pi^2}{4} \int_0^1 \left(x - \frac{1}{2}\right) \sec(\pi x) dx = G.$$

[11]

$$(3.87) \quad \frac{2^{s+2}}{\pi} \int_0^{\pi/2} x \cos^s x \sin(sx) dx - \gamma = \psi(s+1).$$

[110]

$$(3.88) \quad \int_0^{\pi/4} x^m \tan x dx = 2\pi G - \frac{7}{2} \zeta(3), \quad m = 2.$$

[57]

$$(3.89) \quad \int_0^{1/2} x^n \cot \pi x dx = \frac{n!}{2^n} \sum_{k=1, \text{ odd}}^n \frac{(-)^{(k-1)/2}}{\pi^k} \frac{\eta(k)}{(n-k+1)!} + \frac{(-)^n + 1}{2} \frac{4n!(1-2^{-n-1})}{(2\pi)^{n+1}} \zeta(n+1),$$

where $\eta(s) \equiv (1-2^{1-s})\zeta(s)$.

[149, p 6]

$$(3.90) \quad \int_0^\infty \frac{\cos(xy)}{(a^2 + x^2)^{\nu+1/2}} dx = \sqrt{\pi} \left(\frac{y}{2a}\right)^\nu \frac{1}{\Gamma(\nu+1/2)} K_\nu(ay).$$

[149, p 10]

(3.91)

$$\int_0^\infty \frac{x^\nu}{(a^2 + x^2)^{\mu+1}} \cos(xy) dx = \frac{a^{\nu-2\mu-1}}{2} B\left(\frac{1}{2} + \frac{1}{2}\nu, \mu - \frac{1}{2}\nu + \frac{1}{2}\right) {}_1F_2\left(\frac{\nu+1}{2}; \frac{\nu+1}{2} - \mu, \frac{1}{2}, \frac{a^2 y^2}{4}\right)$$

(3.92)

$$+ \sqrt{\pi} \frac{2^{-2\mu+\nu-2}}{\Gamma(1+\mu-\frac{1}{2}\nu)} y^{2\mu-\nu+1} \Gamma\left(\frac{1}{2}\nu - \mu - \frac{1}{2}\right) {}_1F_2\left(\mu+1 - \frac{\nu}{2}; \mu - \frac{\nu}{2} + \frac{3}{2}, \frac{a^2 y^2}{4}\right).$$

[110]

(3.93)

$$\int_0^\infty \frac{x^{2m} \cos(ax) dx}{(\beta^2 + x^2)^{n+1/2}} = \frac{(-1)^m \sqrt{\pi}}{2^n \beta^n \Gamma(n+1/2)} \cdot \frac{d^{2m}}{da^{2m}} \{a^n K_n(a\beta)\}, \quad a > 0, \Re \beta > 0, 0 \leq m \leq n.$$

[149, p 116]

$$(3.94) \quad \int_0^\infty \frac{\sin(xy)}{x(a^2 + x^2)^{\nu+1/2}} dx = \frac{\pi y}{2a^{2\nu}} [K_\nu \mathbf{L}_{\nu-1}(ay) + \mathbf{L}_\nu(ay) K_{\nu-1}(ay)]$$

where \mathbf{L} are Struve functions.

[6]

$$(3.95) \quad \frac{1}{2\pi} \int_0^{\pi/2} \frac{x^2}{\sin x} dx = G - \frac{7}{4\pi} \zeta(3).$$

[42]

$$(3.96) \quad \int_0^{\pi/4} \frac{x^2}{\sin^2 x} dx = G - \frac{1}{16} \pi^2 + \frac{1}{4} \pi \log 2.$$

[6, 42]

$$(3.97) \quad \int_0^{\pi/2} \frac{x \csc x}{\cos x + \sin x} dx = G + \frac{\pi}{4} \log 2.$$

[6, 42]

$$(3.98) \quad -2 \int_0^{\pi/2} \frac{x \cos x}{\cos x + \sin x} dx = G - \frac{\pi^2}{8} - \frac{\pi}{4} \log 2.$$

[6, 42]

$$(3.99) \quad 2 \int_0^{\pi/2} \frac{x \sin x}{\cos x + \sin x} dx = G + \frac{\pi^2}{8} - \frac{\pi}{4} \log 2.$$

[42]

$$(3.100) \quad \frac{3}{4} \int_0^{\pi/6} \frac{x}{\sin x} dx = G - \frac{1}{8} \pi \log(2 + \sqrt{3}).$$

[9]

$$(3.101) \quad \int_0^\infty x^{-p} \cos^{2n+1}(x+b) dx = \frac{\Gamma(1-p)}{2^{2n}} \sum_{k=0}^n \binom{2n+1}{n-k} \frac{\sin[\pi p/2 - (2k+1)b]}{(2k+1)^{1-p}}.$$

[9]

$$(3.102) \quad \int_0^\infty x^{-p} \sin^{2n+1}(x+b) dx = \frac{\Gamma(1-p)}{2^{2n}} \sum_{k=0}^n \binom{2n+1}{n-k} \frac{\cos[\pi p/2 - (2k+1)b]}{(2k+1)^{1-p}}$$

[9]

$$(3.103) \quad \int_0^\infty x^{-p} \cos^{2n+1} x dx = \frac{\Gamma(1-p)}{2^{2n}} \sin\left(\frac{\pi p}{2}\right) \sum_{k=0}^n \frac{\binom{2n+1}{n-k}}{(2k+1)^{1-p}}$$

for $0 < p < 1$.

[9]

$$(3.104) \quad \int_0^\infty x^{-p} \sin^{2n+1} x dx = \frac{\Gamma(1-p)}{2^{2n}} \cos\left(\frac{\pi p}{2}\right) \sum_{k=0}^n (-1)^k \frac{\binom{2n+1}{n-k}}{(2k+1)^{1-p}}$$

for $0 < p < 1$.

[9]

$$(3.105) \quad \int_0^\infty \cos^{2n+1} x^p dx = \frac{1}{2^{2n}} \Gamma\left(\frac{p+1}{p}\right) \cos\left(\frac{\pi}{2p}\right) \sum_{k=0}^n \frac{\binom{2n+1}{n-k}}{(2k+1)^{1/p}}$$

for $p > 1$.

[9]

$$(3.106) \quad \int_0^\infty \sin^{2n+1} x^p dx = \frac{1}{2^{2n}} \Gamma\left(\frac{p+1}{p}\right) \sin\left(\frac{\pi}{2p}\right) \sum_{k=0}^n (-1)^k \frac{\binom{2n+1}{n-k}}{(2k+1)^{1/p}}$$

for $p > 1$.

[9]

$$(3.107) \quad \int_0^{\pi/2} x^p \cos^{2n} x dx = \sum_{j=0}^{\lfloor p/2 \rfloor} a_{n,p,p+1-2j} \pi^{p+1-2j} + \delta_{\text{odd},p} \cdot a_{n,p}^*,$$

where for $p \geq 2$ and $0 \leq j \leq \lfloor p/2 \rfloor$

$$(3.108) \quad a_{n,p,p+1-2j} = \frac{(-1)^j \binom{2n}{n} p!}{2^{2n+p+1} (p+1-2j)!} \sum_{1 \leq k_1 \leq k_2 \leq \dots \leq k_j \leq n} \frac{1}{k_1^2 k_2^2 \dots k_j^2}.$$

and $a_{n,p}^*$ is a similar multinomial sum. A similar form exists for odd powers of the cosine.

[110]

$$(3.109) \quad \int_0^\infty \sin^{2m+1} x \frac{xdx}{a^2 + x^2} = \frac{\pi}{2^{2m+1}} e^{-(2m+1)a} \sum_{k=0}^m (-1)^{m+k} \binom{2m+1}{k} e^{2ka}.$$

[110]

$$(3.110) \quad \int_0^\infty \cos^{2m} x \frac{xdx}{a^2 + x^2} = \frac{\pi}{2^{2m+1}a} \binom{2m}{m} + \frac{\pi}{2^{2m}a} \sum_{k=1}^m \binom{2m}{m+k} e^{-2ka}, \quad a > 0.$$

[69]

$$I(a, b) \equiv \int_0^\infty x^{-a} \left(1 - \frac{\sin^b x}{x^b}\right) dx.$$

then

$$(3.111) \quad I(a, b) = \frac{\pi \sec(\pi a/2)}{2^b \Gamma(a+b)} \sum_{k=0}^{\lfloor (b-1)/2 \rfloor} (-1)^{k+1} \binom{b}{k} (b-2k)^{a+b-1},$$

with special cases

$$(3.112) \quad I(2, b) = \frac{\pi}{2^b (b+1)!} \sum_{k=0}^{\lfloor (b-1)/2 \rfloor} (-1)^{k+1} \binom{b}{k} (b-2k)^{b+1},$$

$$(3.113) \quad I(a, 1) = -\frac{\pi \sec(\pi a/2)}{2\Gamma(1+a)} \int_0^\infty x^{-3/2} \left(1 - \frac{\sin x}{x}\right) dx = \frac{2\sqrt{2\pi}}{3},$$

$$(3.114) \quad I(a, 2) = -\frac{\pi 2^{a-1} \sec(\pi a/2)}{\Gamma(2+a)} \int_0^\infty x^{-3/2} \left(1 - \frac{\sin^2 x}{x^2}\right) dx = \frac{16\sqrt{\pi}}{15},$$

$$(3.115) \quad I(a, 3) = \frac{(3 - 3^{2+a})\pi \sec(\pi a/2)}{8\Gamma(3+a)} \int_0^\infty x^{-3/2} \left(1 - \frac{\sin^3 x}{x^3}\right) dx = \frac{2}{35}(9\sqrt{3} - 1)\sqrt{2\pi}.$$

3.8. Trigonometric Functions and Exponentials. [120]

$$(3.116) \quad \int_0^\infty e^{-tx^2} \cos x^2 dx = \sqrt{\pi/8} \sqrt{\frac{\sqrt{1+t^2} + t}{1+t^2}}.$$

[120]

$$(3.117) \quad \int_0^\infty e^{-tx^2} \sin x^2 dx = \sqrt{\pi/8} \sqrt{\frac{\sqrt{1+t^2} - t}{1+t^2}}.$$

4. DEFINITE INTEGRALS OF ELEMENTARY FUNCTIONS II

[167] In terms of the constant (0.102) we have

$$(4.1) \quad \int_0^1 \frac{\log^2 u}{u} \log(1+u) du = \frac{7\pi^4}{360},$$

$$(4.2) \quad \int_0^1 \frac{\log^2 u}{u} \log(1-u) du = -\frac{\pi^4}{45},$$

$$(4.3) \quad \int_0^1 \frac{\log^2 u}{u} \log \frac{1+u}{1-u} du = \frac{\pi^4}{24},$$

$$(4.4) \quad \int_0^1 \frac{\log u}{u} \log^2(1+u) du = A_4 - \frac{\pi^4}{288},$$

$$(4.5) \quad \int_0^1 \frac{\log u}{u} \log^2(1-u) du = -\frac{\pi^4}{180},$$

$$(4.6) \quad \int_0^1 \frac{\log u}{u} \log^2 \frac{1+u}{1-u} du = 2A_4 - \frac{\pi^4}{60},$$

$$(4.7) \quad \int_0^1 \frac{1}{u} \log^3(1+u) du = \frac{3}{2}A_4 - \frac{\pi^4}{960},$$

$$(4.8) \quad \int_0^1 \frac{1}{u} \log^3(1-u) du = -\frac{\pi^4}{15},$$

[167]

$$(4.9) \quad \int_0^1 \frac{\log u}{u} \log(1+u) du = -\frac{3}{4}\zeta(3),$$

$$(4.10) \quad \int_0^1 \frac{\log u}{u} \log(1-u) du = \zeta(3),$$

$$(4.11) \quad \int_0^1 \frac{\log u}{u} \log \frac{1+u}{1-u} du = -\frac{7}{4}\zeta(3),$$

$$(4.12) \quad \int_0^1 \frac{1}{u} \log^2(1+u) du = \frac{1}{4}\zeta(3),$$

$$(4.13) \quad \int_0^1 \frac{1}{u} \log^2(1-u) du = 2\zeta(3).$$

[110]

$$(4.14) \quad 2 \int_0^{\pi/2} \ln |1 - \sin x| dx = -\pi \ln 2 - 4G.$$

[42]

$$(4.15) \quad -2 \int_0^{\pi/4} \log(2 \sin x) dx = G.$$

A factor 2 is missing in [6].

[42]

$$(4.16) \quad \frac{1}{4} \int_0^{\pi/2} \log \frac{1 + \cos x}{1 - \cos x} dx = G.$$

[42]

$$(4.17) \quad \frac{1}{4} \int_0^{\pi/2} \log \frac{1 + \sin x}{1 - \sin x} dx = G.$$

[110]

$$(4.18) \quad \int_0^{\pi/2} (\ln \tan x)^{2n} dx = (\pi/2)^{2n+1} |E_{2n}|.$$

[110]

$$(4.19) \quad \int_0^\infty \frac{\ln x dx}{(x+a)^2} = \frac{\ln a}{a}, \quad 0 < a.$$

[6, 42]

$$(4.20) \quad 2 \int_0^{\pi/4} \log(2 \cos x) dx = G.$$

[11, 144][175, A115252]

$$(4.21) \quad \int_{\pi/4}^{\pi/2} \ln \ln \tan x dx = \int_0^1 \ln \ln \left(\frac{1}{x} \right) \frac{dx}{1+x^2} = \frac{\pi}{2} \ln \left(\frac{\Gamma(3/4)\sqrt{2\pi}}{\Gamma(1/4)} \right)$$

[142]

$$(4.22) \quad \frac{1}{4} \int_0^1 \frac{x^4 - 6x^2 + 1}{(1+x^2)^3} \log \log(1/x) dx = -(\gamma + \log 4)[\zeta(-1, 1/4) - \zeta(-1, 3/4)] + \zeta'(-1, 1/4) - \zeta'(-1, 3/4).$$

[6]

$$(4.23) \quad - \int_0^1 \frac{\log(x)}{x^2 + 1} dx = G.$$

[56]

$$(4.24) \quad \int_0^1 \frac{\log t}{1 - (1-t)(1-u)} dt = \frac{1}{1-u} \text{Li}_2 \left(-\frac{u-1}{u} \right)$$

[11]

$$(4.25) \quad \int_0^b \frac{\ln t}{(1+t)^{n+1}} dt = \frac{1}{n} [1 - (1+b)^{-n}] \ln b - \frac{1}{n} \ln(1+b) - \frac{1}{n(1+b)^{n-1}} \sum_{j=1}^{n-1} \frac{1}{j!} \binom{n-1}{j} |S_{j+1}^{(2)}| b^j$$

[4]

$$(4.26) \quad \frac{z^p}{(p-1)!} \int_0^1 \frac{t^{z-1}}{(1-t)^z} \log^{p-1} \frac{1}{t} dt = {}_{p+1}F_p \left(\begin{matrix} z, z, \dots, z \\ z+1, \dots, z+1 \end{matrix} \middle| 1 \right).$$

[4]

$$(4.27) \quad \frac{1}{\Gamma(1-z)\Gamma(p)} \int_0^1 \frac{t^{z-1}}{(1-t)^z} \log^{p-1}(t) dt = \left[\begin{matrix} z \\ p \end{matrix} \right].$$

[11]

$$(4.28) \quad \int_0^x \frac{\ln t}{(1+t^2)^{n+1}} dt = \frac{\binom{2n}{n}}{2^{2n}} \left[g_0(x) + p_n(x) \ln x - \sum_{k=0}^{n-1} \frac{\tan^{-1} x + p_k(x)}{2k+1} \right]$$

where

$$(4.29) \quad g_0(x) \equiv \ln x \tan^{-1} x - \int_0^x \frac{\tan^{-1} t}{t} dt.$$

[144]

$$(4.30) \quad \int_0^\infty \frac{\ln^{n-1} x dx}{(x-1)(x+a)} = \frac{(-)^n (n-1)!}{1+a} \{ [1 + (-)^n] \zeta(n) - \sum_{j=0}^{[n/2]} \binom{n}{2j} (2^{2j} - 2) (-)^j B_{2j} \pi^{2j} \log^{n-2j} a \}$$

for $n \geq 2$, $a > 0$.

[177]

$$(4.31) \quad -2t \int_0^1 \frac{(1-x)^{j+1} \log(1-x)}{(1-tx(1-x))^3} dx = \sum_{n \geq 1} \frac{t^n}{C_n(j)} \sum_{r=1}^n \frac{1}{r+j+n}$$

where $C_n(j) = \binom{2n+j}{n}/(n+1)$ are Catalan related numbers. For $t = 2$ the integral becomes a sum of G , $\zeta(2)$, $\pi \ln 2$, π and 1 with rational coefficients, and similar results are given for $t = 1/2$.

[20]

$$(4.32) \quad \int_0^\infty \frac{\log(1+x)}{1+x+x^2} dx = - \int_0^1 \frac{(1+t) \log t}{1+t^3} dt = -\frac{3}{2} L_{-3}(2),$$

where $L_{-3}(s)$ is a Dirichlet series [175, A086724].

[11]

$$(4.33) \quad \int_0^1 \frac{\ln t}{(1+t^2)^{n+1}} dt = -2^{-2n} \binom{2n}{n} \left(G + \sum_{k=0}^{n-1} \frac{\frac{\pi}{4} + p_k(1)}{2k+1} \right)$$

where

$$(4.34) \quad p_k(1) = \sum_{j=1} k \frac{2^j}{2j \binom{2j}{j}}.$$

[6]

$$(4.35) \quad \int_0^1 \left(\frac{2}{x^2 - 4x + 8} - \frac{3}{x^2 + 2x + 2} \right) \log x dx = C.$$

[56]

$$(4.36) \quad (-1)^{p+1} n \int_0^1 (1-t)^{n-1} \log^p t dt = p! \sum_{k=1}^n \binom{n}{k} \frac{(-1)^k}{k^p}.$$

[56][175, A152648]

$$(4.37) \quad \int_0^1 \frac{\log^2 t}{1-t} dt = \sum_{n=1}^\infty \frac{H_n^{(1)}}{n^2} = 2\zeta(3) = 2\text{Li}_3(t) - 2\text{Li}_2(t) \log t - \log(1-t) \log^2 t + c$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.

[56]

$$(4.38) \quad \int_0^1 \frac{\log^3 u}{1-u} du = -6\zeta(4).$$

[196]

$$(4.39) \quad \int_0^{\pi/2} \frac{\ln(2 \cos x)}{x^2 + \ln^2(2 \cos x)} dx = \pi/4.$$

[196]

$$(4.40) \quad \int_0^{\pi/2} \ln[x^2 + \ln^2(2 \cos x)] dx = 0.$$

[196]

$$(4.41) \quad \int_0^{\pi/2} \ln[x^2 + \ln^2(2e^{-a} \cos x)] dx = x \ln \frac{a}{e^b - 1},$$

and

$$(4.42) \quad \int_0^{\pi/2} \ln[x^2 + \ln^2(2e^{-a} \cos x)] \cos 2x dx = \frac{\pi}{2} \left(1 - \frac{1}{a} - e^b + \frac{1}{e^b - 1} \right)$$

where $b = \min(a, \ln 2)$.

[196]

$$(4.43) \quad \int_0^{\pi/2} \ln[x^2 + \ln^2(\cos x)] dx = \frac{\pi}{2} \ln \ln 2.$$

[196]

$$(4.44) \quad \int_0^{\pi/2} \ln[x^2 + \ln^2(\cos x)] \cos 2x dx = -\frac{\pi}{\ln 2}.$$

[196]

$$(4.45) \quad \int_0^{\pi/2} \frac{\ln \cos x}{x^2 + \ln^2(\cos x)} dx = \frac{\pi}{2} \left(1 - \frac{1}{\ln 2}\right).$$

[196]

$$(4.46) \quad \int_0^{\pi/2} \frac{x \sin 2x}{x^2 + \ln^2(\cos x)} dx = \frac{\pi}{4 \ln^2 2}.$$

[196]

$$(4.47) \quad \int_0^{\pi/2} \frac{x \sin 2x}{x^2 + \ln^2(2 \cos x)} dx = \frac{13\pi}{48}.$$

[196]

$$(4.48) \quad \int_{-\pi/2}^{\pi/2} \frac{(1 + e^{-2ix})^\beta}{\ln(1 + e^{-2ix}) - a} dx = -\frac{\pi}{a} + \pi \frac{e^{(\beta+1)a}}{e^a - 1} H(\ln 2 - a),$$

and

$$(4.49) \quad \int_0^{\pi/2} \frac{x \sin x}{x^2 + \ln^2(2e^{-a} \cos x)} dx = \frac{\pi}{4a^2} + \frac{\pi e^a}{4} \left(1 - \frac{1}{(e^a - 1)^2}\right) H(\ln 2 - a),$$

where H is the unit step function.

[11, 196]

$$(4.50) \quad \frac{4}{\pi} \int_0^{\pi/2} \frac{x^2 dx}{x^2 + \ln^2(2 \cos x)} = \frac{1}{2} (1 + \ln(2\pi) - \gamma).$$

[11]

$$(4.51) \quad \frac{4}{\pi} \int_0^{\pi/2} \frac{x^2 dx}{x^2 + \ln^2(2e^{-a} \cos x)} = \frac{\gamma}{a} + \frac{a + \ln(1 - e^{-a}) - \gamma - \ln a}{1 - e^{-a}} + \frac{a}{1 - e^{-a}} \int_0^1 e^{-at} \ln \Gamma(t) dt$$

$$(4.52) \quad = \frac{\gamma}{a} + \frac{a + \ln(1 - e^{-a}) + \Gamma(0, a)}{1 - e^{-a}} + \frac{1}{1 - e^{-a}} \int_0^1 e^{-at} \psi(t+1) dt$$

where $0 < a < \ln 2$.

[11]

$$(4.53) \quad \frac{4}{\pi} \int_0^{\pi/2} \frac{x^2 dx}{x^2 + \ln^2(2e^{-a} \cos x)} = \frac{\gamma}{a} + \int_0^\infty e^{-at} \psi(t+1) dt$$

where $a > \ln 2$.

[11]

$$(4.54) \quad \int_0^{\pi/2} \frac{x^2 \ln(2 \cos x) dx}{(x^2 + \ln^2(2 \cos x))^2} = \frac{7\pi}{192} + \frac{\pi \ln 2\pi}{96} - \frac{\zeta'(2)}{16\pi}.$$

[196]

$$(4.55) \quad \frac{1}{2i} \int_{-\pi/2}^{\pi/2} \frac{x(1 + \exp(-2ix))^\beta}{\ln(1 + \exp(-2ix))} dx = \frac{\pi}{8} [1 + \ln 2\pi - \gamma(2\beta + 1) - 2 \ln \Gamma(\beta + 1)]$$

with $\Re \beta > -1$.

[196]

$$(4.56) \quad \int_{-\pi/2}^{\pi/2} \frac{(1 + \exp(-2ix))^\beta}{\ln(1 + \exp(-2ix))} dx = \frac{\pi}{2} (1 + 2\beta).$$

4.1. Logarithmic functions of compound arguments and powers. [6]

$$(4.57) \quad - \int_0^1 \frac{\log(\frac{1}{\sqrt{2}}(1-x))}{x^2 + 1} = G.$$

[6]

$$(4.58) \quad - \int_0^1 \frac{\log[\frac{1}{2}(1-x^2)]}{x^2 + 1} = G.$$

[32]

$$(4.59) \quad \int_0^\infty \left[\frac{1}{x^4 - x^2 + 1} \right]^r \ln \frac{x^2}{x^4 - x^2 + 1} \frac{dx}{x^2} = \frac{\sqrt{\pi}}{2} \frac{\Gamma(r) \Gamma'(r - 1/2) - \Gamma(r - 1/2) \Gamma'(r)}{\Gamma^2(r)}$$

[32]

$$(4.60) \quad \int_0^\infty \frac{1}{x^4 - x^2 + 1} \ln^2 \frac{x^2}{x^4 - x^2 + 1} \frac{dx}{x^2} = \frac{\pi}{2} \left(\frac{\pi^2}{3} + 4 \ln^2 2 \right).$$

[32]

$$(4.61) \quad \int_0^\infty \left[\frac{x}{x^2 + 1} \right]^{2r} \ln \frac{x}{x^2 + 1} \frac{dx}{x^2} = \frac{\sqrt{\pi}}{4} G(r) [\psi(r - 1/2) - \psi(r) - 2 \ln 2],$$

where $G \equiv 2^{1-2r} B(r - 1/2, 1/2) / \sqrt{\pi}$.

[32]

$$(4.62) \quad \int_0^\infty \left[\frac{x}{x^2 + 1} \right]^{2r} \ln \frac{x}{x^2 + 1} \frac{x^2 + 1}{x^2(x^2 + 1)} dx = \frac{\sqrt{\pi}}{4} G'(r)$$

where $G \equiv 2^{1-2r} B(r - 1/2, 1/2) / \sqrt{\pi}$.

[32]

$$(4.63) \quad \int_0^\infty \left[\frac{x}{x^2 + 1} \right]^{2r} \ln^2 \frac{x}{x^2 + 1} \frac{dx}{x^2} = \frac{\sqrt{\pi}}{8} G''(r)$$

where $G'' \equiv G(r) [\psi'(r - 1/2) - \psi'(r) + (\psi(r - 1/2) - \psi(r) - 2 \ln 2)^2]$

[11]

$$(4.64) \quad \int_0^\infty e^{-ax} \ln x dx = -\frac{\gamma + \ln a}{a}.$$

$$(4.65) \quad \int_0^\infty \frac{\ln x}{e^x + e^{-x} - 1} dx = \int_0^1 \ln \ln \left(\frac{1}{x} \right) \frac{dx}{1 - x + x^2} = \frac{2\pi}{\sqrt{3}} \left(\frac{5}{6} \ln 2\pi - \ln \Gamma \left(\frac{1}{6} \right) \right).$$

[112]

$$(4.66) \quad \frac{a^\nu}{\Gamma(\nu)} \int_0^\infty e^{-ax} x^{\nu-1} \ln^m x dx = \phi^m(a, \nu) + \sum_{j=1}^m \binom{m}{j} \eta_j \phi^{m-j}(a, \nu),$$

where

$$(4.67) \quad \phi(a, \nu) \equiv \psi(\nu) - \ln a,$$

$$(4.68) \quad \eta_j(\nu) \equiv (-1)^j \sum_{\pi_0(j)} (j; 0, k_2, \dots, k_j)^* \zeta^{k_2}(2, \nu) \cdots \zeta^{k_j}(j, \nu),$$

$$(4.69) \quad \zeta(k, \nu) \equiv \sum_{l=0}^\infty \frac{1}{(l + \nu)^k}, \quad k \geq 2.$$

[9]

$$\int_0^\infty \log x \cos^{2n+1} x^2 dx = -\frac{\sqrt{\pi}}{2^{2n+3}} (\pi + 2\gamma + 4 \log 2) \sum_{k=0}^n \binom{2n+1}{n-k} \frac{1}{\sqrt{4k+2}} \\ - \frac{\sqrt{\pi}}{2^{2n+2}} \sum_{k=0}^n \binom{2n+1}{n-k} \frac{\log(2k+1)}{\sqrt{4k+2}}.$$

[11]

$$(4.70) \quad \int_0^{\pi/2} x \ln(2 \cos x) dx = -\frac{7}{16} \zeta(3).$$

[11]

$$(4.71) \quad \int_0^{\pi/2} x^2 \ln(2 \cos x) dx = -\frac{\pi}{4} \zeta(3).$$

[11]

$$(4.72) \quad \int_0^{\pi/2} x^2 \ln^2(2 \cos x) dx = \frac{11\pi}{16} \zeta(4) = \frac{11\pi^5}{1440}.$$

[52]

$$(4.73) \quad \int_0^{\pi/2} x^4 \ln^2(2 \cos x) dx = \frac{5\pi^7}{8064} + \frac{3\pi}{4} \zeta^2(3).$$

[52]

$$(4.74) \quad \int_0^{\pi/2} x^2 \ln^4(2 \cos x) dx = \frac{33\pi^7}{4480} + \frac{3\pi}{2} \zeta^2(3).$$

[52]

(4.75)

$$\begin{aligned} \int_0^{\pi/2} x^3 \ln(\cos x) \sin[(p-1)x] \cos^{p-1} x dx = & -\frac{\pi}{15} 2^{-(p+6)} [60\gamma^4 - 60\gamma^2\pi^2 + \pi^4 + 60\gamma(\pi^2 - 2\gamma^2) \ln 2 \\ & + 60\psi'''(p) + 60\{-(\pi^2 + 6\gamma(-\gamma + \ln 2))\psi^2(p) + (4\gamma - 2\ln 2)\psi^3(p) + \psi^4(p) + 6\gamma \ln 2\psi'(p) \\ & - 3[\psi'(p)]^2 - 2(\gamma + \ln 2)\psi''(p) + (8\gamma - 4\ln 2)\zeta(3) + \psi(p)(4\gamma^3 - 2\gamma\pi^2 + (\pi^2 - 6\gamma^2) \ln 2 \\ & + 6\ln 2\psi'(p) - 2\psi''(p) + 8\zeta(3))\}]. \end{aligned}$$

[6, 42]

$$(4.76) \quad \int_0^{\pi/4} \log(\cot x) dx = - \int_0^{\pi/4} \log(\tan x) dx = G.$$

[75, 134]

$$(4.77) \quad \int_{\pi/4}^{\pi/2} \ln \ln \tan x dx = \frac{\pi}{2} \ln \frac{\Gamma(3/4)\sqrt{2\pi}}{\Gamma(1/4)}.$$

[134, 3]

$$(4.78) \quad \int_0^1 x^j \log \log \frac{1}{x} dx = -\frac{\gamma + \log(j+1)}{j+1}.$$

[3]

(4.79)

$$\int_0^1 \frac{x^{p-1}}{1+x^n} \log \log \frac{1}{x} dx = \frac{1}{2n} [\log(2n) + \gamma] \left(\psi\left(\frac{p}{2n}\right) - \psi\left(\frac{n+p}{2n}\right) \right) + \frac{1}{2n} \left(\zeta'\left(1, \frac{p}{2n}\right) - \zeta'\left(1, \frac{n+p}{2n}\right) \right)$$

for $\Re p > 0$ and $\Re n > 0$, and a similar expression if $(1+x^n)^2$ or $(1+x^n)^3$ are in the denominator.

[3]

(4.80)

$$\int_0^1 \frac{x^{nr-1}}{1+x^n} \log \log \frac{1}{x} dx = \frac{1}{2n} [\log(2n) + \gamma] \left(\psi(r/2) - \psi\left(\frac{r+1}{2}\right) \right) + \frac{1}{2n} (\zeta'(1, r/2) - \zeta'(1, (r+1)/2)).$$

[3]

(4.81)

$$\int_0^1 x^{p-1} \frac{1-x}{1-x^n} \log \log \frac{1}{x} dx = \frac{1}{n} [\log(n) + \gamma] \left(\psi(p/n) - \psi\left(\frac{p+1}{n}\right) \right) + \frac{1}{n} (\zeta'(1, p/n) - \zeta'(1, (p+1)/n)).$$

[134] Let

$$(4.82) \quad R_{m,j}(a) \equiv \int_0^1 \frac{x^j \log \log 1/x}{(x+a)^{m+1}} dx,$$

and (Eulerian numbers)

$$(4.83) \quad A_{m,j} = \sum_{k=0}^j (-)^k \binom{m+1}{k} (j-k)^m$$

and

$$(4.84) \quad E_m \equiv \int_0^1 \frac{T_{m-1}(x) \log \log 1/x}{(x+1)^{m+1}} dx$$

defined via polynomials

$$(4.85) \quad T_m(x) \equiv \sum_{j=0}^m (-)^j A_{m+1,j+1} x^j,$$

then

$$(4.86) \quad E_m = (1 - 2^m) \zeta'(1 - m) - (-)^m [\gamma(2^m - 1) + 2^m \log 2] \frac{B_m}{m}$$

where B_m are the Bermoulli numbers. The $R_{m,j}$ are then recursively

$$(4.87) \quad R_{0,0}(1) = -(\log^2 2)/2; \quad R_{m,0}(1) = \frac{E_m}{b_0(m)} - \sum_{k=1}^{m-1} \frac{b_k(m)}{b_0(m)} R_{m-k,0}(1),$$

$$(4.88) \quad R_{0,0}(a) = -\gamma \log(1 + 1/a) - \text{Li}'_1(-1/a); \quad \text{Li}'_c(x) = - \sum_{n \geq 1} \frac{\log n}{n^c} x^n,$$

and for $m > 0$

$$(4.89) \quad R_{m,0}(a) = -\frac{\gamma}{a^m(1+a)m} - \frac{\gamma}{a^{m+1}m!} \sum_{j=2}^m \frac{S_1(m,j) T_{j-2}(1/a)}{(1+1/a)^j} - \frac{1}{a^m m!} \sum_{j=1}^m S_1(m,j) \text{Li}'_{1-j}(-1/a),$$

where

$$(4.90) \quad b_k(m) \equiv (-)^k \sum_{j=0}^{m-1} \binom{j}{k} A_{m,j+1},$$

and the unsigned Stirling numbers of the first kind are $S_1(m,j)$ as in $(t)_m = \sum_{j=1}^m S_j(m,j) t^j$. For larger parameters m then

$$(4.91) \quad R_{m,0}(a) = \sum_{j=0}^r \alpha_{j,r}(a) R_{m-r+j,j}(a), \quad \alpha_{j,r}(a) \equiv (-)^j \binom{r}{j} a^{-r}.$$

[134] Let

$$(4.92) \quad D_{m,j}(r, \theta) \equiv \int_0^1 \frac{x^j \log \log 1/x}{(x^2 - 2rx \cos \theta + r^2)^{m+1}} dx,$$

then

$$(4.93) \quad D_{0,0}(1, \theta) = \frac{\pi}{2 \sin \theta} \left[(1 - \theta/\pi) \log 2\pi + \log \frac{\Gamma(1 - \theta/2\pi)}{\Gamma(\theta/2\pi)} \right],$$

$$(4.94) \quad D_{0,0}(r, \theta) = -\frac{\gamma}{r \sin \theta} \tan^{-1} \frac{\sin \theta}{r - \cos \theta} + \frac{1}{2ri \sin \theta} (\text{Li}'_1(e^{i\theta}/r) - \text{Li}'_1(e^{-i\theta}/r)),$$

(4.95)

$$D_{0,1}(r, \theta) = -\frac{\gamma}{2} \log \frac{r^2 - 2r \cos \theta + 1}{r^2} - \gamma \cot \theta \tan^{-1} \frac{\sin \theta}{r - \cos \theta} + \frac{1}{2ri \sin \theta} [\Phi'(e^{i\theta}/r, 1, 1) - \Phi'(e^{-i\theta}/r, 1, 1)],$$

$$(4.96) \quad D_{m,j}(r\theta) = -\frac{1}{2rm \sin \theta} \frac{\partial}{\partial \theta} D_{m-1,j-1}(r, \theta), \quad m, j > 0.$$

[134]

$$(4.97) \quad \int_0^1 \frac{\log(1-x)}{x} \log \log 1/x dx = \int_0^\infty \log t \log(1 - e^{-t}) dt = \frac{\gamma \pi^2}{6} - \zeta'(2).$$

[134]

$$(4.98) \quad \int_0^1 \frac{\log(1+x)}{x} \log \log 1/x dx = \frac{\pi^2}{12} (\log 2 - \gamma) + \zeta'(2)/2,$$

and other examples involving the kernel $\log \log 1/x$.

[75]

$$(4.99) \quad \int_0^1 q^n \ln(\sin \pi q) dq = -\frac{\ln 2}{n+1} + n! \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{(-1)^k (\zeta(2k+1))}{(2\pi)^{2k} (n+1-2k)!}.$$

[80, 129] by differentiation of [91, 3.761.4] w.r.t. the parameter:

$$(4.100) \quad \int_0^\infty \frac{\sin x}{x^s} \ln x dx = \frac{\pi}{2} \frac{1}{[\Gamma(s) \sin \frac{\pi s}{2}]^2} \left[\Gamma'(s) \sin \frac{s\pi}{2} + \frac{\pi}{2} \Gamma(s) \cos \frac{s\pi}{2} \right].$$

$$(4.101) \quad \int_0^\infty \frac{\sin x}{x^s} \ln^2 x dx = \pi \frac{1}{[\Gamma(s) \sin \frac{\pi s}{2}]^3} \left\{ \Gamma'(s) \sin \frac{s\pi}{2} + \frac{\pi}{2} \Gamma(s) \cos \frac{s\pi}{2} \right\} \\ - \frac{\pi}{2} \frac{1}{[\Gamma(s) \sin \frac{\pi s}{2}]^3} \left\{ \Gamma''(s) \sin \frac{s\pi}{2} + \pi \Gamma'(s) \cos \frac{s\pi}{2} - \frac{\pi^2}{4} \Gamma(s) \sin \frac{s\pi}{2} \right\}.$$

[108] Define

$$(4.102) \quad s_{n,p} \equiv \frac{(-1)^{n+p-1}}{(n-1)!p!} \int_0^1 t^{-1} \log^{n-1} t \log^p(1-t) dt$$

then

$$(4.103) \quad s_{n,p} = s_{p,n} = \sum_{k=1}^p \frac{(-1)^{k+1}}{k!} \sum_{m_i} \frac{H_p(m_1, \dots, m_k)}{m_1 \dots m_k} \zeta(m_1) \dots \zeta(m_k),$$

where

$$(4.104) \quad H_p(m_1, \dots, m_k) = \sum_{p_i} \binom{m_1}{p_1} \dots \binom{m_k}{p_k},$$

the sum over m_i over all sets of integers which satisfy $m_i \geq 2$, $\sum_{i=1}^k m_i = n+p$, and the sum over p_i over all sets of integers which satisfy $1 \leq p_i \leq m_i - 1$, $\sum_{i=1}^k p_i = p$. Examples with $s_{n,p} = \sum_{k=1}^p \frac{(-1)^{k+1}}{k!} \alpha_k(n,p)/k!$ are $\alpha_1(n,p) = (n+p-1)! \zeta(n+p)/(n!p!)$ or $\alpha_2(n,2) = \sum_{\nu=2}^n \zeta(\nu) \zeta(n-\nu+2)$. The reference provides an explicit table for $n, p \leq 4$.

[109]

$$(4.105) \quad r_{np} \equiv \int_0^{\pi/2} \log^n \cos x \log^p \sin x dx;$$

$$(4.106) \quad r_{10} = -\frac{\pi}{2} \log 2,$$

$$(4.107) \quad r_{11} = \frac{\pi}{2} \left(-\frac{\pi^2}{24} + \log^2 2 \right),$$

$$(4.108) \quad r_{20} = \frac{\pi}{2} \left(\frac{\pi^2}{12} + \log^2 2 \right),$$

$$(4.109) \quad r_{21} = \frac{\pi}{2} \left(-\log^3 2 + \frac{1}{4} \zeta(3) \right),$$

$$(4.110) \quad r_{22} = \frac{\pi}{2} \left(\frac{\pi^4}{160} + \log^4 2 - \zeta(3) \log 2 \right).$$

[56]

$$(4.111) \quad \int_0^1 \frac{\log u}{1-u} \operatorname{Li}_2 \left(\frac{u-1}{u} \right) du = \frac{17}{4} \zeta(4).$$

[56]

$$(4.112) \quad - \int_0^1 \frac{\operatorname{Li}_{q-1}(1-t) \log t}{1-t} dt = \sum_{n=1}^{\infty} \frac{H_n^{(1)}}{n^q},$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.

[56]

$$(4.113) \quad \int \frac{\operatorname{Li}_2(1-t) \log t}{1-t} dt = \frac{1}{2} [\operatorname{Li}_2(1-t)]^2 + c.$$

[56]

$$(4.114) \quad \int_0^1 \frac{\operatorname{Li}_{2p}(1-t) \log t}{1-t} dt = \frac{1}{2} \sum_{j=2}^{2p} (-1)^j \zeta(j) \zeta(2p-j+2),$$

[42]

$$(4.115) \quad -\frac{1}{4} \int_0^1 \frac{\log x}{(x+1)\sqrt{x}} dx = \frac{1}{4} \int_1^{\infty} \frac{\log x}{(x+1)\sqrt{x}} dx = G.$$

[42]

$$(4.116) \quad \frac{1}{\sqrt{2}} \int_0^{\pi/2} \log \left(\frac{1 + \frac{1}{\sqrt{2}} \sin x}{1 - \frac{1}{\sqrt{2}} \sin x} \right) \frac{dx}{1 + \cos^2 x} = G.$$

[42]

$$(4.117) \quad \frac{1}{2} \int_0^{\pi/4} \log \left(\frac{1 + \sin x}{1 - \sin x} \right) \frac{dx}{\cos x \sqrt{\cos 2x}} = G.$$

[42]

$$(4.118) \quad \int_0^{\frac{\sqrt{2}+1}{\sqrt{2}-1}} \frac{(x+1) \log x}{4x \sqrt{6x - x^2 - 1}} dx = G.$$

[42]

$$(4.119) \quad -\int_0^1 \frac{\log x}{1+x^2} dx = \int_1^\infty \frac{\log x}{1+x^2} dx = G.$$

[42]

$$(4.120) \quad -\int_0^1 \log\left(\frac{1-x}{\sqrt{2}}\right) \frac{dx}{1+x^2} = G.$$

[42]

$$(4.121) \quad -\int_0^1 \log\left(\frac{1-x^2}{2}\right) \frac{dx}{1+x^2} = G.$$

[42]

$$(4.122) \quad \int_1^\infty \log\left(\frac{x+1}{\sqrt{2}}\right) \frac{dx}{1+x^2} = G.$$

[42]

$$(4.123) \quad \int_0^\infty \frac{\log(1+x)}{1+x^2} dx = G + \frac{1}{4}\pi \log 2.$$

[42]

$$(4.124) \quad -\int_0^1 \frac{\log(1+x^2)}{1+x^2} dx = G - \frac{1}{2}\pi \log 2.$$

[42]

$$(4.125) \quad -\int_1^{\sqrt{2}} \frac{2 \log x}{x\sqrt{x^2-1}} dx = G - \frac{1}{2}\pi \log 2.$$

[42]

$$(4.126) \quad \int_0^{\pi/2} \log(\cos x + \sin x) dx = G - \frac{1}{4}\pi \log 2.$$

[42]

$$(4.127) \quad -\frac{3}{2} \int_0^{2-\sqrt{3}} \frac{\log x}{1+x^2} dx = G.$$

[42]

$$(4.128) \quad \frac{3}{2} \int_{2+\sqrt{3}}^\infty \frac{\log x}{1+x^2} dx = G.$$

4.2. Inverse Trigonometric Functions. [6, 42]

$$(4.129) \quad \frac{2}{\pi} \int_0^1 \frac{\tan^{-1} x}{x} dx = G - \frac{7}{4\pi} \zeta(3).$$

[42]

$$(4.130) \quad \frac{3}{2} \int_0^{2-\sqrt{3}} \frac{\tan^{-1} x}{x} dx = G - \frac{1}{8}\pi \log(2+\sqrt{3}).$$

[42]

$$(4.131) \quad - \int_0^1 (\tan^{-1} x)^2 dx = G - \frac{1}{16}\pi^2 - \frac{1}{4}\pi \log 2.$$

[6, 42]

$$(4.132) \quad \int_0^1 \frac{\tan^{-1} x}{x} dx = G.$$

[42]

$$(4.133) \quad 2 \int_0^1 \left(\frac{1}{4}\pi - \tan^{-1} x \right) \frac{dx}{1-x^2} = G.$$

[42]

$$(4.134) \quad - \int_0^1 \frac{\sin^{-1} x}{\sqrt{1+x^2}} dx = G - \frac{1}{2}\pi \log(1+\sqrt{2}).$$

[88]

$$(4.135) \quad \int_0^b \tan^{-1} \frac{a}{\sqrt{1+x^2}} \cdot \frac{dx}{\sqrt{1+x^2}} = \frac{1}{2}[g(a, b) + g(b, a)]$$

where

$$(4.136) \quad g(a, b) = \tan^{-1} \frac{b}{a} \ln[4(a^2+1) - \frac{4a\sqrt{a^2+1}}{2a^2+1}] - 2\eta \ln(\sqrt{a^2+1} - a) \\ - \text{Cl}_2(2 \tan^{-1} \frac{b}{a}) + \frac{1}{2} \text{Cl}_2(4 \tan^{-1} \frac{b}{a} - 2\eta) + \frac{1}{2} \text{Cl}_2(2\eta)$$

and

$$(4.137) \quad \eta = \tan^{-1} \frac{ab}{(\sqrt{a^2+1} - a + 1)a^2 + (\sqrt{a^2+1} - a - 1)b^2}.$$

4.3. Multiple Integrals. [20] Let

$$(4.138) \quad C_n = \frac{4}{n!} \int_0^\infty \cdots \int_0^\infty \frac{1}{[\sum_{j=1}^n (u_j + 1/u_j)]^2} \frac{du_1}{u_1} \cdots \frac{du_n}{u_n}$$

then

$$(4.139) \quad C_2 = 1; \quad C_3 = L_{-3}(s); \quad C_4 = 7\zeta(3)/12$$

see (4.32).

[20] Let

$$(4.140) \quad D_n = \frac{4}{n!} \int_0^\infty \cdots \int_0^\infty \frac{\prod_{i<j} \left(\frac{u_i - u_j}{u_i + u_j} \right)^2}{[\sum_{j=1}^n (u_j + 1/u_j)]^2} \frac{du_1}{u_1} \cdots \frac{du_n}{u_n} = \frac{1}{n!} \int d^n x \frac{\prod_{i<j} \tanh^2[(x_i - x_j)/2]}{(\cosh x_1 + \cdots + \cosh x_n)^2}$$

then

$$(4.141) \quad D_1 = 2; \quad D_2 = 1/3; \quad D_3 = 8 + 4\pi^2/3 - 27L_{-3}(2); \quad D_4 = 4\pi^2/9 - 1/6 - 7\zeta(3)/2,$$

see (4.32).

[42]

$$(4.142) \quad \frac{2}{\pi} \int_0^{\pi/2} \int_0^{\pi/2} \tan^{-1}(\sin x \sin y) \frac{dx dy}{\sin x} = G.$$

[42]

$$(4.143) \quad \frac{1}{2} \int_0^1 \int_0^{\pi/2} \frac{d\theta dx}{\sqrt{1-x^2 \sin^2 \theta}} = G.$$

[42]

$$(4.144) \quad \int_0^1 \int_0^{\pi/2} \sqrt{1-x^2 \sin^2 \theta} d\theta dx = G + \frac{1}{2}.$$

[42]

$$(4.145) \quad 8 \int_0^1 \int_0^1 \frac{\tan^{-1}(xy) dx dy}{1+x^2 y^2} = 2\pi G - \frac{7}{2} \zeta(3).$$

[42]

$$(4.146) \quad 4 \int_0^1 \int_0^1 \frac{\tan^{-1} x}{1+x^2 y^2} dx dy = 2\pi G - \frac{7}{2} \zeta(3).$$

[42]

$$(4.147) \quad - \int_0^1 \int_0^1 \frac{\log(1-x^2 y^2)}{xy \sqrt{(1-x^2)(1-y^2)}} dx dy = 2\pi G - \frac{7}{2} \zeta(3).$$

[30]

$$(4.148) \quad \int_0^{\pi/2} \int_0^{\pi/2} \frac{\tan(\phi/2) d\theta d\phi}{\sqrt{1-x \cos^2 \theta \cos^2 \phi}} = \frac{\pi}{4} \int_0^{\pi/2} \frac{d\phi}{\sqrt{1-(1-x) \sin^2 \phi}} + \frac{1}{4} \log x \int_0^{\pi/2} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}}.$$

[6, 42]

$$(4.149) \quad \frac{1}{4} \int_0^1 \int_0^1 \frac{1}{\sqrt{(1-x)(1-y)}} \frac{dx dy}{x+y} = G.$$

[96]

$$(4.150) \quad \int_0^1 \cdots \int_0^1 \prod_{1 \leq i < j \leq n} |t_i - t_j|^{2z} \prod_{j=1}^n t_j^{x-1} (1-t_j)^{y-1} dt_j = \prod_{j=1}^n \frac{\Gamma(x+(j-1)z) \Gamma(y+(j-1)z) \Gamma(jz+1)}{\Gamma(x+y+(n+j-2)z) \Gamma(z+1)},$$

where n is a positive integer, x, y, z are in \mathbb{C} , and $\Re x, \Re y > 0, \Re z > -\max\{1/n, \Re x/(n-1), \Re y/(n-1)\}$.

[56]

$$(4.151) \quad \int_0^1 \int_0^1 \frac{\text{Li}_{q-2}[(1-t)(1-u)] \log t \log u du dt}{(1-t)(1-u)} = \sum_{n=1}^{\infty} \frac{[H_n^{(1)}]^2}{n^q}$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.

[56]

$$(4.152) \quad \int_0^1 \int_0^1 \frac{\log t \log u du dt}{1-(1-t)(1-u)} = \sum_{n=1}^{\infty} \frac{[H_n^{(1)}]^2}{n^2}$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.

[56]

$$(4.153) \quad - \int_0^1 \int_0^1 \frac{\log[1 - (1-t)(1-u)] \log t \log u \, du \, dt}{(1-t)(1-u)} = \sum_{n=1}^{\infty} \frac{[H_n^{(1)}]^2}{n^3}$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.

[56]

$$(4.154) \quad n^2 \int_0^1 (1-t)^{n-2} \log t \, dt \int_0^1 (1-u)^{n-1} \log u \, du = [H_n^{(1)}]^2,$$

where $H_n^{(r)} \equiv \sum_{k=1}^n \frac{1}{k^r}$.

[9]

$$(4.155) \quad \int_0^\infty \int_0^\infty \frac{\cos^{2n+1}(x+y)}{x^p y^q} dx \, ds = -\Gamma(1-p)\Gamma(1-q) \cos \frac{\pi(p+q)}{2} \sum_{k=0}^n \binom{2n+1}{n-k} \frac{(2k+1)^{p+1-2}}{2^{2n}}$$

[9]

$$(4.156) \quad \int_0^\infty \int_0^\infty \frac{\cos(x+y)}{x^p y^q} dx \, dy = -\Gamma(1-p)\Gamma(1-q) \cos \frac{\pi(p+q)}{2}.$$

[9]

$$(4.157) \quad \int_0^\infty \int_0^\infty \frac{\log x \log y}{\sqrt{xy}} \cos(x+y) dx \, dy = (\gamma + 2 \log 2) \pi^2.$$

[9]

$$(4.158) \quad \int_{R_+^n} (\cos ||x||^2) \cdot \prod_{j=1}^n \log x_j \, dV = \frac{(-)^{\Delta_n} \pi^{n/2}}{2^{2n}} \times \begin{cases} \Re \psi_n & n \text{ even} \\ \Im \psi_n & n \text{ odd} \end{cases}$$

where $\Delta_n \equiv n(n+1)/2$ and $\psi_n \equiv (\gamma + 2 \log 2 + \pi i/2)^n e^{\pi i n/4}$.

[95]

$$(4.159) \quad \int_0^1 \int_0^1 \frac{x^{u-1} y^{v-1}}{1-xyz} (-\ln xy)^s dx \, dy = \Gamma(s+1) \frac{\Phi(z, s+1, v) - \Phi(z, s+1, u)}{u-v},$$

[95]

$$(4.160) \quad \int_0^1 \int_0^1 \frac{(xy)^{u-1}}{1-xyz} (-\ln xy)^s dx \, dy = \Gamma(s+1) \Phi(z, s+2, u),$$

where

$$(4.161) \quad \Phi(z, s, u) = \sum_{k=0}^{\infty} \frac{z^k}{(u+k)^s} = \frac{1}{\Gamma(s)} \int_0^\infty \frac{e^{-(u-1)t}}{e^t - z} t^{s-1} dt$$

is the Lerch transcendent. [95]

$$(4.162) \quad \int_0^1 \int_0^1 \frac{1}{1+xyi} dx \, dy = G - \frac{\pi^2 i}{48},$$

[95]

$$(4.163) \quad \int_0^1 \int_0^1 \frac{-x \ln xy}{1+x^2 y^2} dx \, dy = G - \frac{\pi^2}{48},$$

[95]

$$(4.164) \quad \int_0^1 \int_0^1 \frac{-x \ln xy}{1 - x^2 y^2} dx dy = \frac{\pi^2}{12},$$

[95]

$$(4.165) \quad \int_0^1 \int_0^1 \frac{(-\ln xy)^n}{1 - xyz} dx dy = \frac{(n+1)! \operatorname{Li}_{n+2}(z)}{z},$$

[95]

$$(4.166) \quad \int_0^1 \int_0^1 \frac{-1}{(1 - xyz) \ln xy} dx dy = -\frac{\ln(1-z)}{z},$$

[95]

$$(4.167) \quad \int_0^1 \int_0^1 \frac{-1}{(2 - xyz) \ln xy} dx dy = \ln 2,$$

[95]

$$(4.168) \quad \int_0^1 \int_0^1 \frac{1}{2 - xy} dx dy = \frac{\pi^2}{12} - \frac{\ln^2 2}{2}.$$

[95]

$$(4.169) \quad \int_0^1 \int_0^1 \frac{-\ln xy}{2 - xy} dx dy = \frac{7\zeta(3)}{4} - \frac{\pi^2 \ln 2}{6} + \frac{\ln^3 2}{3}.$$

[95]

$$(4.170) \quad \int_0^1 \int_0^1 \frac{-1}{(\varphi - xy) \ln xy} dx dy = \ln \varphi,$$

[95]

$$(4.171) \quad \int_0^1 \int_0^1 \frac{1}{\varphi - xy} dx dy = \frac{\pi^2}{10} - \ln^2 \varphi,$$

[95]

$$(4.172) \quad \int_0^1 \int_0^1 \frac{-1}{(\varphi^2 - xy) \ln xy} dx dy = \ln \varphi,$$

etc where

$$(4.173) \quad \varphi \equiv (1 + \sqrt{5})/2.$$

[95]

$$(4.174) \quad \int_0^1 \int_0^1 \frac{1 - 2xy}{(8 + xy)(9 - xy)} dx dy = \frac{1}{2} \ln^2 \frac{9}{8}.$$

[95]

$$(4.175) \quad \int_0^1 \int_0^1 \frac{52 - 7xy}{(2 + xy)(9 - xy)} dx dy = \frac{\pi^3}{3} + 3 \ln^2 2 + 2 \ln^2 3 - 6 \ln 2 \ln 3.$$

etc [95]

$$(4.176) \quad \int_0^1 \int_0^1 \frac{(-\ln xy)^s}{1 - xy} dx dy = \Gamma(s+2) \zeta(s+2), \Re s > -1.$$

[95]
 (4.177)
$$\int_0^1 \int_0^1 \frac{(-\ln xy)^s}{1+xy} dx dy = \Gamma(s+2)\zeta^*(s+2), \Re s > -2, \quad \zeta^*(s) \equiv (1-2^{1-s})\zeta(s).$$

[95]
 (4.178)
$$\int_0^1 \int_0^1 \frac{(-\ln xy)^s}{1+x^2y^2} dx dy = \Gamma(s+2)\beta(s+2), \Re s > -2.$$

[95]
 (4.179)
$$\int_0^1 \int_0^1 \frac{1}{1-xy} dx dy = \zeta(2).$$

[95]
 (4.180)
$$\int_0^1 \int_0^1 \frac{-\ln xy}{1-xy} dx dy = 2\zeta(3).$$

[95]
 (4.181)
$$\int_0^1 \int_0^1 \frac{-1}{(1+x^2y^2)\ln xy} dx dy = \pi/4.$$

[95, 42]
 (4.182)
$$\int_0^1 \int_0^1 \frac{1}{1+x^2y^2} dx dy = G.$$

[95]
 (4.183)
$$\int_0^1 \int_0^1 \frac{\ln xy}{1+x^2y^2} dx dy = \frac{\pi^3}{16}.$$

[95]
 (4.184)
$$\int_0^1 \int_0^1 \frac{(-\ln xy)^s}{1+x^2y^2z^2} dx dy = \Gamma(s+1) \frac{\chi_{s+2}(z)}{z}, \quad \Re s > -2, \Re z \neq 0.$$

[95]
 (4.185)
$$\int_0^1 \int_0^1 \frac{1}{1-x^2y^2 \tan^2(\pi/8)} dx dy = \frac{\pi^2}{16 \tan \frac{\pi}{8}} - \frac{\ln^2 \tan \frac{\pi}{8}}{4 \tan \frac{\pi}{8}}.$$

[95]
 (4.186)
$$\int_0^1 \int_0^1 \frac{x^{u-1}y^{v-1}}{-\ln xy} dx dy = \frac{1}{u-v} \ln \frac{u}{v}.$$

[95]
 (4.187)
$$\int_0^1 \int_0^1 \frac{(xy)^{u-1}}{-\ln xy} dx dy = \frac{1}{u}.$$

[95]
 (4.188)
$$\int_0^1 \int_0^1 \frac{-x^{u-1}y^{v-1}}{(1+xy)\ln xy} dx dy = \frac{1}{u-v} \ln \frac{\Gamma(u/2)\Gamma(\frac{u+1}{2})}{\Gamma(v/2)\Gamma(\frac{v+1}{2})}, \quad u > 0, v > 0.$$

[95]

$$(4.189) \quad \int_0^1 \int_0^1 \frac{-(xy)^{u-1}}{(1+xy) \ln xy} dx dy = \frac{1}{2} \left[\psi\left(\frac{u+1}{2}\right) - \psi\left(\frac{u}{2}\right) \right].$$

[95][175, A053510]

$$(4.190) \quad \int_0^1 \int_0^1 \frac{1+x}{-(1+xy) \ln xy} dx dy = \ln \pi.$$

[95][175, A094640]

$$(4.191) \quad \int_0^1 \int_0^1 \frac{1-x}{-(1+xy) \ln xy} dx dy = \ln \frac{4}{\pi}.$$

[95]

$$(4.192) \quad \int_0^1 \int_0^1 \frac{-x}{(1+x^2 y^2) \ln xy} dx dy = \ln \frac{\sqrt{2\pi}}{\Gamma^2(3/4)}.$$

[95]

$$(4.193) \quad \int_0^1 \int_0^1 \frac{x^{u-1} y^{v-1}}{1-xy} dx dy = \frac{\psi(u) - \psi(v)}{u-v}; \quad \int_0^1 \int_0^1 \frac{(xy)^{u-1}}{1-xy} dx dy = \psi'(u).$$

[95][175, A073010]

$$(4.194) \quad \int_0^1 \int_0^1 \frac{y}{(1-x^3 y^3)} dx dy = \frac{\pi}{3\sqrt{3}}.$$

[95]

$$(4.195) \quad \int_0^1 \int_0^1 x^{u-1} y^{v-1} (-\ln xy)^s dx dy = \Gamma(s+1) \frac{v^{-s-1} - u^{-s-1}}{u-v}$$

and others of similar shape.

[37]

$$(4.196) \quad W_n(2k) = \sum_{a_1+a_2+\dots+a_n=k} \left(\begin{matrix} k \\ a_1, \dots, a_n \end{matrix} \right)^2,$$

where the sum is over all compositions (unordered partitions) with n terms, and

$$(4.197) \quad W_n(s) \equiv \int_{[0,1]^n} \left| \sum_{k=1}^n e^{2\pi i x_k} \right|^s d^n x.$$

[37]

$$(4.198) \quad W_3(k) = \Re {}_3F_2 \left(\begin{matrix} 1/2, -k/2, -k/2 \\ 1, 1 \end{matrix} \mid 4 \right).$$

[37]

$$(4.199) \quad W_n(s) = n^s \sum_{m \geq 0} (-1)^m \binom{s/2}{m} \sum_{k=0}^m \frac{(-)^k}{n^{2k}} \binom{m}{k} \sum_{a_1+a_2+\dots+a_n=k} \left(\begin{matrix} k \\ a_1, \dots, a_n \end{matrix} \right)^2.$$

5. INDEFINITE INTEGRALS OF SPECIAL FUNCTIONS

5.1. Elliptic Integrals and Functions.

5.2. The Exponential Integral.

5.3. The Sine Integral and Cosine Integral.

5.4. The Error Function and Fresnel Integrals.

5.5. Cylinder Functions.

$$(5.1) \quad \int x^2 Z_{\nu+1}(x) dx = -2x^2 Z_{\nu}(x) + 4 \int x Z_{\nu}(x) dx + \int x^2 Z_{\nu-1}(x) dx,$$

by partial integration of $\int x Z_{\nu} dx$ with [91, 8.471.2], where Z is a Bessel Function.

$$(5.2) \quad \int x^{\mu+1} Z_{\nu-1}(x) dx = x^{\mu+1} Z_{\nu}(x) + (\nu - \mu - 1) \int x^{\mu} Z_{\nu}(x) dx,$$

$\mu \neq -1$, by partial integration of $\int x^{\mu} Z_{\nu} dx$ with [91, 8.472.1], where Z is a Bessel Function. Equivalent formula for spherical Bessel functions $j_n(z) \equiv \sqrt{\pi/(2z)} J_{n+1/2}(z)$:

$$(5.3) \quad (n - m) \int x^m j_n(x) dx = \int x^{m+1} j_{n-1}(x) dx - x^{m+1} j_n(x).$$

$$(5.4) \quad \int x^{\mu+1} Z_{\nu+1}(x) dx = -x^{\mu+1} Z_{\nu}(x) + (\mu + 1 + \nu) \int x^{\mu} Z_{\nu}(x) dx,$$

$\mu \neq -1$, by partial integration of $\int x^{\mu} Z_{\nu} dx$ with [91, 8.472.2], where Z is a Bessel Function.

[154]

$$(5.5) \quad \int \frac{\sin(x) Z_{\nu}(x)}{x^{3/2}} dx = \frac{2[(2\nu + 1) \sin(x) - 2x \cos(x)]}{x^{1/2}(2\nu - 1)(2\nu + 1)} Z_{\nu}(x) - \frac{4x^{1/2} \sin(x)}{(2\nu - 1)(2\nu + 1)} Z_{\nu+1}(x)$$

where Z is a Bessel function J or Y .

[154]

$$(5.6) \quad \int \frac{\cos(x) Z_{\nu}(x)}{x^{3/2}} dx = \frac{2[(2\nu + 1) \cos(x) + 2x \sin(x)]}{x^{1/2}(2\nu - 1)(2\nu + 1)} Z_{\nu}(x) - \frac{4x^{1/2} \cos(x)}{(2\nu - 1)(2\nu + 1)} Z_{\nu+1}(x)$$

where Z is a Bessel function J or Y .

[154]

$$(5.7) \quad \begin{aligned} \int \frac{Z_{\mu}(x) \bar{Z}_{\nu}(x)}{x^2} dx = & -\frac{1 + \mu + \nu + 2\mu\nu + \mu\nu^2 + \mu^2\nu - \mu^2 - \mu^3 - \nu^2 - \nu^3 + 2x^2}{x(-1 + \mu - \nu)(-1 + \mu + \nu)(1 + \mu - \nu)(1 + \mu + \nu)} Z_{\mu} \bar{Z}_{\nu} \\ & + \frac{1}{(1 - \mu - \nu)(1 - \mu + \nu)} Z_{\mu} \bar{Z}_{\nu+1} + \frac{1}{(1 - \mu - \nu)(1 + \mu - \nu)} Z_{\mu+1} \bar{Z}_{\nu} \\ & - \frac{2x}{(1 - \mu - \nu)(1 - \mu + \nu)(1 + \mu - \nu)(1 + \mu + \nu)} Z_{\mu+1} \bar{Z}_{\nu+1}. \end{aligned}$$

[154]

(5.8)

$$\begin{aligned} \int \frac{Z_\mu(x)\bar{Z}_\nu(x)}{x^3} dx = & -\frac{(2+\mu+\nu)(4\mu+4\nu+\mu\nu^2+\mu^2\nu-\mu^3-\nu^3+4x^2)}{x^2(\mu+\nu)[\nu^2-(2-\mu)^2][\nu^2-(2+\mu)^2]} Z_\mu\bar{Z}_\nu \\ & - \frac{4\mu\nu^2+2\mu^2\nu^2-4\mu^2-4\nu^3-\mu^4+4\nu^2-\nu^4+8x^2}{x(\mu^2-\nu^2)[\nu^2-(2-\mu)^2][\nu^2-(2+\mu)^2]} Z_\mu\bar{Z}_{\nu+1} \\ & + \frac{4\mu^2\nu+2\mu^2\nu^2+4\mu^2-\mu^4-4\nu^2-4\nu^3-\nu^4+8x^2}{x(\mu^2-\nu^2)[\nu^2-(2-\mu)^2][\nu^2-(2+\mu)^2]} Z_{\mu+1}\bar{Z}_\nu \\ & - \frac{4}{[-\nu^2+(2-\mu)^2][-\nu^2+(2+\mu)^2]} Z_{\mu+1}\bar{Z}_{\nu+1}. \end{aligned}$$

[154]

(5.9)

$$\int \frac{Z_\nu^2(x)}{x^2} dx = \frac{1+2\nu+2x^2}{(4\nu^2-1)x} Z_\nu^2(x) - \frac{2}{-1+2\nu} Z_\nu(x)Z_{\nu+1}(x) - \frac{2x}{1-4\nu^2} Z_{\nu+1}^2(x).$$

[154]

(5.10)

$$\begin{aligned} \int \frac{Z_\nu^2(x)}{x^4} dx = & \frac{-9-6\nu+x^2(6+16\nu+8\nu^2)+36\nu^2+24\nu^3+16x^4}{3x^3(1-4\nu^2)(9-4\nu^2)} Z_\nu^2(x) \\ & - \frac{2(-3+4\nu+4\nu^2+8x^2)}{3x^2(1-2\nu)(9-4\nu^2)} Z_\nu(x)Z_{\nu+1}(x) - \frac{2(1-4\nu^2-8x^2)}{3x(1-4\nu^2)(9-4\nu^2)} Z_{\nu+1}^2(x). \end{aligned}$$

[154]

(5.11)

$$\int x^2 Z_{1/3}^3(x) dx = \left(-\frac{4}{9}x - \frac{16}{81x}\right) Z_{1/3}^3(x) - \frac{4x}{3} Z_{1/3}(x) Z_{4/3}^2(x) + \left(\frac{8}{9} + x^2\right) Z_{1/3}^2(x) Z_{4/3}(x) + \frac{2}{3} x^2 Z_{4/3}^3(x).$$

[154]

(5.12)

$$\int \frac{Z_1^4(x)}{x} dx = \frac{x^2}{4} Z_2^4(x) + \left(\frac{3}{4} + \frac{x^2}{4}\right) Z_1^4(x) - \frac{3x}{2} Z_1(x) Z_2^3(x) + 6\left(\frac{1}{2} + \frac{x^2}{12}\right) Z_1^2(x) Z_2^2(x) + 4\left(-\frac{3x}{8} - \frac{1}{2x}\right) Z_1^3(x) Z_2(x).$$

[154]

(5.13)

$$\begin{aligned} \int \frac{Z_3^4(x)}{x^3} dx = & \left(\frac{1}{24} + \frac{1}{2x^2} + \frac{2}{x^4} + \frac{x^2}{378}\right) Z_3^4(x) + \left(\frac{5}{216} + \frac{2}{27x^2} + \frac{x^2}{378}\right) Z_4^4(x) \\ & + 4\left(-\frac{x}{108} - \frac{5}{54x} - \frac{1}{3x^3}\right) Z_3(x) Z_4^3(x) + 6\left(\frac{7}{216} + \frac{1}{3x^2} + \frac{4}{3x^4} + \frac{x^2}{1134}\right) Z_3^2(x) Z_4^2(x) \\ & + 4\left(-\frac{x}{108} - \frac{1}{8x} - \frac{1}{x^3} - \frac{4}{x^5}\right) Z_3^3(x) Z_4(x). \end{aligned}$$

where Z and \bar{Z} are Bessel functions J or Y .

[154]

(5.14)

$$\begin{aligned} \int x^l Z_\mu(x)\bar{Z}_\nu(x) dx = & A_{00}(x)Z_\mu(x)\bar{Z}_\nu(x) + A_{01}(x)Z_\mu(x)\bar{Z}_{\nu+1}(x) + A_{10}(x)Z_{\mu+1}(x)\bar{Z}_\nu(x) \\ & + A_{11}(x)Z_{\mu+1}(x)\bar{Z}_{\nu+1}(x), \end{aligned}$$

where Z and \bar{Z} are Bessel functions J or Y , where

$$A_{00} = \frac{x}{2(\mu + \nu)} D^3 A_{11} + \frac{3 + \mu + \nu}{2(\mu + \nu)} D^2 A_{11} + \frac{-7 - 3\mu - 3\nu - 2\mu\nu + \mu^2 + \nu^2 - 4x^2}{2x(\mu + \nu)} D A_{11} \\ + \frac{(-2 - \mu - \nu)(-4 - 2\mu\nu + \mu^2 + \nu^2 - 2x^2)}{2x^2(\mu + \nu)} A_{11} + \frac{x^{l+1}}{\mu + \nu},$$

$$A_{01} = \frac{-x^2}{2(\mu^2 - \nu^2)} D^3 A_{11} + \frac{3x}{2(\mu^2 - \nu^2)} D^2 A_{11} - \frac{7 - 3\mu^2 - \nu^2 + 4x^2}{2(\mu^2 - \nu^2)} D A_{11} \\ + \frac{4 + \mu\nu^2 - 3\mu^2 - \mu^3 - \nu^2 + 2x^2}{x(\mu^2 - \nu^2)} A_{11} + \frac{x^{l+2}}{\mu^2 - \nu^2},$$

$$A_{10} = \frac{x^2}{2(\mu^2 - \nu^2)} D^3 A_{11} - \frac{3x}{2(\mu^2 - \nu^2)} D^2 A_{11} + \frac{7 - \mu^2 - 3\nu^2 + 4x^2}{2(\mu^2 - \nu^2)} D A_{11} \\ - \frac{4 + \mu^2\nu - \mu^2 - 3\nu^2 - \nu^3 + 2x^2}{x(\mu^2 - \nu^2)} A_{11} - \frac{x^{l+2}}{\mu^2 - \nu^2},$$

$$A_{11}(x) = x^{l+3} \sum_{n=0}^{n'} d_n x^{2n},$$

$$d_0 = \frac{2(l+1)}{(l+3)^4 - 8(l+3)^3 + 2(12 - \mu^2 - \nu^2)(l+3)^2 - 8(l+3)(4 - \mu^2 - \nu^2) + [(2 - \mu)^2 - \nu^2][(2 + \mu)^2 - \nu^2]},$$

$$\{(3 + 2n + l)^4 - 8(3 + 2n + l)^3 + 2(12 - \mu^2 - \nu^2)(3 + 2n + l)^2 - 8(3 + 2n + l)(4 - \mu^2 - \nu^2) + [(2 - \mu)^2 - \nu^2][(2 + \mu)^2 - \nu^2]\} d_n = -4(1 + 2n + l)(2n + l) d_{n-1}, n' \geq n > 0, \\ \text{and } d_n = 0 \text{ if } n > n'.$$

$$n' = \begin{cases} 0, & l = -1 \\ \frac{|l|}{2} - 1, & l < -1, \text{even} \\ \frac{|l|-3}{2}, & l < -1, \text{odd} \\ \infty, & l \geq 0 \end{cases}$$

[154]

$$(5.15) \quad \int x^l Z_\nu^2(x) dx = A_{00}(x) Z_\nu^2(x) + 2A_{01}(x) Z_\nu(x) Z_{\nu+1}(x) + A_{11}(x) Z_{\nu+1}^2(x)$$

where

$$(5.16) \quad A_{00}(x) = \frac{1}{2} D^2 A_{11}(x) - \frac{3 + 2\nu}{2x} D A_{11}(x) + \frac{(2 + 2\nu + x^2)}{x^2} A_{11}(x),$$

$$(5.17) \quad A_{01}(x) = \frac{1}{2} D A_{11}(x) - \frac{1 + \nu}{x} A_{11}(x),$$

$$(5.18) \quad A_{11}(x) = xy(x),$$

$$(5.19) \quad y = \sum_{n=0}^{(l-1)/2} b_n x^{2n+1}, \quad \frac{b_{l-1}}{2} = 1/2l,$$

$$(5.20) \quad \frac{b_{l-1}}{2} = 1/2l, \quad b_n = \frac{2(n+1)[\nu^2 - (n+1)^2]}{2n+1} b_{n+1},$$

if $0 \leq n \leq (l-3)/2$, and if $l \geq 3$ a positive odd integer.

[198, 136,1]

(5.21)

$$\int z^{-\mu-\nu-1} \mathcal{C}_{\mu+1}(z) \mathcal{D}_{\nu+1}(z) dz = -\frac{z^{-\mu-\nu}}{2(\mu+\nu+1)} \{ \mathcal{C}_{\mu}(z) \mathcal{D}_{\nu}(z) + \mathcal{C}_{\mu+1}(z) \mathcal{D}_{\nu+1}(z) \}$$

where \mathcal{C} and \mathcal{D} are arbitrary Bessel functions.

[198, 136,2]

$$(5.22) \quad \int z^{\mu+\nu+1} \mathcal{C}_{\mu}(z) \mathcal{D}_{\nu}(z) dz = \frac{z^{\mu+\nu+2}}{2(\mu+\nu+1)} \{ \mathcal{C}_{\mu}(z) \mathcal{D}_{\nu}(z) + \mathcal{C}_{\mu+1}(z) \mathcal{D}_{\nu+1}(z) \}$$

where \mathcal{C} and \mathcal{D} are arbitrary Bessel functions.

[198]

$$(\rho + \mu + \nu) \int z^{\rho-1} \mathcal{C}_{\mu}(z) \mathcal{D}_{\nu}(z) dz + (\rho - \mu - \nu - 2) \int z^{\rho-1} \mathcal{C}_{\mu+1}(z) \mathcal{D}_{\nu+1}(z) dz = z^{\rho} \{ \mathcal{C}_{\mu}(z) \mathcal{D}_{\nu}(z) + \mathcal{C}_{\mu+1}(z) \mathcal{D}_{\nu+1}(z) \}$$

where \mathcal{C} and \mathcal{D} are arbitrary Bessel functions.

[198, 136,5]

$$(\mu + \nu) \int \mathcal{C}_{\mu}(z) \mathcal{D}_{\nu}(z) \frac{dz}{z} - (\mu + \nu + 2n) \int \mathcal{C}_{\mu+n}(z) \mathcal{D}_{\nu+n}(z) \frac{dz}{z} = \mathcal{C}_{\mu}(z) \mathcal{D}_{\nu}(z) + 2 \sum_{m=1}^{n-1} \mathcal{C}_{\mu+m}(z) \mathcal{D}_{\nu+m}(z) + \mathcal{C}_{\mu+n}(z) \mathcal{D}_{\nu+n}(z)$$

where \mathcal{C} and \mathcal{D} are arbitrary Bessel functions.

[198, 137,1]

(5.23)

$$\int \mathcal{C}_n(z) \mathcal{D}_n(z) \frac{dz}{z} = -\frac{1}{2n} \left[\mathcal{C}_0(z) \mathcal{D}_0(z) + 2 \sum_{m=1}^{n-1} \mathcal{C}_m(z) \mathcal{D}_m(z) + \mathcal{C}_n(z) \mathcal{D}_n(z) \right]$$

where \mathcal{C} and \mathcal{D} are arbitrary Bessel functions.

[198, 138]

$$(5.24) \quad (\mu + 2) \int z^{\mu+2} \mathcal{C}_{\nu}^2(z) dz = (\mu + 1) \left\{ \nu^2 - \frac{1}{4}(\mu + 1)^2 \right\} \int z^{\mu} \mathcal{C}_{\nu}^2(z) dz + \frac{1}{2} \left[z^{\mu+1} \left\{ z \mathcal{C}'_{\nu}(z) - \frac{1}{2}(\mu + 1) \mathcal{C}_{\nu}(z) \right\}^2 + z^{\mu+1} \left\{ z^2 - \nu^2 + \frac{1}{4}(\mu + 1)^2 \right\} \mathcal{C}_{\nu}^2(z) \right]$$

where \mathcal{C} and \mathcal{D} are arbitrary Bessel functions.

With [1, 10.121], then partial integration for a product of three spherical Bessel functions

$$(5.25) \quad (n + m + l + 2) \int j_n(x) j_m(x) j_l(kx) dx = -(2n - 1) j_{n-1}(x) j_m(x) j_l(kx) + (n - 3 - m - l) \int j_{n-2}(x) j_m(x) j_l(kx) dx + (2n - 1) \int j_{n-1}(x) j_{m-1}(x) j_l(kx) dx + (2n - 1) k \int j_{n-1}(x) j_m(x) j_{l-1}(kx) dx.$$

$$(5.26) \quad \int x J_1(x) dx = \frac{\pi}{2} x [J_1(x) \mathbf{H}_0(x) - J_0(x) \mathbf{H}_1(x)]$$

where \mathbf{H} are Struve functions [1, §12].

5.6. Orthogonal Polynomials. [154]

$$(5.27) \quad \int \ln(1 \pm x) P_\nu(x) dx = \left[-\frac{x}{\nu} \ln(1 \pm x) \pm \frac{1}{\nu(\nu+1)} - \frac{x}{\nu^2} \right] P_\nu + \left[\frac{1}{\nu} \ln(1 \pm x) + \frac{1}{\nu^2(\nu+1)} \right] P_{\nu+1}.$$

where P are Legendre functions.

[154]

$$(5.28) \quad \int P_\mu(x) \bar{P}_\nu(x) dx = -\frac{x}{1+\mu+\nu} P_\mu(x) \bar{P}_\nu(x) - \frac{1+\nu}{(\mu-\nu)(1+\mu+\nu)} P_\mu(x) \bar{P}_{\nu+1}(x) \\ + \frac{1+\mu}{(\mu-\nu)(1+\mu+\nu)} P_{\mu+1}(x) \bar{P}_\nu(x).$$

[154]

$$(5.29) \quad \int x [P_\nu(x)]^2 dx = -\frac{1+\nu}{2\nu} \left[\frac{x^2+\nu}{1+\nu} [P_\nu(x)]^2 - 2x P_\nu(x) P_{\nu+1}(x) + [P_{\nu+1}(x)]^2 \right].$$

[154]

$$(5.30) \quad \int [P_{1/2}(x)]^2 dx = \frac{9}{2} x [P_{3/2}(x)]^2 + \frac{x(-7+16x^2)}{2} [P_{1/2}(x)]^2 - 3(-1+2x)(1+2x) P_{1/2}(x) P_{3/2}(x).$$

[154]

$$(5.31) \quad \int [P_{3/2}(x)]^2 dx = \frac{x(93-480x^2+512x^4)}{18} x [P_{3/2}(x)]^2 + \frac{25x(-3+8x^2)}{18} [P_{5/2}(x)]^2 \\ - \frac{5}{9} (3-42x^2+64x^4) P_{3/2}(x) P_{5/2}(x).$$

[154]

$$(5.32) \quad \int x^5 P_{1/3}(x) P_{2/3}(x) dx = \left(-\frac{335}{2352} - \frac{512x^2}{392} + \frac{235x^4}{336} + \frac{x^6}{3} \right) P_{1/3}(x) P_{2/3}(x) \\ + \left(\frac{685x}{784} - \frac{573x^3}{1176} - \frac{5x^5}{48} \right) P_{1/3}(x) P_{5/3}(x) + \left(\frac{235x}{196} - \frac{295x^3}{588} - \frac{2x^5}{21} \right) P_{2/3}(x) P_{4/3}(x) \\ + \left(-\frac{5}{6} + \frac{65x^2}{196} + \frac{25x^4}{588} \right) P_{4/3}(x) P_{5/3}(x).$$

[154]

$$(5.33) \quad \int x [P_{1/3}(x)]^3 dx = \left(\frac{125x^4}{12} - \frac{14}{3} x^2 - \frac{5}{12} \right) [P_{1/3}(x)]^3 + (-4+20x^2) P_{1/3}(x) [P_{4/3}(x)]^2 \\ + (9x-25x^3) [P_{1/3}(x)]^2 P_{4/3}(x) - \frac{16}{3} x [P_{4/3}(x)]^3.$$

[154]

$$(5.34) \quad \int x[P_{1/2}(x)]^4 dx = \left(-\frac{5}{16} - \frac{19}{4}x^2\right)[P_{1/2}(x)]^4 + \frac{81}{4}xP_{1/2}(x)[P_{3/2}(x)]^3 \\ + 6\left(-\frac{9}{16} - \frac{9}{2}x^2\right)[P_{1/2}(x)]^2[P_{3/2}(x)]^2 + 4\left(\frac{33x}{16} + 3x^3\right)[P_{1/2}(x)]^3P_{3/2}(x) - \frac{81}{16}[P_{3/2}(x)]^4.$$

[154]

$$(5.35) \quad \int x^l P_\mu(x) P_\nu(x) dx = A_{00}(x)P_\mu(x)P_\nu(x) + A_{01}(x)P_\mu(x)P_{\nu+1}(x) \\ + A_{10}(x)P_{\mu+1}(x)P_\nu(x) + A_{11}(x)P_{\mu+1}(x)P_{\nu+1}(x),$$

[154]

$$(5.36) \quad \int e^{-x^2} H_\nu(x) dx = -e^{-x^2} H_{\nu-1}(x).$$

[154]

$$(5.37) \quad \int H_\nu(x) x^{-(\nu+3)} dx = \left[\frac{2x^{-\nu}}{(\nu+1)(\nu+2)} - \frac{x^{-(\nu+2)}}{\nu+2} \right] H_\nu(x) - \frac{2\nu}{(\nu+1)(\nu+2)} x^{-(\nu+1)} H_{\nu-1}(x).$$

[154]

$$(5.38) \quad \int x H_\nu(x) dx = \frac{1+2x^2}{2(\nu+2)} H_\nu(x) - \frac{\nu x}{\nu+2} H_{\nu-1}(x).$$

[154]

$$(5.39) \quad \int e^{-x^2} H_\mu(x) \bar{H}_\nu dx = \frac{e^{-x^2}}{2(\mu-\nu)} [-H_\mu(x) \bar{H}_{\nu+1}(x) + H_{\mu+1}(x) \bar{H}_\nu(x)].$$

[154]

$$(5.40) \quad \int x e^{-x^2} H_\mu(x) \bar{H}_\nu dx = \frac{e^{-x^2}}{2} \left[-\frac{1+\mu+\nu}{(1-\mu+\nu)(1+\mu-\nu)} H_\mu(x) \bar{H}_{\nu+1}(x) + \frac{x}{1+\mu-\nu} H_{\mu+1}(x) \bar{H}_\nu(x) \right. \\ \left. + \frac{x}{1-\mu+\nu} H_\mu(x) \bar{H}_{\nu+1}(x) - \frac{1}{(1-\mu+\nu)(1+\mu-\nu)} H_{\mu+1}(x) \bar{H}_{\nu+1}(x) \right].$$

[154]

$$(5.41) \quad \int x^2 e^{-x^2} H_\mu(x) \bar{H}_\nu dx = -e^{-x^2} \frac{H_\mu(x) \bar{H}_\nu(x) x(\mu+\nu)}{(2-\mu+\nu)(2+\mu-\nu)} \\ + H_{\mu+1}(x) \bar{H}_\nu(x) \frac{2+\mu+3\nu+2\mu x^2-2\nu x^2-\mu^2 x^2-\nu^2 x^2+2\mu\nu x^2}{2(\mu-\nu)(2-\mu+\nu)(2+\mu-\nu)} \\ + H_\mu(x) \bar{H}_{\nu+1}(x) \frac{2+3\mu+\nu-2\mu x^2+2\nu x^2-\mu^2 x^2-\nu^2 x^2+2\mu\nu x^2}{2(\mu-\nu)(2-\mu+\nu)(2+\mu-\nu)} \\ - H_{\mu+1}(x) \bar{H}_{\nu+1}(x) \frac{x}{(2-\mu+\nu)(2+\mu-\nu)}.$$

[154]

(5.42)

$$\int e^{-3x^2} x^2 H_{2/3}^3(x) dx = e^{-3x^2} \left[-\frac{1}{12} x(5+6x^2) H_{2/3}^3(x) + \frac{1}{8} (1+6x^2) H_{2/3}^2(x) H_{5/3}(x) - \frac{3}{8} x H_{2/3}(x) H_{5/3}^2(x) + \frac{1}{16} H_{5/3}^3(x) \right].$$

where H and \bar{H} are Hermite functions.

[154]

(5.43)

$$\int x e^{-(\nu+1)x} L_\nu(x) dx = \frac{e^{-(\nu+1)x}}{\nu+1} [-(1+x)L_\nu(x) + L_{\nu-1}(x)].$$

[154]

(5.44)

$$\int x(1+x)^{-(\nu+3)} L_\nu(x) dx = \frac{(1+x)^{-(\nu+1)}}{\nu+2} \left[\left(\frac{x-\nu}{\nu+1} - \frac{x}{1+x} \right) L_\nu(x) + \frac{\nu}{\nu+1} L_{\nu-1}(x) \right].$$

[154]

(5.45)

$$\int e^{-x} L_\mu(x) \bar{L}_\nu(x) dx = e^{-x} \left[L_\mu(x) \bar{L}_\nu(x) + \frac{1+\nu}{\mu-\nu} L_\mu(x) \bar{L}_{\nu+1}(x) - \frac{1+\mu}{\mu-\nu} L_{\mu+1}(x) \bar{L}_\nu(x) \right].$$

[154]

(5.46)

$$\begin{aligned} \int x e^{-x} L_\mu(x) \bar{L}_\nu(x) dx &= e^{-x} \left[-\frac{1+\mu+\nu-x+2\mu\nu+\mu^2x+\nu^2x-2\mu\nu x}{(1-\mu+\nu)(1+\mu-\nu)} L_\mu(x) \bar{L}_\nu(x) \right. \\ &+ \frac{(1+\nu)(1+2\mu-\mu x+\nu x)}{(\mu-\nu)(1-\mu+\nu)} L_\mu(x) \bar{L}_{\nu+1}(x) - \frac{(1+\mu)(1+2\nu+\mu x-\nu x)}{(\mu-\nu)(1+\mu-\nu)} L_{\mu+1}(x) \bar{L}_\nu(x) \\ &\left. - \frac{2(1+\mu)(1+\nu)}{(1-\mu+\nu)(1+\mu-\nu)} L_{\mu+1}(x) \bar{L}_{\nu+1}(x) \right]. \end{aligned}$$

[154]

(5.47)

$$\begin{aligned} \int e^{-3x} x L_{2/3}^3(x) dx &= e^{-3x} \left\{ L_{2/3}^3(x) \left[\frac{125}{24} - \frac{625x}{24} + \frac{853x^2}{16} - \frac{675x^3}{16} + \frac{225x^4}{16} - \frac{27x^5}{16} \right] \right. \\ &+ L_{5/3}^3(x) \left[-\frac{125}{24} + \frac{125x}{12} - \frac{125x^2}{16} \right] + 3 \left[\frac{125}{24} - \frac{125x}{8} + \frac{275x^2}{16} - \frac{75x^3}{16} \right] L_{2/3}(x) L_{5/3}^2(x) \\ &\left. + 3 \left[-\frac{125}{24} + \frac{125x}{6} - \frac{515x^2}{16} + \frac{135x^3}{8} - \frac{45x^4}{16} \right] L_{2/3}^2(x) L_{5/3}(x) \right\}. \end{aligned}$$

where L and \bar{L} are Laguerre functions.

6. DEFINITE INTEGRALS OF SPECIAL FUNCTIONS

6.1. Elliptic Integrals and Functions. [6]

(6.1)

$$\frac{1}{2} \int_0^1 K(x^2) dx = G.$$

[6]

$$(6.2) \quad \int_0^1 E(x^2) dx = G + \frac{1}{2}.$$

6.2. **The Exponential Integral and Related Functions.** [34]

$$(6.3) \quad \int_0^\infty t^{2\beta} e^{-3t^2} \operatorname{erf}(t) dt = \frac{\Gamma(1+\beta)}{\sqrt{\pi} 3^{\beta+1}} {}_2F_1\left(\frac{1}{2}, 1+\beta; \frac{3}{2}; -\frac{1}{3}\right).$$

6.3. **The Gamma Function and Related Functions.** [11][175, A075700]

$$(6.4) \quad \int_0^1 \ln \Gamma(t) dt = \frac{1}{2} \ln 2\pi.$$

[11]

$$(6.5) \quad \int_0^1 t \ln \Gamma(t) dt = \frac{\zeta'(2)}{2\pi^2} + \frac{1}{6} \ln 2\pi - \frac{\gamma}{12}.$$

[11]

$$(6.6) \quad \int_0^\infty 2^{-t} \ln \Gamma(t) dt = 2 \int_0^1 2^{-t} \ln \Gamma(t) dt - \frac{\gamma + \ln \ln 2}{\ln 2}.$$

[11]

$$(6.7) \quad \int_0^\infty 2^{-t} t \ln \Gamma(t) dt = 2 \int_0^1 2^{-t} (t+1) \ln \Gamma(t) dt - \frac{(\gamma + \ln \ln 2)(1 + 2 \ln 2) - 1}{\ln^2 2}.$$

[75]

$$(6.8) \quad \int_0^1 \ln \Gamma(q) \cos((2n+1)\pi q) dq \frac{2}{\pi^2} \left(\frac{\gamma + 2 \ln \sqrt{2\pi}}{(2n+1)^2} + 2 \sum_{k=1}^\infty \frac{\ln k}{4k^2 - (2n+1)^2} \right).$$

[75]

$$(6.9) \quad \int_0^1 B_{2m}(q) \ln \Gamma(q) dq = (-)^{m+1} \frac{(2m)! \zeta(2m+1)}{2(2\pi)^{2m}}.$$

[75]

$$(6.10) \quad \int_0^1 B_{2m-1}(q) \ln \Gamma(q) dq = \frac{B_{2m}}{2m} \left[\frac{\zeta'(2m)}{\zeta(2m)} - A \right],$$

where $A = 2 \ln \sqrt{2\pi} + \gamma$.

[75]

$$(6.11) \quad \int_0^1 q^n \ln \Gamma(q) dq = \frac{1}{n+1} \sum_{k=1}^{\lfloor (n+1)/2 \rfloor} (-)^k \binom{n+1}{2k-1} \frac{(2k!)}{k(2\pi)^{2k}} [A \zeta(2k) - \zeta'(2k)] \\ - \frac{1}{n+1} \sum_{k=1}^{\lfloor n/2 \rfloor} (-)^k \binom{n+1}{2k} \frac{(2k!)}{2(2\pi)^{2k}} \zeta(2k+1) + \frac{\ln \sqrt{2\pi}}{n+1},$$

where $A = 2 \ln \sqrt{2\pi} + \gamma$.

[75]

$$(6.12) \quad \int_0^1 (q - 1/2) \ln \Gamma(q) dq = \frac{1}{12} \left(\frac{6\zeta'(2)}{\pi^2} - 2 \ln \sqrt{2\pi} - \gamma \right).$$

[75]

$$(6.13) \quad \int_0^{1/2} \ln \Gamma(q+1) dq = \frac{\gamma}{8} + \frac{3 \ln \sqrt{2\pi}}{4} - \frac{13 \ln 2}{24} - \frac{3\zeta'(2)}{4\pi^2} - \frac{1}{2}.$$

[75]

(6.14)

$$\int_0^1 q^n \psi^{(m)}(q) dq = (-1)^m \frac{n!}{(n-m)!} \left[\frac{\gamma}{n-m+1} + (n-m)! \sum_{k=0}^{m-2} \frac{\Gamma(m-k)\zeta(m-k)}{(n-k)!} \right. \\ \left. + \sum_{k=0}^{n-m-1} (-)^k \binom{n-m}{k} [H_k \zeta(-k) + \zeta'(-k)] \right],$$

where H_k are harmonic sums.

[75]

$$(6.15) \quad \int_0^1 q^n \psi(q) dq = \zeta'(0) + \sum_{k=1}^{n-1} (-)^k \binom{n}{k} [H_k \zeta(-k) + \zeta'(-k)],$$

where H_k are harmonic sums.

[16]

$$(6.16) \quad \int_0^\infty \frac{t^{\alpha+z} \psi(\alpha+z+1)}{\Gamma(\alpha+z+1)} dz = \frac{t^\alpha}{\Gamma(\alpha+1)} + \nu(t, \alpha) \ln t, \quad \Re \alpha > -1.$$

$$(6.17) \quad \int_0^\infty \frac{\psi(\alpha+z+1)}{\Gamma(\alpha+z+1)} dz = \frac{1}{\Gamma(\alpha+1)}, \quad \Re \alpha > -1.$$

$$(6.18) \quad \int_1^\infty \frac{\psi(z)}{\Gamma(z)} dz = 1.$$

$$(6.19) \quad \int_0^\infty \frac{t^z \psi(z+1)}{\Gamma(z+1)} dz = 1 + \nu(t) \ln t, \quad t \neq 1.$$

[16]

(6.20)

$$\int_0^\infty \frac{t^{\alpha+z}}{\Gamma(\alpha+z+1)} \{ \psi(\alpha+z+1)^2 - \psi^{(1)}(\alpha+z+1) \} dz = \frac{2t^\alpha \ln t}{\Gamma(\alpha+1)} + (\ln t)^2 \nu(t, \alpha) + L^{-1} \left\{ \frac{\ln s}{s^{\alpha+1}} \right\}, \quad \Re \alpha > -1$$

where L^{-1} is the inverse Laplace transform.

$$(6.21) \quad \int_0^\infty \frac{t^z}{\Gamma(z+1)} \{ \psi(z+1)^2 - \psi^{(1)}(z+1) \} dz = -\gamma + \ln t (1 + \nu(t) \ln t).$$

$$(6.22) \quad \int_1^\infty \frac{1}{\Gamma(z)} \{ \psi^{(1)}(z) - \psi(z)^2 \} dz = \gamma.$$

(6.23)

$$\int_0^\infty \frac{t^{\alpha+z}}{\Gamma(\alpha+z+1)} \{ \psi(\alpha+z+1)^2 - \psi^{(1)}(\alpha+z+1) \} dz = \frac{t^\alpha}{\Gamma(\alpha+1)} [\psi(\alpha+1) + \ln t] + (\ln t)^2 \nu(t, \alpha).$$

(6.24)

$$\int_0^\infty \frac{t^{\alpha+z} z^\beta}{\Gamma(\alpha+z+1)} \frac{\psi(\alpha+z+1)}{\Gamma(\beta+1)} dz = \mu(t, \beta-1, \alpha) + \ln t \mu(t, \beta, \alpha), \quad \Re \alpha > -1, \Re \beta > -1.$$

$$(6.25) \quad \int_0^\infty \frac{t^{\alpha+z} z^\beta}{\Gamma(\alpha+z+1)} \frac{\psi(\alpha+z+1)^2 - \psi^{(1)}(\alpha+z+1)}{\Gamma(\beta+1)} dz$$

$$= \mu(t, \beta-2, \alpha) + 2 \ln t \mu(t, \beta-1, \alpha) + (\ln t)^2 \mu(t, \beta, \alpha), \quad \Re \alpha > -1, \Re \beta > 1.$$

[16] Let $L\{f(t)\} \equiv \int_0^\infty e^{-st} f(t) dt = F(s)$ be the Laplace transform and

$$(6.26) \quad \nu(z) \equiv \int_0^\infty \frac{z^t dt}{\Gamma(t+1)},$$

$$(6.27) \quad \nu(z, \alpha) \equiv \int_0^\infty \frac{z^{\alpha+t} dt}{\Gamma(\alpha+t+1)},$$

$$(6.28) \quad \mu(z, \beta) \equiv \int_0^\infty \frac{z^t t^\beta dt}{\Gamma(\beta+1)\Gamma(t+1)},$$

$$(6.29) \quad \mu(z, \beta, \alpha) \equiv \int_0^\infty \frac{z^{\alpha+t} t^\beta dt}{\Gamma(\beta+1)\Gamma(\alpha+t+1)},$$

then

$$(6.30) \quad L\{\nu(t)\} = \frac{1}{s \ln s}.$$

[16]

$$(6.31) \quad L\{\nu(t, \alpha)\} = \frac{1}{s^{1+\alpha} \ln s}.$$

[16]

$$(6.32) \quad L\{\mu(t, \beta)\} = \frac{1}{s(\ln s)^{\beta+1}}.$$

[16]

$$(6.33) \quad L\{\mu(t, \beta, \alpha)\} = \frac{1}{s^{\alpha+1}(\ln s)^{\beta+1}}.$$

Above $\Re \alpha > -1, \Re \beta > -1, \Re s > 1$. [16]

$$(6.34) \quad \frac{F(\ln s)}{s \ln s} = L\left\{\int_0^\infty \nu(t, x) f(x) dx\right\}.$$

[16]

$$(6.35) \quad \frac{F(\ln s)}{s^{\alpha+1} \ln s} = L\left\{\int_0^\infty \nu(t, \alpha+x) f(x) dx\right\}.$$

[16]

$$(6.36) \quad \frac{F(\ln s)}{s(\ln s)^{\beta+1}} = L\left\{\int_0^\infty \mu(t, \beta, x) f(x) dx\right\}.$$

[16]

$$(6.37) \quad \frac{F(\ln s)}{s^{\alpha+1}(\ln s)^{\beta+1}} = L\left\{\int_0^\infty \mu(t, \beta, \alpha+x) f(x) dx\right\}.$$

[16]

$$(6.38) \quad \int_0^\infty x^{\beta-1} \nu(t, \alpha+x) dx = \Gamma(\beta) \mu(t, \beta, \alpha), \quad \Re \alpha > -1, \Re \beta > 0.$$

[16]

$$(6.39) \quad \int_0^\infty x^{\beta-1} \nu(t, x) dx = \Gamma(\beta) \mu(t, \beta).$$

[16]

$$(6.40) \quad \int_0^\infty \nu(t, \alpha + x) dx = \mu(t, 1, \alpha).$$

[16]

$$(6.41) \quad \int_0^\infty \nu(t, x) dx = \mu(t, 1).$$

[16]

$$(6.42) \quad \int_0^\infty t^{\lambda-1} \mu(t, \beta, \alpha + x) dx = \Gamma(\lambda) \mu(t, \beta + \lambda, \alpha), \quad \Re \alpha > -1, \Re \beta > -1, \Re \lambda > 0.$$

[16]

$$(6.43) \quad \frac{1}{3\pi} \int_0^\infty \left(\frac{x}{t}\right)^{1/2} K_{1/2}(\phi) \mu(x, \beta, \alpha) dx = 3^{\beta+1} \mu(t, \beta, \alpha/3), \phi \equiv 2(x^3/27t)^{1/2},$$

and others,

6.4. Cylinder Functions. [147]

(6.44)

$$2\pi(-1)^{(n-m)/2} \int_0^\infty dk J_{n+1}(2\pi k) J_m(2\pi kr) = R_n^m(r) = \sum_{s=0}^{(n-m)/2} (-1)^s \binom{n-s}{s} \binom{n-2s}{(n-m)/2-s} r^{n-2s},$$

for $n \geq 0$, $0 \leq m \leq n$, $n-m$ even.

[27]

$$(6.45) \quad \int_0^\infty J_\mu(ct \sin \phi) J_\nu(ct \sin \Phi) K_\rho(ct \cos \phi \cos \Phi) dt = \dots$$

[27]

(6.46)

$$\int_0^\infty J_\mu(ct \sin \phi \sin \Phi) J_\nu(ct \cos \phi \sin \Phi) J_\rho(ct) dt = \frac{\Gamma(\frac{1+\mu+\nu+\rho}{2}) \sin^\mu \phi \cos^\mu \Phi \cos^\nu \phi \sin^\nu \Phi}{c^{\mu+\nu+1} \Gamma(\mu+1) \Gamma(\nu+1) \Gamma(\frac{1-\mu-\nu+\rho}{2})}$$

(6.47)

$$\times {}_2F_1\left(\frac{1+\mu+\nu-\rho}{2}, \frac{1+\mu+\nu+\rho}{2}; \mu+1; \sin^2 \phi\right) {}_2F_1\left(\frac{1+\mu+\nu-\rho}{2}, \frac{1+\mu+\nu+\rho}{2}; \nu+1; \sin^2 \Phi\right)$$

where ϕ and Φ are positive angles whose sum is acute.

6.4.1. Cylinder Functions combined with x and x^2 . [203]

$$(6.48) \quad \int_0^1 x J_\nu(\lambda_n x) J_\nu(\lambda_m x) dx = \frac{1}{2} \{ (1 - \nu^2/\lambda_n^2) J_\nu^2(\lambda_n) + J_\nu'^2(\lambda_n) \} \delta_{nm}$$

for $\{\lambda_n\}$ ($n = 1, 2, \dots$) a sequence of successive positive roots of the equation $x J_\nu'(x) + H J_\nu(x) = 0$, where H is a real number and $\nu \geq -1$.

[125]

$$(6.49) \quad \int_0^1 x^{2+l+2n} j_l(2\pi\sigma x) dx = \frac{1}{2\pi\sigma} \sum_{k=0}^n \frac{(-n)_k}{(\pi\sigma)^k} j_{l+k+1}(2\pi\sigma); \quad l, n = 0, 1, 2, 3, \dots$$

[152]

$$(6.50) \quad \int_0^\infty k^2 j_l(kr) j_l(kr') dk = \frac{\pi \delta(r - r')}{2r^2}.$$

[93, 136]

$$(6.51) \quad \frac{4p_1 p_2 p_3}{\pi} \int_0^\infty x^2 j_1(p_1 x) j_2(p_2 x) j_3(p_3 x) dx = \Delta(p_1, p_2, p_3) (-1)^{(l_1 + l_2 + l_3)/2} \\ \times \frac{1}{2} \sum_{k_1=0}^{l_1} \sum_{k_2=0}^{l_2} \sum_{k_3=0}^{l_3} \frac{(-1)^{k_1+k_2+k_3}}{(k_1+k_2+k_3)!} \prod_{i=1}^3 \frac{(l_i+k_i)!}{k_i!(l_i-k_i)!} (2p_i)^{-k_i} \\ \times [(-1)^{l_1+k_1} (p_2+p_3-p_1)^{k_1+k_2+k_3} + (-1)^{l_2+k_2} (p_3+p_1-p_2)^{k_1+k_2+k_3} \\ + (-1)^{l_3+k_3} (p_1+p_2-p_3)^{k_1+k_2+k_3} - (p_1+p_2+p_3)^{k_1+k_2+k_3}]$$

supposed that p_1 , p_2 and p_3 can be the sides of a plane triangle, that is where $\Delta(\cdot) = 1$ if they form a non-degenerate triangle, $\Delta(\cdot) = 1/2$ if they form a degenerate triangle, and $\Delta(\cdot) = 0$ otherwise.

$$(6.52) \quad \int_0^\infty x j_0(ax) j_0(bx) j_1(cx) dx = \frac{\pi}{8abc^2} (c^2 - (a-b)^2)$$

for $c > 0$, $a > 0$, $|c-a| < b < c+a$.

[102]

$$(6.53) \quad \int_0^\infty J_{n_1}(k_1\rho) J_{n_2}(k_2\rho) J_{n_3}(k_3\rho) \rho d\rho = \frac{\Delta}{6\pi A} [\cos(n_1\alpha_2 - n_2\alpha_1) + \cos(n_2\alpha_3 - n_3\alpha_2) + \cos(n_3\alpha_1 - n_1\alpha_3)]$$

if $n_1 + n_2 + n_3 = 0$, Δ as above, the area of the triangle of k_1 , k_2 and k_3 given by $2A = k_1 k_3 \sin \theta_{13}$, and α three external angles in that triangle.

6.4.2. Cylinder Functions and Rational Functions. [82]

$$(6.54) \quad \int_0^\infty \frac{J_\nu(x)}{x^2 + a^2} dx = \frac{i}{a} [S_{0,\nu}(ia - e^{-i\nu\pi/2} K_\nu(a))] = \frac{1}{a} [is_{0,\nu}(ia) + \frac{\pi}{2} \sec \frac{\nu\pi}{2} I_\nu(a)].$$

[180]

$$(6.55) \quad \int_0^\infty x^{1-2n} J_\nu(ax) J_\nu(bx) \frac{dx}{x^2 + c^2} = (-1)^n c^{-2n} \left\{ I_\nu(bc) K_\nu(ac) \right. \\ \left. - \frac{1}{2} \left(\frac{b}{a} \right)^\nu \frac{\pi}{\sin \pi \nu} \sum_{p=0}^{n-1} \frac{(ac/2)^{2p}}{p! \Gamma(1-\nu+p)} \sum_{k=0}^{n-1-p} \frac{(bc/2)^{2k}}{k! \Gamma(1+\nu+k)} \right\}$$

for $0 < b < a$, $\Re c > 0$, $\Re \nu > n-1$, $n = 1, 2, \dots$. For $0 < a < b$, the arguments a and b should be interchanged.

6.4.3. *Cylinder Functions and Powers.* [110, 114]

(6.56)

$$\int_0^1 x^\mu J_\nu(ax) dx = 2^\mu \frac{\Gamma(\frac{1}{2} + \frac{1}{2}\mu + \frac{1}{2}\nu)}{a^{\mu+1} \Gamma(\frac{\nu}{2} + \frac{1}{2} - \frac{1}{2}\mu)} + a^{-\mu} \{(\mu + \nu - 1)J_\nu(a)S_{\mu-1, \nu-1}(a) - J_{\nu-1}(a)S_{\mu, \nu}(a)\}$$

$$[a > 0, \Re(\mu + \nu) > -1].$$

$$(6.57) \quad \int_0^u z^\nu K_\nu(x) dx = \frac{2^\nu - 1}{1 + 2\nu} [(1 + 2\nu)\Gamma(\nu) {}_1F_2(1/2; 3/2, 1 - \nu; u^2/4) + 2(u/2)^{1+2\nu} \Gamma(-\nu) {}_1F_2(\nu + 1/2; 1 + \nu, 3/2 + \nu; u^2/4)].$$

[147]

$$(6.58) \quad \int_0^\infty x^{-P} [1 - J_0(bx)] dx = \frac{\pi b^{P-1}}{2^P \Gamma^2(\frac{P+1}{2}) \sin[\pi(P-1)/2]}.$$

[72, p 22]

(6.59)

$$\int_0^1 x^\mu J_\nu(xy) \sqrt{xy} dx = y^{-\mu-1} [(\nu + \mu - \frac{1}{2}) y J_\nu(y) S_{\mu-1/2, \nu-1}(y) - y J_{\nu-1}(y) S_{\mu+1/2, \nu}(y) + 2^{\mu+1/2} \frac{\Gamma(\frac{\mu}{2} + \frac{\nu}{2} + \frac{3}{4})}{\Gamma(\frac{\nu}{2} - \frac{\mu}{2} + \frac{1}{4})}]$$

for $\Re(\mu + \nu) > -3/2$.

[82]

$$(6.60) \quad \int_0^\infty \frac{x^{\rho-1} J_\nu(ax)}{(x^2 + k^2)^{\mu+1}} dx = \frac{a^\nu k^{\rho+\nu-2\mu-2} \Gamma(\rho/2 + \nu/2) \Gamma(\mu + 1 - \rho/2 - \nu/2)}{2^{\nu+1} \Gamma(\mu + 1) \Gamma(\nu + 1)} \\ \times {}_1F_2\left(\frac{\rho + \nu}{2}; \frac{\rho + \nu}{2} - \mu, \nu + 1; \frac{a^2 k^2}{4}\right) + \frac{a^{2\mu+2-\rho} \Gamma(\nu/2 + \rho/2 - \mu - 1)}{2^{2\mu+3-\rho} \Gamma(\mu + 2 + \nu/2 - \rho/2)} \\ \times {}_1F_2\left(\mu + 1; \mu + 2 + \frac{\nu - \rho}{2}, \mu + 2 - \frac{\nu + \rho}{2}; \frac{a^2 k^2}{4}\right), \quad a > 0, -\Re\nu < \Re\rho < 2\Re\mu + \frac{7}{2}, \Re k > 0.$$

[204]

$$\int_0^\infty \frac{J_{n+1}(k) J_{n'+1}(k)}{k(k^2 + k_0^2)^{1+\gamma/2}} dk =$$

$$k_0^{n+n'-\gamma} \frac{\Gamma(\frac{\gamma-n-n'}{2}) \Gamma(\frac{n+n'}{2} + 1)}{2^{n+n'+3} \Gamma(n+2) \Gamma(n'+2) \Gamma(1 + \gamma/2)}$$

$$\times {}_3F_4\left(\begin{matrix} \frac{n+n'}{2} + 1, \frac{n+n'}{2} + 2, \frac{n+n'+3}{2} \\ n+2, n'+2, n+n'+3, 1 + \frac{n+n'-\gamma}{2} \end{matrix} \mid k_0^2\right)$$

$$+ \frac{\Gamma(\frac{n+n'-\gamma}{2}) \Gamma(3 + \gamma)}{2^{3+\gamma} \Gamma(3 + \frac{\gamma+n+n'}{2}) \Gamma(2 + \frac{\gamma+n-n'}{2}) \Gamma(2 + \frac{\gamma+n'-n}{2})} {}_3F_4\left(\begin{matrix} 2 + \frac{\gamma}{2}, 1 + \frac{\gamma}{2}, \frac{3+\gamma}{2} \\ 2 + \frac{\gamma+n-n'}{2}, 2 + \frac{\gamma+n'-n}{2}, 3 + \frac{\gamma+n+n'}{2}, 1 + \frac{\gamma-n-n'}{2} \end{matrix} \mid k_0^2\right)$$

[122, p50]
(6.61)

$$\int_0^\infty J_{\nu+n}(at)J_{\nu-n-1}(bt)dt = \begin{cases} \frac{b^{\nu-n-1}\Gamma(\nu)}{a^{\nu-n}\Gamma(\nu-n)} {}_2F_1(\nu, -n; \nu-n; \frac{b^2}{a^2}), & 0 < b < a, \\ (-1)^n/(2a), & 0 < b = a, \\ 0, & 0 < a < b. \end{cases}$$

where $n = 0, 1, 2, \dots$, $\Re \nu > 0$.

[122, p50]

(6.62)
$$\int_0^\infty J_\nu(at)J_{\nu+1}(bt)dt = \begin{cases} a^\nu b^{-\nu-1}, & 0 < b < a \\ 1/(2a), & 0 < b = a \\ 0, & 0 < a < b \end{cases}$$

where a, b real positive, $\Re \nu > -1$.

[122, p50]

(6.63)
$$\int_0^\infty J_\mu(at)J_\nu(at)dt = \frac{2}{\pi} \frac{\sin(\frac{\nu-\mu}{2}\pi)}{\nu^2 - \mu^2},$$

$\Re(\nu + \mu) > 0$, $a > 0$.

[147]

(6.64)
$$\int_0^\infty x^{-P} \left(1 - \frac{4J_1^2(x)}{x^2}\right) dx = \frac{\pi\Gamma(P+2)}{2^P\Gamma^2(\frac{P+3}{2})\Gamma(\frac{P+5}{2})\Gamma(\frac{P+1}{2})\sin[\pi(P-1)/2]}.$$

[79]

(6.65)
$$\int_0^\infty t^{\rho-\mu-\nu-3} J_\mu(at)J_\nu(bt)J_\rho(ct)dt = \frac{2^{\rho-\mu-\nu-3}a^\mu b^\nu \Gamma(\rho-1)}{c^{\rho-2}\Gamma(\mu+1)\Gamma(\nu+1)} \left(1 - \frac{\rho-1}{\mu-1} \frac{a^2}{c^2} - \frac{\rho-1}{\nu-1} \frac{b^2}{c^2}\right).$$

[190] Let

(6.66)
$$G_{lmn} \equiv \int_0^\infty dx x^\alpha J_l(x)J_m(x)J_n(\beta x)$$

with $\Re \alpha < 3/2$ and $\Re(\alpha + l + m + n + 1) > 0$, then for $\beta/2 < 1$

(6.67)
$$\begin{aligned} G_{lmn} = & \frac{\Gamma(\frac{\alpha+n}{2})(\beta/2)^{-\alpha}}{2\Gamma(\frac{-l+m+1}{2})\Gamma(\frac{l-m+1}{2})\Gamma(\frac{-\alpha+n+2}{2})} {}_4F_3\left(\begin{matrix} \frac{l+m+1}{2}, \frac{l-m+1}{2}, \frac{-l-m+1}{2}, \frac{-l+m+1}{2} \\ \frac{-\alpha+n+2}{2}, \frac{1}{2}, \frac{-\alpha-n+2}{2} \end{matrix} \mid \beta^2/4\right) \\ & - \frac{\Gamma(\frac{l+m+2}{2})\Gamma(\frac{\alpha+n-1}{2})(\beta/2)^{-\alpha+1}}{\Gamma(\frac{-l+m}{2})\Gamma(\frac{l+m}{2})\Gamma(\frac{l-m}{2})\Gamma(\frac{-\alpha+n+3}{2})} {}_4F_3\left(\begin{matrix} \frac{l+m+2}{2}, \frac{l-m+2}{2}, \frac{-l-m+2}{2}, \frac{-l+m+2}{2} \\ \frac{-\alpha+n+3}{2}, \frac{3}{2}, \frac{-\alpha-n+3}{2} \end{matrix} \mid \beta^2/4\right) \\ & + \frac{2^{\alpha+n}\Gamma(-\alpha-n)\Gamma(\frac{\alpha+l+m+n+1}{2})(\beta/2)^n}{\Gamma(\frac{-\alpha-l+m-n+1}{2})\Gamma(\frac{-\alpha+l+m-n+1}{2})\Gamma(\frac{-\alpha+l-m-n+1}{2})\Gamma(n+1)} \\ & \times {}_4F_3\left(\begin{matrix} \frac{\alpha+l+m+n+1}{2}, \frac{\alpha+l-m+n+1}{2}, \frac{\alpha-l-m+n+1}{2}, \frac{\alpha-l+m+n+1}{2} \\ n+1, \frac{n+\alpha+2}{2}, \frac{\alpha+n+1}{2} \end{matrix} \mid \beta^2/4\right), \end{aligned}$$

and for $\beta/2 > 1$

(6.68)
$$G_{lmn}(\alpha, \beta) = \frac{2^{-l-m-1}\Gamma(\frac{\alpha+l+m+n+1}{2})(\beta/2)^{-\alpha-l-m-1}}{\Gamma(\frac{-\alpha-l-m+n+1}{2})\Gamma(l+1)\Gamma(m+1)} {}_4F_3\left(\begin{matrix} \frac{\alpha+l+m+n+1}{2}, \frac{l+m+1}{2}, \frac{l+m+2}{2}, \frac{\alpha+l+m-n+1}{2} \\ m+1, l+1, m+l+1 \end{matrix} \mid 4/\beta^2\right)$$

[187]
 (6.69)
$$\int_0^\infty t \left(\frac{t}{\sqrt{u^2 + t^2}} - 1 \right) J_0(\gamma t) dt = \frac{u^2}{2} [I_1(u\gamma/2)K_1(u\gamma/2) - I_0(u\gamma/2)K_0(u\gamma/2)].$$

6.4.4. *Cylinder Functions and Exponentials.*

6.4.5. *Cylinder Functions, Exponentials and Powers.* With

$$l_1(a, b, c) \equiv \frac{1}{2} \left[\sqrt{(a+b)^2 + c^2} - \sqrt{(a-b)^2 + c^2} \right]$$

$$l_2(a, b, c) \equiv \frac{1}{2} \left[\sqrt{(a+b)^2 + c^2} + \sqrt{(a-b)^2 + c^2} \right]$$

at $a > 0, b > 0, c > 0$ [79]

(6.70)
$$\int_0^\infty e^{-cx} J_1(ax) J_{1/2}(bx) \frac{dx}{x^{3/2}} = \frac{\sqrt{2}}{\sqrt{\pi ba}} \left[\frac{l_1}{2} \sqrt{a^2 - l_1^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{l_1}{a} \right) + c(\sqrt{b^2 - l_1^2} - b) \right].$$

(6.71)
$$\int_0^\infty e^{-cx} J_1(ax) J_{1/2}(bx) \frac{dx}{\sqrt{x}} = \frac{\sqrt{2}}{\sqrt{\pi ba}} (b - \sqrt{b^2 - l_1^2}).$$

(6.72)
$$\int_0^\infty e^{-cx} J_1(ax) J_{1/2}(bx) \sqrt{x} dx = \frac{\sqrt{2}}{\sqrt{\pi ba}} \frac{l_1 \sqrt{a^2 - l_1^2}}{l_2^2 - l_1^2}.$$

(6.73)
$$\int_0^\infty e^{-cx} J_1(ax) J_{3/2}(bx) \sqrt{x} dx = \frac{2l_1^2 \sqrt{b^2 - l_1^2}}{\sqrt{2\pi} b^{3/2} a (l_2^2 - l_1^2)}.$$

(6.74)
$$\int_0^\infty e^{-cx} J_1(ax) J_{3/2}(bx) \frac{dx}{\sqrt{x}} = \frac{1}{\sqrt{2\pi} b^{3/2} a} \left(-l_1 \sqrt{a^2 - l_1^2} + a^2 \sin^{-1} (l_1/a) \right).$$

(6.75)
$$\int_0^\infty e^{-cx} J_1(ax) J_{5/2}(bx) \frac{dx}{\sqrt{x}} = \frac{2^{-1/2} c}{\sqrt{\pi} b^{5/2} a} \left(l_1 \sqrt{a^2 - l_1^2} + \frac{2a^2 l_1}{\sqrt{a^2 - l_1^2}} - 3a^2 \sin^{-1} (l_1/a) \right).$$

and similar combinations of even and odd-indexed $J(ax)$ and $J(bx)$.

[7]

(6.76)
$$\int_0^\infty x^\nu e^{-x/2} J_\nu(\mu x) L_n^{2\nu}(x) dx = 2^\nu \Gamma(\nu + \frac{1}{2}) \frac{1}{\sqrt{\pi \mu}} (\sin \theta)^{\nu + \frac{1}{2}} C_n^{\nu + \frac{1}{2}}(\cos \theta),$$

$\mu \geq 0, \nu > -\frac{1}{2}, \cos \theta \equiv \frac{\mu^2 - 1/4}{\mu^2 + 1/4}, C_n^\lambda(x)$ ultraspherical polynomial.

[150]

(6.77)
$$\int_0^\infty \frac{x^{\nu+1}}{a^2 + x^2} J_\nu(xy) dx = \frac{a^\nu}{\sqrt{y}} K_\nu(ay), \quad 1 < \Re \nu < 3/2.$$

[150]

(6.78)
$$\int_0^\infty \frac{x^{\nu+1}}{(a^2 + x^2)^\mu} J_\nu(xy) dx = \frac{2^{1-\mu} a^{\nu-\mu+1} y^{\mu-1}}{\Gamma(\mu)} K_{\nu-\mu+1}(ay), \quad \Re \nu > -1, \Re(2\mu-\nu) > 1/2.$$

[150]

$$(6.79) \quad \int_0^\infty \frac{x^{1-\nu}}{(a^2+x^2)^\mu} J_\nu(xy) dx = a^{-\mu-\nu+1} y^{\mu-1} \left[2^{-\mu} \frac{\Gamma(1-\mu)}{1-\nu} I_{\nu+\mu-1}(ay) \right. \\ \left. - 2 \frac{1-\mu}{\Gamma(\nu)} e^{-i\frac{\pi}{2}(\nu-\mu+1)} s_{-\mu+\nu, -\mu-\nu+1}(iay) \right], \quad \Re(\nu+2\mu) > 1/2.$$

[34]

$$(6.80) \quad \int_0^\infty x^{\beta-1} e^{-x} I_n(x) dx = \sum_{j=0}^\infty \frac{\Gamma(\beta+2j+n)}{2^{2j+n} j! (j+n)!}.$$

[34]

$$(6.81) \quad \int_0^\infty x^{2n+1} e^{-x^2/4a} I_0(x) dx = 2^{2n+1} a^{n+1} n! e^a L_n(-a) = 2^{2n+1} a^{n+1} n! {}_1F_1(n+1; 1; a).$$

$$(6.82) \quad \int_0^\infty x^3 e^{-x^2/4a} I_0(x) dx = 8a^2(1+a)e^a.$$

[2]

$$(6.83) \quad \int_0^\infty x e^{-x^2/z} J_2(x) Y_2(x) dx = -\frac{2}{\pi} + \frac{4}{\pi z} - \frac{z K_2(z/2)}{2\pi \exp(z/2)}.$$

[2]

$$(6.84) \quad \int_0^\infty x e^{-x^2/z} I_3(x) K_3(x) dx = -\frac{32+16z+3z^2}{2z^2} + \frac{\exp(z/2) z K_3(z/2)}{4}.$$

[2]

$$(6.85) \quad \int_0^\infty x^3 e^{-x^2/z} J_2(x) Y_2(x) dx = -\frac{4}{\pi} + \frac{z^2(2+z)K_0(z/2)}{4\pi \exp(z/2)} + \frac{z(8+4z+z^2)K_1(z/2)}{4\pi \exp(z/2)}.$$

[2]

$$(6.86) \quad \int_0^\infty x^5 e^{-x^2/z} I_3(x) K_3(x) dx = -32 + \frac{1}{8} e^{z/2} z^2 (32 - 16z + 5z^2 - z^3) K_0(z/2) \\ + \frac{1}{8} e^{z/2} z (128 - 64z + 24z^2 - 6z^3 + z^4) K_1(z/2).$$

[68] Let

$$(6.87) \quad I(\mu, \nu, \lambda) \equiv \int_0^\infty J_\mu(at) J_\nu(bt) e^{-ct} t^\lambda dt,$$

then

$$(6.88) \quad I(n, n; 0) = \frac{(-)^n k}{\pi \sqrt{ab}} \int_0^{\pi/2} \frac{\cos(2n\psi) d\psi}{\sqrt{1-k^2 \sin^2 \psi}},$$

$$(6.89) \quad I(n, n; 1) = \frac{(-)^n c k^3}{4\pi (ab)^{3/2}} \int_0^{\pi/2} \frac{\cos(2n\psi) d\psi}{(1-k^2 \sin^2 \psi)^{3/2}},$$

in particular

$$(6.90) \quad I(0, 0; 0) = \frac{k}{2\sqrt{ab}} F_0(k),$$

$$(6.91) \quad I(1, 1; 0) = \frac{1}{k\sqrt{ab}} [(1 - k^2/2)F_0(k) - E_0(k)],$$

$$(6.92) \quad I(0, 0; 1) = \frac{ck^3 E_0(k)}{8k'^2(ab)^{3/2}},$$

and more results on $I(n+1, n; \pm 1)$ and $I(n+1, n; 0)$ in terms of Elliptic Integrals. Associated recurrences:

$$(6.93) \quad a[I(\mu+1, \nu; \lambda) + I(\mu-1, \nu; \lambda)] = 2\mu I(\mu, \nu; \lambda-1);$$

$$(6.94) \quad b[I(\mu, \nu+1; \lambda) + I(\mu, \nu-1; \lambda)] = 2\nu I(\mu, \nu; \lambda-1);$$

$$(6.95) \quad aI(\mu+1, \nu; \lambda) - bI(\mu, \nu-1; \lambda) = C_{\mu, \nu} + (\mu - \nu + \lambda)I(\mu, \nu; \lambda-1) - cI(\mu, \nu; \lambda),$$

with

$$(6.96) \quad C_{\mu, \nu} \equiv \begin{cases} \frac{a^\mu b^\nu}{2^{\mu+\nu} \Gamma(\mu+1) \Gamma(\nu+1)}, & \text{if } \lambda + \mu + \nu = 0; \\ 0 & \text{if } \lambda + \mu + \nu > 0. \end{cases}$$

6.4.6. *Cylinder and Trigonometric Functions and Powers.* [79]

$$(6.97) \quad \int_0^\infty \sin(cx) x^{\nu-\mu-4} J_\mu(ax) J_\nu(bx) dx = \frac{\Gamma(\nu) a^\mu b^{-\nu} c}{2^{\mu-\nu+3} \Gamma(\mu+1)} \left(\frac{b^2}{\nu-1} - \frac{a^2}{\mu+1} - \frac{2c^2}{3} \right).$$

[79]

$$(6.98) \quad \int_0^\infty \cos(cx) x^{\nu-\mu-3} J_\mu(ax) J_\nu(bx) dx = \frac{\Gamma(\nu) a^\mu b^{-\nu}}{2^{\mu-\nu+3} \Gamma(\mu+1)} \left(\frac{b^2}{\nu-1} - \frac{a^2}{\mu+1} - 2c^2 \right).$$

[165]

$$(6.99) \quad \int_0^\infty J_{2n}(a\sqrt{t^2+2bt}) e^{-pt} dt = (-1)^n \left\{ \frac{e^{b(p-\sqrt{p^2+a^2})}}{\sqrt{p^2+a^2}} + \frac{2e^{bp}}{a} \right. \\ \left. \times \sum_{k=1}^n (-1)^k (2k-1) I_{k-1/2}[b(\sqrt{p^2+a^2}-a)/2] K_{k-1/2}[b(\sqrt{p^2+a^2}+a)/2] \right\}.$$

[165]

$$(6.100) \quad \int_0^\infty J_{2n}(a\sqrt{t^2+2bt}) e^{-pt} dt = (-1)^n \left\{ \frac{e^{b(p-\sqrt{p^2+a^2})}}{\sqrt{p^2+a^2}} + 2n \sum_{\lambda=1}^n \frac{(n-1+\lambda)!}{\lambda!(n-\lambda)!} \frac{1}{b^{2\lambda-1} a^{2\lambda}} \right. \\ \left. \times \sum_{\mu=0}^{\lambda-1} \frac{(2\lambda-2-\mu)!}{\mu!(\lambda-1-\mu)!} (2b\sqrt{p^2+a^2})^\mu \left(e^{b(p-\sqrt{p^2+a^2})} - \sum_{\nu=0}^{2\lambda-2-\mu} \frac{[b(p-\sqrt{p^2+a^2})]^\nu}{\nu!} \right) \right\}.$$

[34]

$$(6.101) \quad \int_0^\infty x^{\beta-1} e^{-x} I_0(2\sqrt{ax}) dx = \sum_{k=0}^\infty \frac{a^k}{k!^2} \Gamma(\beta+k).$$

7. DEFINITE INTEGRALS OF SPECIAL FUNCTIONS II.

7.1. Associated Legendre Functions and Powers. [159]

$$\begin{aligned} & \int_{-1}^1 P_{l+\alpha}^{-\alpha}(x) P_{k+\beta}^{-\beta}(x) (1-x^2)^{-p-1} dx = 2^{-(\alpha+\beta)} \\ & \times \frac{\Gamma[\frac{1}{2}(k+1)]\Gamma(-\frac{1}{2}k-\beta)\Gamma[\frac{1}{2}(\alpha+\beta)-p]\Gamma[\frac{1}{2}(\alpha-\beta-p)]\Gamma[\frac{1}{2}(l-k)]\Gamma[\frac{1}{2}(l+k+1)+\beta]}{\Gamma(\beta+1)\Gamma(-\beta)\Gamma(\alpha+1)\Gamma(-\frac{1}{2}k)\Gamma[\frac{1}{2}(k+1)+\beta]\Gamma[\frac{1}{2}(l-k+\alpha-\beta)-p]\Gamma[\frac{1}{2}(l+k+\alpha+\beta+1)-p]} \\ & \times {}_4F_3\left(\begin{matrix} \frac{1}{2}(\alpha-\beta)+p+1, & \frac{1}{2}(\alpha-\beta)-p, & -\frac{1}{2}(l-1), & -\frac{1}{2}l \\ \frac{1}{2}(k-l)+1, & -\frac{1}{2}(l+k-1)-\beta, & \alpha+1 & \end{matrix}; 1\right) \end{aligned}$$

for $\Re[\frac{1}{2}(\alpha+\beta)-p] > 0$, for k, l both even or both odd.
[159]

$$\begin{aligned} (7.1) \quad & \int_{-1}^1 P_l^m(x) P_k^n(x) (1-x^2)^{-p-1} dx = (-1)^{(l-k+m-n)/2} 2^{-(m+n)} \\ & \times \frac{(l+m)! \Gamma[\frac{1}{2}(m+n)-p] \Gamma[\frac{1}{2}(m-n)-p] \Gamma[\frac{1}{2}(l+k+1-m+n)]}{(l-m)! m! [\frac{1}{2}(k-l+m-n)]! \Gamma[\frac{1}{2}(l-k)-p] \Gamma[\frac{1}{2}(l+k+1)-p]} \\ & \times {}_4F_3\left(\begin{matrix} \frac{1}{2}(m-n)+p+1, & \frac{1}{2}(m-n)-p, & -\frac{1}{2}(l-m-1), & -\frac{1}{2}(l-m) \\ \frac{1}{2}(k-l+m-n)+1, & -\frac{1}{2}(l+k-1-m+n), & m+1 & \end{matrix}; 1\right) \end{aligned}$$

for $\Re[\frac{1}{2}(m+n)-p] > 0$, for $k-n \geq l-m$.
[159]

$$(7.2) \quad \int_{-1}^1 P_l^m(x) P_k^m(x) (1-x^2)^{-1} dx = \frac{1}{m} \frac{(l+m)!}{(l-m)!}, \quad k \geq l \geq m > 0.$$

[159]

$$\begin{aligned} & \int_{-1}^1 P_l^m(x) P_k^m(x) (1-x^2)^{-2} dx = \frac{(l+m)!}{2(l-m)!(m-1)m(m+1)} \\ & \times [l(l+1) + (m-1)(m+1) + \frac{1}{2}(k-l)(m+1)(k+l+1)], \quad k \geq l \geq m > 1. \end{aligned}$$

[159]

$$(7.3) \quad \int_{-1}^1 P_l^m(x) P_k^m(x) dx = \frac{2}{2l+1} \frac{(l+m)!}{(l-m)!} \delta_{kl}$$

[159]

$$\begin{aligned} & \int_{-1}^1 P_l^m(x) P_k^m(x) (1-x^2) dx = \\ & (-1)^{(k-l)/2} \frac{(l+m)! [l(l+1) + (m-1)(m+1) + \frac{1}{2}(k-l)(m+1)(k+l+1)]}{(l-m)! [\frac{1}{2}(k-l)+1]! \Gamma[\frac{1}{2}(l-k)+2] (k+l-1)(k+l+1)(k+l+3)} \end{aligned}$$

for $k \geq l \geq m \geq 0$, zero for $k-l \geq 4$.

7.2. Associated Legendre functions, powers, and trigonometric functions.

[26]

$$(7.4) \quad \int_0^z P_n[\cos(z-t)]P_n^{-m}(\cos t)\frac{dt}{\sin t} = \frac{P_n^{-m}(\cos z)}{m}, \quad \Re m > 0.$$

$$(7.5) \quad \int_0^z \sin^m t P_{n-m-1}[\cos(z-t)]P_n^{-m}(\cos t)dt = \frac{\Gamma(m+1/2)}{2^{1/2}\Gamma(m+1)} \sin^{m+1/2} z P_{n-1/2}^{-m-1/2}(\cos z), \quad \Re m > -1/2.$$

$$(7.6) \quad \int_0^z \sin^{m+1} t P_{n-m-2}[\cos(z-t)]P_n^{-m}(\cos t)dt = \frac{\Gamma(m+3/2)}{2^{1/2}\Gamma(m+2)} \sin^{m+3/2} z P_{n-1/2}^{-m-1/2}(\cos z), \quad \Re m > -1.$$

$$(7.7) \quad \int_0^z \sin^{-1-k} t \sin^k(z-t)P_n^{-m}(\cos t)P_n^{-k}[\cos(z-t)] = \frac{2^k \Gamma(m-k)\Gamma(k+1/2)}{\sqrt{\pi}\Gamma(k+m+1)} \sin^k z P_n^{-m}(\cos z), \quad \Re m > \Re k > -1/2.$$

$$(7.8) \quad \int_0^z \sin^m t \sin^{-m}(z-t)P_n^{-m}(\cos t)P_{n-1}^m[\cos(z-t)] = \frac{\sin[(m+n)z]}{(m+n)\cos m\pi}, \quad -1/2 < \Re m < 1/2.$$

$$(7.9) \quad \int_0^z P_{n+1}^{-m}(\cos t) \sin[n(z-t)]dt = 2^{3/2} \sin^{1/2} z \sum_{r=0}^{\infty} \frac{(-1)^r (m+n+2)_{2r} (3/2)_r \Gamma(m+r+3/2)}{r! \Gamma(m+r+1)} \\ \times \frac{m+2r+3/2}{(m+2r+1)(m+2r+2)} P_{n-1/2}^{-(m+3/2+2r)}(\cos z).$$

[146]

$$(7.10) \quad \int_0^\pi d\theta \sin^{|m|+1} \theta \exp(\pm iR \cos \theta) P_n^{|m|}(\cos \theta) = 2(\pm i)^{n+|m|} \frac{(n+|m|)!}{(n-|m|)!} \frac{j_n(R)}{R^{|m|}}.$$

[146, (23)]

$$(7.11) \quad \int_0^\pi d\theta \sin \theta \exp(iR \cos \alpha \cos \theta) P_n^m(\cos \theta) J_m(R \sin \alpha \sin \theta) = 2i^{n-m} P_n^m(\cos \alpha) j_n(R).$$

[15, §11.4,13][123]

$$(7.12) \quad \int_0^\infty e^{-c^2 x^2} H_{2m}(ax) H_{2k}(bx) dx = \\ (-)^{m+k} 2^{2m+2k-1} \frac{\Gamma(m+1/2)\Gamma(k+1/2)}{\pi c^{2m+2k+1}} (c^2-a^2)^m (c^2-b^2)^k {}_2F_1(-m, -k; \frac{1}{2}; \frac{a^2 b^2}{(c^2-a^2)(c^2-b^2)}),$$

$$c^2 - a^2 - b^2 > 0.$$

[15, §11.4,14]

$$(7.13) \quad \int_0^\infty e^{-c^2 x^2} H_{2m+1}(ax) H_{2k+1}(bx) dx = (-)^{m+k} 2^{2m+2k+1} \frac{\Gamma(m+3/2)\Gamma(k+3/2)}{\Gamma(3/2)} \\ \times \frac{ab(c^2-a^2)^m (c^2-b^2)^k}{c^{2m+2k+3}} {}_2F_1(-m, -k; \frac{3}{2}; \frac{a^2 b^2}{(c^2-a^2)(c^2-b^2)}).$$

[15, §11.4,11]

$$(7.14) \quad \int_{-\infty}^{\infty} e^{-2x^2} [H_n(x)]^2 dx = 2^{n-1/2} \Gamma(n+1/2).$$

[15, §11.4,20]

$$(7.15) \quad \int_{-\infty}^{\infty} e^{-(a^2+b^2)x^2} H_{2m}(ax) H_{2k}(bx) dx = (-)^{m+k} 2^{2(m+k)} \Gamma(m+k+\frac{1}{2}) \frac{a^{2k} b^{2m}}{(a^2+b^2)^{m+k+1/2}}.$$

[15, §11.4,21]

$$(7.16) \quad \int_{-\infty}^{\infty} e^{-(a^2+b^2)x^2} H_{2m+1}(ax) H_{2k+1}(bx) dx = (-)^{m+k} 2^{2(m+k+1)} \Gamma(m+k+\frac{3}{2}) \frac{a^{2k+1} b^{2m+1}}{(a^2+b^2)^{m+k+3/2}}.$$

[111]

$$(7.17) \quad \int_0^{\infty} x^{\nu+1} e^{-\alpha x^2} L_m^{\nu-\sigma}(\alpha x^2) L_n^{\sigma}(\alpha x^2) J_{\nu}(xy) dx = \frac{(-1)^{m+n}}{2\alpha} \left(\frac{y}{2\alpha}\right)^{\nu} \exp\left(-\frac{y^2}{4\alpha}\right) L_m^{\sigma-m+n}\left(\frac{y^2}{4\alpha}\right) L_n^{\nu-\sigma+m-n}\left(\frac{y^2}{4\alpha}\right),$$

for $y > 0$, $\Re \alpha > 0$, $\Re \nu > -1$.

(7.18)

$$\int_0^{\infty} x^{(\alpha+\beta)/2} e^{-x} L_m^{\alpha}(x) L_n^{\beta}(x) J_{\alpha+\beta}(2\sqrt{ax}) dx = (-1)^{m+n} a^{(\alpha+\beta)/2} L_m^{\beta-m+n}(a) L_n^{\alpha+m-n}(a); \quad \Re(\alpha+\beta) > -1.$$

7.3. Hypergeometric Functions. [73, p 238]

(7.19)

$$\frac{1}{\Gamma(2\lambda+2\nu)} \int_0^{\infty} dt e^{-pt} t^{2\lambda+2\nu-1} {}_1F_2(\nu; \lambda+\nu, \lambda+\nu+\frac{1}{2}; -\frac{1}{4}a^2t^2) = \frac{1}{p^{2\lambda}} \frac{1}{(p^2+a^2)^{\nu}}; \quad \Re(\lambda+\nu) > 0.$$

previous formula at $\lambda = 0$ with [1, 9.1.69] gives [91, 6.623.1]

(7.20)

$$\begin{aligned} \frac{1}{\Gamma(2\nu)} \int_0^{\infty} dt e^{-pt} t^{2\nu-1} {}_0F_1(; \nu+\frac{1}{2}; -\frac{1}{4}a^2t^2) &= \frac{\Gamma(\nu+1/2)}{\Gamma(2\nu)} \left(\frac{a}{2}\right)^{1/2-\nu} \int_0^{\infty} e^{-pt} t^{\nu-1/2} J_{\nu-1/2}(at) \\ &= \frac{1}{(p^2+a^2)^{\nu}}; \quad \Re(\nu) > 0. \end{aligned}$$

[133]

(7.21)

$$\int_0^{\infty} J_{\mu}^2(\omega\rho) {}_3F_2\left(\begin{matrix} 3/2, (\sigma+\nu)/2, (\sigma-\nu)/2 \\ 1-\mu, 1+\mu \end{matrix} \middle| -4\omega^2\right) \omega d\omega = -\frac{\mu\rho^{\sigma-2} K_{\nu}(\rho)}{2^{\sigma-1}\Gamma[(\sigma+\nu)/2]\Gamma[(\sigma-\nu)/2]}$$

for $\Re \sigma > 1 + |\Re \nu|$, $\mu = l + 1/2$ with $l \in \mathbb{N}$, $\Re(\sigma+\nu) > 0$, $\Re \rho > 0$.

[133]

(7.22)

$$\int_0^{\infty} J_{\mu}^2(\omega\rho) J_{\mu+1}(\omega\rho) {}_3F_2\left(\begin{matrix} 3/2, (\sigma+\nu)/2, (\sigma-\nu)/2 \\ 1-\mu, 2+\mu \end{matrix} \middle| -4\omega^2\right) \omega^2 d\omega = -\frac{\mu(\mu+1)\rho^{\sigma-3} K_{\nu}(\rho)}{2^{\sigma-1}\Gamma[(\sigma+\nu)/2]\Gamma[(\sigma-\nu)/2]}$$

for $\Re \sigma > 2 + |\Re \nu|$, $\mu = l + 1/2$ with $l \in \mathbb{N}$, $\Re(\sigma+\nu) > 0$, $\Re \rho > 0$.

8. SPECIAL FUNCTIONS

8.1. **The exponential integral and related functions.** [30] If

$$(8.1) \quad \int_0^x \frac{\sin u}{u} du = \frac{\pi}{2} - r \cos(x - \theta); \quad \int_0^x \frac{1 - \cos u}{u} du = \gamma + \log x - r \sin(x - \theta)$$

then

$$(8.2) \quad r \cos \theta \sim \sum_{k \geq 0} \frac{(-)^k (2k)!}{x^{2k+1}}; \quad r \sin \theta \sim \sum_{k \geq 1} \frac{(-)^{k+1} (2k-1)!}{x^{2k}}; \quad r \sim \sum_{k \geq 1} \frac{(-)^{k+1} (2k-1)!}{k x^{2k}}$$

for $x \rightarrow \infty$.

8.2. **The error function and Fresnel integrals.** [34]

$$(8.3) \quad 2\sqrt{\pi} e^{b^2} \operatorname{erf}(b) = \sum_{j=0}^{\infty} \frac{j! b^{2j+1}}{(2j+1)! 2^{2j+2}}.$$

8.3. **The gamma function.**

$$(8.4) \quad \frac{1}{a+r} = \frac{1}{a} \frac{(a)_r}{(a+1)_r}.$$

[174, 2.2.3.1]

$$(8.5) \quad (a)_{m-r} = \frac{(-1)^r (a)_m}{(1-a-m)_r}.$$

[174, (2.4.5.2.)][64, 166]

$$(8.6) \quad (a)_{2r} = (a/2)_r \left(\frac{a+1}{2}\right)_r 2^{2r}; \quad (a)_{qr} = (a/q)_r \left(\frac{a+1}{q}\right)_r \cdots \left(\frac{a+q-1}{q}\right)_r q^{qr}$$

$$(8.7) \quad \Gamma(n+1-k) = \frac{(-1)^k \Gamma(n+1)}{(-n)_k}.$$

[174, (I.4)]

$$(8.8) \quad (a+kn)_n = \frac{(a)_{(k+1)n}}{(a)_{kn}}.$$

[174, (I.6)]

$$(8.9) \quad (a-kn)_n = \frac{(-1)^n (1-a)_{kn}}{(1-a)_{(k-1)n}}.$$

[174, (I.9)][166]

$$(8.10) \quad (a)_{N-n} = \frac{(-1)^n (a)_N}{(1-a-N)_n}.$$

[174, (I.11)]

$$(8.11) \quad (a+kn)_{N-n} = \frac{(a)_N (a+N)_{(k-1)n}}{(a)_{kn}}.$$

[174, (I.13)]

$$(8.12) \quad (a-kn)_{N-n} = \frac{(-1)^n (a)_N (1-a)_{kn}}{(1-a-N)_{(k+1)n}}.$$

[70, (§1.2)]

$$(8.13) \quad \frac{\Gamma(n + \frac{1}{2} + z)\Gamma(n + \frac{1}{2} - z)}{\Gamma^2(n + \frac{1}{2})} = \frac{1}{\cos(\pi z)} \prod_{l=1}^n \left[1 - \frac{4z^2}{(2l-1)^2} \right],$$

for $n = 1, 2, 3, \dots$

[194][175, A073006]

$$(8.14) \quad \Gamma(2/3) = \frac{2\pi}{\sqrt{3}} \frac{1}{\Gamma(1/3)}.$$

[194][175, A068465]

$$(8.15) \quad \Gamma(3/4) = \pi\sqrt{2} \frac{1}{\Gamma(1/4)}.$$

[194][175, A175379]

$$(8.16) \quad \Gamma(1/6) = \frac{\sqrt{3}}{\sqrt{\pi}2^{1/3}} \Gamma^2(1/3).$$

[194]

$$(8.17) \quad \Gamma(3/5) = \frac{\pi\sqrt{2}\sqrt{\phi^*}}{\sqrt{5}} \frac{1}{\Gamma(2/5)},$$

where $\phi^* = 5 - \sqrt{5}$.

[194][175, A203145]

$$(8.18) \quad \Gamma(5/6) = \frac{\pi^{3/2}2^{4/3}}{\sqrt{3}} \frac{1}{\Gamma^2(1/3)}.$$

[194]

$$(8.19) \quad \Gamma(4/5) = \frac{\pi\sqrt{2}\sqrt{\phi}}{\sqrt{5}} \frac{1}{\Gamma(1/5)},$$

where $\phi = 5 + \sqrt{5}$.

[194][175, A203143]

$$(8.20) \quad \Gamma(3/8) = \sqrt{\pi}\sqrt{\sqrt{2}-1} \frac{\Gamma(1/8)}{\Gamma(1/4)}.$$

[194][175, A203144]

$$(8.21) \quad \Gamma(5/8) = \sqrt{\pi}2^{3/4} \frac{\Gamma(1/4)}{\Gamma(1/8)}.$$

[194][175, A203146]

$$(8.22) \quad \Gamma(7/8) = \pi^{3/4}\sqrt{\sqrt{2}+1} \frac{1}{\Gamma(1/8)}.$$

[194]

$$(8.23) \quad \Gamma(1/10) = \frac{\sqrt{\phi}}{\sqrt{\pi}2^{7/10}} \Gamma(1/5)\Gamma(2/5),$$

where $\phi = 5 + \sqrt{5}$.

[194]

$$(8.24) \quad \Gamma(3/10) = \frac{\sqrt{\pi}\phi^*}{2^{3/5}\sqrt{5}} \frac{\Gamma(1/5)}{\Gamma(2/5)},$$

where $\phi^* = 5 - \sqrt{5}$.

[194]

$$(8.25) \quad \Gamma(7/10) = \sqrt{\pi} 2^{3/5} \frac{\Gamma(2/5)}{\Gamma(1/5)}.$$

[194]

$$(8.26) \quad \Gamma(9/10) = \frac{\pi^{3/2} 2^{7/10} \sqrt{\phi}}{\sqrt{5}} \frac{1}{\Gamma(2/5)\Gamma(1/5)}.$$

[194]

$$(8.27) \quad \Gamma(1/12) = \frac{3^{3/8} \sqrt{\sqrt{3}+1}}{\sqrt{\pi} 2^{1/4}} \Gamma(1/3) \Gamma(1/4).$$

[194]

$$(8.28) \quad \Gamma(5/12) = \frac{\sqrt{\pi} 2^{1/4} \sqrt{\sqrt{3}-1}}{3^{1/8}} \frac{\Gamma(1/4)}{\Gamma(1/3)}.$$

[164]

$$(8.29) \quad (N + \frac{1}{2}, K)(N + \frac{1}{2}, L) = \sum_{j=\max(K,L)}^{\min(N,K+L)} \binom{2j-K-L}{j-K} (j + \frac{1}{2}, K+L-j)(N + \frac{1}{2}, j),$$

where $(n + \frac{1}{2}, j) = \frac{(n+j)!}{j!(n-j)!}$ is Hankel's symbol.

[164]

$$(8.30) \quad (-)^N \frac{(N + \frac{1}{2}, K)(N + \frac{1}{2}, L)}{\binom{K+L}{K}} = \sum_{j=\max(K,L)}^{\min(N,K+L)} (-)^j \binom{2j-K-L}{j-K} \frac{(j + 1/2, K+L-j)(N + 1/2, j)}{\binom{K+L}{j}},$$

where $(n + \frac{1}{2}, j) = \frac{(n+j)!}{j!(n-j)!}$ is Hankel's symbol.

[104]

$$(8.31) \quad \log \frac{\Gamma(z + 1/2)}{\sqrt{z}\Gamma(z)} = - \sum_{r=1}^k \frac{(1 - 2^{-2r})B_{2r}}{r(2r-1)z^{2r-1}} + O(z^{-2k+1/2}).$$

[104]

$$(8.32) \quad \log \frac{\Gamma(z + 3/4)}{\sqrt{z}\Gamma(z + 1/4)} = - \sum_{r=1}^k \frac{E_{2r}}{4r(4z)^{2r}} + O(z^{-2k+1/2}).$$

[104]

$$(8.33) \quad \frac{1}{z} \left[\frac{\Gamma(z + 3/4)}{\Gamma(z + 1/4)} \right]^2 = 1 + \frac{2u}{1+} \frac{9u}{1+} \frac{25u}{1+} \frac{49u}{1+} \dots$$

where $u = 1/(64z^2)$.

8.4. The psi function ψ . [6]

$$(8.34) \quad \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} [\psi(k+1) + \gamma] = G - \frac{\pi}{2} \log 2.$$

[6]

$$(8.35) \quad \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} [\psi(k+3/2) + \gamma] = G - \frac{\pi}{4} \log 2.$$

[6]

$$(8.36) \quad \frac{(-1)^n}{q^n(n-1)!} \sum_{k=1}^{q-1} e^{2\pi i k p/q} \psi^{n-1}(k/q) = \text{Li}_n(e^{2\pi i p/q}) - \frac{\zeta(n)}{q^n}.$$

[6]

$$(8.37) \quad \frac{1}{4} \sum_{k=0}^{\infty} \frac{2^k k!^2}{(2k+1)!} [\psi(k+3/2) + \gamma] = G - \frac{\pi}{4} \log 2.$$

[6]

$$(8.38) \quad q \sum_{k=0}^{q-1} \text{Li}_2(e^{2\pi i k/q} t) = \text{Li}_2(t^q).$$

8.5. Integral representations of the functions J_ν and N_ν . [185, La. 4.13]

$$(8.39) \quad J_{\mu+\nu+1}(t) = \frac{t^{\nu+1}}{2^\nu \Gamma(\nu+1)} \int_0^1 J_\mu(ts) s^{\mu+1} (1-s^2)^\nu ds.$$

[71, 7.2.7][118]

$$(8.40) \quad \Gamma(\nu+1)\Gamma(\mu+1)J_\nu(z)J_\mu(z) = (z/2)^{\nu+\mu} {}_2F_3\left(\frac{1+\nu+\mu}{2}, 1+\frac{\nu+\mu}{2}; 1+\nu, 1+\mu, 1+\nu+\mu; -z^2\right).$$

[24]

$$(8.41) \quad \begin{aligned} \frac{z}{2} J_\mu(z \cos \phi \cos \Phi) J_\nu(z \sin \phi \sin \Phi) &= (\cos \phi \cos \Phi)^\mu (\sin \phi \sin \Phi)^\nu \\ &\times \sum_{n=0}^{\infty} (-1)^n (\mu + \nu + 2n + 1) J_{\mu+\nu+2n+1}(z) \frac{\Gamma(\mu + \nu + n + 1) \Gamma(\nu + n + 1)}{n! \Gamma(\mu + n + 1) \Gamma^2(\nu + 1)} \\ &\times F(-n, \mu + \nu + n + 1; \nu + 1; \sin^2 \phi) F(-n, \mu + \nu + n + 1; \nu + 1; \sin^2 \Phi), \end{aligned}$$

where ν and μ are not negative integers.

[63]

$$(8.42) \quad J_\mu(Xz)J_\nu(xz) = \frac{X^\mu x^\nu}{\pi} \int_{-\pi/2}^{\pi/2} e^{i(\mu-\nu)\theta} (\lambda_1/\lambda_2)^{\mu+\nu} J_{\mu+\nu}(z\lambda_1\lambda_2) d\theta$$

where $\lambda_1 = +\sqrt{(e^{i\theta} + e^{-i\theta})}$, $\lambda_2 = (X^2 e^{i\theta} + x^2 e^{-i\theta})^{1/2}$.

[71, p 99]

$$(8.43) \quad (z/2)^{\gamma-\mu-\nu} J_\mu(\alpha z) J_\nu(\beta z) = \frac{\alpha^\mu \beta^\nu}{\Gamma(\mu+1)\Gamma(\nu+1)} \sum_{m=0}^{\infty} \frac{(\gamma+2m)\Gamma(\gamma+m)}{m!} \\ \times F_4(-m, \gamma+m; \mu+1, \nu+1; \alpha^2, \beta^2) J_{\gamma+2m}(z) \\ (8.44) \\ = \frac{\alpha^\mu \beta^\nu}{\Gamma(\nu+1)} \sum_{m=0}^{\infty} (\gamma+2m) J_{\gamma+2m}(z) \sum_{n=0}^{\infty} \frac{(-1)^n \Gamma(\gamma+m+n) \alpha^{2n}}{n!(m-n)!\Gamma^2(n+\mu+1)} {}_2F_1(-n, -n-\mu; \nu+1; \frac{\beta^2}{\alpha^2}).$$

Application of [81, (3.5)] to [1, (9.1.14)]

$$(8.45) \quad J_\rho(cz) J_\nu(z) J_\mu(z) = \frac{c^\rho (z/2)^{\nu+\mu+\rho}}{\Gamma(\rho+1)\Gamma(\nu+1)\Gamma(\mu+1)} \sum_{n \geq 0} \frac{(-c^2 z^2/4)^n}{(\rho+1)_n n!} \\ \times {}_4F_3 \left(-n, -n-\rho, 1 + \frac{\nu+\mu}{2}, \frac{1+\nu+\mu}{2}; \nu+1, \mu+1, \nu+\mu+1; \frac{4}{c^2} \right).$$

[31]

$$(8.46) \quad x^{-\alpha} I_\alpha(2\sqrt{x}) = \sum_{r=0}^{\infty} \frac{x^r}{r! \Gamma(r+\alpha+1)}.$$

[31][175, A002426]

$$(8.47) \quad \exp(t) I_0(2t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} c_n; \quad c_n \equiv \sum_{k=0}^{[n/2]} \frac{n!}{(k!)^2 (n-2k)!} = i^n \sqrt{3^n} P_n \left(-\frac{i}{\sqrt{3}} \right).$$

[31]

$$(8.48) \quad \exp(yt) \exp(xt^2)^{-\alpha/2} I_\alpha(2t\sqrt{x}) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \Pi_n^\alpha(x, y); \quad \Pi_n^\alpha(x, y) \equiv n! \sum_{k=0}^{[n/2]} \frac{x^k y^{n-2k}}{(n-2k)! k! \Gamma(k+\alpha+1)}.$$

$$(8.49) \quad \frac{\exp(t)}{t} I_1(t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \Pi_n^1(1, 1).$$

[146]

$$(8.50) \quad j_s(R) = \frac{R^s}{2^{s+1} s!} \int_0^\pi d\theta \sin \theta \cos(R \cos \theta) (\sin \theta)^{2s}.$$

[175, A122848]

$$(8.51) \quad \frac{d^s}{dt^s} t^\nu Z_\nu(t) = \sum_{l=\lfloor (s+1)/2 \rfloor}^s \alpha_{l,s} t^{\nu-(s-l)} Z_{\nu-l}(t).$$

$$(8.52) \quad \frac{d^s}{dt^s} t^\nu K_\nu(t) = (-)^s \sum_{l=\lfloor (s+1)/2 \rfloor}^s \alpha_{l,s} t^{\nu-(s-l)} K_{\nu-l}(t)$$

with Bessel polynomial coefficients [97]

$$\alpha_{l,s} = \frac{s!}{(s-l)!(2l-s)! 2^{s-l}}.$$

[164]

$$(8.53) \quad \sum_{k=1}^n \frac{2k-1}{\sqrt{\pi}} K_{k-1/2}(x) K_{k-1/2}(y) = \sum_{k=1}^n \frac{n(n-1+k)!}{k!(n-k)!} \left(\frac{x+y}{2xy} \right)^{k-1/2} K_{k-1/2}(x+y).$$

[164]

$$(8.54) \quad \frac{1}{\sqrt{\pi}} K_{n+1/2}(x) K_{n+1/2}(y) = \sum_{\mu=0}^n \frac{(n+\mu)!}{\mu!(n-\mu)!} \left(\frac{x+y}{2xy} \right)^{\mu+1/2} K_{k+1/2}(x+y).$$

[87]

$$(8.55) \quad \sum_{n=1}^{\infty} (-1)^n \frac{J_{2m}(n\pi)}{a^2 - n^2} = \frac{\pi J_{2m}(a\pi)}{2a \sin a\pi}.$$

[87]

$$(8.56) \quad \sum_{n=1}^{\infty} (-1)^n \frac{n J_{2m-1}(n\pi)}{a^2 - n^2} = \frac{\pi J_{2m-1}(a\pi)}{2 \sin a\pi}.$$

[168]

$$(8.57) \quad z^m [K_{\nu+m}(z) - K_{\nu-m}(z)] = \sum_{j=0}^{m-1} (-1)^{m-j-1} b_m(j) z^j K_{\nu+j}(z),$$

$$(8.58) \quad z^m [I_{\nu-m}(z) - I_{\nu+m}(z)] = \sum_{j=0}^{m-1} b_m(j) z^j I_{\nu+j}(z),$$

$$(8.59) \quad z^m [J_{\nu-m}(z) - (-1)^m I_{\nu-m}(z)] = \sum_{j=0}^{m-1} b_m(j) (-z)^j J_{\nu+j}(z),$$

with $m = 1, 2, 3, \dots$ and

$$b_m(j) = 2^{m-j} \binom{m}{j} \nu(\nu-1) \cdots (\nu+j-m+1) = \Gamma(\nu+1) \frac{2^{m-j} \binom{m}{j}}{\Gamma(\nu+j-m+1)}.$$

[71, §7.10.1]

$$(8.60) \quad f(z) = \frac{1}{z^\nu} \sum_{n=0}^{\infty} a_n J_{\nu+n}(z); \quad a_n = (\nu+n) 2^{\nu+n} \sum_{s=0}^{\lfloor n/2 \rfloor} \frac{\Gamma(\nu+n-s)}{s! 2^{2s}} b_{n-2s},$$

where $f(z) = \sum_{n=0}^{\infty} b_n z^n$.

[71, §7.10.1]

$$(8.61) \quad f(z) = \frac{1}{z^\nu} \sum_{n=0}^{\infty} a_n z^n J_{\nu+n}(z); \quad a_n = \sum_{s=0}^n \frac{\Gamma(\nu+s+1)}{(n-s)!} 2^{2s-n+\nu} b_s,$$

$$(8.62) \quad \Gamma(\nu+n+1) b_n = \sum_{s=0}^{\infty} (-1)^s 2^{-\nu-n-s} \frac{a_{n-s}}{s!}.$$

where $f(z) = \sum_{l=0}^{\infty} b_l z^{2l}$.

$$(8.63) \quad [z^{2n}] \frac{1}{z^{2\nu}} \sum_{r=0}^{\infty} A_r J_{\nu+r}^2(z) = \frac{(-)^n \Gamma(\nu+n+1/2)}{n! \sqrt{\pi} \Gamma(2\nu+n+1) \Gamma(\nu+n+1)} \sum_{r=0}^n \frac{A_r (-n)_r}{(2\nu+n+1)_r}.$$

$$(8.64) \quad [z^{2\nu}] \frac{1}{z^{2\nu}} \sum_{r=0}^{\infty} \frac{(2\nu)_r (\nu+1)_r (a_3)_r (a_4)_r}{r! (\nu)_r (1+2\nu-a_3)_r (1+2\nu-a_4)_r} J_{\nu+r}^2(z) = \frac{1}{4\nu \Gamma^2(1+\nu)} {}_2F_3 \left(\begin{matrix} \nu+1/2, 1+2\nu-a_3-a_4 \\ \nu+1, 1+2\nu-a_3, 1+2\nu-a_4 \end{matrix} \mid -z^2 \right)$$

$$(8.65) \quad z^\nu J_\nu(2z) = \frac{2\sqrt{\pi}}{\Gamma(\nu+1/2)} \sum_{r=0}^{\infty} (-)^r (\nu+r) \frac{\Gamma(2\nu+r)}{r!} J_{\nu+r}^2(z).$$

$$(8.66) \quad z^{-1/2} J_{2\nu+1/2}(2z) = -\frac{\Gamma(\nu+1)}{\Gamma(\nu+1/2)} \sum_{r=0}^{\infty} (\nu+r) \frac{\Gamma(2\nu+r) \Gamma(r-1/2)}{r! \Gamma(2\nu+r+3/2)} J_{\nu+r}^2(z).$$

$$(8.67) \quad [z^{2n}] \frac{1}{z^{2\nu}} \sum_{r=0}^{\infty} A_r J_{\nu+2r}^2(z) = \frac{(-)^n \Gamma(\nu+n+1/2)}{n! \sqrt{\pi} \Gamma(2\nu+n+1) \Gamma(\nu+n+1)} \sum_{r=0}^n \frac{A_r (-n)_{2r}}{(2\nu+n+1)_{2r}}.$$

$$(8.68) \quad [z^{2\nu}] \frac{1}{z^{2\nu}} \sum_{r=0}^{\infty} \frac{(\nu)_r (1+\nu/2)_r (a_3)_r}{r! (\nu/2)_r (1+\nu-a_3)_r} J_{\nu+2r}^2(z) = \frac{1}{4\nu \Gamma^2(1+\nu)} {}_1F_2 \left(\begin{matrix} 1/2+\nu-a_3 \\ \nu+1, 1+2\nu-2a_3 \end{matrix} \mid -z^2 \right).$$

$$(8.69) \quad \frac{J_\lambda(2z \sin \frac{1}{2}\phi)}{(2z \sin \frac{1}{2}\phi)^{\lambda-2\nu}} = \sum_{n=0}^{\infty} \frac{2^{1-\lambda+2\nu} \sqrt{\pi} \Gamma(\nu+1)}{n! \Gamma(\nu+1/2) \Gamma(\lambda+1)} \sin^{2\nu} \frac{1}{2}\phi (\nu+n) \Gamma(2\nu+n) J_{\nu+n}^2(z) {}_3F_2 \left(\begin{matrix} \nu+1, 2\nu+n, -n \\ \nu+1/2, \lambda+ \end{matrix} \mid \sin^2 \frac{1}{2}\phi \right)$$

$$(8.70) \quad J_0(z \cos a) + J_0(z \sin a) \equiv 2 \sum_{n=0}^{\infty} b_{2n}(a) J_{2n}(z);$$

where

$2n$	b_{2n}
0	1
2	1
4	$1 + 6 \cos^4 a - 6 \cos^2 a = \frac{1}{4} + \frac{3}{4} \cos(4a)$
6	$1 + 6 \cos^4 a - 6 \cos^2 a = \frac{1}{4} + \frac{3}{4} \cos(4a)$
8	$1 + 70 \cos^8 a - 20 \cos^2 a - 140 \cos^6 a + 90 \cos^4 a = \frac{9}{64} + \frac{35}{64} \cos(8a) + \frac{5}{16} \cos(4a)$
10	$1 + 70 \cos^8 a - 20 \cos^2 a - 140 \cos^6 a + 90 \cos^4 a = \frac{9}{64} + \frac{35}{64} \cos(8a) + \frac{5}{16} \cos(4a)$
12	$1 - 42 \cos^2 a + 420 \cos^4 a - 1680 \cos^6 a + 3150 \cos^8 a - 2772 \cos^{10} a + 924 \cos^{12} a$ $= \frac{25}{256} + \frac{104}{512} \cos(4a) + \frac{63}{256} \cos(8a) + \frac{231}{512} \cos(12a)$

Apparently, the b_{2n} count numbers of Delannoy paths, A109983 in [175].

[122, p31]

$$(8.71) \quad e^{ik \cos \varphi} = 2^\nu \Gamma(\nu) \sum_{m=0}^{\infty} (\nu + m) i^m J_{\nu+m}(k\rho) (k\rho)^{-\nu} C_m^{(\nu)}(\cos \varphi),$$

$\nu \neq 0, -1, -2, -3, \dots$

[82]

(8.72)

$$S_{\mu,\nu}(z) \sim z^{\mu-1} \sum_{m=0}^{\infty} (-1)^m \left(\frac{1-\mu+\nu}{2} \right)_m \left(\frac{1-\mu-\nu}{2} \right)_m (z/2)^{-2m}, \quad |z| \rightarrow \infty, |\arg z| < \pi.$$

The series terminates and is equal to $S_{\mu,\nu}(z)$ when $\mu \pm \nu$ is a positive odd integer.

[158]

(8.73)

$$\left(\frac{\sin \beta}{\sin \alpha} \right)^{m+n} P_{m+n}^m(\cos \alpha) = \sum_{r=0}^n \binom{2m+n}{r} \left(\frac{\sin(\beta-\alpha)}{\sin \alpha} \right)^r P_{n+m-r}^m(\cos \beta).$$

8.6. Orthogonal Polynomials. [197]

(8.74)

$$L_m^{(\alpha)} L_n^{(\alpha)} = \sum_{M=|m-n|}^{m+n} \frac{L_M^{(\alpha)}(z)(-)^{m+n-M} 2^{m+n-M} M!}{(M-m)!(M-n)!(m+n-M)!} {}_3F_2 \left(\begin{matrix} \alpha + M + 1, \frac{M-m-n}{2}, \frac{M-m-n+1}{2} \\ M-m+1, M-n+1 \end{matrix} \mid 1 \right).$$

[54]

$$(8.75) \quad \sum_{n \geq 0} t^n L_n^{a+bn}(x) = \frac{(1+v)^{a+1}}{1-bv} \exp(-xv),$$

where $v = t(1+v)^{b+1}$, $v(0) = 0$.

[54]

$$(8.76) \quad \sum_{n \geq 0} t^n L_n^{v+bn}(x(1+an)) = \frac{(1-z)^{1-v}}{1-z(b+2-ax) + z^2(b+1)} e^{xz/(z-1)},$$

where $t = z(1-z)^b \exp[axz/(1-z)]$ and $|t| < 1$.

[54]

(8.77)

$$\sum_{n \geq 0} \frac{t^n}{v+bn+n} L_n^{v+bn}(x(1+an)) = \frac{\exp[xz/(z-1)]}{v(1-z)^v} {}_1F_1 \left(\begin{matrix} 1 \\ \frac{v+1+b}{1+b} \end{matrix} \mid \frac{xz(1+b-av)}{(1-z)(1+b)} \right),$$

where $t = z(1-z)^b \exp[axz/(1-z)]$ and $|t| < 1$.

[54]

$$(8.78) \quad \sum_{n \geq 0} \frac{t^n}{1+an} L_n^{v+avn-n}(x(1+an)) = \frac{1}{(1-z)^v} e^{xz/(z-1)},$$

where $t = z(1-z)^{av-1} \exp[axz/(1-z)]$ and $|t| < 1$.

[54]

$$(8.79) \quad \sum_{n \geq 0} \frac{t^n}{1+an} L_n^{v+bn}(x(1+an)) = \frac{\exp[xz/(z-1)]}{(1-z)^v} {}_2F_1 \left(\begin{matrix} 1, \frac{1+b-av}{a} \\ \frac{a+1}{a} \end{matrix} \mid z \right),$$

where $t = z(1-z)^b \exp[axz/(1-z)]$ and $|t| < 1$.

[54]

$$(8.80) \quad \sum_{n \geq 0} \frac{t^n}{1+n} L_n^{v+bn}(x(1+an)) = \frac{\exp[x(1-a-z)/(1-z)]}{(1-z)^v z(v-b)} \left\{ {}_1F_1 \left(\begin{matrix} v-b \\ v-b+1 \end{matrix} \mid \frac{x(a-1)}{1-z} \right) \right. \\ \left. - (1-z)^{v-b} {}_1F_1 \left(\begin{matrix} v-b \\ v-b+1 \end{matrix} \mid x(a-1) \right) \right\},$$

where $t = z(1-z)^b \exp[axz/(1-z)]$ and $|t| < 1$.

[54]

$$(8.81) \quad \sum_{n \geq 0} \frac{(1+bn)^{n/2}}{n!} t^n H_n[x(1+an)/(1+bn)^{1/2}] = \frac{e^{-z^2-2xz}}{1+2bz^2+2axz},$$

where $t = (-z)e^{bz^2+2axz}$ and $|2axz \exp[bz^2+2axz+1]| < 1$.

[54]

$$(8.82) \quad \sum_{n \geq 0} \frac{(1+bn)^{n/2-1}}{n!} t^n H_n[x(1+an)/(1+bn)^{1/2}] = e^{-z^2-2xz} {}_1F_1 \left(\begin{matrix} 1 \\ 1+1/b \end{matrix} \mid 2xz(b-a)/b \right)$$

where $t = (-z)e^{bz^2+2axz}$ and $|2axz \exp[bz^2+2axz+1]| < 1$.

[54]

$$(8.83) \quad \sum_{n \geq 0} \frac{(1+an)^{n/2-1}}{n!} t^n H_n[x(1+an)] = e^{-z^2-2xz}$$

where $t = (-z)e^{az^2+2axz}$ and $|2axz \exp[az^2+2axz+1]| < 1$.

[83]

$$(8.84) \quad \int_0^1 f(t) t^r dt = c_r, \quad (r = 0, 1, \dots, n) \rightsquigarrow f(t) = \sum_{i=0}^n (2i+1) \sum_{r=0}^i \left\{ (-)^r \frac{(i+r)!}{(i-r)!} \frac{1}{(r!)^2} c_r \right\} P_i(1-2t).$$

9. SPECIAL FUNCTIONS II.

9.1. Hypergeometric Functions. [29]

$$(9.1) \quad {}_2F_1(a, b; c; z) = (a)_m (b)_m z^m {}_2F_1(a+m, b+m; 1+m; z).$$

[29]

$$(9.2) \quad {}_2F_1(a, b; c; z) = (1-z)^{c-a-b} {}_2F_1(c-a, c-b; c; z).$$

[29]

$$(9.3) \quad {}_2F_1(a, b; c; z) = (a)_m (b)_m z^m (1-z)^{1-a-b-m} {}_2F_1(1-b, 1-a; 1+m; z).$$

[29]

$$(9.4) \quad {}_2F_1(a, b; c; z) = (1-z)^{-a} {}_2F_1(a, c-b; c; z/(z-1)).$$

[145]

$$(9.5) \quad \frac{(z/2)^c}{\Gamma(1+c)} {}_2F_1(a, b; c+1; -z^2/(4ab)) = \sum_{\nu \geq 0} \frac{1}{\nu!} {}_3F_0(-\nu, a, b; 1/(ab))(z/2)^\nu J_{c+\nu}(z).$$

[29]

$$(9.6) \quad {}_2F_1(a, b; c; z) = (1-z)^{-a} \left(\frac{z}{z-1} \right)^{1-c} \left\{ \frac{\Gamma(|m|)}{\Gamma(a+\bar{m})\Gamma(c-a-\underline{m})} \left(\frac{1}{1-z} \right)^{\underline{m}} \right. \\ \sum_{n=0}^{|m|-1} \frac{(1-a-\bar{m})_n (1-c+a+\underline{m})_n}{(1-|m|)_n \Gamma(1+n)} \left(\frac{1}{1-z} \right)^n - \frac{(-)^m}{\Gamma(a+\underline{m})\Gamma(c-a-\bar{m})} \left(\frac{1}{1-z} \right)^{\bar{m}} \\ \left. \sum_{n=0}^{\infty} \frac{(1-a-\underline{m})_n (1-c+a+\bar{m})_n}{\Gamma(1+|m|+n)\Gamma(1+n)} \left(\frac{1}{1-z} \right)^n \right. \\ \left. [-\ln(1-z) - \pi \cot \pi(c-a) - \pi \cot \pi a + \psi(1-a-\underline{m}+n) + \psi(1-c+a+\bar{m}+n) - \psi(1+|m|+n) - \psi(1+n)] \right\},$$

where $\bar{m} \equiv \max(0, m)$, $\underline{m} \equiv \min(0, m)$.

[163]

(9.7)

$${}_1F_1(1/2; 9/2; z) = -\frac{525 + 280z + 140z^2}{128z^3} e^z + \frac{525 + 630z + 420z^2 + 280z^3}{256z^{7/2}} \sqrt{\pi} \operatorname{erfi}(\sqrt{z}).$$

[173]

(9.8)

$${}_1F_1\left(\frac{1}{2} + \frac{1}{2}a - b, 1 + a - b; x\right) = e^{x/2} \sum_{r=0}^{\infty} \frac{(a)_r (b)_r (-x/4)^r}{r! (a/2)_r (1+a-b)_r} {}_0F_1\left(\frac{1}{2}a + r + 1; (x/4)^2\right).$$

[174, 1.7.7]

(9.9)

$${}_2F_1(a, -m; c, 1) = \frac{(c-a)_m}{(c)_m}.$$

[47]

(9.10)

$${}_2F_1(a, -n; c; p) = \frac{(a)_n}{(c)_n} (-p)^n {}_2F_1(1-c-n, -n; 1-a-n; 1/p).$$

[193]

(9.11)

$${}_2F_1(a+n, b; a-b; -1) = P(n) \frac{\Gamma(a-b)\Gamma(\frac{a+1}{2})}{\Gamma(a)\Gamma(\frac{a+1}{2}-b)} + Q(n) \frac{\Gamma(a-b)\Gamma(\frac{a}{2})}{\Gamma(a)\Gamma(\frac{a}{2}-b)}$$

where

$$P(n) = \frac{1}{2^{n+1}} {}_3F_2(-n/2, -(n+1)/2, a/2-b; 1/2, a/2; 1); \\ Q(n) = \frac{n+1}{2^{n+1}} {}_3F_2(-(n-1)/2, -n/2, (a+1)/2-b; 3/2, (a+1)/2; 1).$$

[13]

(9.12)

$${}_2F_1(-2n, b; -2n+2r-b; -1) = \frac{(1/2)_n (b+1-r)_n}{(b/2+1-r)_n (b/2+1/2-r)_n} \sum_{i=0}^{r-1} \frac{2^{2i} i! \binom{r+i-1}{2i}}{(b-r+1)_i} \binom{n}{i}.$$

[193]

(9.13)

$${}_2F_1(-a, 1/2; 2a+3/2+n; 1/4) = \frac{2^{n+3/2}}{3^{n+1}} \frac{\Gamma(a+5/4+n/2)\Gamma(a+3/4+n/2)\Gamma(a+1/2)}{\Gamma(a+7/6+n/3)\Gamma(a+5/6+n/3)\Gamma(a+1/2+n/3)} K(n) \\ - (-3)^{n-2} 2^{3/2} \frac{\Gamma(a+5/4+n/2)\Gamma(a+3/4+n/2)\Gamma(a+1)}{\Gamma(a+3/2)\Gamma(a+1/2+n/2)\Gamma(a+1+n/2)} L(n)$$

where $K(n)$ and $L(n)$ are defined in the reference.

[30]

(9.14)

$$\pi {}_2F_1(1/2, 1/2; 1; 1-x) = \log \frac{16}{x} {}_2F_1(1/2, 1/2; 1; x) - 4 \sum_{k \geq 1} \frac{(1/2)_k^2}{(k!)^2} \sum_{j=1}^k \frac{x^k}{(2j-1)(2j)}.$$

[30]

(9.15)

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; \frac{1+x}{2}\right) = \frac{\sqrt{\pi}}{\Gamma^2(3/4)} {}_2F_1(1/4, 1/4; 1/2; x^2) + \frac{\Gamma^2(3/4)}{\pi^{3/2}} {}_2F_1(3/4, 3/4; 3/2; x^2).$$

[30]

(9.16)

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; \frac{1}{2} + \frac{x}{1+x^2}\right) = \frac{\sqrt{\pi}}{\Gamma^2(3/4)} \sqrt{1+x^2} {}_2F_1(1/4, 1/2; 3/4; x^4) + \frac{\Gamma^2(3/4)}{\pi^{3/2}} x(1+x^2)^{3/2} {}_2F_1(1/2, 3/4; 5/4; x^4).$$

[30]

(9.17)

$${}_2F_1(n, -n; 1/2; x^2) = \cos(2n \sin^{-1} x).$$

[30]

(9.18)

$$2nx {}_2F_1\left(\frac{1}{2} + n, \frac{1}{2} - n; 3/2; x^2\right) = \sin(2n \sin^{-1} x).$$

[30]

(9.19)

$${}_2F_1\left(\frac{1}{2} + n, \frac{1}{2} - n; 1/2; x^2\right) = (1-x^2)^{-1/2} \cos(2n \sin^{-1} x).$$

[163]

(9.20)

$${}_2F_1(-3/2, -1/2; 1/2, z) = \frac{2+z}{2} \sqrt{1-z} + \frac{3\sqrt{z}}{2} \arcsin \sqrt{z}.$$

[163]

(9.21)

$$\frac{z^{\mu+1}}{2^\mu \sqrt{\pi} \Gamma(\frac{3}{2} + \mu)} {}_2F_1(1, 3/2; 3/2 + \mu, z^2/4) = L_\mu(z).$$

[163]

(9.22)

$${}_2F_1(-3/2, 1/2; 3/2, z) = \frac{5-2z}{8} \sqrt{1-z} + \frac{3}{8\sqrt{z}} \arcsin \sqrt{z}.$$

[163]

(9.23)

$${}_2F_1(-3/2, -1/2; 3/2, z) = \frac{13+2z}{16} \sqrt{1-z} + \frac{3+12z}{16\sqrt{z}} \arcsin \sqrt{z}.$$

[163]
 (9.24) ${}_1F_2(-3/2; -1/2, 1/2; z) = (1 + 2z) \cosh(2\sqrt{z}) + \sqrt{z} \sinh(2\sqrt{z}) - 4z^{3/2} \text{Shi}(2\sqrt{z}).$
 [163]

(9.25) ${}_1F_2(-3/2; -1/2, 2; z) = -\frac{4 + 24z - 28z^2}{15z\pi} K(\sqrt{z}) + \frac{4 + 56z + 4z^2}{15z\pi} E(\sqrt{z}).$
 [195]

(9.26) ${}_2F_1\left(1/4, -1/12; 2/3; \frac{x(4+x)^3}{4(2x-1)^3}\right) = (1-2x)^{-1/4}..$
 [195]

(9.27) ${}_2F_1\left(5/4, -1/12; 5/3; \frac{x(4+x)^3}{4(2x-1)^3}\right) = \frac{1+x}{(1+\frac{1}{4}x)^2} (1-2x)^{-1/4}.$
 [195]

(9.28) ${}_2F_1\left(1/4, 7/12; 4/3; \frac{x(4+x)^3}{4(2x-1)^3}\right) = \frac{1}{1+\frac{1}{4}x} (1-2x)^{3/4}.$
 [195]

(9.29) ${}_2F_1\left(1/4, -5/12; 1/3; \frac{x(4+x)^3}{4(2x-1)^3}\right) = (1+\frac{5}{2}x)(1-2x)^{-5/4}.$
 [195]

(9.30) ${}_2F_1\left(1/2, -1/6; 2/3; \frac{x(2+x)^3}{(2x+1)^3}\right) = (1+2x)^{-1/2}.$
 [195]

(9.31) ${}_2F_1\left(1/2, 5/6; 2/3; \frac{x(2+x)^3}{(2x+1)^3}\right) = \frac{1}{(1-x)^2} (1+2x)^{3/2}.$
 [195]

(9.32) ${}_2F_1\left(1/6, 5/6; 4/3; \frac{x(2+x)^3}{(2x+1)^3}\right) = \frac{1}{1+\frac{1}{2}x} (1+2x)^{1/2} (1+x)^{1/3}.$
 [195]

(9.33) ${}_2F_1\left(1/6, -1/6; 1/3; \frac{x(2+x)^3}{(2x+1)^3}\right) = (1+2x)^{-1/2} (1+x)^{1/3}.$
 [195]

(9.34) ${}_2F_1\left(7/24, -1/24; 3/4; \frac{108x(x-1)^4}{(x^2+14x+1)^3}\right) = (1+14x+x^2)^{-1/8}.$
 [195]

(9.35) ${}_2F_1\left(7/24, 23/24; 7/4; \frac{108x(x-1)^4}{(x^2+14x+1)^3}\right) = \frac{1+2x-\frac{1}{11}x^2}{(1-x)^2} (1+14x+x^2)^{7/8}.$
 [195]

(9.36) ${}_2F_1\left(5/24, 13/24; 5/4; \frac{108x(x-1)^4}{(x^2+14x+1)^3}\right) = \frac{1}{1-x} (1+14x+x^2)^{5/8}.$

[195]

$$(9.37) \quad {}_2F_1\left(5/24, -11/24; 1/4; \frac{108x(x-1)^4}{(x^2+14x+1)^3}\right) = \frac{1-22x-11x^2}{(1+14x+x^2)^{11/8}}.$$

[195]

$$(9.38) \quad {}_2F_1(19/60, -1/60; 4/5; \varphi_1(x)) = (1-228x+494x^2+228x^3+x^4)^{-1/20}.$$

(9.39)

$${}_2F_1(19/60, 59/60; 4/5; \varphi_1(x)) = \frac{(1+66x-11x^2)(1-228x+494x^2+228x^3+x^4)^{19/20}}{(1+x^2)(1+522x-10006x^2-522x^3+x^4)}.$$

$$(9.40) \quad {}_2F_1(11/60, 31/60; 6/5; \varphi_1(x)) = \frac{(1-228x+494x^2+228x^3+x^4)^{11/20}}{1+11x-x^2}.$$

$$(9.41) \quad {}_2F_1(11/60, -29/60; 1/5; \varphi_1(x)) = \frac{1+435x-6670x^2-3335x^4-87x^5}{(1-228x+494x^2+228x^3+x^4)^{29/20}}.$$

$$(9.42) \quad {}_2F_1(13/60, -7/60; 3/5; \varphi_1(x)) = \frac{1-7x}{(1-228x+494x^2+228x^3+x^4)^{7/20}}.$$

(9.43)

$${}_2F_1(13/60, 53/60; 3/5; \varphi_1(x)) = \frac{(1+119x+187x^2+17x^3)(1-228x+494x^2+228x^3+x^4)^{13/20}}{(1+x^2)(1+522x-10006x^2-522x^3+x^4)}.$$

(9.44)

$${}_2F_1(17/60, 37/60; 7/5; \varphi_1(x)) = \frac{(1+\frac{1}{7}x)(1-228x+494x^2+228x^3+x^4)^{17/20}}{(1+11x-x^2)^2}.$$

$$(9.45) \quad {}_2F_1(17/60, -23/60; 2/5; \varphi_1(x)) = \frac{(1+107x-391x^2+1173x^3+46x^4)}{(1-228x+494x^2+228x^3+x^4)^{23/20}}.$$

Where

$$(9.46) \quad \varphi_1(x) = \frac{1728x(x^2-11x-1)^5}{(x^4+228x^3+494x^2-228x+1)^3}.$$

[195]

$$(9.47) \quad {}_2F_1(7/20, -1/20; 4/5; \varphi_2(x)) = \frac{(1+x)^{7/20}}{(1-x)^{1/20}(1-4x-x^2)^{1/4}}.$$

(9.48)

$${}_2F_1(7/20, 19/20; 4/5; \varphi_2(x)) = \frac{(1+3x)(1+x)^{7/20}(1-x)^{19/20}(1-4x-x^2)^{7/4}}{(1+x^2)(1+22x-6x^2-22x^3+x^4)}.$$

Where

$$(9.49) \quad \varphi_2(x) = \frac{64x(x^2-x-1)^5}{(x^2-1)(x^2+4x-1)^5}.$$

[151, 51]

$$(9.50) \quad {}_2F_2(a, d; b, c; x) = e^x \sum_{n \geq 0} \frac{(c-d)_n}{(c)_n n!} (-x)^n {}_2F_2(b-a, d; b, c+n; -x).$$

[140]

$$(9.51) \quad {}_{r+1}F_{r+1}(a, (f_r+1); b, (f_r); y) = e^y {}_{r+1}F_{r+1}(b-a-r, (\xi_r+1); b, (\xi_r); -y),$$

where ξ_r are nonvanishing zeros of an associated parametric polynomial of Q degree r ,

$$(9.52) \quad Q_r(t) = \sum_{j=0}^r s_{r-j} \sum_{l=0}^j \left\{ \begin{matrix} j \\ l \end{matrix} \right\} (a)_l (t)_l (b-a-r-t)_{r-l},$$

and the s_{r-j} are determined by

$$(9.53) \quad (f_1 + x) \cdots (f + r + x) = \sum_{j=0}^r s_{r-j} x^j.$$

[11]

$$(9.54) \quad {}_3F_2(1, 1, 2-t; 2, 3; 1) = \frac{2(1-\gamma-\psi(t+1))}{1-t}.$$

[166]

$$(9.55) \quad {}_3F_2(-n, -a, -b; c, 2-n-a-b-c; 1) = \frac{(c+b-1)_n (c+a)_n}{(c+a+b-1)_n (c)_n} \left[1 - \frac{a}{(c+b-1)(a+c+n-1)} \right].$$

for $n = 1, 2, 3, \dots$

[113][174, (2.3.1.3)]

$$(9.56) \quad {}_3F_2(-m, b, c; e, -m+b+c-e+1; 1) = \frac{(e-b)_m (e-c)_m}{(e)_m (e-b-c)_m}; \quad m = 0, 1, 2, \dots, \quad e, e-b-c \neq 0, -1, -2, \dots$$

[64]

$$(9.57) \quad {}_3F_2(a, b/2, (b+1)/2; c/2, (c+1)/2; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-b)\Gamma(c-a)} {}_2F_1(a, b; c-a; -1), \quad \Re c > \Re b > 0, \Re(c-a-b) > 0.$$

[64]

$$(9.58) \quad {}_3F_2(a, b/2, (b+1)/2; c/2, (c+1)/2; 1/2) = 2^a \sum_{k=0}^{\infty} \binom{-a}{k} \frac{(c-b)_k}{(c)_k} {}_2F_1(-k, b; c+k; -1), \quad \Re c > \Re b > 0.$$

[47]

$$(9.59) \quad {}_0F_1(a; px) {}_0F_1(c'; qx) = \sum_{n \geq 0} \frac{(px)^n}{n!(c)_n} {}_2F_1(1-c-n, -n; c'; q/p).$$

[47]

$$(9.60) \quad {}_1F_1(a; c; px) {}_1F_1(a'; c'; qx) = \sum_{n \geq 0} \frac{(a)_n (px)^n}{n!(c)_n} {}_3F_2(a', 1-c-n, -n; c', 1-a-n; -q/p).$$

[47]

$$(9.61) \quad {}_2F_0(a, b; px) {}_2F_0(a', b'; qx) = \sum_{n \geq 0} \frac{(a)_n (b)_n (px)^n}{n!} {}_3F_2(a', b', -n; 1-a-n, 1-b-n; -q/p).$$

[47]
(9.62)

$${}_2F_1(a, b; c; px) {}_2F_1(a', b'; c'; qx) = \sum_{n \geq 0} \frac{(a)_n (b)_n (px)^n}{n! (c)_n} {}_4F_3(a', b', 1-c-n, -n; c', 1-a-n, 1-b-n; q/p).$$

[202]
(9.63)

$$(a+b+1) {}_3F_2[-c, -a, 1; b+1, \frac{1-a-c}{2}; \frac{1}{2}] = (b+1) {}_3F_2[-\frac{a}{2}, \frac{1-a}{2}, 1; \frac{1-a-b}{2}, \frac{1-a-c}{2}; 1]$$

where a , b and c are positive integers of the same parity.

[64]
(9.64)

$${}_3F_2(-n, b/q, (b+1)/2; c/2, (c+1)/2; 1) = \frac{(c-b)_n}{(c)_n} {}_2F_1(-n, b; c+n; -1), \quad \Re c > \Re b > 0.$$

[113]

$$\begin{aligned} & {}_3F_2(a, b, c; e, a+b+c-e+1; 1) + \frac{\Gamma(e-1)\Gamma(a-e+1)\Gamma(b-e+1)\Gamma(c-e+1)\Gamma(a+b+c-e+1)}{\Gamma(1-e)\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(a+b+c-2e+2)} \\ & \quad \times {}_3F_2(a-e+1, b-e+1, c-e+1; 2-e, a+b+c-2e+2; 1) \\ & = \frac{\Gamma(a-e+1)\Gamma(b-e+1)\Gamma(c-e+1)\Gamma(a+b+c-e+1)}{\Gamma(1-e)\Gamma(b+c-e+1)\Gamma(a+c-e+1)\Gamma(a+b-e+1)}. \end{aligned}$$

[113]
(9.65)

$${}_3F_2(a, b, c; e, a+b+c-e+1; 1) = \frac{\Gamma(e)\Gamma(a+b+c-e+1)}{\Gamma(a)\Gamma(b+1)\Gamma(c+1)} {}_3F_2(e-a, b+c-e+1, 1; b+1, c+1; 1); \quad \Re a > 0.$$

[160][139]

$${}_3F_2(a, b, c; f, e; 1)$$

(9.66)

$$= \frac{\Gamma(f)\Gamma(e)\Gamma(f+e-a-b-c)}{\Gamma(e-b+f-c)\Gamma(f+e-a-c)} {}_3F_2(e+f-a-b-c, f-c, e-c; e-b+f-c, e+f-a-c; 1)$$

(9.67)

$$= \frac{\Gamma(f)\Gamma(e)\Gamma(f+e-a-b-c)}{\Gamma(e-b+f-c)\Gamma(e-b+f-a)} {}_3F_2(e+f-a-b-c, f-b, e-b; e-b+f-c, e-b+f-a; 1)$$

(9.68)

$$= \frac{\Gamma(e)\Gamma(f+e-a-b-c)}{\Gamma(e-a)\Gamma(e-b+f-c)} {}_3F_2(f-c, f-b, a; e-b+f-c, f; 1)$$

(9.69)

$$= \frac{\Gamma(f)\Gamma(f+e-a-b-c)}{\Gamma(f-a)\Gamma(e-b+f-c)} {}_3F_2(e-c, e-b, a; e-b+f-c, e; 1)$$

(9.70)

$$= \frac{\Gamma(e)\Gamma(f)\Gamma(f+e-a-b-c)}{\Gamma(e+f-a-c)\Gamma(e-b+f-a)} {}_3F_2(e+f-a-b-c, f-a, e-a; e+f-a-c, e-b+f-a; 1)$$

(9.71)

$$= \frac{\Gamma(e)\Gamma(f+e-a-b-c)}{\Gamma(e-b)\Gamma(e+f-a-c)} {}_3F_2(f-c, f-a, b; e+f-a-c, f; 1)$$

(9.72)

$$= \frac{\Gamma(f)\Gamma(f+e-a-b-c)}{\Gamma(f-b)\Gamma(e+f-a-c)} {}_3F_2(e-c, e-a, b; e+f-a-c, e; 1)$$

(9.73)

$$= \frac{\Gamma(e)\Gamma(f+e-a-b-c)}{\Gamma(e-c)\Gamma(e-b+f-a)} {}_3F_2(f-b, f-a, c; e+f-a-b, f; 1)$$

(9.74)

$$= \frac{\Gamma(f)\Gamma(f+e-a-b-c)}{\Gamma(f-c)\Gamma(e-b+f-a)} {}_3F_2(e-b, e-a, c; e+f-a-b, e; 1).$$

[157, (6)]

(9.75)

$${}_3F_2(-n, \alpha, \beta; \gamma, \delta; 1) = \frac{\Gamma(\gamma)\Gamma(\gamma+n-\alpha)}{\Gamma(\gamma+n)\Gamma(\gamma-\alpha)} {}_3F_2(-n, \alpha, \delta-\beta; 1+\alpha-\gamma-n, \delta; 1).$$

[30]

$$(9.76) \quad {}_3F_2(-2\alpha, -2\beta, \gamma; \gamma+1/2, 2\gamma; x) = {}_2F_1^2(-\alpha, -\beta; \gamma+1/2; x).$$

[30]

$$(9.77) \quad {}_3F_2(\alpha, \beta, \gamma; \delta, \epsilon; 1) = \frac{\Gamma(\delta)\Gamma(\delta-\alpha-\beta)}{\Gamma(\delta-\alpha)\Gamma(\delta-\beta)} {}_3F_2(\alpha, \beta, \epsilon-\gamma; \alpha+\beta-\delta+1, \epsilon; 1) \\ + \frac{\Gamma(\delta)\Gamma(\epsilon)\Gamma(\alpha+\beta-\gamma)\Gamma(\delta+\epsilon-\alpha-\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\epsilon-\gamma)\Gamma(\delta+\epsilon-\alpha-\beta)} {}_3F_2(\delta-\alpha, \delta-\beta, \delta+\epsilon-\alpha-\beta-\gamma; \delta-\alpha-\beta+1, \delta+\epsilon-\alpha-\beta; 1).$$

[174, (2.2.3.2)]

(9.78)

$${}_{A+1}F_B[(a), -m; (b); z] = \frac{((a))_m (-z)^m}{((b))_m} {}_{B+1}F_A[1-(b), 1-m; 1-(a); \frac{(-1)^{A+B}}{z}].$$

[174, (4.3.5.1)][200][105]

(9.79)

$${}_4F_3(x, y, z, -n; u, v, w; 1) = \frac{(v-z)_n (w-z)_n}{(v)_n (w)_n} {}_4F_3(u-x, u-y, z, -n; 1-v+z-n, 1-w+z-n, w; 1),$$

if $u + v + w = 1 + x + y + z - n$.

[174]

(9.80)

$$[\delta(\delta+b_1-1)(\delta+b_2-1)\cdots(\delta+b_B-1)-z(\delta+a_1)(\delta+a_2)\cdots(\delta+a_A)] {}_A F_B((a), (b), z) = 0,$$

where $\delta = z \frac{d}{dz}$.

[47]

$$(9.81) \quad \frac{(\delta+h)_m}{(\delta+k)_m} {}_r F_s \left(\begin{matrix} a, \dots \\ c, \dots \end{matrix} \middle| x \right) = \frac{(h)_m}{(k)_m} {}_{r+2} F_{s+2} \left(\begin{matrix} a, \dots, h+m, k \\ c, \dots, h, k+m \end{matrix} \middle| x \right).$$

[47]

$$(9.82) \quad \sum_{n \geq 0} \frac{(a)_n}{n!} F \left(\begin{matrix} -n, A, \dots \\ C, \dots \end{matrix} \middle| p \right) x^n = (1-x)^{-a} F \left(\begin{matrix} a, A, \dots \\ C, \dots \end{matrix} \middle| -\frac{px}{1-x} \right).$$

[47]

(9.83)

$$\sum_{n \geq 0} \frac{(a)_n}{n!} F \left(\begin{matrix} -n, a+n, A, \dots \\ C, \dots \end{matrix} \middle| p \right) x^n = (1-x)^{-a} F \left(\begin{matrix} a/2, (1+a)/2, A, \dots \\ C, \dots \end{matrix} \middle| -\frac{4px}{(1-x)^2} \right).$$

[47]

$$(9.84) \quad \sum_{n \geq 0} \frac{(a)_n}{n!} F \left(\begin{matrix} -n, A, \dots \\ 1-a-n, C, \dots \end{matrix} \middle| p \right) x^n = (1-x)^{-a} F \left(\begin{matrix} A, \dots \\ C, \dots \end{matrix} \middle| px \right).$$

[47]

$$(9.85) \quad [\delta(\delta+c-1) - px(\delta+a)(\delta-n)] F_n = 0,$$

where $F_n \equiv \frac{(c)_n}{n!} {}_2 F_1(a, -n; c; px)$.

[47]

$$(9.86) \quad nF_n - [2n+c-2-p(n+a-1)]F_{n-1} + (1-p)(n+c-2)F_{n-2} = 0$$

where $F_n \equiv \frac{(c)_n}{n!} {}_2 F_1(a, -n; c; p)$.

[47]

$$(9.87) \quad [\delta(\delta+c-1)(\delta+c'-1) - (\delta+a)(\delta+a')(\delta-n)] F_n = 0,$$

where $F_n \equiv \frac{(c)_n}{n!} {}_3 F_2(a, a', -n; c, c'; x)$.

[47]

(9.88)

$$n(n+c'-1)F_n - [2(n-1)^2 + (2c+2c'-a-a'-1)(n-1) + cc' - aa']F_{n-1} + (n+c-2)(n+c+c'-a-a'-2)F_{n-2} = 0$$

where $F_n \equiv \frac{(c)_n}{n!} {}_3 F_2(a, a', -n; c, c'; 1)$.

[174]

$$(9.89) \quad \left[\sum_{\nu=1}^B z^{\nu-1} (a_\nu z - b_\nu) \frac{d^\nu}{dz^\nu} + a_0 + z^B (1-z) \frac{d^{B+1}}{dz^{B+1}} \right] {}_{B+1} F_B((a), (b), z) = 0.$$

[163]

$$(9.90) \quad {}_2 F_2(-3/2, -1/2; -5/2, 1; z) = \frac{5-4z}{5} e^{z/2} I_0(z/2) + \frac{4z}{5} e^{z/2} I_1(z/2).$$

[163]

$${}_2F_3(-1/2, 1; 1/4, 1/2, 3/4; z) = 1 + z^{1/4}\sqrt{2}\sqrt{\pi}e^{2\sqrt{z}}\operatorname{erf}(\sqrt{2}z^{1/4}) \\ - z^{1/4}\sqrt{2}\sqrt{\pi}e^{-2\sqrt{z}}\operatorname{erfi}(\sqrt{2}z^{1/4}) - 2\sqrt{z}\pi\operatorname{erf}(\sqrt{2}z^{1/4})\operatorname{erfi}(\sqrt{2}z^{1/4}).$$

[163]

$${}_1F_2(3/2; 5/2, 5; z) = -\frac{432 - 24z + 96z^2}{5z^3}I_0(2\sqrt{z}) + \frac{432 + 192z + 48z^2}{5z^{7/2}}I_1(2\sqrt{z}) \\ - \frac{48}{5z}\pi(I_0(2\sqrt{z})L_1(2\sqrt{z}) - I_1(2\sqrt{z})L_0(2\sqrt{z})).$$

[163]

$${}_3F_2(-1/2, 1, 2; 3, 4; z) = -\frac{480 + 3472z - 2100z^2}{525z^3} + \frac{480 + 3712z - 1024z^2 + 192z^3}{525z^3}\sqrt{1-z} \\ - \frac{32}{5z^2}\log\left(\frac{1}{2} + \frac{\sqrt{1-z}}{2}\right).$$

[30]

$$(9.91) \quad {}_2F_3(-\beta, \beta + \gamma; \gamma; \gamma/2, \frac{1+\gamma}{2}; x^2/4) = {}_1F_1(-\beta; \gamma; -x){}_1F_1(-\beta; \gamma; x).$$

[30]

$$(9.92) \quad {}_2F_3(1, n; n+1; (n+1)/2, 2+n2; x^2/4) = {}_1F_1(1; n+1; -x){}_1F_1(1; n+1; x).$$

[30]

(9.93)

$${}_4F_1(-\alpha, -\beta, -\frac{\alpha+\beta}{2} - \frac{\alpha+\beta-1}{2}; -\alpha-\beta; 4x^2) = {}_2F_0(-\alpha, -\beta; x){}_2F_0(-\alpha, -\beta; -x).$$

if α or β a nonnegative integer.

[174]

(9.94)

$${}_{A+1}F_{B+1}(c, (a); d, (b); z) = \frac{\Gamma(d)}{\Gamma(c)\Gamma(d-c)} \int_0^1 t^{c-1}(1-t)^{d-c-1} {}_A F_B((a), (b), tz) dt.$$

[53, 64]

(9.95)

$${}_{q+1}F_q\left(\begin{matrix} a, \frac{b}{q}, \frac{b+1}{q}, \dots, \frac{b+q-1}{q} \\ \frac{c}{q}, \frac{c+1}{q}, \dots, \frac{c+q-1}{q} \end{matrix} \middle| x\right) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1}(1-t)^{c-b-1}(1-xt^q)^{-a} dt \\ = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \sum_{k \geq 0} \binom{c-b-1}{k} \frac{(-)^k}{b+k} {}_2F_1(a, (b+k)/q; 1+(b+k)/q; x)$$

[64]

(9.96)

$${}_{p+k}F_{q+k}\left(\begin{matrix} a_1, \dots, a_p, \frac{\alpha}{k}, \frac{\alpha+1}{k}, \dots, \frac{\alpha+k-1}{k} \\ b_1, \dots, b_q, \frac{\alpha+\beta}{k}, \frac{\alpha+\beta+1}{k}, \dots, \frac{\alpha+\beta+k-1}{k} \end{matrix} \middle| ct^k\right) = \frac{t^{1-\alpha-\beta}}{B(\alpha, \beta)} \int_0^t x^{\alpha-1}(t-x)^{\beta-1} {}_pF_q\left(\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} \middle| cx^k\right) dx.$$

$$(9.97) \quad {}_{r+2}F_{r+1} \left(\begin{matrix} a, b, & (f_r + 1) \\ c, & (f_r) \end{matrix} \mid x \right) = (1-x)^{-a} {}_{r+2}F_{r+1} \left(\begin{matrix} a, \lambda, & (\xi_r + 1) \\ c, & (\xi_r) \end{matrix} \mid \frac{x}{x-1} \right)$$

$$(9.98) \quad {}_{r+2}F_{r+1} \left(\begin{matrix} a, b, & (f_r + 1) \\ c, & (f_r) \end{matrix} \mid x \right) = (1-x)^{c-a-b-r} {}_{r+2}F_{r+1} \left(\begin{matrix} \lambda, \lambda', & (\xi_r + 1) \\ c, & (\xi_r) \end{matrix} \mid x \right)$$

$$(9.99) \quad {}_{p+1}F_p \left(\begin{matrix} a, a, \dots, a \\ a+1, \dots, a+1 \end{matrix} \mid 1 \right) = \frac{a^p}{(a-1)!} \sum_{k=0}^{a-1} (-1)^{a-k-1} \zeta(p-k) \left[\begin{matrix} a \\ k+1 \end{matrix} \right]$$

for a a positive integer, where the $\left[\begin{matrix} n \\ k \end{matrix} \right]$ is Stirling numbers of the first kind.

$$(9.100) \quad {}_{p+1}F_p(a_1, \dots, a_{p+1}; b_1 \dots b_p \mid z\zeta) = (1-z)^{-a_1} \sum_{k \geq 0} \frac{(a_1)_k}{k!} {}_{p+1}F_p(-k, a_2, \dots, a_{p+1}; b_1, \dots, b_p \mid \zeta) \left(\frac{z}{z-1} \right)^k.$$

$$(9.101) \quad {}_{p+1}F_p \left(\begin{matrix} a, a, \dots, a \\ a+1, \dots, a+1 \end{matrix} \mid 1 \right) = \frac{(-1)^{p-1} \pi a^p}{\sin(a\pi)(p-1)!} w(a, p-1)$$

where

$$(9.102) \quad w(n, m) = \frac{1}{(n-1)!} \sum_{i=m+1}^n \left[\begin{matrix} n \\ i \end{matrix} \right] (i-m)_m (-1)^{n-i} n^{i-m-1},$$

recursively

$$(9.103) \quad w(n, 0) = 1, \quad w(n, m) = \sum_{k=0}^{m-1} (1-m)_k H_{n-1}^{(k+1)} w(n, m-1-k)$$

and the Harmonic numbers defined in (0.88).

$$(9.104) \quad {}_{q+1}F_q \left(\begin{matrix} a_1, \dots, a_{q+1} \\ b_1, \dots, b_q \end{matrix} \mid 1 \right) + {}_{q+1}F_q \left(\begin{matrix} a_1, \dots, a_{q+1} \\ b_1, \dots, b_q \end{matrix} \mid -1 \right) \\ = 2 {}_{2q+2}F_{2q+1} \left(\begin{matrix} a_1/2, a_1/2+1/2, \dots, a_{q+1}/2+1/2 \\ b_1/2, b_1/2+1/2, \dots, b_q/2+1/2, 1/2 \end{matrix} \mid 1 \right).$$

$$(9.105) \quad {}_{q+1}F_q \left(\begin{matrix} a_1, \dots, a_{q+1} \\ b_1, \dots, b_q \end{matrix} \mid 1 \right) - {}_{q+1}F_q \left(\begin{matrix} a_1, \dots, a_{q+1} \\ b_1, \dots, b_q \end{matrix} \mid -1 \right) \\ = 2 \frac{a_1 a_2 \dots a_{q+1}}{b_1 b_2 \dots b_q} {}_{q+1}F_q \left(\begin{matrix} a_1/2+1/2, a_1/2+1, \dots, a_{q+1}/2+1/2, a_{q+1}/2+1 \\ b_1/2+1/2, b_1/2+1, \dots, b_q/2+1, 3/2 \end{matrix} \mid 1 \right).$$

[78]

(9.106)

$${}_{q+1}F_q \left(\begin{matrix} a, a+2, a+4, \dots, a+2q \\ a+1, a+3, a+5, \dots, a+2q-1 \end{matrix} \mid 1 \right) + {}_{q+1}F_q \left(\begin{matrix} a, a+2, a+4, \dots, a+2q \\ a+1, a+3, a+5, a+2q-1 \end{matrix} \mid -1 \right) \\ = {}_2F_1(a/2, a+q+1/2; 1/2; 1).$$

[78]

(9.107)

$${}_{q+1}F_q \left(\begin{matrix} a, a+2, a+4, \dots, a+2q \\ a+1, a+3, a+5, \dots, a+2q-1 \end{matrix} \mid 1 \right) - {}_{q+1}F_q \left(\begin{matrix} a, a+2, a+4, \dots, a+2q \\ a+1, a+3, a+5, a+2q-1 \end{matrix} \mid -1 \right) \\ = \frac{2a(a+2)(a+4) \cdots (a+2q)}{(a+1)(a+3) \cdots (a+2q-1)} {}_2F_1(a/2+1/2, a/2+q+1; 3/2; 1).$$

[78]

(9.108)

$${}_{q+1}F_q \left(\begin{matrix} a, a+2, a+4, \dots, a+2q \\ a+1, a+3, a+5, \dots, a+2q-1 \end{matrix} \mid 1 \right) \\ = \frac{\sqrt{\pi}\Gamma(-a-q)}{\Gamma(1/2-a)\Gamma(-a/2-q)} + \frac{2(a/2)_q}{(a/2+1/2)_{q-1}\Gamma(1-a/2)\Gamma(1/2-a/2-q)},$$

if $\Re(a+q) < 0$, and a similar expression for argument -1 .

[139]

(9.109) ${}_3F_2(a, m, b; c, m-n; 1) = \Gamma(a+n-m+1)$

$$\times \left(\sum_{L=0}^{m-1} \frac{(-1)^L \Gamma(-b+c-a-n+L) \Gamma(1-b+L)}{\Gamma(m-L) \Gamma(-b-n+1+L) \Gamma(-b+c-m+1+L) \Gamma(L+1)} \right) \\ \times \frac{\Gamma(c) \Gamma(m-n) (-1)^n}{\Gamma(c-a) \Gamma(a)}, \quad 0 < n < m.$$

(9.110) ${}_3F_2(a, b, -n; c, m-n; 1) = \Gamma(a+n-m+1)$

$$\times \left(\sum_{L=0}^{m-1} \frac{(-1)^L \Gamma(c-a+L) \Gamma(1+b-m+n+L)}{\Gamma(m-L) \Gamma(b-m+1+L) \Gamma(c+n-m+1+L) \Gamma(L+1)} \right) \\ \times \frac{\Gamma(c) \Gamma(m-n) (-1)^n}{\Gamma(c-a) \Gamma(a)}, \quad 0 < n < m.$$

(9.111)

$${}_3F_2(a, -n, b; c, a-c+b-n+m; 1) = \Gamma(c-b+n-m+1) \Gamma(-a+c-b-m+1) \\ \times \left(\sum_{L=0}^{m-1} \frac{(-1)^L \Gamma(b+L) \Gamma(1-a+c-m+n+L)}{\Gamma(m-L) \Gamma(-a+c-m+1+L) \Gamma(c+n-m+1+L) \Gamma(L+1)} \right) \\ \times \frac{\Gamma(c) \Gamma(m)}{\Gamma(n+1-a+c-b-m) \Gamma(b) \Gamma(c-b)}, \quad 0 < n, m.$$

(9.112)

$${}_3F_2(a, 1, b; n+1, c; 1) = \frac{\Gamma(a-n)\Gamma(b-n)\Gamma(n+1)\Gamma(c)\Gamma(-b+c-a+n)\Gamma(1-b)\Gamma(1+n)\Gamma(c)}{\Gamma(b)\Gamma(c-b)\Gamma(a)\Gamma(c-a)\Gamma(a)} \\ \times \left(\sum_{L=0}^{n-1} \frac{(-1)^L \Gamma(a-1-L)}{\Gamma(c-1-L)\Gamma(n-L)\Gamma(2+L-b)} \right) \\ , \quad 0 < n, m.$$

and 60 others.

[\[64\]](#)

$$(9.113) \quad {}_4F_3(a, b/3, (b+1)/3, (b+2)/3; c/3, (c+1)/3, (c+2)/3; 1)$$

$$= \frac{\Gamma(c)\Gamma(c-b-a)}{\Gamma(c-a)\Gamma(c-b)} \sum_{k=0}^{\infty} \frac{(a)_k (-1)^k (b)_k}{k! (c-a)_k} {}_2F_1(-k, b+k; c-a+k; -1).$$

[\[105\]](#)

$$(9.114) \quad {}_4F_3\left(\frac{\alpha}{2}, \frac{\alpha+1}{2}, \beta+n, -n; 1+\alpha, \frac{\beta}{2}, \frac{\beta+1}{2}; 1\right) = \frac{(\beta-\alpha)_n}{(\beta)_n}.$$

[\[105\]](#)

$$(9.115) \quad {}_4F_3\left(\alpha, -\alpha, -\frac{m}{2}, \frac{1-m}{2}; \frac{1}{2}, \beta, 1-m-\beta; 1\right) = \frac{(\alpha+\beta)_m + (\beta-\alpha)_m}{2(\beta)_m}.$$

[\[105\]](#)

$$(9.116) \quad {}_4F_3\left(a, b, \frac{1}{2}-a-b-n, -n; a+b-\frac{1}{2}, 1-a-n, 1-b-n; 1\right) = \frac{(2a)_n (2b)_n (a+b)_n}{(2a+2b-1)_n (a)_n (b)_n}.$$

[\[105\]](#)

$$(9.117) \quad {}_4F_3\left(a-\frac{1}{2}, b-\frac{1}{2}, \frac{1}{2}-a-b-n, -n; a+b-\frac{1}{2}, \frac{1}{2}-a-n, \frac{1}{2}-b-n; 1\right) = \frac{(2a)_n (2b)_n (a+b)_n}{(2a+2b-1)_n (a+\frac{1}{2})_n (b+\frac{1}{2})_n}.$$

[\[105\]](#)

$$(9.118) \quad {}_4F_3\left(a+1, b, \frac{1}{2}-a-b-n, -n; a+b+\frac{1}{2}, 1-a-n, 1-b-n; 1\right) = \frac{(2a+1)_n (2b)_n (a+b)_n}{(2a+2b)_n (a)_n (b)_n}.$$

[\[105\]](#)

$$(9.119) \quad {}_4F_3\left(a+\frac{1}{2}, b-\frac{1}{2}, \frac{1}{2}-a-b-n, -n; a+b+\frac{1}{2}, \frac{1}{2}-a-n, \frac{1}{2}-b-n; 1\right) = \frac{(2a+1)_n (2b)_n (a+b)_n}{(2a+2b)_n (a+\frac{1}{2})_n (b+\frac{1}{2})_n}.$$

[\[105\]](#)

$$(9.120) \quad {}_4F_3\left(a, b, \frac{1}{2}-a-b-n, -n; a+b+\frac{1}{2}, 1-a-n, 1-b-n; 1\right) = \frac{(2a)_n (2b)_n (a+b)_n}{(2a+2b)_n (a)_n (b)_n}.$$

[105]
(9.121)

$${}_4F_3(a, b, a+b-\frac{1}{2}-n, -n; a+b+\frac{1}{2}, \frac{1}{2}+a-n, \frac{1}{2}+b-n; 1) = \frac{(1/2)_n(a-b+\frac{1}{2})_n(b-a+\frac{1}{2})_n}{(a+b+\frac{1}{2})_n(\frac{1}{2}-a)_n(\frac{1}{2}-b)_n}.$$

[105]
(9.122)

$${}_4F_3(a, -a, \frac{1}{2}-n, -n; \frac{1}{2}, \frac{1}{2}+b-n, \frac{1}{2}-b-n; 1) = \frac{(\frac{1}{2}+a+b)_n(\frac{1}{2}-a-b)_n + (\frac{1}{2}+a-b)_n(\frac{1}{2}-a+b)_n}{2(\frac{1}{2}+b)_n(\frac{1}{2}-b)_n}.$$

[105]
(9.123)

$${}_4F_3(a, -a, -\frac{1}{2}-n, -n; \frac{1}{2}, b-n, -b-n; 1) = \frac{(a+b)_{n+1}(1-a-b)_n + (b-a)_{n+1}(1+a-b)_n}{2b(1+b)_n(1-b)_n}.$$

[105]
(9.124)

$${}_4F_3(\frac{1}{2}+a, \frac{1}{2}-a, -\frac{1}{2}-n, -n; \frac{1}{2}, \frac{1}{2}+b-n, \frac{1}{2}-b-n; 1) = \frac{(a+b)_{n+1}(1-a-b)_n + (b-a)_{n+1}(1+a-b)_n}{2b(\frac{1}{2}+b)_n(\frac{1}{2}-b)_n}.$$

[105]

$$(9.125) \quad {}_4F_3(a, -a, -\frac{1}{2}-n, -n; \frac{1}{2}, \frac{1}{2}+b-n, \frac{1}{2}-b-n; 1)$$

$$= \frac{(a+b)(\frac{1}{2}+a+b)_n(\frac{1}{2}-a-b)_n + (b-a)(\frac{1}{2}+b-a)_n(\frac{1}{2}+a-b)_n}{2b(\frac{1}{2}+b)_n(\frac{1}{2}-b)_n}.$$

[105]
(9.126)

$${}_4F_3(a, -a, \frac{1}{2}-n, -n; \frac{1}{2}, 1+b-n, 1-b-n; 1) = \frac{(a+b)_n(-a-b)_n + (a-b)_n(b-a)_n}{2(b)_n(-b)_n}.$$

[30]
(9.127)

$${}_4F_3(\alpha, \beta, \frac{\alpha+\beta}{2}, \frac{\gamma+\delta}{2}; \gamma, \delta, \alpha+\beta; x) = {}_2F_1(\alpha, \beta; \gamma; \frac{1-\sqrt{1-x}}{2}) {}_2F_1(\alpha, \beta; \delta; \frac{1-\sqrt{1-x}}{2})$$

if $\alpha + \beta + 1 = \gamma + \delta$.

[45]

(9.128)

$$\frac{1}{\Gamma(b_1)\Gamma(b_2)\Gamma(b_3)} {}_4F_3\left(\begin{matrix} a_1, a_2, a_3, a_4 \\ b_1, b_2, b_3 \end{matrix} \mid 1\right) = \frac{\Gamma(s)}{\Gamma(a_1+s)\Gamma(a_2+s)\Gamma(a_3)\Gamma(a_4)} \\ \times \sum_{k=0}^{\infty} \frac{(b_1+b_3-a_3-a_4)_k(b_2+b_3-a_3-a_4)_k(s)_k}{(a_1+s)_k(a_2+s)_k k!} {}_3F_2\left(\begin{matrix} b_3-a_3, b_3-a_4, -k \\ b_1+b_3-a_3-a_4, b_2+b_3-a_3-a_4 \end{matrix} \mid 1\right).$$

[78]

$$(9.129) \quad {}_4F_3(a/2, a/2+1/2, b/2, b/2+1/2; 1/2+a/2-b/2, 1+a/2-b/2, 1/2; 1) \\ = \frac{\Gamma(1+a-b)\Gamma(1-2b)}{\Gamma(1-b)\Gamma(1+a-2b)} + \frac{\Gamma(1+a-b)\Gamma(1+a/2)}{\Gamma(1+a)\Gamma(1+a/2-b)}.$$

where $\Re b < 1/2$.

[78]

$$(9.130) \quad \frac{2ab}{1+a-b} {}_4F_3(a/2+1/2, a/2+1, b/2+1/2, b/2+1; 1+a/2-b/2, 3/2+a/2-b/2, 3/2; 1) \\ = \frac{\Gamma(1+a-b)\Gamma(1-2b)}{\Gamma(1-b)\Gamma(1+a-2b)} + \frac{\Gamma(1+a-b)\Gamma(1+a/2)}{\Gamma(1+a)\Gamma(1+a/2-b)}.$$

where $\Re b < 1/2$.

[45]

$$(9.131) \quad \frac{1}{\prod_{j=1}^p \Gamma(b_j)} {}_{p+2}F_{p+1} \left(\begin{matrix} a_1, a_2, \dots, a_{p+1}, -m \\ b_1, b_2, \dots, b_p, 1-s-m \end{matrix} \mid 1 \right) \\ = \frac{(a_1+s)_m (a_2+s)_m \prod_{j=3}^{p+1} (a_j)_m}{(s)_m \prod_{j=1}^p \Gamma(b_j+m)} \sum_{k=0}^m \frac{(s)_k (-m)_k}{(a_1+s)_k (a_2+s)_k k!} \\ \times (S)_k {}_{p+1}F_p \left(\begin{matrix} 1-b_1-m, 1-b_2-m, \dots, 1-b_p-m, -k \\ 1-a_3-m, 1-a_4-m, \dots, 1-a_{p+1}-m, 1-S-k \end{matrix} \mid 1 \right)$$

where $s = \sum_{j=1}^p b_j - \sum_{j=1}^{p+1} a_j$ and $S = a_1 + a_2 + s + m - 1$ and s not a negative integer or zero.

[45]

$$(9.132) \quad \frac{1}{\Gamma(b_1)\Gamma(b_2)} {}_4F_3 \left(\begin{matrix} a_1, a_2, a_3, -m \\ b_1, b_2, 1-s-m \end{matrix} \mid 1 \right) = \frac{(a_1+s)_m (a_2+s)_m (a_3)_m}{(s)_m \Gamma(b_1+m) \Gamma(b_2+m)} {}_4F_3 \left(\begin{matrix} b_1-a_3, b_2-a_3, s, -m \\ a_1+s, a_2+s, 1-a_3-m \end{matrix} \mid 1 \right)$$

where $s = b_1 + b_2 - a_1 - a_2 - a_3$ not a negative integer or zero, $m = 0, 1, 2, \dots$

[30]

$$(9.133) \quad {}_0F_2(m+n+1, n+1; x) {}_0F_2(m+1, 1-n; -x) = 1 + \sum_{k \geq 1} \frac{\alpha_k \binom{2m+n+k+2}{2}_k (2x)^k}{(m+n+1)_k (m+1)_k k!}$$

where

$$(9.134) \quad \alpha_k = \begin{cases} \frac{n}{(n^2-1^2)(n^2-3^2)\dots(n^2-k^2)}, & \text{if } k \text{ odd;} \\ \frac{1}{(n^2-2^2)(n^2-4^2)\dots(n^2-k^2)}, & \text{if } k \text{ even} \end{cases}$$

[54]

$$(9.135) \quad \sum_{k \geq 0} t^k \frac{(\alpha+1+sk)_k}{k!} {}_rF_r \left(\begin{matrix} \Delta(-k, r) \\ \Delta(-\alpha-sk-k, r) \end{matrix} \mid x(\beta+sk) \right) = (1-z)^{\alpha+1} \frac{1}{1+sz+rsy} e^{-\beta y},$$

where $t = (-z)(1-z)^{-s-1} \exp(sy)$, $x = (-y)[1-1/z]^r$, $(a)_k = \Gamma(a+k)/\Gamma(a)$ and $\Delta(-k, r) \equiv -k/r, (-k+1)/r, \dots, (-k+r-1)/r, |t| < 1$.

[54]

(9.136)

$$\begin{aligned} & \sum_{k \geq 0} t^k \frac{(\alpha + 1 + sk)_k}{(\alpha + l + 1 + sk)k!((\alpha + l + 1 + s'' + sk)/s'')_{l'}} {}_rF_r \left(\begin{matrix} \Delta(-k, r) \\ \Delta(-\alpha = sk - k, r) \end{matrix} \mid x(\beta + sk) \right) \\ &= \frac{(1-z)^{\alpha+l+1} e^{-\beta y}}{l'!} \sum_{q=0}^{l'} \sum_{k=0}^{l+s''q} \frac{(-z)^k (1-z)^{s''q-k} (-l')_q (-l-s''q)_k}{k!q!(\alpha + l + 1 + s''q + sk)} \\ & \quad \times {}_1F_1 \left(\begin{matrix} 1 \\ (\alpha + l + 1 + s' + s''r + sk)/s' \end{matrix} \mid y(\beta - \alpha - l - 1 - s''q) \right), \end{aligned}$$

where $t = (-z)(1-z)^{-s-1} e^{sy}$, $x = -y[1 - 1/z]^r$ and $|t| < 1$.

[54]

$$(9.137) \quad \sum_{p \geq 0} t^p \frac{(\beta + s'p)^p}{p!} {}_rF_0 \left(\begin{matrix} \Delta(-p, r) \\ - \end{matrix} \mid \frac{xr^r(\alpha + s'p)}{(\beta + s'p)^r} \right) = \frac{e^{-\beta y - \alpha z}}{1 + s'y + rs'z},$$

where $t = -ye^{s'y+s'z}$, $x = -z/y^r$ and $|s'y \exp(s'y + s'z + 1)| < 1$.

[54]

(9.138)

$$\sum_{p \geq 0} t^p \frac{(\beta + s'p)^p}{p!(\alpha + s'p)} {}_rF_0 \left(\begin{matrix} \Delta(-p, r) \\ - \end{matrix} \mid \frac{xr^r(\alpha + s'p)}{(\beta + s'p)^r} \right) = \frac{e^{-\beta y - \alpha z}}{\alpha} {}_1F_1 \left(\begin{matrix} 1 \\ \alpha/s' + 1 \end{matrix} \mid y(\beta - \alpha) \right)$$

where $t = -ye^{s'y+s'z}$, $x = -z/y^r$ and $|s'y \exp(s'y + s'z + 1)| < 1$.

[170]

$$(9.139) \quad F_1 \left(\begin{matrix} a; b, b \\ 1 + a - b \end{matrix} \mid e^{2\pi i/3}, e^{-2\pi i/3} \right) = \frac{\Gamma(1 + a - b)\Gamma(1 + a/3)}{\Gamma(1 + a)\Gamma(1 + a/3 - b)}.$$

[170]

$$(9.140) \quad F_D \left(\begin{matrix} a; b, b, \dots b \\ c \end{matrix} \mid \omega_{1,n}, \dots, \omega_{n-1,n} \right) = \frac{\Gamma(a - b + 1)\Gamma(1 + a/n)}{\Gamma(1 + a)\Gamma(1 + a/n - b)},$$

where $\omega_{k,n} \equiv e^{2k\pi i/n}$.

[170]

(9.141)

$$F_D \left(\begin{matrix} 2mb - a; b, b, \dots b \\ 2mb \end{matrix} \mid x_1, \dots, x_{2m} \right) = \frac{1}{2m} \frac{\Gamma(a/(2m))\Gamma(2mb)\Gamma((2mb - a)/(2m))}{\Gamma(a)\Gamma(b)\Gamma(2mb - a)},$$

where $x_k = 1 + e^{(2k-1)\pi i/(2m)}$, for $k = 1, \dots, 2m$, and $2m$ is an even integer, $a > 0$, $b > 0$, $nb > a$.

[170]

$$\begin{aligned} (9.142) \quad & F_D \left(\begin{matrix} (2m-1)b - a; b, b, \dots b \\ (2m-1)b \end{matrix} \mid y_1, \dots, y_{2m-1} \right) \\ &= \frac{1}{2m-1} \frac{\Gamma(a/(2m-1))\Gamma((2m-1)b)\Gamma(((2m-1)b - a)/(2m-1))}{\Gamma(a)\Gamma(b)\Gamma((2m-1)b - a)}, \end{aligned}$$

where $y_k = 1 + e^{(2k-1)\pi i/(2m-1)}$, and $2m-1$ is an odd integer, $a > 0$, $b > 0$, $nb > a$.

[18, p 20]

$$(9.143) \quad \alpha F_4(\alpha + 1, \beta; \gamma, \gamma'; x, y) - \beta F_4(\alpha, \beta + 1; \gamma, \gamma'; x, y) \\ = (\alpha - \beta) F_4(\alpha, \beta; \gamma, \gamma'; x, y).$$

[18, p 21]

$$(9.144) \quad \frac{\beta}{\gamma} x F_4(\alpha + 1, \beta + 1; \gamma + 1, \gamma'; x, y) + \frac{\beta}{\gamma'} y F_4(\alpha + 1, \beta + 1; \gamma, \gamma' + 1; x, y) \\ = F_4(\alpha + 1, \beta; \gamma, \gamma'; x, y) - F_4(\alpha, \beta; \gamma, \gamma'; x, y).$$

[18, p 26]

$$(9.145) \quad F_4(\alpha, \beta; \gamma, \gamma'; x, y) = \sum \frac{(\alpha)_m (\beta)_m}{(\gamma)_m m!} \frac{\Gamma(\gamma') \Gamma(\beta - \alpha)}{\Gamma(\gamma' - \alpha - m) \Gamma(\beta + m)} (-y)^{-\alpha - m} \\ \times {}_2F_1(\alpha + m, \alpha + m + 1 - \gamma'; \alpha + 1 - \beta; \frac{1}{y}) x^m \\ + \sum \frac{(\alpha)_m (\beta)_m}{(\gamma)_m m!} \frac{\Gamma(\gamma') \Gamma(\alpha - \beta)}{\Gamma(\gamma' - \beta - m) \Gamma(\alpha + m)} (-y)^{-\beta - m} \\ \times {}_2F_1(\beta + m, \beta + m + 1 - \gamma'; \beta + 1 - \alpha; \frac{1}{y}) x^m.$$

[18]

$$(9.146) \quad F_4(\alpha, \beta; \gamma, \gamma'; x, y) = \frac{\Gamma(\gamma') \Gamma(\beta - \alpha)}{\Gamma(\gamma' - \alpha) \Gamma(\beta)} (-y)^{-\alpha} F_4(\alpha, \alpha + 1 - \gamma'; \gamma, \alpha + 1 - \beta; \frac{x}{y}, \frac{1}{y}) \\ + \frac{\Gamma(\gamma') \Gamma(\alpha - \beta)}{\Gamma(\gamma' - \beta) \Gamma(\alpha)} (-y)^{-\beta} F_4(\beta + 1 - \gamma', \beta; \gamma, \beta + 1 - \alpha; \frac{x}{y}, \frac{1}{y}).$$

[27, 85]

$$(9.147) \quad F_4[\alpha, \beta; \gamma, 1 + \alpha + \beta - \gamma; x(1 - y), y(1 - x)] = F(\alpha, \beta; \gamma; x) F(\alpha, \beta; 1 + \alpha + \beta - \gamma; y).$$

$$(9.148) \quad F_4 \left[\alpha, \beta; \gamma, \beta; -\frac{x}{(1-x)(1-y)}, -\frac{y}{(1-x)(1-y)} \right] = (1-x)^\alpha (1-y)^\alpha F_1[\alpha; \gamma - \beta, 1 + \alpha - \gamma; \gamma; x, xy].$$

$$(9.149) \quad F_4 \left[\alpha, \beta; \alpha, \beta; -\frac{x}{(1-x)(1-y)}, -\frac{y}{(1-x)(1-y)} \right] = (1-xy)^{-1} (1-x)^\beta (1-y)^\alpha.$$

$$(9.150) \quad F_4 \left[\alpha, \beta; \beta, \beta; -\frac{x}{(1-x)(1-y)}, -\frac{y}{(1-x)(1-y)} \right] = (1-x)^\alpha (1-y)^\alpha F[\alpha, 1 + \alpha - \beta; \beta; xy].$$

$$(9.151) \quad F_4 \left[\alpha, \beta; 1 + \alpha - \beta, \beta; -\frac{x}{(1-x)(1-y)}, -\frac{y}{(1-x)(1-y)} \right] = (1-y)^\alpha F[\alpha, \beta; 1 + \alpha - \beta; -\frac{x(1-y)}{1-x}].$$

[105]

$$(9.152) \quad F_{q;1;1}^{p;2;2} \left[\begin{array}{ccc} \alpha_1, \dots, \alpha_p : & a_1, b_1 & a_2, b_2 \\ \gamma_1, \dots, \gamma_q : & c_1 & c_2 \end{array} \mid X, X \right] =_{p+3} F_{q+2}(\alpha_1, \dots, \alpha_p, \beta_1, \beta_2, \beta_3; \gamma_1, \dots, \gamma_q, \delta_1, \delta_2; X)$$

$$\begin{array}{c|c|c|c|c}
a_1 & b_1 & c_1 & a_2 & b_2 & c_2 & \beta_1 & \beta_2 & \beta_3 & \delta_1 & \delta_2 \\
\hline
a & b & a+b-\frac{1}{2} & a & b & a+b+\frac{1}{2} & 2a & 2b & a+b & 2a+2b-1 & a+b+\frac{1}{2} \\
a-\frac{1}{2} & b-\frac{1}{2} & a+b-\frac{1}{2} & a+\frac{1}{2} & b+\frac{1}{2} & a+b+\frac{1}{2} & 2a & 2b & a+b & 2a+2b-1 & a+b+\frac{1}{2}
\end{array}$$

and more of that format.

[183]

(9.153)

$$F_{1:1;1}^{1:2;2} \left[\begin{array}{c} \alpha : \quad -M, \beta : \quad -N, \beta' : \\ \beta + \beta' : \quad \alpha - \beta' - M + 1 \quad \alpha - \beta - N + 1 \end{array} \mid 1, 1 \right] = \frac{(\beta + \beta' - \alpha)_{M+N} (\beta')_M (\beta)_N}{(\beta + \beta')_{M+N} (\beta' - \alpha)_M (\beta - \alpha)_N}$$

for $M, N = 0, 1, 2, \dots$

[183]

$$\begin{aligned}
(9.154) \quad & F_{1:1;1}^{1:2;2} \left[\begin{array}{c} \beta + \beta' - \alpha : \quad \beta, \gamma : \quad \beta', \gamma' : \\ \beta + \beta' : \quad \delta \quad \delta' \end{array} \mid 1, 1 \right] \\
&= \frac{\Gamma(\delta)\Gamma(\delta')\Gamma(\alpha - \beta - \gamma + \delta)\Gamma(\alpha - \beta' - \gamma' + \delta')}{\Gamma(\delta - \gamma)\Gamma(\delta' - \gamma')\Gamma(\alpha - \beta + \delta)\Gamma(\alpha - \beta' + \delta')} F_{1:1;1}^{1:2;2} \left[\begin{array}{c} \alpha : \quad \beta', \gamma : \quad \beta, \gamma' : \\ \beta + \beta' : \quad \alpha - \beta + \delta \quad \alpha - \beta' + \delta' \end{array} \mid 1, 1 \right].
\end{aligned}$$

[183]

(9.155)

$$\begin{aligned}
F_{1:s;v}^{1:r;u} \left[\begin{array}{c} \alpha : \quad (a_r) : \quad (c_u) : \\ \gamma : \quad (b_s) : \quad (d_v); \end{array} \mid x, y \right] &= \sum_{n=0}^{\infty} \frac{(\alpha)_n (\gamma - \alpha)_n \prod_{j=1}^r (a_j)_n \prod_{j=1}^u (c_j)_n}{(\gamma + n - 1)_n (\gamma)_{2n} \prod_{j=1}^s (b_j)_n \prod_{l=1}^v (d_l)_n} \frac{(xy)^n}{n!} \\
&\times {}_{r+1}F_{s+1} \left(\begin{array}{c} (a_r) + n, \quad \alpha + n \\ (b_s) + n, \quad \gamma + 2n \end{array} \mid x \right) {}_{u+1}F_{v+1} \left(\begin{array}{c} (c_u) + n, \quad \alpha + n; \\ (d_v) + n, \quad \gamma + 2n; \end{array} \mid y \right).
\end{aligned}$$

[85]

$$\begin{aligned}
(9.156) \quad & {}_3F_2 \left(\begin{array}{c} -n, n+a, b \\ c, d \end{array} \mid 1 \right) {}_3F_2 \left(\begin{array}{c} -n, n+a, e \\ c, f \end{array} \mid 1 \right) \\
&= \frac{(-)^n (a-c+1)_n}{(c)_n} F_{2:1;1}^{2:2;2} \left[\begin{array}{c} -n, n+a : \quad b, e; \quad d-b, f-e \\ d, f : \quad c; \quad a-c+1 \end{array} \mid 1, 1 \right].
\end{aligned}$$

[85]

$$\begin{aligned}
(9.157) \quad & F_{2:1;1}^{2:2;2} \left[\begin{array}{c} a, b : \quad -x, y+e; \quad -y, x+d \\ d, e : \quad c; \quad b \end{array} \mid 1, 1 \right] \\
&= \frac{(d-a)_x (e-a)_y}{(d)_x (e)_y} F_{2:1;0}^{2:2;2} \left[\begin{array}{c} a, -x : \quad 1+a-c, -y; \quad c-b \\ c, 1+a-d-x : \quad 1+a-e-y; \quad - \end{array} \mid 1, 1 \right]
\end{aligned}$$

if $x, y = 0, 1, \dots$

[85]

(9.158)

$$\begin{aligned}
F_{2:1;1}^{2:2;2} \left[\begin{array}{c} a, b : \quad -x, y+e; \quad -y, x+d \\ d, e : \quad c; \quad c' \end{array} \mid 1, 1 \right] &= \sum_{r=0}^{\min(x,y)} \frac{(a)_r (b)_r (a+b-c-c'+1)_r}{r! (c)_r (c')_r} \frac{(-x)_r (-y)_r}{(d)_r (e)_r} \\
&\times {}_3F_2 \left(\begin{array}{c} r+a, r+b, r-x \\ r+c, r+c \end{array} \mid 1 \right) {}_3F_2 \left(\begin{array}{c} r+a, r+b, r-y \\ r+c', r+e \end{array} \mid 1 \right)
\end{aligned}$$

if $x, y = 0, 1, \dots$

9.2. The Confluent Hypergeometric Function.

9.3. The Meijer G-Function. [2]

$$(9.159) \quad G_{p+1,q+2}^{m+1,n+1} \left(z \mid \begin{matrix} a, a_p \\ a, b_q, b \end{matrix} \right) = (-)^{a-b} G_{p,q+1}^{m+1,n} \left(z \mid \begin{matrix} a_p \\ b_q, b \end{matrix} \right) \\ - (-1)^{a-b} \sum_{k=1}^{a-b} \text{Res}_{s=k-a} \left[\frac{\Gamma(b+s) \prod_{i=1}^m \Gamma(b_i+s) \prod_{i=1}^n \Gamma(1-a_i-s)}{\prod_{i=n+1}^p \Gamma(a_i+s) \prod_{i=m+1,q} \Gamma(1-b_i-s)} z^{-s} \right]$$

if $a-b > 0$.

[2]

$$(9.160) \quad G_{p+1,q+2}^{m+1,n+1} \left(z \mid \begin{matrix} a, a_p \\ a, b_q, b \end{matrix} \right) = (-)^{a-b} G_{p,q+1}^{m+1,n} \left(z \mid \begin{matrix} a_p \\ b_q, b \end{matrix} \right)$$

if $a-b \leq 0$.

[2]

$$(9.161) \quad G_{p+2,q+1}^{m,n+1} \left(z \mid \begin{matrix} a, a_p, b \\ b_q, b \end{matrix} \right) = (-)^{a-b} G_{p+1,q}^{m,n} \left(z \mid \begin{matrix} a_p, a \\ b_q, b \end{matrix} \right)$$

if $a-b$ is an integer.

9.4. The MacRobert E-function.

9.5. Riemann and Hurwitz zeta functions. [188, (2.4.1)]

$$(9.162) \quad \zeta(s) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1}}{e^x - 1} dx, \quad \sigma > 1.$$

[184, 5]

$$(9.163) \quad \zeta(3) = \frac{5}{2} \sum_{k \geq 1} \frac{(-)^{k-1}}{k^3 \binom{2k}{k}} = -\frac{4\pi^2}{7} \sum_{k \geq 0} \frac{\zeta(2k)}{(2k+1)(2k+2)2^{2k}}.$$

[153]

$$(9.164) \quad \zeta(4) = \sum_{n=1}^{\infty} \frac{1}{n!^4} \left(\frac{4n+1}{2n^2} L_n + \frac{7n^3}{4} L_{n-1} \right)$$

where $L_0 = 0$, $L_1 = 1/3$ and

$$(9.165) \quad 4(4n+3)(4n+5)L_{n+1} + 2(n+1)^3(6n^3+9n^2+5n+1)L_n - n^6(n+1)^3L_{n-1} = 0.$$

[142]

$$(9.166) \quad \zeta'(2k) = (-)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} \{2k\zeta'(1-2k) - [\psi(2k) - \log(2\pi)]B_{2k}\}.$$

[142]

$$(9.167) \quad \zeta'(1-2k, p/q) = \frac{[\psi(2k) - \log(2\pi q)]B_{2k}(p/q)}{2k} - \frac{[\psi(2k) - \log(2\pi)]B_{2k}(p/q)}{q^{2k}2k} \\ + (-)^{k+1} \frac{\pi}{(2\pi q)^{2k}} \sum_{n=1}^{q-1} \sin\left(\frac{2\pi pn}{q}\right) \psi^{(2k-1)}(n/q) \\ + (-)^{k+1} \frac{2(2k-1)!}{(2\pi q)^{2k}} \sum_{n=1}^{q-1} \cos\left(\frac{2\pi pn}{q}\right) \zeta'(2k, n/q) + \frac{\zeta'(1-2k)}{q^{2k}}.$$

[142]

$$(9.168) \quad \zeta'(1-2k, 1/2) = -\frac{B_{2k} \log 2}{k4^k} - \frac{(2^{2k-1} - 1)\zeta'(1-2k)}{2^{2k-1}}.$$

[142]

$$(9.169) \quad \zeta'(1-2k, 1/3) = -\frac{(9^k - 1)B_{2k}\pi}{\sqrt{3}(3^{2k-1} - 1)8k} - \frac{B_{2k} \log 3}{3^{2k-1}4k} \\ + (-)^k \frac{\psi^{(2k-1)}(1/3)}{2\sqrt{3}(6\pi)^{2k-1}} - \frac{(3^{2k-1} - 1)\zeta'(1-2k)}{2 \times 3^{2k-1}}.$$

[153]

$$(9.170) \quad \sum_{k=1}^{\infty} \frac{k}{k^4 - x^2 k^2 - y^4} = \sum_{n,m=0}^{\infty} \binom{n+m}{n} \zeta(2n+4m+3) x^{2n} y^{4m} \\ = \frac{1}{2} \sum_{n=1}^{\infty} \frac{(-)^{n-1} r(n)}{n \binom{2n}{n}} \frac{\prod_{m=1}^{n-1} ((m^2 - x^2)^2 + 4y^4)}{\prod_{m=n}^{2n} (m^4 - x^2 m^2 - y^4)}$$

where

$$(9.171) \quad r(n) = 205n^6 - 160n^5 + (32 - 62x^2)n^4 + 40x^2n^3 + (x^4 - 8x^2 - 25y^4)n^2 + 10y^4n + y^4(x^2 - 2).$$

[5]

$$(9.172) \quad \sum_{k \geq 1} \frac{(-)^k}{k} [\zeta(nk) - 1] = \log \left(\prod_{j=0}^{n-1} \Gamma[2 - (-)^{(2j+1)/n}] \right).$$

[5]

$$(9.173) \quad \sum_{k=2}^{\infty} (-)^k [\zeta(k) - 1] k^n = -1 + \frac{1 - 2^{n+1}}{n+1} B_{n+1} - \sum_{k=1}^n (-)^k k! \zeta(k+1) S(n+1, k+1),$$

with S the Stirling numbers of the second kind.

[5]

$$(9.174) \quad \sum_{k=2}^{\infty} [\zeta(k) - 1] k^n = 1 + \sum_{k=1}^n k! \zeta(k+1) S(n+1, k+1),$$

[6]

$$(9.175) \quad -\sum_{k=1}^{\infty} \frac{k}{4^{2k}} \zeta(2k+1) = G - 1.$$

[184]

$$(9.176) \quad \sum_{k \geq 0} \frac{\zeta(2k)}{(2k+1)4^{2k}} = -\frac{G}{\pi} - \frac{1}{4} \log 2.$$

[184]

$$(9.177) \quad \sum_{k \geq 0} \frac{\zeta(2k)}{2k+1} \left(\frac{3}{4}\right)^{2k} = \frac{G}{3\pi} - \frac{1}{4} \log 2.$$

[6]

$$(9.178) \quad \frac{1}{16} \sum_{k=1}^{\infty} \frac{3^k - 1}{4^k} (k+1) \zeta(k+2) = G.$$

[5]

$$(9.179) \quad \sum_{k=1}^{\infty} \frac{\zeta(2k+1) - 1}{k+2} = \frac{2}{3} - \frac{\gamma}{2} + \log 2 + 6\zeta'(-1).$$

[5]

$$(9.180) \quad \sum_{k=2}^{\infty} \frac{\zeta(k)}{(k+1)(k+2)} = \frac{1-\gamma}{6} - 2\zeta'(-1).$$

[5]

$$(9.181) \quad \sum_{k=2}^{\infty} \frac{k^2}{k+1} [\zeta(k) - 1] = \frac{3}{2} - \frac{\gamma}{2} + \frac{\pi^2}{6} - \frac{1}{2} \log(2\pi).$$

[5]

$$(9.182) \quad \sum_{k=2}^{\infty} [\zeta(4k) - 1] = \frac{7}{8} - \frac{\pi}{4} \coth \pi$$

[142]

$$(9.183) \quad 4 \sum_{k=0}^{\infty} \frac{1 - \zeta(2k)}{(2k+1)3^{2k}} = \log(192) - \pi \frac{2\sqrt{3}}{9} + \frac{\sqrt{3}}{3\pi} \psi^{(1)}(1/3).$$

[5]

$$(9.184) \quad \sum_{k=1}^{\infty} [\zeta(4k) - 1] z^{4k} = \frac{3z^4 - 1}{2(z^4 - 1)} - \frac{\pi z}{4} [\cot(\pi z) + \coth(\pi z)], \quad |z| < 2.$$

[5]

$$(9.185) \quad \sum_{k=1}^{\infty} [\zeta(2k) - 1] \sin k = -\frac{1}{2} \cot(1/2) + \frac{\pi}{2} \frac{\sin(1/2) \sin[2\pi \cos(1/2)] - \cos(1/2) \sinh[2\pi \sin(1/2)]}{\cos[2\pi \cos(1/2)] - \cosh[2\pi \sin(1/2)]}$$

[5]

$$(9.186) \quad \sum_{k=1}^{\infty} \binom{p+k}{k} \zeta(p+k+1, a) z^k = \frac{(-)^p}{p!} [\psi^{(p)}(a) - \psi^{(p)}(a-z)].$$

[5]

$$(9.187) \quad \sum_{k=1}^{\infty} \frac{t^k}{k^2} \zeta(2k) = \log[\pi \sqrt{t} \csc(\pi \sqrt{t})].$$

[6]

$$(9.188) \quad \frac{1}{8} \sum_{k=2}^{\infty} \frac{k}{2^k} \zeta(k+1, 3/4) = G.$$

[6]

$$(9.189) \quad -\frac{1}{8} \sum_{k=2}^{\infty} \frac{k}{2^k} \zeta(k+1, 5/4) = G - 1.$$

[3]

(9.190)

$$\zeta'(1, p/q) - \zeta'(1, 1 - p/q) = \pi \cot \frac{\pi p}{q} [\log(2\pi q) + \gamma] - 2\pi \sum_{j=1}^{q-1} \log(\Gamma(j/q)) \sin \frac{2\pi j p}{q}.$$

[75]

$$(9.191) \quad \int_0^1 \sin(2\pi q) \zeta(z, q) dq = \frac{(2\pi)^z}{4\Gamma(z)} \csc \frac{z\pi}{2}.$$

[75]

$$(9.192) \quad \int_0^1 \sin(2k\pi q) \zeta(z, q) dq = \frac{(2\pi)^z k^{z-1}}{4\Gamma(z)} \csc \frac{z\pi}{2}.$$

[75]

$$(9.193) \quad \int_0^1 \cos(2k\pi q) \zeta(z, q) dq = \frac{(2\pi)^z k^{z-1}}{4\Gamma(z)} \sec \frac{z\pi}{2}.$$

[75]

$$(9.194) \quad \int_0^1 \zeta(z', q) \zeta(z, q) dq = -\zeta(z + z' - 1) B(1 - z, 1 - z') \frac{\cos \frac{\pi(z-z')}{2}}{\cos \frac{\pi(z+z')}{2}}.$$

[75]

$$(9.195) \quad \int_0^1 q^n \zeta(z, q) dq = -n! \sum_{j=1}^n \frac{\zeta(z-j)}{(z-j)_j (n-j+1)!}.$$

[75]

$$(9.196) \quad \int_0^1 \ln(\sin \pi q) \zeta(z, q) dq = -\frac{\Gamma(1-z)}{(2\pi)^{1-z}} \sin \frac{\pi z}{2} \zeta(2-z).$$

[95]

$$(9.197) \quad \Phi(z, s, u) = 2^{-s} [\Phi(z^2, s, u/2) + z\Phi(z^2, s, (u+1)/2)].$$

[57]

$$(9.198) \quad \zeta(r, s) \equiv \sum_{m < n} \frac{1}{n^s m^r}.$$

[57]

$$(9.199) \quad \zeta(r, s) + \zeta(s, r) = \zeta(r)\zeta(s) - \zeta(r+s).$$

[57]

$$(9.200) \quad \zeta(r, s) = -\frac{1}{2} \zeta(r+s) + \sum_{j=1, j \text{ odd}} \left(\binom{j-1}{s-1} + \binom{j-1}{r-1} \right) \zeta(j) \zeta(r+s-j).$$

for r even and s odd.

[57]

$$(9.201) \quad \zeta(1, s) = \frac{s}{2} \zeta(s+1) - \frac{1}{2} \sum_{j=1}^{s-2} \zeta(j+1) \zeta(s-j).$$

[57]

$$(9.202) \quad \zeta(r, s) = -\frac{1}{2} \zeta(r+s) + \sum_{k=3, \text{odd}}^{\infty} \Phi_k \sum_{j=0, \text{even}}^{k-1} \binom{k}{j} \eta(r-j) \eta(s-k+j),$$

where

$$(9.203) \quad \Phi_k \equiv -\frac{2}{\pi} \sum_{d=1, \text{dodd}}^{k-2} (-1)^{(d-1)/2} \frac{\pi^d}{d!} \zeta(k-d+1),$$

and

$$(9.204) \quad \eta(s) = (1 - 2^{1-s}) \zeta(s).$$

9.6. Bernoulli Numbers and Polynomials. [95]

$$(9.205) \quad B_m(x) = \sum_{n=0}^m \frac{1}{n+1} \sum_{k=0}^n (-1)^k \binom{n}{k} (x+k)^m.$$

[48]

$$(9.206) \quad \sum_{k=0}^m \frac{\binom{m}{k}}{\binom{n+k+s}{s}} y^{m-k} B_{n+k+s}(x) = \sum_{k=0}^n \frac{\binom{n}{k}}{\binom{m+k+s}{s}} (-y)^{n-k} B_{m+k+s}(x+y) \\ + \sum_{j=0}^{s-1} \sum_{i=0}^{s-1-j} \binom{s-1-j}{i} \binom{s-1}{j} \frac{(-1)^{n+1+i} s y^{m+n+s-j} B_j(x)}{(m+n+1+i) \binom{m+n+l}{n}}$$

and

$$(9.207) \quad \sum_{k=0}^m \binom{m}{k} \binom{n+k}{s} y^{m-k} B_{n+k-s}(x) = \sum_{k=0}^n \binom{n}{k} \binom{m+k}{s} (-y)^{n-k} B_{m+k-s}(x+y).$$

[48]

$$(9.208) \quad \sum_{j=0}^k \binom{k+1}{j} (k+j+1) B_{k+j} = 0, \quad k \geq 1.$$

[57]

$$(9.209) \quad B_s(t) B_r(t) = \sum_{j>0, j \equiv r+s \pmod{2}} \frac{1}{j} \left(r \binom{s}{j-r} + s \binom{r}{j-s} \right) B_{r+s-j}(t) + \frac{1}{2} ((-1)^{r+s} + 1) \frac{(-)^r B_{r+s}}{\binom{r+s}{s}}$$

[95]

$$(9.210) \quad E_m(x) = \sum_{n=0}^m \frac{1}{2^n} \sum_{k=0}^n (-1)^k \binom{n}{k} (x+k)^m.$$

REFERENCES

1. Milton Abramowitz and Irene A. Stegun (eds.), *Handbook of mathematical functions*, 9th ed., Dover Publications, New York, 1972. MR 0167642 (29 #4914)
2. V. S. Adamchik, *The evaluation of integrals of Bessel functions via g-function identities*, J. Comput. Appl. Math. **64** (1995), no. 3, 283–290. MR 1365430
3. ———, *A class of logarithmic integrals*, Proc. 1997 Intl. Symp. Symbol. Algebr. Comput., 1997, pp. 1–8. MR 1809963
4. ———, *On Stirling numbers and Euler sums*, J. Comput. Appl. Math **79** (1997), no. 1, 119–130. MR 1437973 (97m:11025)
5. V. S. Adamchik and H. M. Srivastava, *Some series of the Zeta and related functions*, Analysis (Munich) **18** (1998), no. 2, 131–144. MR 1625172 (99d:11096)
6. Victor Adamchik, *33 representations for catalan's constant*, <http://www.cs.cmu.edu/~adamchik/articles/catalan/catalan.htm>, 2007.
7. A. D. Alhaidari, *Evaluation of integrals involving orthogonal polynomials: Laguerre polynomial and Bessel function example*, Appl. Math. Lett **20** (2007), 38–42. MR 2273125
8. Tewodros Amdeberhan, M. L. Glasser, M. C. Jones, V. H. Moll, R. Posey, and D. Varela, *The Cauchy-Schlömilch transformation*, arXiv:math/1004.2445 (2010).
9. Tewodros Amdeberhan, Luis A. Medina, and Victor H. Moll, *The integrals in Gradshteyn and Ryzhik. part 5: Some trigonometric integrals*, arXiv:math/0705.2379 (2007).
10. Tewodros Amdeberhan and Victor H. Moll, *A formula for a quartic integral: A survey of old proofs and some new ones*, arXiv:math/0707.2118 (2007).
11. ———, *The laplace transform of the digamma function: an integral due to Glasser, Manna and Oloa*, arXiv:math/0707.3663 (2007).
12. S. Amghibech, *On sums involving binomial coefficients*, J. Integer Seq. **10** (2007), # 07.2.1. MR 2276785
13. Moa Apagodu and Doron Zeilberger, *Searching for strange hypergeometric identities by sheer brute force*, Integers: El. J. Combin. Number Theory **8** (2008), #A36. MR 2438527
14. Alexander Apelblat, *Repeating use of integral transforms—a new method for evaluation of some infinite integrals*, IMA J. Appl. Math. **27** (1981), no. 4, 481–496. MR 0637508
15. ———, *Table of definite and infinite integrals*, Physical Sciences Data, vol. 13, Elsevier, 1983. MR 0902582
16. ———, *Some integrals of gamma, polygamma and Volterra functions*, IMA J. Appl. Math. **34** (1985), no. 2, 173–186. MR 0795510
17. ———, *Tables of integrals and series*, 1 ed., Harri Deutsch, Thun, 1996.
18. Paul Appell and Joseph Kampé de Fériet, *Fonctions hypergéométriques et hypersphériques. Polynomes d'Hermite*, Gauthier-Villars, Paris, 1926.
19. Nigel Backhouse, *Pancake functions and approximations to π* , Math. Gaz. **79** (1995), no. 485, 371–374.
20. D. H. Bailey, J. M. Borwein, and R. E. Crandall, *Integrals of the Ising class*, J. Phys. A: Math. Gen. **39** (2006), 12271–12302. MR 2261886
21. David Bailey, Jonathan M. Borwein, and Roland Girgensohn, *Experimental evaluation of Euler sums*, Exper. Math. **3** (1994), no. 1, 17. MR 1302815
22. David Bailey, Peter Borwein, and Simon Plouffe, *On the rapid computation of various polylogarithmic constants*, Math. Com. **66** (1997), 903–913. MR 1415794
23. David H. Bailey, *A compendium of BBP-type formals for mathematical constants*, 16 October 2009.
24. W. N. Bailey, *A note on Bateman's expansion in Bessel functions*, Proc. Cambr. Phil. Soc. **25** (1929), no. 1, 48–49.
25. ———, *Some series of squares of Bessel functions*, Proc. Cambr. Phil. Soc. **26** (1930), no. 1, 82–87.
26. ———, *Some series and integrals involving associated legendre functions (ii)*, Proc. Cambr. Phil. Soc. **27** (1931), no. 3, 381–386.
27. ———, *Some infinite integrals involving Bessel functions*, Proc. Lond. Math. Soc. **40** (1936), no. 1, 37–48, E: [103].
28. Eugenio P. Balanzario and Jorge Sánchez-Ortiz, *A generating function for a class of generalized Bernoulli polynomials*, Raman. J. **19** (2009), 9–18. MR 2501233

29. W. Becken and P. Schmelcher, *The analytic continuation of the gaussian hypergeometric function ${}_1F_1(a, b; c; z)$ for arbitrary parameters*, J. Comput. Appl. Math. **126** (2000), no. 1–2, 449–478. MR 1806771
30. Bruce C. Berndt, *Chapter 11 of Ramanujan's second notebook*, Bull. Lond. Math. Soc. **15** (1983), no. 4, 273–320. MR 0703753 (85a:01043)
31. P. Blasiak, G. Dattoli, A. Horzela, K. A. Penson, and K. Zhukovsky, *Motzkin numbers, central trinomial coefficients and hybrid polynomials*, arXiv:math.CO/0802.0075 (2008).
32. George Boros and Victor H. Moll, *An integral with three parameters*, SIAM Rev. **40** (1998), no. 4, 972–980. MR 1659712
33. ———, *Sums of arctangents and some formulas of Ramanujan*, Sci. Ser. A Math. Sci (N.S.) **11** (2005), 13–24. MR 2196063 (2006j:33001)
34. ———, *A summation method due to Carr: Part 1*, Scientia. Series A: Math. Sci. (2006), no. 12, 21–38. MR 2257096
35. David Borwein and J. M. Borwein, *On an intriguing integral and some series related to $\zeta(4)$* , Proc. Am. Math. Soc. **123** (1995), 1191–1198. MR 1231029
36. David Borwein, Jonathan M. Borwein, and David M. Bradley, *Parametric Euler sum identities*, J. Math. Anal. Appl. **316** (2006), 328–338. MR 2201764
37. Jonathan Borwein, Dirk Nuyens, Armin Straub, and James Wan, *Some arithmetic properties of short random walk integrals*, Ramanujan J. **26** (2011), no. 1, 109–132. MR 2837721
38. Jonathan Michael Borwein, David J. Broadhurst, and Joel Kamnitzer, *Central binomial sums, multiple Clausen values and Zeta values*, Exp. Math. **10** (2001), no. 1, 25–34. MR 1821569 (2002k:11105)
39. Jonathan Michael Borwein and Roland Girgensohn, *Evaluations of binomial series*, Tech. Report 02-188, CECM, 2002.
40. Khristo N. Boyadzhiev, *Series with central binomial coefficients, Catalan numbers, and harmonic numbers*, J. Integer Seq. **15** (2012), 12.1.7. MR 2872464
41. David M. Bradley, *A class of series acceleration formulae for Catalan's constant*, Ramanujan J. **3** (1999), no. 2, 159–173. MR 1703281
42. ———, *Representations of Catalan's constant*, 2001.
43. David K. Brice, *Three-parameter formula for the electronic stopping cross section at non-relativistic velocities*, Phys. Rev. A **6** (1972), no. 5, 1791–1805.
44. D. J. Broadhurst, *Polylogarithmic ladders, hypergeometric series and the ten millionth digits of $\zeta(3)$ and $\zeta(5)$* , arXiv:math/9803067 [math.CA] (1998).
45. Wolfgang Bühring, *Transformation formulas for terminating Saalschützian hypergeometric series of unit argument*, J. Appl. Math. Stoch. Anal. **8** (1995), no. 2, 189–194. MR 1330122
46. Marc Chamberland, *Binary BBP-formulae for logarithms and generalized Gaussian-Mersenne primes*, J. Integer Seq. **6** (2003), 03.3.7. MR 2046407 (2005a:11201)
47. T. W. Chaundy, *An extension of hypergeometric functions (i)*, Quart. J. Math. **14** (1943), 55–78. MR 0010749
48. Kwang-Wu Chen, *Identities from the binomial transform*, J. Number Theory **124** (2007), 142–150. MR 2320995
49. William Y. C. Chen, Quing-Hu Hou, and Yan-Ping Mu, *A telescoping method for double summations*, J. Comp. Appl. Math. **196** (2006), 553–566. MR 2249445
50. Junesang Choi and Arjun Kumar Rathie, *Generalizations of two summation formulas for the generalized hypergeometric function of higher order due to Exton*, Commun. Korean Math. Soc. **25** (2010), no. 3, 385–389.
51. Wenchang Chu and Wenlong Zhang, *Transformations of kummer-type for ${}_2F_2$ -series and their q -analogues*, J. Comp. Appl. Math. **216** (2008), 467–473. MR 2412920
52. Mark W. Coffey, *On some log-cosine integrals related to $\zeta(3)$, $\zeta(4)$ and $\zeta(6)$* , J. Comp. Appl. Math. **159** (2003), 205–215. MR 20005956
53. Mark W. Coffey and S. J. Johnston, *Some results involving series representations of hypergeometric functions*, J. Comp. Appl. Math. **233** (2009), no. 3, 674–679. MR 2583002
54. M. E. Cohen, *Some classes of generating functions for the laguerre and hermite polynomials*, Math. Comp. **31** (1977), no. 238, 511–518. MR 0442323
55. Lin Cong, *On Bernoulli numbers and its properties*, arXiv:math.HO/0408082 (2004).
56. Donal F. Connors, *Some series and integrals involving the Riemann zeta function, binomial coefficients and the harmonic numbers*, arXiv:0710.4023 (2007).

57. Richard E. Crandall and J. P. Buhler, *On the evaluation of Euler sums*, *Exper. Math.* **3** (1994), no. 4, 275–285. MR 1341720
58. S. A. Cruz, C. Cisneros, and I. Alvarez, *Individual orbital contribution to the electronic stopping cross section in the low-velocity region*, *Phys. Rev. A* **17** (1978), no. 1, 132–140, (E): [59].
59. ———, *Erratum: Individual orbital contribution to the electronic stopping cross section in the low-velocity region*, *Phys. Rev. A* **20** (1979), no. 2, 628–629.
60. Djurdje Cvijović and Jacek Klinowski, *Closed-form summation of some trigonometric series*, *Math. Comp.* **64** (1995), no. 209, 205–210. MR 1270616
61. Guang-ming Dai, *Model compensation of atmospheric turbulence with the use of Zernike polynomials and Karhunen-Loève functions*, *J. Opt. Soc. Am. A* **12** (1995), no. 10, 2182–2193.
62. Ayhan Dil and Veli Kurt, *Polynomials related to harmonic numbers and evaluation of harmonic number series ii*, *Appl. An. Discr. Math.* **5** (2011), 212–229. MR 2883425
63. A. L. Dixon and W. L. Ferrar, *Integrals for the product of two Bessel functions*, *Quart. J. Math.* **4** (1933), no. 1, 193–208.
64. K. A. Driver and S. J. Johnston, *An integral representation of some hypergeometric functions*, *El. Trans. Num. Analysis* **25** (2006), 115–120. MR 2280366
65. Herbert Bristol Dwight, *Tables of integrals and other mathematical data*, 3rd ed., Macmillan, New York, 1957. MR 0129577 (23 #B2613)
66. F. J. Dyson, N. E. Frankel, and M. L. Glasser, *Lehmer’s second interesting series*, arXiv:1009.4274 [math-ph] (2010).
67. Askar Dzhumadil’daev and Damir Yeliussizov, *Power sums of binomial coefficients*, *J. Integer Seq.* **16** (2013), # 13.1.4. MR 3022328
68. G. Eason, B. Noble, and I. N. Sneddon, *On certain integrals of Lipschitz-Hankel type involving products of Bessel functions*, *Phil. Trans. Roy. Soc. Lond. A* **247** (1955), no. 935, 529–551. MR 0069961
69. H. El-Rabii, *On the evaluation of the integral $\int_0^\infty x^{-a}(1 - \sin^b x/x^b)dx$* , *Scientia A: Math. Sci.* **14** (2007), 21–25. MR 2330700
70. Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, and Francesco G. Tricomi (eds.), *Higher transcendental functions*, vol. 1, McGraw-Hill, New York, London, 1953. MR 0058756 (15,419i)
71. Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, and Francesco G. Tricomi (eds.), *Higher transcendental functions*, vol. 2, McGraw-Hill, New York, London, 1953.
72. Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, and Francesco G. Tricomi (eds.), *Tables of integral transforms*, vol. 2, McGraw-Hill, New York, London, 1954. MR 0065685 (16,468c)
73. Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, and Francesco G. Tricomi (eds.), *Tables of integral transforms*, vol. 1, McGraw-Hill, New York, London, 1954. MR 0061695 (15,868a)
74. Olivier Espinosa, *On the evaluation of Matsubara sums*, *Math. Comp.* **79** (2010), no. 271, 1709–1725. MR 2630009
75. Olivier Espinosa and Victor H. Moll, *On some integrals involving the Hurwitz zeta function: Part 1*, *Raman. J.* **6** (2002), no. 2, 159–188. MR 1908196
76. ———, *The evaluation of Tornheim double sums. part 2*, arXiv:0811.0557 [math.NT] (2008).
77. ———, *The evaluation of Tornheim double sums. part 2*, *Raman. J.* **22** (2010), no. 1, 55–99. MR 2610609
78. Harold Exton, *Some new summation formulae for the generalised hypergeometric function of higher order*, *J. Comp. Appl. Math.* **79** (1997), no. 2, 183–187, E:[50]. MR 1450279
79. Valery I. Fabrikant, *Computation of infinite integrals involving three Bessel functions by introduction of a new formalism*, *Z. Angew. Math. Mech.* **83** (2003), no. 6, 363–374. MR 1986960
80. G. M. Fichtenholz, *Differential- und Integralrechnung*, vol. II, Deutscher Verlag der Wissenschaften, Berlin, 1964.
81. Jerry L. Fields and Jet Wimp, *Expansions of hypergeometric functions in series of other hypergeometric functions*, *Math. Comp.* **15** (1961), no. 76, 390–395. MR 23 #A3289
82. George Fikioris, *Table errata 634*, *Math. Comp.* **67** (1998), no. 224, 1753–1754. MR 1625064
83. C. Fox, *The solution of a moment problem*, *J. Lond. Math. Soc.* **s1-13** (1938), no. 1, 12–14.

84. J. M. Gandhi, *A conjectured representation of Genocchi numbers*, Amer. Math. Monthly **77** (1970), no. 5, 505. MR 1535914
85. George Gasper, *Products of terminating ${}_3F_2(1)$ series*, Pac. J. Math. **56** (1975), no. 1, 87–95. MR 0374508
86. M. L. Glasser, *Reduction formulas for multiple series*, Math. Comp. **28** (1974), no. 125, 265–266. MR 0328414
87. ———, *A sum of Bessel functions*, SIAM Rev. **22** (1980), no. 4, 508.
88. ———, *Problems and solutions*, SIAM Rev. **26** (1984), no. 2, 276–277.
89. Xavier Gourdon and Pascal Sebah, *The logarithmic constant: $\log 2$* , 2008, <http://numbers.computations.free.fr/Constants/Log2/log2.html>.
90. Boris Gourévitch and Jesús Guillera Goyanes, *Construction of binomial sums for π and polylogarithmic constants inspired by BBP formulas*, Appl. Math. E-Notes **7** (2007), 237–246.
91. I. Gradstein and I. Ryzhik, *Summen-, Produkt- und Integraltafeln*, 1st ed., Harri Deutsch, Thun, 1981. MR 0671418 (83i:00012)
92. R. L. Graham and John Riordan, *The solution of a certain recurrence*, Am. Math. Monthly **73** (1966), no. 5, 604–608. MR 0227029
93. I. P. Grant and H. M. Quiney, *A class of Bessel function integrals with application in particle physics*, J. Phys. A: Math. Gen. **26** (1993), no. 24, 7547–7562. MR 1257782
94. Jesus Guillera, *History of the formulas and algorithms for π* , arXiv:0807.0872 [math.HO] (2009).
95. Jesús Guillera and Jonathan Sondow, *Double integrals and infinite products for some classical constants via analytic continuations of Lerch's transcendent*, Raman. J. **16** (2008), no. 3, 247–270. MR 2429900
96. Robert A. Gustafson, *A generalization of Selberg's beta integral*, Bull. Am. Math. Soc. **22** (1990), no. 1, 97–105. MR 1001607
97. Hyuk Han and Seunghyn Seo, *Combinatorial proofs of inverse relations and log-concavity for Bessel numbers*, Eur. J. Combinat. **29** (2008), no. 7, 1544–1554. MR 2431746
98. Leetsch Charles Hsu and Evelyn L. Tan, *A refinement of de Bruyn's formula for $\sum a^k k^p$* , Fib. Quart. **38** (2000), no. 1, 56–59. MR 1738647 (2000k:11030)
99. Simon Hubbert and Stefan Müller, *Interpolation with circular basis functions*, Numer. Algorithms **42** (2006), 75–90. MR 2249568
100. Robert Israel, priv. commun., 4 January 2009.
101. ———, SeqFan List, 4 July 2009.
102. A. D. Jackson and L. C. Maximon, *Integrals of products of Bessel functions*, SIAM J. Math. Anal. **3** (1972), no. 3, 446–460. MR 0311958 (47 #520)
103. Augustus J. E. M. Janssen, *Zernike circle polynomials and infinite integrals involving the product of Bessel functions*, arXiv:1007.0667 [math-ph] (2010).
104. W. B. Jordan, *Problem 76-2, an infinite sum*, SIAM Rev. **26** (1984), no. 2, 276–278.
105. Per W. Karlsson, *Some reduction formulae for double power series and Kempé de Fériet functions*, Nederl. Akad. Wetensch. Indag. Math. **46** (1984), no. 1, 31–36. MR 0748975 (85h:3304)
106. Emrah Kilic and Dursun Tasci, *Factorizations and representations of the backward second-order linear recurrences*, J. Comput. Appl. Math. **201** (2007), no. 1, 182–197. MR 2293547 (2008e:65399)
107. Peter Kirschenhofer, *A note on alternating sums*, El. J. Combinat. **3** (1996), no. 2, #R7. MR 1392492
108. K. S. Kölbig, *Closed expressions for $\int_0^1 t^{-1} \log^{n-1} t \log^p(1-t) dt$* , Math. Comp. **39** (1982), no. 160, 647–654. MR 0669656
109. ———, *On the integral $\int_0^{\pi/2} \log^n \cos x \log^p \sin x dx$* , Math. Comp. **40** (1983), no. 162, 565–570. MR 0689472
110. ———, *Table errata 617*, Math. Comp. **64** (1995), no. 209, 449–460. MR 1270626
111. K. S. Kölbig and H. Scherb, *On a Hankel transform integral containing an exponential function and two Laguerre polynomials*, J. Comput. Appl. Math. **71** (1996), no. 2, 357–363. MR 1399902
112. K. S. Kölbig and W. Strampp, *Some infinite integrals with powers of logarithms and the complete Bell polynomials*, J. Comput. Appl. Math. **69** (1996), 39–47. MR 1391610

113. Tom H. Koornwinder, *Identities of nonterminating series by Zeilberger's algorithm*, arXiv:math.CA/9805010 (1998).
114. Ernst D. Krupnikov, *Table errata 601*, Math. Comp. **41** (1983), no. 164, 782–783. MR 0717727
115. Jaume Oliver Lafont, *Bbp binary and ternary formulas*, 7 June 2010, priv. commun.
116. ———, *Generalized mercator series*, 13 September 2011, priv. commun.
117. Adeline Lambert, *Table errata 628*, Math. Comp. **66** (1997), no. 217, 463–463. MR 1388890
118. L. J. Landau and N. J. Luswili, *Asymptotic expansion of a Bessel function integral using hypergeometric functions*, J. Comp. Appl. Math. **132** (2001), no. 2, 387–397. MR 1840636
119. Derrick Henry Lehmer, *Interesting series involving the central binomial coefficient*, Amer. Math. Monthly **92** (1985), no. 7, 449–457, E: [127]. MR 0801217 (87c:40002)
120. I. E. Leonard, *More on fresnel integrals*, Math. Monthly **95** (1988), no. 5, 431–433. MR 0937530
121. S. K. Lucas, *Approximations to π derived from integrals with nonnegative integrands*, Amer. Math. Monthly **116** (2009), no. 2, 166–172. MR 2478060
122. Wilhelm Magnus and Fritz Oberhettinger, *Formeln und Sätze für die speziellen Funktionen der Mathematischen Physik*, 2nd ed., Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, vol. 52, Springer, Berlin, Heidelberg, 1948.
123. Richard J. Mathar, *Mutual conversion of three flavors of Gaussian Type Orbitals*, Int. J. Quant. Chem. **90** (2002), no. 1, 227–243, E: [130].
124. ———, *Orthogonal set of basis functions over the binocular pupil*, arXiv:physics/0706.3682 [optics] (2007).
125. ———, *Karhunen-Loève basis functions of Kolmogorov turbulence in the sphere*, Baltic Astronomy **17** (2008), no. 3/4, 383–398, E: [131].
126. ———, *Series of reciprocal powers of k -almost primes*, arXiv:0803.0900 [math.NT] (2008).
127. ———, *Corrigenda to “interesting series involving the central binomial coefficient”* [Am. Math. Monthly vol 92 (1985)], arXiv:0905.0215 [math.CA] (2009).
128. ———, *A java Math.BigDecimal implementation of core mathematical functions*, arXiv:0908.3030 [math.NA] (2009).
129. ———, *Numerical evaluation of the oscillatory integral over $e^{i\pi x}x^{1/x}$ between 1 and infinity*, arXiv:0912.3844 [math.CA] (2009).
130. ———, *Erratum: Mutual conversion of three flavors of Gaussian Type Orbitals*, Int. J. Quant. Chem. **110** (2010), no. 4, 962.
131. ———, *Karhunen-Loève basis functions of Kolmogorov turbulence in the sphere (erratum)*, Baltic Astronomy **19** (2010), no. 1/2, 143–144.
132. ———, *Tightly circumscribed regular polygons*, arXiv:1301.6293 [math.MG] (2013).
133. Harry A. Mavromatis, *Two integrals involving products of two Bessel and a generalized hypergeometric function*, J. Comp. Appl. Math. **59** (1995), no. 3, 381–384. MR 1346420
134. Luis A. Medina and Victor H. Moll, *A class of logarithmic integrals*, Ramanujan J. **20** (2009), 91–125. MR 2546186
135. Luis A. Medina, Victor H. Moll, and Eric S. Rowland, *Iterated primitives of logarithmic powers*, arXiv:0911.1325 [math.NT] (2009).
136. R. Mehrem, J. T. Londergan, and M. H. Macfarlane, *Analytic expressions for integrals of products of spherical Bessel functions*, J. Phys. A: Math. Gen. **24** (1991), 1435–1453. MR 1121820
137. István Mező, *Summation of hyperharmonic series*, arXiv:0811.0042 [math.CO] (2008).
138. István Mező and Ayhan Dil, *Euler-seidel method for certain combinatorial numbers and a new characterization of fibonacci sequence*, Cent. Eur. J. Math **7** (2009), no. 2, 310–321. MR 2506968
139. Michael Milgram, *On hypergeometric $3f_2(1)$* , arXiv:math.CA/0603096 (2006).
140. Allen R. Miller, *Certain summation and transformation formulae for generalized hypergeometric series*, J. Comput. Appl. Math. **231** (2009), 964–972. MR 2549757
141. Allen R. Miller and R. B. Paris, *Euler-type transformations for the generalized hypergeometric function*, Z. Angew. Math. Phys. **62** (2010), no. 1, 31–45.
142. Jeff Miller and Victor S. Adamchik, *Derivatives of the Hurwitz zeta function for rational arguments*, J. Comput. Appl. Math. **100** (1998), no. 2, 201–206. MR 1569109
143. Victor H. Moll, *The evaluation of integrals: A personal story*, Not. AMS **49** (2002), no. 3. MR 1879857

144. ———, *The integrals in Gradshteyn and Ryzhik. part 1: A family of logarithmic integrals*, Scientia A **14** (2007), 1–6. MR 2330697
145. Bengt Nagel, *An expansion of the hypergeometric function in bessel functions*, J. Math. Phys. **42** (2001), no. 12, 5910. MR 1866696
146. A. A. R. Neves, L. A. Padilha, A. Fontes, E. Rodrigues, C. H. B. Cruz, L. C. Barbosa, and C. L. Cesar, *Analytical results for a Bessel function times Legendre polynomial class integrals*, J. Phys. A: Math. Gen. **39** (2006), no. 18, L293–L296. MR 2243193
147. Robert J. Noll, *Zernike polynomials and atmospheric turbulence*, J. Opt. Soc. Am. **66** (1976), no. 3, 207–211.
148. N. E. Nørlund, *Hypergeometric functions*, Acta Math. **94** (1955), no. 1, 289–349. MR 0074585
149. Fritz Oberhettinger, *Tabellen zur Fourier-Transformation*, Springer, Berlin, Heidelberg, 1957. MR 0081997
150. ———, *Tables of Bessel Transforms*, Springer, Berlin, Heidelberg, 1972. MR 0352888
151. R. B. Paris, *A kummer-type transformation of a ${}_2F_2$ hypergeometric function*, J. Comput. Appl. Math. **173** (2005), no. 2, 379–382. MR 2102904
152. P. J. E. Peebles, *Statistical analysis of catalogs of extragalactic objects. I. Theory*, Astroph. J. **185** (1973), 413–440.
153. Kh. Hessami Pilehrood and Tatiana Hessami Pilehrood, *Simultaneous generation for zeta values by the Markov-WZ method*, Discr. Math. Theor. Comp. Sci. **10** (2008), no. 3, 115–124. MR 2457055
154. Jean C. Piquette, *Table of special function integrals*, ACM SIGSAM Bulletin **24** (1990), no. 4, 8–21.
155. Alexander Povolotsky, priv. commun., 9 January 2009.
156. Helmut Prodinger and Wojciech Szpankowski, *A note on binomial recurrences arising in the analysis of algorithms*, Inf. Proc. Lett. **46** (1993), no. 6, 309–311. MR 1231833
157. V. Rajeswari and K. Srinivasa Rao, *Four sets of ${}_3F_1(1)$ functions, Hahn polynomials and recurrence relations for the $3-j$ coefficients*, J. Phys. A: Math. Gen. **22** (1989), no. 19, 4113–4123. MR 1016765
158. S. K. Rangarajan, *On “a new formula for $p_{m+n}^m(\cos \alpha)$ ”*, Quart. J. Math. **15** (1964), no. 1, 32–34. MR 0159968
159. M. A. Rashid, *Evaluation of integrals involving powers of $1 - x^2$ and two associated Legendre functions of Gegenbauer polynomials*, J. Phys. A: Math. Gen. **29** (1986), 2505–2512. MR 0857047 (87m:33009)
160. Arjun K. Rathie, Y. S. Yong Sup Kim, and Hyeong Kee Song, *Another method for a new two-term relation for the hypergeometric function ${}_3F_2$ due to Exton*, J. Comp. Appl. Math. **167** (2004), no. 2, 485–487. MR 2064704
161. John Riordan, *Combinatorial identities*, John Wiley, New York, 1968. MR 0231725 (38 #53)
162. Theodore J. Rivlin, *The Chebyshev Polynomials*, Pure and Applied Mathematics, John Wiley, New York, London, 1974. MR 0450850 (56 #9142)
163. Kelly Roach, *Hypergeometric function representations*, Proc. of the 1996 Intl. Symp. on Symbolic and algebraic computation (Y. N. Lakshman, ed.), ACM, 1996, pp. 301–308.
164. Klaus Rottbrand, *Finite-sum rules for Macdonald’s functions and Hankel’s symbols*, Int. Transf. Special Func. **10** (2000), no. 2, 115–124. MR 1812510
165. Klaus Rottbrand and Christian Weddigen, *Closed-form representations of the Laplace transforms of Bessel functions $j_{2n}(a\sqrt{t^2 + 2b})$* , Int. Transf. Special Func. **9** (2000), no. 3, 217–228. MR 1782973
166. Ranjan Roy, *Binomial identities and hypergeometric series*, Amer. Math. Monthly **94** (1987), no. 1, 36–46. MR 0873603 (88f:05012)
167. George Rutledge and R. D. Douglas, *Table of definite integrals*, Am. Math. Monthly **45** (1938), no. 8, 525–530. MR 1524383
168. Stefan G. Samko, *On some index relations for Bessel functions*, Integr. Transf. Spec. Func. **11** (2001), no. 4, 397–402. MR 1887291
169. Richard J. Sasiela and John D. Shelton, *Transverse spectral filtering and Mellin transform techniques applied to the effect of outer scale on tilt and tilt anisoplanatism*, J. Opt. Soc. Am. A **10** (1993), no. 4, 646–660.
170. Giovanni Mingari Scarpello and Daniele Ritelli, *On computing some special values of hypergeometric functions*, arXiv:1212.0251 (2012).

171. Glenn M. Schmiege, *Table errata 392.*, Math. Comp. **20** (1966), no. 95, 468–468.
172. E. B. Shanks, *Iterated sums of powers of the binomial coefficients*, Amer. Math. Monthly **58** (1951), no. 6, 404–407. MR 0047594
173. L. J. Slater, *Expansions of generalized Whittaker functions*, Proc. Camb. Phil. Soc. **50** (1954), no. 4, 628–631. MR 0064210
174. Lucy Joan Slater, *Generalized hypergeometric functions*, Cambridge University Press, 1966. MR 0201688
175. Neil J. A. Sloane, *The On-Line Encyclopedia Of Integer Sequences*, Notices Am. Math. Soc. **50** (2003), no. 8, 912–915, <http://oeis.org/>. MR 1992789 (2004f:11151)
176. Donald R. Snow, *Formulas for sums of powers of integers by functional equations*, Aequat. Math. **18** (1978), no. 1–2, 269–286. MR 0522512
177. Anthony Sofo, *Catalan related sums*, J. Inequal. Pure Appl. Math. **10** (2009), no. 3, # 69. MR 2551092
178. ———, *Integrals and polygamma representations for binomial sums*, J. Integer Seq. **13** (2010), # 10.2.8. MR 2594742
179. ———, *Extensions of Euler harmonic sums*, Applic. Anal. Discr. Math. **6** (2012), 317–328. MR 3012679
180. G. Solt, *Table errata 607.*, Math. Comput. **47** (1986), no. 176, 768–768. MR 0856719
181. Michael Z. Spivey, *The humble sum of remainders function*, Math. Mag. **78** (2005), no. 4, 300–305.
182. ———, *Combinatorial sums and finite differences*, Discr. Math. **307** (2007), no. 24, 3130–3146. MR 2370116 (2008j:05013)
183. H. M. Srivastava, *A certain class of q -series transformations*, J. Math. Anal. Applic. **107** (1985), no. 2, 498–508. MR 0787729
184. H. M. Srivastava, M. L. Glasser, and V. S. Adamchik, *Some definite integrals associated with the Riemann zeta function*, Zeitsch. f. Analys. Anwend. **19** (2000), no. 3, 831–846. MR 1784133
185. Elias M. Stein and Guido Weiss, *Introduction to fourier analysis in euclidean spaces*, Princeton University Press, Princeton, 1971. MR 0304972
186. Lajos Takács, *A sum of binomial coefficients*, Math. Comp. **32** (1978), no. 144, 1271–1273. MR 0502014
187. Münevver Tezer, *On the numerical evaluation of an oscillating infinite series*, J. Comp. Appl. Math. **28** (1989), 383–390. MR 1038858
188. E. C. Titchmarsh and D. R. Heath-Brown, *The theory of the Riemann zeta-function*, 2 ed., Oxford Science Publications, 1986. MR 0882550 (88c:11049)
189. Tiberiu Trif, *Combinatorial sums and series involving inverses of binomial coefficients*, Fib. Quart. **38** (2000), no. 1, 79–83. MR 1738651
190. Glenn A. Tyler, *Analysis of propagation through turbulence: evaluation of an integral involving the product of three Bessel functions*, J. Opt. Soc. Am. A **7** (1990), no. 7, 1218–1223.
191. Alfred J. van der Poorten and Xuan Chuong Tran, *Periodic continued fractions in elliptic function fields*, Lect. Not. Comp. Sci. **2369** (2002), 390–403. MR 2041099
192. H. van Haeringen and L. P. Kok, *Table errata 589*, Math. Comput. **39** (1982), no. 160, 747–757. MR 0669666
193. Raimundas Vidūnas, *A generalization of Kummer’s identity*, Rocky Mount. J. Math. **32** (2002), no. 2, 919–936. MR 1934920
194. ———, *Expressions for values of the gamma function*, arXiv:math.CA/0403510 (2004).
195. ———, *Darboux evaluations of algebraic gauss hypergeometric functions*, arXiv:math/0504264 [math.CA] (2005).
196. Niaz M. Vildanov, *Generalization of the Glasser - Manna - Oloa integral and some new integrals of similar type*, arXiv:1007.3460 [math.CA] (2010).
197. G. N. Watson, *A note on the polynomials of hermite and laguerre*, J. Lond. Math. Soc. **s1-13** (1938), 29–32.
198. ———, *A treatise on the theory of Bessel functions*, 2nd ed., Cambridge University Press, 1958.
199. Kurt Wegschaidter, *Computer generated proofs of binomial multi-sum identities*, Master’s thesis, Johannes Kepler Universität Linz, 1997.
200. F. J. W. Whipple, *Well-poised series and other generalized hypergeometric series*, Proc. Lond. Math. Soc. **25** (1926), no. 2, 525–544.

- 201. Kenneth S. Williams, *On Salié's sum*, J. Numb. Theory **3** (1971), 316–317. MR 0286742
- 202. James A. Wilson and O. G. Ruehr, *Identity for hypergeometric functions (problem 81-15)*, SIAM Rev. **24** (1982), no. 3, 346.
- 203. G. Milton Wing, *The mean convergence of orthogonal series*, Am. J. Math. **72** (1950), no. 4, 792–808. MR 0037923
- 204. D. M. Winker, *Effect of a finite outer scale on the Zernike decomposition of atmospheric optical turbulence*, J. Opt. Soc. Am. A **8** (1991), no. 10, 1568–1573, E: [169].
- 205. Roman Wituła and Damian Słota, *Partial fractions decompositions of some rational functions*, Appl. Math. Comput. **197** (2008), 328–336. MR 2396331
- 206. Jin-Hua Yang and Feng-Zhen Zhao, *Certain sums involving inverses of binomial coefficients and some integrals*, J. Integer Sequ. **10** (2007), no. 07.8.7. MR 2335861

URL: <http://www.mpia.de/~mathar>

E-mail address: mathar@mpia.de

HOESCHSTR. 7, 52372 KREUZAU, GERMANY