


# Pillar[5]arene glyco(mimetic)rotaxanes for the functional interrogation of multivalency responsive glycosidases

Iwona Nierengarten, Manuel González-Cuesta, Jaime Rodríguez-Guerra Pedregal, Uwe Hahn, Sonia Romero-Tellez, Jean-Didier Maréchal, Laura Masgrau, \* José Manuel García Fernández, \* Jean-François Nierengarten\* and Carmen Ortiz Mellet\*

**Abstract:** Multiconjugation of a pillar[5]arene scaffold and rotaxation of a glycomimetic-exposing axel has been exploited as a tactic to alter molecular volume and rigidity in hetero(multivalent) constructs purposely conceived to investigate the mechanisms at play in multivalent enzyme inhibition (MEI). With the help of a newly developed glycosidase binding mode screening tool, using reporter lectins and computational docking, the influence of the glycosidase tertiary and quaternary structure in MEI has been evidenced. The results provide a rational for the activation of biomimetic lectin-like associations in the intriguing case of multivalency-responsive glycosidases with deeply located catalytic pockets.

The observation that the inhibitory potency of the iminosugar-type glycomimetic 1-deoxynojirimycin (DNJ) towards the Jack bean  $\alpha$ -mannosidase (JbM) is enhanced by 2000-fold when displayed in twelve copies coating a C<sub>60</sub>-fullerene ball (**1**)<sup>[1]</sup> shook the foundations of the generally accepted key-and-lock model for enzyme inhibition by substrate analogues (Figure 1). The phenomenon, generically termed multivalent enzyme inhibition (MEI), has been further confirmed for a variety of glycomimetic architectures,<sup>[2,3]</sup> depicting a scenario that is reminiscent of that portraying sugars binding to their cognate protein receptors (lectins). The fact that the active site of JbM, a member of the glycosyl hydrolase (GH) family GH38 in the Carbohydrate Active Enzyme

(CAZy) classification,<sup>[4]</sup> is relatively open and that the enzyme is multimeric in solution reinforced this notion, which is consistent with recent X-ray evidence.<sup>[5]</sup> However, this thesis cannot apply for multivalency responsive enzymes that are either monomeric and/or have deeply buried catalytic pockets, which includes human enzymes of biological and therapeutic relevance such as lysosomal  $\beta$ -glucosidase,<sup>[6]</sup> hexosaminidases and O-glucosaminidase<sup>[7]</sup> or heparanase.<sup>[8]</sup> Most disturbing, sugar glycosides, the putative substrates of glycosidases, were found to turn into inhibitors when multivalently presented at the surface of nanomaterials, resulting in lectin/glycosidase binding promiscuity.<sup>[9]</sup> The impact of these discoveries in the glycoscience is palpable, promoting a reformulation of multivalency from a “safe” tactic to achieve biologically useful lectin affinities to a “multi-switch” tool with the potential to elicit lectin/glycosidase cross-talk behaviors.<sup>[10]</sup>



**Figure 1.** Structures of the iminosugar 1-deoxynojirimycin (DNJ) and the C<sub>60</sub> fullerene/DNJ dodecavalent conjugate **1**.

Advancing our understanding on the mechanisms underpinning MEI events beyond the  $\alpha$ -mannosidase case is critical to substantiate this paradigm shift and uncover the biological and technological repercussions. The inhibition of enzymes featuring narrow catalytic sites by bulky multivalent conjugates is a counterintuitive phenomenon whose interpretation requires the design of precise (macro)molecular probes and purposely-conceived binding site-informing screening tools. Herein we report a new class

[\*] Dr. I. Nierengarten, U. Hahn, Prof. J.-F. Nierengarten  
Laboratoire de Chimie des Matériaux Moléculaires  
Université de Strasbourg et CNRS (UMR 7509)  
Ecole Européenne de Chimie, Polymères et Matériaux  
25 rue Becquerel, 67087 Strasbourg, France  
E-mail: E-mail: [nierengarten@unistra.fr](mailto:nierengarten@unistra.fr)  
M. González-Cuesta, Dr. M. I. García Moreno, Prof. C. Ortiz Mellet  
Department of Organic Chemistry, Faculty of Chemistry  
University of Sevilla  
c/ Profesor García González 1, 41012 Sevilla, Spain  
E-mail: [mellet@us.es](mailto:mellet@us.es)  
J. Rodríguez-Guerra Pedregal, S. Romero-Tellez, Dr. J.-D.  
Maréchal, Dr. L. Masgrau  
Departament de Química, Universitat Autònoma de Barcelona,  
08193 Bellaterra, Spain  
E-mail: [Laura.Masgrau@uab.cat](mailto:Laura.Masgrau@uab.cat)  
Prof. J. M. García Fernández  
Instituto de Investigaciones Químicas (IIQ), CSIC– Universidad de  
Sevilla, Américo Vespucio 49, E-41092 Sevilla, Spain  
E-mail: [jogarcia@iiq.csic.es](mailto:jogarcia@iiq.csic.es)

Supporting information for this article is available on the WWW  
under <http://dx.doi.org/10.1002/anie.2011xxxxxx>.

