

# Canadian Journal of Forest Research Revue canadienne de recherche forestière

# Long-term compositional changes following partial disturbance revealed by the re-survey of logging concession limits in the northern temperate forest of eastern Canada

Journal:	Canadian Journal of Forest Research		
Manuscript ID	cjfr-2016-0047.R1		
Manuscript Type: Article			
Date Submitted by the Author: 12-Apr-2016			
Complete List of Authors:	Danneyrolles, Victor; Universite du Quebec en Abitibi Temiscamingue Arseneault, Dominique; Universite du Quebec a Rimouski Bergeron, Yves; Universite du Quebec en Abitibi Temiscamingue		
Keyword:	forest succession, historical ecology, ecosystem-based management, preindustrial forests, presettlement forests		

SCHOLARONE™ Manuscripts

- 1 Long-term compositional changes following partial disturbance revealed by the re-
- 2 survey of logging concession limits in the northern temperate forest of eastern Canada
- 3 Victor Danneyrolles Dominique Arseneault Yves Bergeron
- 4 **Danneyrolles, V.** (corresponding author, victor.danneyrolles@uqat.ca, 438-820-6903)
- 5 <sup>2</sup>Arseneault, D. (dominique arseneault@uqar.ca)
- 6 <sup>1</sup>Bergeron, Y. (yves.bergeron@uqat.ca)
- 7 <sup>1</sup> Centre d'étude de la forêt (CEF) and Chaire industrielle CRSNG-UQAT-UQAM en
- 8 Aménagement Forestier Durable, Université du Québec en Abitibi-Témiscamingue, 445
- 9 boulevard de l'Université, Rouyn-Noranda, QC J9X 5E4, Canada
- <sup>2</sup> Groupe BOREAS, Centre d'étude de la forêt (CEF) and Chaire de Recherche sur la Forêt
- Habitée, Université du Québec à Rimouski, Rimouski, QC G5L 3A1, Canada
- 12 Manuscript word count: 4860 words

#### Abstract

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

Land use changes that are linked to European settlement of North America have transformed north-eastern temperate forest landscapes. Many studies report a regional increase of young early-successional forests due to high disturbance rates since the preindustrial era (fire, landclearing, clear-cuts). In this study, we document specific compositional changes to present day mature forest landscapes, which have only been managed with partial cutting (highgrading and diameter-limit cuts) since the preindustrial era in south-western Quebec. We resurveyed 108 forest observations that were extracted from logbooks of former logging concession limits (surveyed between 1870 and 1890). Results highlight an increase in midto late-successional shade-tolerant taxa (Betula alleghaniensis, Thuja occidentalis, Acer saccharum) at the expense of preindustrial dominant conifers (Abies balsamea, Pinus strobus). Former logging activities and spruce budworm outbreaks appear to be the main drivers of these changes, which were also strongly structured across the topographic gradient. To some extent, these results highlight the relevance of partial cutting management since it has allowed long-term maintenance of a mid- to late-successional forest composition, while also pointing the need for *Pinus strobus* restoration. We conclude that by allowing sitespecific comparisons, the re-survey of historical observations greatly improve the analytical strengths of historical reconstruction.

- 31 **Keywords**: Forest succession, historical ecology, ecosystem-based management,
- 32 preindustrial forests, presettlement forests.

#### Résumé

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

La colonisation européenne de l'est de l'Amérique du Nord a engendré une profonde transformation des paysages forestiers. De nombreuses études ont décrit une augmentation des jeunes forêts de début de succession comme conséquence des forts taux de perturbation liés à la colonisation (feux, défrichement, coupes à blanc). Dans cette étude, nous documentons les changements de composition spécifiques à des paysages forestiers aujourd'hui matures, qui n'ont été exploités que par des coupes partielles (coupes d'écrémage et coupes à diamètre limite) depuis l'époque préindustrielle. Nous avons réarpenté 108 observations de composition issues de rapports d'arpentage d'anciennes limites de concessions forestières (arpentées entre 1870 et 1890). Les résultats indiquent une augmentation de taxons de milieux à fin de succession (Betula alleghaniensis, Thuja occidentalis, Acer saccharum) au détriment des conifères dominant à l'époque préindustrielle (Abies balsamea, Pinus strobus). Ces changements de composition sont principalement la conséquence des coupes et des épidémies de tordeuse des bourgeons de l'épinette, et se sont aussi largement structurés le long du gradient topographique. Ceci témoigne de la pertinence de l'aménagement par coupes partielles, puisqu'il a permis le maintient d'une composition de milieux à fin de succession. Cependant, la restauration des populations de *Pinus strobus* parait aussi nécessaire. Nous concluons qu'en permettant la comparaison de sites appariés, le réarpentage d'observations historiques améliore remarquablement la compréhension des changements de composition survenus depuis l'époque préindustrielle.

- Mots clefs: Succession forestière, écologie historique, aménagement écosystémique, forêts
- 54 préindustrielles, forêts précoloniales.

#### Introduction

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

Global change is responsible for major shifts in forest ecosystems characteristics. In northeastern North America, land use changes that are linked to European settlement have transformed forest landscapes (Whitney 1994). Modern landscapes are generally considered to be younger, and composed of an increased proportion of early-successional species compared to preindustrial landscapes (Mladenoff et al. 1993; Foster et al. 1998; Dupuis et al. 2011; Thompson et al. 2013). Knowledge of forest composition prior to Euro-American settlement is therefore used as a reference for restoration and forest management (Egan & Howell 2001; Foster et al. 2003). In northern temperate forests, several studies have reported that preindustrial landscapes were dominated by old uneven-aged forests (Lorimer & White 2003; Boucher et al. 2009), the dynamics of which were mainly controlled by natural partial disturbances (gaps, insect outbreaks). In order to maintain forest landscapes within their range of natural variability, ecosystem-based management of these forests should be based mainly upon different types of partial cutting (Seymour et al. 2002; Raymond et al. 2009). In south-western Quebec, a substantial proportion of the landscape has only been managed through partial cutting (highgrading and diameter-limit cuts) since the preindustrial era. These present day mature forest landscapes appear to have followed distinct post-industrial compositional trajectories, compared with those of highly disturbed landscapes, which were converted into younger early-successional forests (Danneyrolles et al. 2016). The knowledge of long-term compositional changes after these types of partial cutting management could help in assessing the relevance of such practices, and in developing management tools adapted to these mature forest landscapes. Reconstructions of post-industrial compositional changes that are methodologically based on regional-scale comparisons between historical data-sets (i.e. historical land survey data) and

modern forest inventories commonly reveal general trends (eg. Friedman & Reich 2005; Thompson et al. 2013). However, these reconstructions are not sufficiently reliable to properly characterize more precise compositional changes that concern only specific landscapes. Since the main objective of this study was to document specific compositional changes that occurred in landscapes having only been managed by partial cutting, our reconstruction is based on the re-survey of historical observations (i.e. historical and modern observations that were paired at the same location). Yet, few studies have used this original methodological approach (Siccama 1971; Fahey & Lorimer 2014), even though it may allow a deeper understanding of long-term compositional changes. More particularly, we addressed two main questions: (1) what were the main drivers of these compositional changes? (2) How have environmental gradients, particularly the topographic gradient, structured these compositional changes?

### Study area

The study covers an area of about 4000 km² along the south-west boundary of the Canadian province of Quebec (Fig. 1). The forests in the study area form the northern end of the Great Lakes St. Lawrence mixed forest region (Rowe 1972), which also corresponds to the limits between the maple-yellow birch and balsam fir-yellow birch bioclimatic domains of the provincial classification (Robitaille & Saucier 1998). The principal surface deposits are either undifferentiated or rocky glacial till. The closest weather station (Barrage Témiscamingue, 46°42'N, 76°06'W; 181 m asl.) has recorded a mean annual temperature of 4.9°C and annual total precipitation of 937 mm for the period 1981-2010.

Elevation varies from 290m to 415m (asl.). Several factors that are related to topography influence forest composition (drainage, nutrient availability, soil depth and micro-climate; MacHattie & McCormack 1961). Natural stand-replacing fire rotation has been estimated to

be about 200 years in duration within the balsam fir-yellow birch domain (Grenier et al. 2005), and 500 years within the maple-yellow birch domain (Drever et al. 2006). Outbreaks of spruce budworm (*Choristoneura fumiferana*) also have been identified as a major source of disturbance of forest dynamics over the last century (Bouchard et al. 2006a,b).

First Nations have inhabited the study area for at least 5000 years (Riopel 2002). However, patterns of land use and management by Algonquin tribes during preindustrial times remain unknown. Logging did not begin until 1840, and was mainly focused on high-grading cutting of tall pines trees (*Pinus* spp.) until 1917 (Riopel 2002). Following the construction of a paper mill in the city of Témiscaming in 1917, diameter-limit cuts of smaller diameter spruces (*Picea* spp.) and balsam fir (*Abies balsamea* [L.] Mill.) for wood pulp became the main logging activity until the end of the 20<sup>th</sup> century (Lienert 1966), although high-grading cutting of tall trees for lumber remained frequent.

#### Material and methods

# Site selection and field measurements

This study is based on the re-survey of 108 historical observations of forest composition that were extracted from 10 logbooks reporting the first survey of logging concession boundaries, which were realized by four different surveyors between 1870 and 1890. These logbooks contain point observations mentioning taxa lists (e.g., "Pine, spruce, birch, cedar and a few maples"), which were usually 10 chains (about 200 m) apart along the survey lines. Point observations have been geo-referenced using historical maps with a resolution of about 20 meters. These 10 logbooks were selected for their consistent methodology.

A total of 108 observations that were to be re-surveyed had been selected according to four criteria. (1) Observations had to be spaced at least 200 m apart to minimize spatial auto-correlation. (2) To restrict sampling to the most ecologically accurate observations, only taxa

lists mentioning at least two taxa were retained (90 % of selected observations mentioned
three to five taxa). (3) To document dynamic trajectories of forests that had not experienced
stand-replacing disturbance since preindustrial times, observations that were retained had to
be located in present-day mature forests. These forests corresponded to a dominant age class
of 90 year, 120 year, or as uneven-aged in the modern forest maps (Berger 2008). Their old
ages had to be confirmed in the field through the presence of old trees and irregular
structure. (4) Only observations that were located within 4 km of the road network were
finally retained.

In the field, each observation point was registered using a GPS with a 10 m precision. These locations determined the centres of 11.28 m diameter circular plots of (400 m²). Within each plot, all stems greater than 10 cm in diameter at breast height (DBH, 1.3 m) were inventoried at the species level, and their DBH measured. Evidence of former cuts (stumps) or spruce budworm outbreaks (numerous standing or lying dead stems) were recorded if they were recognizable either inside of or in proximity to the plots.

#### Data-base construction and analysis

For all historical and modern observations, taxa were classified according to 5 abundance ranks (r). Despite the possible existence of bias in these types of ranked list, for example giving more prominence to highly visible or commercially valuable taxa, a recent study demonstrated that taxa positions within lists are well correlated with their relative basal area (Terrail et al. 2014). Thus, for historical observations, taxa were ranked according to their position in the taxa list. The dominant taxon (first position in the list) was ranked as 4; the following two taxa in the list were ranked as 3 and 2, and rank 1 was assigned to the remaining taxa in the list. For example, the observation "Pine, spruce, yellow birch, cedar and a few maples" would have been transcribed as pines (r = 4), spruces (r = 3), yellow

birch (r = 2), cedar (r = 1), maples (r = 1) and (r = 0) for non-cited taxa. The same process was used to assign ranks to taxa that were inventoried in the re-surveyed plots, according to their relative basal area. Some species from the modern plots were grouped at the genus level to match the taxa that were mentioned by surveyors, while rarely mentioned taxa (<5%) within historical and modern observations were grouped in as "others". Taxa that represented less than 5% of the total basal area of a plot were removed from the list, to obtain an equivalent number of taxa for historical and modern observations.

Several simple metrics were used to document changes in forest composition. First, frequency and dominance indices were computed (Scull & Richardson 2007; Dupuis et al. 2011). The frequency index represents the proportion of taxon occurrences regardless of rank (i.e.,  $r \neq 0$ ), while the dominance index represents the proportion of taxon dominance (i.e., r = 4). These indices were computed for each taxon and for each period, within the whole data-set and for three topographic positions (lower slopes, middle slopes and upper slopes; Fig. 1). Second, a simple relative rank difference ( $D_i$ ) was computed according to the formula:

$$D_i = R1880_{ip} - R2014_{ip}$$

where  $R1880_{ip}$  corresponds to the rank of taxon i within plot p during the period 1870-1890, and  $R2014_{ip}$  is the rank of taxon i within plot p in 2014. Paired sample tests (Wilcoxon signed-rank tests) were used to assess the significance of rank differences ( $R1880_{ip}$  vs.  $R2014_{ip}$ ) for the entire data-set and by topographic position.

#### Results

During the preindustrial period (1870-1890), the 108 stands were dominated by balsam fir (*A. balsamea*), pines (*Pinus* spp.) and yellow birch (*Betula alleghaniensis* Britten) with dominance indices of 46 %, 17 % and 17 %, respectively (Table 1). Although less dominant,

eastern white cedar (Thuja occidentalis L.), spruces (Picea spp.) and paper birch (Betula
papyrifera Marshall) were frequent (41 %; 37 % and 28 %, respectively; Table 1). Evidence
of former partial cuts (stumps) and spruce budworm outbreaks (numerous standing and lying
dead stems) were found in all stands. Unfortunately, an obvious distinction between these
two kinds of disturbance was not always possible, and most stands had likely been subjected
to both kinds of disturbance during the 20 <sup>th</sup> century.
These stands have experienced major composition changes since the end of the 19 <sup>th</sup> century.
Overall balsam fir dominance decreased from 46 % to 7 % (Table 1), while its ranking
within plots significantly decreased by an average of 1.6 ranks (Fig. 2). Balsam fir
preindustrial dominance was high at the three topographic positions (Fig. 3) but its ranks
mostly decreased at mid- and upper slope positions (Fig. 4). Consequently, balsam fir has
maintained a greater dominance on lower slopes (Fig. 3), with a high density of small stems
(Fig. 5). Despite these changes, balsam fir is still one of the most frequent taxa in the study
region, with an overall frequency of occurrence greater than 50% (Table 1).
The taxon "pines" corresponds mainly to white pine (Pinus strobus L.; Table 2). This taxon
experienced a sharp decrease in dominance (from 17 % to 3 %) and frequency (57 % to 7 %;
Table 1) since preindustrial times. Pine rankings within plots significantly decreased for the
entire data-set (Fig. 2) at all topographic positions (Fig. 4), especially on lower slopes where
pine was very dominant during preindustrial times.
Overall yellow birch dominance increased from 17% to 37%, while its frequency also
increased from 62 % to 83 % (Table 1). Its rank within plots significantly increased for the
whole data set (Fig. 2), particularly on lower and mid-slopes where it was less abundant
during preindustrial times (Fig. 3 and 4). Yellow birch is currently characterized by low
densities of large stems (Fig. 5). Its dominance and frequency are constant along the

- 200 topographic gradient (Fig. 3).
- Dominance and frequency of white cedar have increased from 8 % to 28 % and from 41 % to
- 202 60 % respectively (Table 1). White cedar ranking within plots has significantly increased for
- the entire data-set (Fig. 3), especially on lower and middle slopes positions (Fig. 4), where
- 204 cedar is currently the most abundant taxon.
- The taxon "maples" corresponds mainly to sugar maple (*Acer saccharum* Marshall; Table 2).
- This taxon experienced the strongest increase in dominance (from 2 % to 19 %) and
- frequency (7 % to 76 %; Table 1). Maple ranking within plots significantly increased for the
- 208 whole data-set (Fig. 2), especially on middle and upper slopes (Fig. 4) where maples
- 209 currently are remarkably abundant.
- 210 Spruces and paper birch have not experienced significant changes in their abundance in the
- whole data-set (Fig. 2) and at all topographic positions, except for spruces, which decreased
- significantly on lower slopes (Fig. 3).

# Discussion

- Our results attest to major compositional shifts in mature stands over the last 130 years.
- 215 Industrial logging started towards the end of the 19th century with high-grading cutting of
- tall pines, then intensified from 1917 (Témiscaming paper mill construction) onward with
- diameter-limit cutting of smaller spruces and balsam fir (Lienert 1966; Riopel 2002). These
- former management practices were mainly focused on harvest, and did not aim to favour
- 219 particular stand characteristics. Intensive exploitation of pines was undoubtedly the main
- factor explaining their disappearance from these stands. Diameter-limit cuts at the beginning
- of the 20th century may also explain, in part, the decline in balsam fir abundances. The
- 222 widespread mortality in balsam fir populations caused by the two major spruce budworm
- 223 outbreaks of the 20th century (1910-1930 and 1970-1990; Bouchard et al. 2006a,b) has been

superimposed on the impacts of logging activities. Surprisingly, the abundance of spruces
has not significantly decreased in these stands (except on lower slopes), while they
represented the main target species for the paper industry. This likely reflect the fact that
spruces were not particularly dominant in these stands in preindustrial times, and thus their
diminution is not as marked as for balsam fir or pines.
Parallel with the decrease in abundance of balsam fir and pines, our results highlight a strong
increase in mid- to late-successional shade-tolerant taxa (yellow birch, white cedar, sugar
maple). As such, our results contrast with the increase in early-successional intolerant taxa
(eg. poplars, paper birch) that has been documented in highly disturbed area within the
region (Pinto et al. 2008; Danneyrolles et al. 2016) and more generally in north-eastern
North America (eg. Friedman & Reich 2005; Boucher et al. 2006; Bouchard et al. 2006a;
Dupuis et al. 2011; Thompson et al. 2013). Canopy openings created by partial cuts and
spruce budworm outbreaks are clearly the main driver that favoured the recruitment of
shade-tolerant taxa (Heitzman et al. 1997; Gasser et al. 2010; Duchesne & Prévost 2013;
Larouche & Ruel 2015). Moreover, the large amount of stumps and rotten wood generated
by these disturbances may have favoured the regeneration of yellow birch (Lambert et al.
2016) and sugar maple (Caspersen & Saprunoff 2005).
The compositional changes were also significantly structured across the topographic
gradient. Lower slopes remained dominated by conifers (white cedar, balsam fir, spruces),
while upper slopes became dominated by hardwood species (sugar maple and yellow birch).
These divergent trajectories could be explained by distinct site characteristics. First, the
colder and moister micro-climate of poorly drained lower slopes may have favoured the
increase in white cedar, where it can multiply by layering (Hofmeyer et al. 2009), while also
allowing balsam fir and spruces to maintain an important abundance. Conversely, well-
drained upper slopes, which lie outside of cold air drainage areas, were particularly

favourable to the development of sugar maple (Barras & Kellman 1998). A second possible explanation for these divergent trajectories is that lower slopes were dominated by conifers in preindustrial times, while yellow birch dominated a large proportion of upper slopes. Thus, the greater abundance of deciduous litter could have promoted the establishment of sugar maples at mid- and upper slope positions (Barras et Kellman 1998; Caspersen et Saprunoff 2005), while conifer-dominated lower slopes have been more resistant to the development of hardwoods.

#### **Conclusions**

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

Our study documents the long-term cumulative effects of partial cutting management and spruce budworm outbreak on north-eastern temperate forest composition. Although these former management practices (high-grading and diameter-limit cuts) focused on harvest and did not aim to favour particular stand characteristics, they allowed mid- to late-successional forest composition to persist. Thus, to some extent, our results highlight the relevance of partial cutting management in north-eastern temperate forests. Nevertheless, these former management practices have led to significant compositional changes, which bring out an important avenue for restoration and management of these mature forest landscapes. Particularly, white pine populations that had been present in the area for thousands of years (Liu 1990), and which were largely removed by the early 20th century logging activities, should be restored. Stands and seed-trees preservation could ensure the establishment of white pine trees in adjacent cuts areas (Uprety et al. 2013). However, the strong expansion of sugar maple have led to stable deciduous stands, given that maple litter tends to inhibit regeneration of conifers (Barras & Kellman 1998). Thus, these stands should be carefully managed in order to maintain or reintroduce white pine. Irregular shelterwood systems can allow the maintenance of mid-tolerant species, such as white pine or yellow birch, in uneven-aged stands dominated by shade-tolerant species

274	(Raymond et al. 2009; Suffice et al. 2015). Moreover, enrichment planting within patch cuts
275	(Fahey & Lorimer 2013; Hébert et al. 2013) could be used to reintroduce white pine in stand
276	from which it was completely removed.
277	More broadly, our study also demonstrate the analytical strength of site-specific comparisons
278	through the resurvey of historical observations (Fahey & Lorimer 2014). This
279	methodological approach was highly relevant to characterize compositional changes with
280	precise spatial resolution and allowed us to distinguish different dynamics trajectories across
281	the topographic gradient. Such an approach can be used to complement regional-scale
282	comparisons between historical land survey data and modern forest inventories (eg.
283	Friedman & Reich 2005; Thompson et al. 2013) in order to refine our knowledge of post-
284	industrial dynamics trajectories across disturbance or environmental gradients.

### Acknowledgements

285

286

287

288

289

290

291

292

293

294

We thank Sébastien Dupuis for his help in constructing the geo-referenced database from the survey records, Chloé Bilodeau for her invaluable help with field-work, and William F.J. Parsons for carefully editing the manuscript. This project was financially supported by the Natural Science and Engineering Research Council of Canada (NSERC), by the Fonds de recherche du Québec Nature et Technologies (FRQNT), by the forest product company TEMBEC Inc. (Témiscaming, QC), by the Conference régionale des élus de l'Abitibi-Témiscamingue (CREAT) and by the Ministère des Forêts, de la Faune et des Parcs of Québec (MFFP).

### References

- Barras, N., & Kellman, M. 1998. The supply of regeneration micro-sites and segregation of tree species in a hardwood/boreal forest transition zone. *Journal of Biogeography* 25: 871–881.
- Berger, J.P. (Coordinator) 2008. Norme de stratification écoforestière. Quatrième inventaire écoforestier. Ministère des Ressources naturelles et de la Faune du Québec, Québec,

Canada.

- Bouchard, M., Kneeshaw, D., & Bergeron, Y. 2006a. Forest dynamics after successive spruce budworm outbreaks in mixedwood forests. *Ecology* 87: 2319–2329.
- Bouchard, M., Kneeshaw, D., & Bergeron, Y. 2006b. Tree recruitment pulses and long-term species coexistence in mixed forests of western Québec. *Ecoscience* 13: 82–88.
- Boucher, Y., Arseneault, D., & Sirois, L. 2006. Logging-induced change (1930-2002) of a preindustrial landscape at the northern range limit of northern hardwoods, eastern Canada. *Canadian Journal of Forest Research* 36: 505–517.
- Boucher, Y., Arseneault, D., Sirois, L., & Blais, L. 2009. Logging pattern and landscape changes over the last century at the boreal and deciduous forest transition in Eastern Canada. *Landscape Ecology* 24: 171–184.
- Caspersen, J.P., & Saprunoff, M. 2005. Seedling recruitment in a northern temperate forest: the relative importance of supply and establishment limitation. *Canadian Journal of Forest Research* 35: 978–989.
- Danneyrolles, V., Arseneault, D., & Bergeron, Y. 2016. Pre-industrial landscape composition patterns and post-industrial changes at the temperate-boreal forest interface in western Quebec, Canada. *Journal of Vegetation Science*. doi: 10.1111/jvs.12373
- Drever, C.R., Messier, C., Bergeron, Y., & Doyon, F. 2006. Fire and canopy species composition in the Great Lakes-St. Lawrence forest of Témiscamingue, Québec. *Forest Ecology and Management* 231: 27–37.
- Duchesne, L., & Prévost, M. 2013. Canopy disturbance and intertree competition: implications for tree growth and recruitment in two yellow birch–conifer stands in Quebec, Canada. *Journal of Forest Research* 18: 168–178.
- Dupuis, S., Arseneault, D., & Sirois, L. 2011. Change from pre-settlement to present-day forest composition reconstructed from early land survey records in eastern Québec, Canada. *Journal of Vegetation Science* 22: 564–575.
- Egan, D., & Howell, E.A. 2001. *The historical ecology handbook: a restorationist's guide to reference ecosystems*. Island Press, Washington, D.C.
- Fahey, R.T., & Lorimer, C.G. 2013. Restoring a midtolerant pine species as a component of late-successional forests: Results of gap-based planting trials. *Forest Ecology and Management* 292: 139–149.
- Fahey, R.T., & Lorimer, C.G. 2014. Habitat associations and 150 years of compositional change in white pine-hemlock-hardwood forests based on resurvey of public land survey corners. *The Journal of the Torrey Botanical Society* 141: 277–293.
- Foster, D.R., Motzkin, G., & Slater, B. 1998. Land-Use History as Long-Term Broad-Scale Disturbance: Regional Forest Dynamics in Central New England. *Ecosystems* 1: 96–119.
- Foster, D.R., Swanson, F., Aber, J., Burke, I., Brokaw, N., Tilman, D., & Knapp, A. 2003. The

- Importance of Land-Use Legacies to Ecology and Conservation. *BioScience* 53: 77–88.
- Friedman, S.K., & Reich, P.B. 2005. Regional legacies of logging: departure from presettlement forest conditions in northern Minnesota. *Ecological Applications* 15: 726–744.
- Gasser, D., Messier, C., Beaudet, M., & Lechowicz, M.J. 2010. Sugar maple and yellow birch regeneration in response to canopy opening, liming and vegetation control in a temperate deciduous forest of Quebec. *Forest Ecology and Management* 259: 2006–2014.
- Grenier, D.J., Bergeron, Y., Kneeshaw, D., & Gauthier, S. 2005. Fire frequency for the transitional mixedwood forest of Timiskaming, Quebec, Canada. *Canadian Journal of Forest Research* 35: 656–666.
- Hébert, F., Roy, V., Auger, I., & Gauthier, M.-M. 2013. White spruce (*Picea glauca*) restoration in temperate mixedwood stands using patch cuts and enrichment planting. *The Forestry Chronicle* 89: 392–400.
- Heitzman, E., Pregitzer, K.S., & Miller, R.O. 1997. Origin and early development of northern white-cedar stands in northern Michigan. *Canadian Journal of Forest Research* 27: 1953–1961.
- Hofmeyer, P.V., Kenefic, L.S., & Seymour, R.S. 2009. Northern white-cedar ecology and silviculture in the northeastern united states and southeastern Canada: a synthesis of knowledge. *Northern Journal of Applied Forestry* 26: 21–27.
- Lambert, J.-B., Ameztegui, A., Delagrange, S., & Messier, C. 2016. Birch and conifer deadwood favour early establishment and shade tolerance in yellow birch juveniles growing in sugar maple dominated stands. *Canadian Journal of Forest Research* 46: 114–121.
- Larouche, C., & Ruel, J.-C. 2015. Development of Northern White-Cedar Regeneration Following Partial Cutting, with and without Deer Browsing. *Forests* 6: 344–359.
- Lienert, A. 1966. *The story of the (Kipawa) Noranda woods division*. Canadian Paper International, Rouyn-Noranda, Canada.
- Liu, K.-B. 1990. Holocene paleoecology of the boreal forest and Great Lakes-St. Lawrence forest in northern Ontario. *Ecological Monographs* 60: 179.
- Lorimer, C.G., & White, A.S. 2003. Scale and frequency of natural disturbances in the northeastern US: implications for early successional forest habitats and regional age distributions. *Forest Ecology and Management* 185: 41–64.
- MacHattie, L.B., & McCormack, R.J. 1961. Forest Microclimate: A Topographic Study in Ontario. *Journal of Ecology* 49: 301–323.
- Mladenoff, D.J., White, M.A., Pastor, J., & Crow, T.R. 1993. Comparing Spatial Pattern in Unaltered Old-Growth and Disturbed Forest Landscapes. *Ecological Applications* 3: 294–306.

- Pinto, F., Romaniuk, S., & Ferguson, M. 2008. Changes to preindustrial forest tree composition in central and northeastern Ontario, Canada. *Canadian Journal of Forest Research* 38: 1842–1854.
- Raymond, P., Bédard, S., Roy, V., Larouche, C., & Tremblay, S. 2009. The irregular shelterwood system: review, classification, and potential application to forests affected by partial disturbances. *Journal of Forestry* 107: 405–413.
- Riopel, M. 2002. Le Témiscamingue: son histoire et ses habitants. Fides, Saint-Laurent, Canada.
- Robitaille, A., & Saucier, J.-P. 1998. *Paysages régionaux du Québec méridional*. Gouvernement du Québec, Ministère des ressources naturelles, Québec.
- Rowe, J.S. 1972. Forest regions of Canada. Fisheries and Environment Canada, Canadian Forest Service, Ottawa.
- Scull, P., & Richardson, J.L. 2007. A method to use ranked timber observations to perform forest composition reconstructions from land survey data. *American Midland Naturalist* 158: 446–460.
- Seymour, R.S., White, A.S., & de Maynadier, P.G. 2002. Natural disturbance regimes in northeastern North America—evaluating silvicultural systems using natural scales and frequencies. *Forest Ecology and Management* 155: 357–367.
- Siccama, T.G. 1971. Presettlement and Present Forest Vegetation in Northern Vermont with Special Reference to Chittenden County. *American Midland Naturalist* 85: 153–172.
- Suffice, P., Joanisse, G., Imbeau, L., Mazerolle, M.J., & Lessard, G. 2015. Short-term effects of irregular shelterwood cutting on yellow birch regeneration and habitat use by snowshoe hare. *Forest Ecology and Management* 354: 160–169.
- Terrail, R., Arseneault, D., Fortin, M.-J., Dupuis, S., & Boucher, Y. 2014. An early forest inventory indicates high accuracy of forest composition data in pre-settlement land survey records. *Journal of Vegetation Science* 25: 691–702.
- Thompson, J.R., Carpenter, D.N., Cogbill, C.V., & Foster, D.R. 2013. Four centuries of change in northeastern united states forests. *PLoS ONE* 8: e72540.
- Uprety, Y., Asselin, H., Bergeron, Y., & Mazerolle, M.J. 2013. White pine (Pinus strobus L.) regeneration dynamics at the species' northern limit of continuous distribution. *New Forests* 45: 131–147.
- Whitney, G.G. 1994. From coastal wilderness to fruited plain: a history of environmental change in temperate North America, 1500 to the present. Cambridge University Press, Cambridge; New York.

**Table 1.** Dominance and frequency indices of each taxon in the 1870-1890 preindustrial period and in 2014 for the whole data-set (n = 108).

	1870-1890		2	2014	
TAXA	Dom.	Freq.	Dom.	Freq.	
Balsam fir	45.9	87.2	7.3	51.4	
Pines	16.5	56.9	2.8	7.3	
Spruces	7.3	37.6	2.8	32.1	
W. cedar	8.3	41.3	27.5	59.6	
Y. birch	17.4	62.4	36.7	82.6	
Maples	1.8	7.3	19.3	51.4	
P. birch	2.8	25.5	1.8	29.4	
Others	0.0	6.4	1.8	12.8	



**Table 2.** Details of species of pines, spruces and maples that were cited by surveyors (1870-1890) and identified on the field (2014). In many cases, surveyors cited these taxa at the genus level, so the data have been regrouped by genus level for the remaining analyses.

3	U	I
J	v	

298

299

	1870-1890		2	2014	
TAXA	Dom.	Freq.	Dom.	Freq.	
white pine	16.5	27.5	2.8	7.3	
red pine	X	X	X	0.9	
pines	X	29.4	X	X	
ALL pines	16.5	56.9	2.8	7.3	
sugar maple	X	X	19.3	40.7	
red maple	X	X	X	30.6	
maples	1.8	7.3	X	X	
ALL maples	1.8	7.3	19.3	51.3	
white spruce black spruce spruces	X	X	0.9	28.7	
	X	X	1.9	8.3	
	7.3	37.6	X	X	
ALL spruces	7.3	37.6	2.8	32.1	




Figure 1. Study area, localization of re-surveyed transect portions, and illustration of the three topographic positions.






Figure 2. Relative rank difference  $(D_i)$  of each taxon for the whole data-set (n = 108).

The asterisks indicate significant (P < 0.001) rank differences (Wilcoxon signed-rank test).




**Figure 3.** Dominance (dark grey) and frequency (light grey) of each taxon during the 1870-1890 period and in 2014, for lower slope (low; n = 31), mid-slope (mid; n = 46) and upper slope (upp; n = 31) positions.

308



**Figure 4.** Relative rank difference  $(D_i)$  of each taxa for lower slope (low; n = 31), mid-slope (mid; n = 46) and upper slope (upp; n = 31) positions. The asterisks indicate significant (P < 0.01) rank differences (Wilcoxon signed-rank tests).

311



313 Figure 5. Overall structural characteristics of present-day forests (total), and of the

four currently most abundant taxa by topographic position.