

Type I and Type II Errors

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

Decision	Reality	
	H_0 True ($\mu = 66.7$)	H_0 False ($\mu > 66.7$)
Reject H_0 (Decide $\mu > 66.7$)	Type I error	Correct
Don't reject H_0 (Decide $\mu = 66.7$)	Correct	Type II error

Type I and Type II Errors

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

Decision	Reality	
	H_0 True ($\mu = 66.7$)	H_0 False ($\mu > 66.7$)
Reject H_0 (Decide $\mu > 66.7$)	Type I error	Correct
Don't reject H_0 (Decide $\mu = 66.7$)	Correct	Type II error

Type I and Type II Errors

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

Decision	Reality	
	H_0 True ($\mu = 66.7$)	H_0 False ($\mu > 66.7$)
Reject H_0 (Decide $\mu > 66.7$)	Type I error	Correct
Don't reject H_0 (Decide $\mu = 66.7$)	Correct	Type II error

Finding the Type II Error Probabilities

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

Finding the Type II Error Probabilities

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

Finding the Type II Error Probabilities

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

For $\alpha = 0.05$: $k_\alpha = 68.26$

Decision rule: Reject H_0 if $\bar{x} > 68.26$

Finding the Type II Error Probabilities

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

For $\alpha = 0.05$: $k_\alpha = 68.26$

Decision rule: Reject H_0 if $\bar{x} > 68.26$

What is the **Type II error probability** if the true value is $\mu = 70$?

$$\mathbf{P}(\text{Do not reject } H_0 | \mu = 70) \longrightarrow \mathbf{P}(\bar{X} < 68.26 | \mu = 70)$$

Finding the Type II Error Probabilities

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

For $\alpha = 0.05$: $k_\alpha = 68.2604$ Decision rule: Reject H_0 if $\bar{X} > 68.26$

What is the **Type II error probability** if the true value is $\mu = 70$?

$$\mathbf{P}(\bar{X} < 68.26 | \mu = 70)$$

Finding the Type II Error Probabilities

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**


$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

For $\alpha = 0.05$: $k_\alpha = 68.2604$ Decision rule: Reject H_0 if $\bar{X} > 68.26$

What is the **Type II error probability** if the true value is $\mu = 70$?

If $\mu = 66.7$ $\bar{X} \sim \mathcal{N}\left(66.7, \frac{3^2}{10}\right)$

$$\mathbf{P}(\bar{X} < 68.26 | \mu = 70)$$



Finding the Type II Error Probabilities

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**


$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

For $\alpha = 0.05$: $k_\alpha = 68.2604$ Decision rule: Reject H_0 if $\bar{X} > 68.26$

What is the **Type II error probability** if the true value is $\mu = 70$?

$$\text{If } \mu = 70 \quad \bar{X} \sim \mathcal{N}\left(70, \frac{3^2}{10}\right)$$

$$\mathbf{P}(\bar{X} < 68.26 | \mu = 70)$$



Finding the Type II Error Probabilities

The mean height for 18 y/o in the US in the 70s was **66.7** in.


$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

For $\alpha = 0.05$: $k_\alpha = 68.2604$ Decision rule: Reject H_0 if $\bar{X} > 68.26$

What is the **Type II error probability** if the true value is $\mu = 70$?

$$\text{If } \mu = 70 \quad \bar{X} \sim \mathcal{N}\left(70, \frac{3^2}{10}\right)$$

$$\mathbf{P}(\bar{X} < 68.26 | \mu = 70) = 0.0333$$



Finding the Type II Error Probabilities

The mean height for 18 y/o in the US in the 70s was **66.7** in.


$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$

For $\alpha = 0.05$: $k_\alpha = 68.2604$ Decision rule: Reject H_0 if $\bar{X} > 68.26$

What is the **Type II error probability** if the true value is $\mu = 70$?

$$\text{If } \mu = 70 \quad \bar{X} \sim \mathcal{N}\left(70, \frac{3^2}{10}\right)$$

$$\beta = P(\bar{X} < 68.26 | \mu = 70) = 0.0333$$



Finding the Type II Error Probabilities

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10 \quad \sigma = 3$$


For $\alpha = 0.05$: $k_\alpha = 68.2604$ Decision rule: Reject H_0 if $\bar{X} > 68.26$

What is the **Type II error probability** if the true value is $\mu = 70$?

$$\text{If } \mu = 70 \quad \bar{X} \sim \mathcal{N}\left(70, \frac{3^2}{10}\right)$$

$$\beta = P(\bar{X} < 68.26 | \mu = 70) = 0.0333$$

This probability doesn't depend on your observed sample!



Power of the Test

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

Power of the test

$$\mathbf{P}(\text{Reject } H_0 | \mu \in H_1)$$

Power of the Test

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

Decision	Reality		Power of the test $P(\text{Reject } H_0 \mu \in H_1)$
	H_0 True ($\mu = 66.7$)	H_0 False ($\mu > 66.7$)	
Reject H_0 (Decide $\mu > 66.7$)	Type I error	Correct	
Don't reject H_0 (Decide $\mu = 66.7$)	Correct	Type II error	

Power of the Test

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

Decision	Reality		Power of the test $P(\text{Reject } H_0 \mu \in H_1)$
	H_0 True ($\mu = 66.7$)	H_0 False ($\mu > 66.7$)	
Reject H_0 (Decide $\mu > 66.7$)	Type I error	Correct	
Don't reject H_0 (Decide $\mu = 66.7$)	Correct	Type II error	

Power of the Test



Power of the Test

Type II error: $\overbrace{\mathbf{P}(\text{Do not reject } H_0 | \mu \in H_1)}$

Power of the Test

Type II error: $\overbrace{\mathbf{P}(\text{Do not reject } H_0 | \mu \in H_1)}^{\beta}$

Power of the test: $\overbrace{\mathbf{P}(\text{Reject } H_0 | \mu \in H_1)}^{1 - \beta}$

Power of the Test

$$\left. \begin{array}{l} \text{Type II error: } \mathbf{P} \left(\text{Do not reject } H_0 \mid \mu \in H_1 \right) \\ \text{Power of the test: } \mathbf{P} \left(\text{Reject } H_0 \mid \mu \in H_1 \right) \end{array} \right\} \overset{\beta}{\overbrace{\phantom{\text{Type II error: } \mathbf{P} \left(\text{Do not reject } H_0 \mid \mu \in H_1 \right)}}^{\text{Complementary probabilities}}}$$

Power of the Test

$$\left. \begin{array}{l} \text{Type II error: } \mathbf{P} \left(\text{Do not reject } H_0 \mid \mu \in H_1 \right) \\ \text{Power of the test: } \mathbf{P} \left(\text{Reject } H_0 \mid \mu \in H_1 \right) \end{array} \right\} \begin{array}{l} \beta \\ 1 - \beta \end{array}$$

Complementary probabilities

Power of the Test

$$\left. \begin{array}{l} \text{Type II error: } \mathbf{P} \left(\text{Do not reject } H_0 \mid \mu \in H_1 \right) \\ \text{Power of the test: } \mathbf{P} \left(\text{Reject } H_0 \mid \mu \in H_1 \right) \end{array} \right\} \begin{array}{l} \beta \\ 1 - \beta \end{array}$$

Complementary probabilities

Power of the test = 1 – Type II error probability


$$= 1 - \mathbf{P} \left(\text{Do not reject } H_0 \mid \mu \in H_1 \right)$$

Power of the Test

$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$


Power of the Test

$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$




Power of the Test

$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$




Power of the Test

$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$




Power of the Test

$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$




Power of the Test

$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$




Power of the Test

$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$




Power of the Test


$H_0 : \mu = 66.7$ vs. $H_1 : \mu > 66.7$




Power of the Test




Power of the Test




Power of the Test



Power of the Test






Power of the Test



Power of the Test

$$\mu = 70$$

Power






Type I error

Power of the Test

$$\mu = 70$$

Power






Type I error

Power of the Test

$$\mu = 70$$


Power



Type I error


Power of the Test

$$\mu = 70$$



Power of the Test




$$\mu = 70$$



Power of the Test

$$\mu = 70$$

Type II error



Type I error



DeepLearning.AI

Hypothesis Testing

Interpreting results

Steps for Performing Hypothesis Testing

Steps for Performing Hypothesis Testing

1. State your hypotheses.

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline → $H_0 : \mu = 66.7$

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline $\rightarrow H_0 : \mu = 66.7$
- Alternative hypothesis: the statement you want to prove $\rightarrow H_1 : \mu > 66.7$

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline $\rightarrow H_0 : \mu = 66.7$
- Alternative hypothesis: the statement you want to prove $\rightarrow H_1 : \mu > 66.7$

2. Design your test

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline $\rightarrow H_0 : \mu = 66.7$
- Alternative hypothesis: the statement you want to prove $\rightarrow H_1 : \mu > 66.7$

2. Design your test

- Decide the test statistic to work with. $\rightarrow \bar{X}$

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline $\rightarrow H_0 : \mu = 66.7$
- Alternative hypothesis: the statement you want to prove $\rightarrow H_1 : \mu > 66.7$

2. Design your test

- Decide the test statistic to work with. $\rightarrow \bar{X}$
- Decide the significance level $\rightarrow \alpha = 0.05$

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline $\rightarrow H_0 : \mu = 66.7$
- Alternative hypothesis: the statement you want to prove $\rightarrow H_1 : \mu > 66.7$

2. Design your test

- Decide the test statistic to work with. $\rightarrow \bar{X}$
- Decide the significance level $\rightarrow \alpha = 0.05$

3. Compute the observed statistic (based on your sample) $\rightarrow \bar{x} = 68.442$

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline $\rightarrow H_0 : \mu = 66.7$
- Alternative hypothesis: the statement you want to prove $\rightarrow H_1 : \mu > 66.7$

2. Design your test

- Decide the test statistic to work with. $\rightarrow \bar{X}$
- Decide the significance level $\rightarrow \alpha = 0.05$

3. Compute the observed statistic (based on your sample) $\rightarrow \bar{x} = 68.442$

4. Reach a conclusion:

Steps for Performing Hypothesis Testing

1. State your hypotheses.

- Null hypothesis: the baseline $\rightarrow H_0 : \mu = 66.7$
- Alternative hypothesis: the statement you want to prove $\rightarrow H_1 : \mu > 66.7$

2. Design your test

- Decide the test statistic to work with. $\rightarrow \bar{X}$
- Decide the significance level $\rightarrow \alpha = 0.05$

3. Compute the observed statistic (based on your sample) $\rightarrow \bar{x} = 68.442$

4. Reach a conclusion:

- If the p -value is less than the significance level reject H_0
 $\rightarrow P(\bar{X} > 68.442 | \mu = 66.7) > ? 0.05$

Important Remarks - Interpreting Tests

Important Remarks - Interpreting Tests

- Type I error:
 - Reject null hypothesis when it is true \rightarrow Reject H_0 when $\mu = 66.7$

Important Remarks - Interpreting Tests

- Type I error:
 - Reject null hypothesis when it is true \rightarrow Reject H_0 when $\mu = 66.7$
- Type II error:
 - Do not reject null hypothesis, when it was false \rightarrow Do not reject H_0 when $\mu > 66.7$

Important Remarks - Interpreting Tests

- Type I error:
 - Reject null hypothesis when it is true \rightarrow Reject H_0 when $\mu = 66.7$
- Type II error:
 - Do not reject null hypothesis, when it was false \rightarrow Do not reject H_0 when $\mu > 66.7$
- Significance level (α):
 - It is the maximum probability of incurring in a Type I error

Important Remarks - Interpreting Tests

- Type I error:
 - Reject null hypothesis when it is true \rightarrow Reject H_0 when $\mu = 66.7$
- Type II error:
 - Do not reject null hypothesis, when it was false \rightarrow Do not reject H_0 when $\mu > 66.7$
- Significance level (α):
 - It is the maximum probability of incurring in a Type I error
- Errors:
 - $\downarrow \alpha \uparrow \beta$

Important Remarks - Interpreting Tests

Important Remarks - Interpreting Tests

- p -values:

Important Remarks - Interpreting Tests

- *p*-values:

- If $\mathbf{P}(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$

Important Remarks - Interpreting Tests

- *p*-values:

- If $\mathbf{P}(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- The *p*-value represents the probability of H_0 being true.

Important Remarks - Interpreting Tests

- *p*-values:

- If $\mathbf{P}(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- The *p*-value represents the probability of H_0 being true.

Important Remarks - Interpreting Tests

Important Remarks - Interpreting Tests

- *p*-values:

- If $\mathbf{P}(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$

Important Remarks - Interpreting Tests

- *p*-values:

- If $\mathbf{P}(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- A small *p*-value indicates that the probability of seeing the observed data by chance is small

Important Remarks - Interpreting Tests

- ***p*-values:**

- If $P(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- A small *p*-value indicates that the probability of seeing the observed data by chance is small

Important Remarks - Interpreting Tests

- ***p*-values:**

- If $P(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- A small *p*-value indicates that the probability of seeing the observed data by chance is small

- **Test conclusions**

Important Remarks - Interpreting Tests

- ***p*-values:**

- If $P(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- A small *p*-value indicates that the probability of seeing the observed data by chance is small

- **Test conclusions**

- $\text{Reject } H_0 \rightarrow H_1 \text{ true}$

Important Remarks - Interpreting Tests

- ***p*-values:**

- If $P(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- A small *p*-value indicates that the probability of seeing the observed data by chance is small

- **Test conclusions**

- **Reject $H_0 \rightarrow H_1$ true**
- **Do not reject $H_0 \rightarrow H_0$ true**

Important Remarks - Interpreting Tests

- ***p*-values:**

- If $P(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- A small *p*-value indicates that the probability of seeing the observed data by chance is small

- **Test conclusions**

- Reject $H_0 \rightarrow H_1$ true
- Do not reject $H_0 \rightarrow H_0$ true



Important Remarks - Interpreting Tests

- ***p*-values:**

- If $P(\text{Reject } H_0 | H_0) < \alpha \longrightarrow \text{Reject } H_0 \text{ and then accept } H_1$
- A small *p*-value indicates that the probability of seeing the observed data by chance is small

- **Test conclusions**

- Reject $H_0 \rightarrow H_1$ true
- Do not reject $H_0 \rightarrow H_0$ true



You can only say that there is not enough evidence




DeepLearning.AI

Hypothesis Testing


t-Distribution

t -Distribution: Motivation

t -Distribution: Motivation




t -Distribution: Motivation



$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

t -Distribution: Motivation




$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

\downarrow

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

t -Distribution: Motivation



$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

This is fine if you know μ and σ

What if σ is unknown?

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$



$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$



$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{10}} \sim \mathcal{N}(0, 1^2)$$

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$



$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{10}} \sim \mathcal{N}(0, 1^2) \text{ (Standardization)}$$

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$



If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{10}} \sim \mathcal{N}(0, 1^2) \text{ (Standardization)}$$

Z statistic

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{10}} \sim \mathcal{N}(0, 1^2) \text{ (Standardization)}$$

Z statistic

What if σ is unknown?

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{10}} \sim \mathcal{N}(0, 1^2) \text{ (Standardization)}$$

Z statistic

What if σ is unknown?

Replace σ with its estimate

$$\frac{\bar{X} - \mu}{S / \sqrt{10}}$$

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{10}} \sim \mathcal{N}(0, 1^2) \text{ (Standardization)}$$

Z statistic

What if σ is unknown?

Replace σ with its estimate

$$\frac{\bar{X} - \mu}{S / \sqrt{10}}$$

T statistic

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

$$S = \sqrt{\frac{1}{10-1} \sum_{i=1}^{10} (X_i - \bar{X})^2}$$

If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{10}} \sim \mathcal{N}(0, 1^2) \text{ (Standardization)}$$

Z statistic

What if σ is unknown?

Replace σ with its estimate

$$\frac{\bar{X} - \mu}{S / \sqrt{10}}$$

T statistic

t -Distribution: Motivation

$$X_i \stackrel{i.i.d}{\sim} \mathcal{N}(\mu, \sigma^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{10}\right)$$

$$S = \sqrt{\frac{1}{10-1} \sum_{i=1}^{10} (X_i - \bar{X})^2}$$

If μ, σ are known

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{10}} \sim \mathcal{N}(0, 1^2) \text{ (Standardization)}$$

Z statistic

What if σ is unknown?

Replace σ with its estimate

$$\frac{\bar{X} - \mu}{S / \sqrt{10}} \sim \mathcal{N}(0, 1^2)$$



t -Distribution


t-Distribution

$\frac{\bar{X} - \mu}{S/\sqrt{10}}$ follows a *t* distribution

t -Distribution

$\frac{\bar{X} - \mu}{S/\sqrt{10}}$ follows a t distribution

What does it look like?




t -Distribution

$\frac{\bar{X} - \mu}{S/\sqrt{10}}$ follows a t distribution

What does it look like?

Still bell-shaped




t-Distribution

$\frac{\bar{X} - \mu}{S/\sqrt{10}}$ follows a *t* distribution

What does it look like?

Still bell-shaped

It has heavier tails that account for the uncertainty introduced with the std estimation



t -Distribution

t-Distribution

Parameters:

- Degrees of freedom

t-Distribution


Parameters:

- Degrees of freedom (ν)

t -Distribution

Parameters:

- Degrees of freedom (ν)




Controls how heavy
the tails are

t -Distribution

Parameters:

- Degrees of freedom (ν)



Controls how heavy
the tails are

$$X \sim t_{\nu}$$


t -Distribution

Parameters:

- Degrees of freedom (ν)


Controls how heavy
the tails are

$$X \sim t_{\nu}$$




t -Distribution and T -Statistic

t -Distribution and T -Statistic




t -Distribution and T -Statistic



$$n = 10$$


$$T = \frac{\bar{X} - \mu}{S/\sqrt{10}} \sim t_{\nu}$$

t -Distribution and T -Statistic



$$n = 10$$
$$\nu = 10 - 1$$
$$T = \frac{\bar{X} - \mu}{S/\sqrt{10}} \sim t_9$$

t -Distribution and T -Statistic




$$n = 10$$
$$T = \frac{\bar{X} - \mu}{S/\sqrt{10}} \sim t_9$$

$\nu = 10 - 1$

Degrees of freedom (ν) = sample size - 1
 $= (n - 1)$

t -Distribution and T -Statistic




$$n = 10$$
$$\nu = 10 - 1$$
$$T = \frac{\bar{X} - \mu}{S/\sqrt{10}} \sim t_9$$

Degrees of freedom (ν) = sample size - 1
 $= (n - 1)$

As n increases, this looks more
like a $\mathcal{N}(0, 1^2)$

t -Distribution and T -Statistic



$$n = 10$$
$$T = \frac{\bar{X} - \mu}{S/\sqrt{10}} \sim t_9$$

$\nu = 10 - 1$

Degrees of freedom (ν) = sample size - 1
= $(n - 1)$

As n increases, this looks more like a $\mathcal{N}(0, 1^2)$

T -statistic is used when

- The population has a Gaussian distribution
- But you don't know the variance




DeepLearning.AI

Hypothesis Testing

t-Tests

Example: Heights

Example: Heights



$$\bar{x} = 68.442$$

Example: Heights

Example: Heights

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**



3 questions

3 sets of hypothesis

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$



$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu < 66.7$$



$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu \neq 66.7$$

Example: Heights

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

Example: Heights

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$\sigma = 3$$

$$n = 10$$

$$H_0 : \mu = 66.7$$

$$\text{If } H_0 \text{ is true: } \bar{X} \sim \mathcal{N}\left(66.7, \frac{3^2}{10}\right)$$

Null hypothesis



Example: Heights


The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**



~~$n = 10$~~

$$H_0 : \mu = 66.7$$

If H_0 is true: $\bar{X} \sim$




Null hypothesis

66.7

μ

Example: Heights

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**



~~n = 10~~

$$H_0 : \mu = 66.7$$

Null hypothesis

66.7

μ

If H_0 is true: $\bar{X} \sim$

$$\text{If } H_0 \text{ is true: } T = \frac{\bar{X} - 66.7}{S/\sqrt{10}} \sim t_9$$

Example: Heights


The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**




~~$n = 10$~~

$$H_0 : \mu = 66.7$$

If H_0 is true: $\bar{X} \sim$



Null hypothesis



$$\text{If } H_0 \text{ is true: } T = \frac{\bar{X} - 66.7}{S/\sqrt{10}} \sim t_9$$

Example: Heights


The mean height for 18 y/o in the US in the 70s was 66.7 in.




~~n = 10~~

$$H_0 : \mu = 66.7$$

If H_0 is true: $\bar{X} \sim$




Null hypothesis



$$\text{If } H_0 \text{ is true: } T = \frac{\bar{X} - 66.7}{S/\sqrt{10}} \sim t_9$$

Right-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7 in.**




Right-Tailed Test for Gaussian Data (Unknown σ)

The **mean** height for 18 y/o in the US in the 70s was **66.7 in.**

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7$$

$$\alpha = 0.05$$



Right-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.


$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10$$

$$\alpha = 0.05$$

$$\bar{x} = 68.442$$

$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$



Right-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10$$

$$\bar{x} = 68.442$$


$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$

$$\alpha = 0.05$$

p-value:

$$P\left(\frac{\bar{X} - 66.7}{S/\sqrt{10}} > 1.771 \mid \mu = 66.7\right) ?$$



Null hypothesis

Right-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10$$

$$\alpha = 0.05$$

$$\bar{x} = 68.442$$


$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$

p-value:

$$P\left(\frac{\bar{X} - 66.7}{S/\sqrt{10}} > 1.771 \mid \mu = 66.7\right)$$

$$= 0.0552$$



Right-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu > 66.7 \quad n = 10$$

$$\bar{x} = 68.442$$

$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$


$$\alpha = 0.05$$

p-value:

$$P\left(\frac{\bar{X} - 66.7}{S/\sqrt{10}} > 1.771 \mid \mu = 66.7\right)$$

$$= 0.0552 > \alpha$$


Conclusion: do not reject H_0
(with a 5% significance level)



Two-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$n = 10 \quad \bar{x} = 68.442 \quad t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$
$$s = 3.113$$



Two-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.


$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu \neq 66.7 \quad n = 10$$

$$\bar{x} = 68.442$$

$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$

$$\alpha = 0.05$$



Two-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu \neq 66.7$$


$$n = 10$$

$$\bar{x} = 68.442$$

$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$

$$\alpha = 0.05$$



p-value:

$$P\left(\left|\frac{\bar{X} - 66.7}{S/\sqrt{10}}\right| > 1.771 \mid \mu = 66.7\right)?$$

Two-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu \neq 66.7$$


$$n = 10$$

$$\bar{x} = 68.442$$

$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$

$$\alpha = 0.05$$



p-value:

$$P\left(\left|\frac{\bar{X} - 66.7}{S/\sqrt{10}}\right| > |1.771| \middle| \mu = 66.7\right)?$$

Two-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu \neq 66.7$$


$$n = 10$$

$$\bar{x} = 68.442$$

$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$

$$\alpha = 0.05$$



p-value:

$$P\left(\left|\frac{\bar{X} - 66.7}{S/\sqrt{10}}\right| > |1.771| \mid \mu = 66.7\right) = 0.1103$$

Two-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu \neq 66.7$$


$$n = 10$$

$$\bar{x} = 68.442$$

$$s = 3.113$$

$$t = \frac{68.442 - 66.7}{3.113/\sqrt{10}} = 1.771$$

$$\alpha = 0.05$$



p-value:

$$P\left(\left|\frac{\bar{X} - 66.7}{S/\sqrt{10}}\right| > |1.771| \mid \mu = 66.7\right) = 0.1103 > \alpha$$


Conclusion: do not reject H_0
(with a 5% significance level)

Left-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$n = 10 \quad \bar{x} = 64.252 \quad t = \frac{64.252 - 66.7}{3.113/\sqrt{10}}$$
$$s = 3.113$$

$$\alpha = 0.05$$



Left-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.


$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu < 66.7 \quad n = 10$$

$$\bar{x} = 64.252$$

$$s = 3.113$$

$$t = \frac{64.252 - 66.7}{3.113/\sqrt{10}}$$

$$\alpha = 0.05$$



Left-Tailed Test for Gaussian Data (Unknown σ)


The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu < 66.7 \quad n = 10$$

$$\bar{x} = 64.252 \\ s = 3.113$$

$$t = \frac{64.252 - 66.7}{3.113/\sqrt{10}} = -2.487$$

$$\alpha = 0.05$$



Left-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu < 66.7$$

$$n = 10$$

$$\bar{x} = 64.252$$


$$s = 3.113$$

$$t = \frac{64.252 - 66.7}{3.113/\sqrt{10}} = -2.487$$

$$\alpha = 0.05$$

p-value:

$$P\left(\frac{\bar{X} - 66.7}{S/\sqrt{10}} < -2.487 \mid \mu = 66.7\right) ?$$



Left-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu < 66.7$$

$$n = 10$$

$$\bar{x} = 64.252$$

$$s = 3.113$$


$$t = \frac{64.252 - 66.7}{3.113/\sqrt{10}} = -2.487$$

$$\alpha = 0.05$$

p-value:

$$P\left(\frac{\bar{X} - 66.7}{S/\sqrt{10}} < -2.487 \mid \mu = 66.7\right)$$

$$= 0.0173$$



Left-Tailed Test for Gaussian Data (Unknown σ)

The mean height for 18 y/o in the US in the 70s was **66.7** in.

$$H_0 : \mu = 66.7 \text{ vs. } H_1 : \mu < 66.7$$

$$n = 10$$

$$\bar{x} = 64.252$$


$$s = 3.113$$

$$t = \frac{64.252 - 66.7}{3.113/\sqrt{10}} = -2.487$$

$$\alpha = 0.05$$

p-value:

$$P\left(\frac{\bar{X} - 66.7}{S/\sqrt{10}} < -2.487 \mid \mu = 66.7\right)$$



$$= 0.0173 < \alpha$$

Conclusion: reject H_0
(with a 5% significance level)




DeepLearning.AI

Hypothesis Testing

Two sample t-test


Independent Two-Sample t -Test

Independent Two-Sample t -Test




Height of 18 y/o in the US

Independent Two-Sample t -Test




Height of 18 y/o in the US




Height of 18 y/o in Argentina

Independent Two-Sample t -Test




$$n_X = 10 \quad \bar{x} = 68.442$$
$$s_X = 3.113$$

Height of 18 y/o in the US




Height of 18 y/o in Argentina

Independent Two-Sample t -Test



$$n_X = 10 \quad \bar{x} = 68.442$$
$$s_X = 3.113$$


Height of 18 y/o in the US



$$n_Y = 9 \quad \bar{y} = 65.949$$
$$s_Y = 3.106$$

Height of 18 y/o in Argentina


Independent Two-Sample t -Test



$$n_X = 10 \quad \bar{x} = 68.442$$
$$s_X = 3.113$$

Height of 18 y/o in the US

$$\mu_{US} \neq \mu_{Ar}$$




$$n_Y = 9 \quad \bar{y} = 65.949$$
$$s_Y = 3.106$$

Height of 18 y/o in Argentina



Independent Two-Sample t -Test: Hypothesis

Independent Two-Sample t -Test: Hypothesis





$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} > \mu_{Ar}$$

Independent Two-Sample t -Test: Hypothesis





$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} > \mu_{Ar}$$





$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} < \mu_{Ar}$$



Independent Two-Sample t -Test: Hypothesis



$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} > \mu_{Ar}$$





$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} < \mu_{Ar}$$





$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} \neq \mu_{Ar}$$



Independent Two-Sample t -Test: Hypothesis



$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$





$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} < \mu_{Ar}$$





$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} \neq \mu_{Ar}$$



Independent Two-Sample t -Test: Hypothesis



$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$




$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} < 0$$




$$H_0 : \mu_{US} = \mu_{Ar} \text{ vs. } H_1 : \mu_{US} \neq \mu_{Ar}$$


Independent Two-Sample t -Test: Hypothesis



$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$



$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} < 0$$



$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} \neq 0$$

Independent Two-Sample t -Test: Assumptions

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}^2)$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg}^2)$$

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}^2)$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg}^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i$$

$$\bar{Y} = \frac{1}{9} \sum_{i=1}^9 Y_i$$

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}^2)$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg}^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i$$

$$\bar{X} - \bar{Y} \sim ?$$

$$\bar{Y} = \frac{1}{9} \sum_{i=1}^9 Y_i$$

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}^2)$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg}^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\quad, \quad \right)$$

$$\bar{Y} = \frac{1}{9} \sum_{i=1}^9 Y_i$$

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}^2)$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg}^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \right)$$

$$\bar{Y} = \frac{1}{9} \sum_{i=1}^9 Y_i$$

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}^2)$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg}^2)$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right)$$

$$\bar{Y} = \frac{1}{9} \sum_{i=1}^9 Y_i$$

Independent Two-Sample t -Test: Assumptions

- All people in the sample from the two groups are different
- Each person in both samples are independent
- Populations are normally distributed

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US})$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$

$$\bar{X} = \frac{1}{10} \sum_{i=1}^{10} X_i$$

$$\bar{Y} = \frac{1}{9} \sum_{i=1}^9 Y_i$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}\right)$$

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US})$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right)$$

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}) \quad Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$
$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right) \xrightarrow{\hspace{1cm}} \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0, 1^2)$$

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}) \quad Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$
$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right) \rightarrow \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0, 1^2)$$

You don't know σ_{US} , σ_{Arg}



Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}) \quad Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$
$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right) \rightarrow \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0, 1^2)$$

You don't know σ_{US} , σ_{Arg}



Replace it with the sample standard deviation

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}) \quad Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$
$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right) \rightarrow \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0, 1^2)$$

You don't know σ_{US} , σ_{Arg}



Replace it with the sample standard deviation

$$\frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{S_X^2}{10} + \frac{S_Y^2}{9}}}$$

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}) \quad Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$
$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right) \rightarrow \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0, 1^2)$$

You don't know σ_{US} , σ_{Arg}



Replace it with the sample standard deviation

$$T = \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{S_X^2}{10} + \frac{S_Y^2}{9}}} \sim t_{\nu}$$

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US}) \quad Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}\right) \rightarrow \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0, 1^2)$$

You don't know σ_{US} , σ_{Arg}




Replace it with the sample standard deviation


$$T = \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{\nu}$$

Degrees of freedom = $\frac{\left(\frac{s_X^2}{n_X} + \frac{s_Y^2}{n_Y}\right)^2}{\frac{\left(\frac{s_X^2}{n_X}\right)^2}{n_X - 1} + \frac{\left(\frac{s_Y^2}{n_Y}\right)^2}{n_Y - 1}}$

Independent Two-Sample t -Test




$$n_X = 10 \quad \bar{x} = 68.442$$
$$s_X = 3.113$$




$$n_Y = 9 \quad \bar{y} = 65.949$$
$$s_Y = 3.106$$

Degrees of freedom = $\frac{\left(\frac{n_X - 1}{2} + \frac{n_Y - 1}{2} \right)^2}{\frac{\left(n_X - 1 \right)^2}{n_X - 1} + \frac{\left(n_Y - 1 \right)^2}{n_Y - 1}}$

Independent Two-Sample t -Test



$$n_X = 10 \quad \bar{x} = 68.442$$
$$s_X = 3.113$$



$$n_Y = 9 \quad \bar{y} = 65.949$$
$$s_Y = 3.106$$

Degrees of freedom = $\frac{\overbrace{\left(\frac{2}{3.113} + \frac{2}{3.106} \right)^2}^{16.8}}{\underbrace{\left(\frac{2}{10} \right)^2}_{10-1} + \underbrace{\left(\frac{2}{9} \right)^2}_{9-1}}$

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US})$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}\right) \rightarrow \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0,1)$$

You don't know σ_{US} , σ_{Arg}



Replace it with the sample standard deviation

$$T = \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Degrees of freedom = $\frac{\left(\frac{s_X^2}{n_X} + \frac{s_Y^2}{n_Y}\right)^2}{\frac{\left(\frac{s_X^2}{n_X}\right)^2}{n_X - 1} + \frac{\left(\frac{s_Y^2}{n_Y}\right)^2}{n_Y - 1}}$

Independent Two-Sample t -Test: Statistic

$$X \sim \mathcal{N}(\mu_{US}, \sigma_{US})$$

$$Y \sim \mathcal{N}(\mu_{Arg}, \sigma_{Arg})$$

$$\bar{X} - \bar{Y} \sim \mathcal{N}\left(\mu_{US} - \mu_{Arg}, \sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}\right) \rightarrow \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{\sigma_{US}^2}{10} + \frac{\sigma_{Arg}^2}{9}}} \sim \mathcal{N}(0,1)$$

You don't know σ_{US} , σ_{Arg}



Replace it with the sample standard deviation


$$T = \frac{(\bar{X} - \bar{Y}) - (\mu_{US} - \mu_{Arg})}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Degrees of freedom

$$\text{Degrees of freedom} = \frac{\left(\frac{s_X^2}{n_X} + \frac{s_Y^2}{n_Y}\right)^2}{\frac{\left(\frac{s_X^2}{n_X}\right)^2}{n_X - 1} + \frac{\left(\frac{s_Y^2}{n_Y}\right)^2}{n_Y - 1}}$$

Independent Two-Sample t -Test: Right Tailed Test


Independent Two-Sample t -Test: Right Tailed Test



$$n_X = 10$$

$$n_Y = 9$$

Independent Two-Sample t -Test: Right Tailed Test




$$\bar{x} = 68.442$$

$$n_X = 10$$

$$s_X = 3.113$$

$$n_Y = 9$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$n_Y = 9$$

$$s_X = 3.113$$

$$s_Y = 3.106$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$n_X = 10$$

$$\bar{y} = 65.949$$


$$n_Y = 9$$

$$s_X = 3.113$$

$$s_Y = 3.106$$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$n_X = 10$$

$$\bar{y} = 65.949$$

$$n_Y = 9$$


$$s_X = 3.113$$

$$s_Y = 3.106$$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$

$$\alpha = 0.05$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$

$$\bar{y} = 65.949$$

$$s_Y = 3.106$$


$$n_Y = 9$$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$


$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{S_X^2}{10} + \frac{S_Y^2}{9}}} \sim t_{16.8}$$

Null hypothesis



Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis




$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$

$$\bar{y} = 65.949$$


$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis

p -Value:




$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{S_X^2}{10} + \frac{S_Y^2}{9}}} \sim t_{16.8}$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis




p-Value: $P(T > 1.7459 \mid \mu_{US} - \mu_{Ar} = 0)$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis




p-Value: $P(T > 1.7459 \mid \mu_{US} - \mu_{Ar} = 0)$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis



p-Value: $P(T > 1.7459 \mid \mu_{US} - \mu_{Ar} = 0)$


$$= 0.0495$$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} > 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Independent Two-Sample t -Test: Right Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis




p-Value: $P(T > 1.7459 \mid \mu_{US} - \mu_{Ar} = 0)$

$$= 0.0495 < 0.05$$

\Rightarrow Reject H_0 (and accept H_1)
(with a 5% significance level)

Independent Two-Sample t -Test: Two Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$

$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$


$$t = 1.7459$$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} \neq 0$$


$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Null hypothesis



Independent Two-Sample t -Test: Two Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$

$$\bar{y} = 65.949$$


$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis

p -Value:




$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} \neq 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Independent Two-Sample t -Test: Two Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$

$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$


$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} \neq 0$$

$$\alpha = 0.05$$


$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Null hypothesis

p -Value: $\mathbf{P}(|T| > 1.7459 | \mu_{US} - \mu_{Ar} = 0)$



Independent Two-Sample t -Test: Two Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis




p-Value: $\mathbf{P}(|T| > 1.7459 | \mu_{US} - \mu_{Ar} = 0)$

$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} \neq 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

Independent Two-Sample t -Test: Two Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis



$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} \neq 0$$


$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{s_X^2}{10} + \frac{s_Y^2}{9}}} \sim t_{16.8}$$

$$p\text{-Value: } \mathbf{P}(|T| > 1.7459 | \mu_{US} - \mu_{Ar} = 0)$$

$$= 0.0991$$

Independent Two-Sample t -Test: Two Tailed Test



$$\bar{x} = 68.442$$

$$s_X = 3.113$$

$$n_X = 10$$


$$\bar{y} = 65.949$$

$$s_Y = 3.106$$

$$n_Y = 9$$

$$t = 1.7459$$

Null hypothesis



$$H_0 : \mu_{US} - \mu_{Ar} = 0 \text{ vs. } H_1 : \mu_{US} - \mu_{Ar} \neq 0$$

$$\alpha = 0.05$$

$$\text{If } H_0 \text{ is true: } T = \frac{(\bar{X} - \bar{Y}) - 0}{\sqrt{\frac{S_X^2}{10} + \frac{S_Y^2}{9}}} \sim t_{16.8}$$

$$p\text{-Value: } \mathbf{P}(|T| > 1.7459 | \mu_{US} - \mu_{Ar} = 0)$$

$$= 0.0991 > 0.05$$

\Rightarrow Do not reject H_0
(with a 5% significance level)





DeepLearning.AI

Hypothesis Testing


Paired t-test

Paired t -Test and Two-Sample t -Test


Paired t -Test and Two-Sample t -Test




Paired t -Test and Two-Sample t -Test




Paired t -Test and Two-Sample t -Test




Paired t -Test and Two-Sample t -Test




Paired t -Test and Two-Sample t -Test




Paired t -Test and Two-Sample t -Test



Paired t -Test and Two-Sample t -Test




Paired t -Test: Statistic



Now you're interested in the difference between pair of samples


Paired t -Test: Statistic



Now you're interested in the difference between pair of samples

$$(X_1 - Y_1)$$

Paired t -Test: Statistic



Now you're interested in the difference between pair of samples

$$(X_1 - Y_1) \quad (X_2 - Y_2) \quad (X_3 - Y_3) \quad (X_4 - Y_4) \quad (X_5 - Y_5) \quad (X_6 - Y_6) \quad (X_7 - Y_7) \quad (X_8 - Y_8) \quad (X_9 - Y_9) \quad (X_{10} - Y_{10})$$