

Buffon's Needle Problem for a Rectangular Grid

Author(s): Robert Fakler

Source: *The Mathematics Teacher*, MARCH 1995, Vol. 88, No. 3 (MARCH 1995), pp. 205-208

Published by: National Council of Teachers of Mathematics

Stable URL: <https://www.jstor.org/stable/27969276>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

National Council of Teachers of Mathematics is collaborating with JSTOR to digitize, preserve and extend access to *The Mathematics Teacher*

Buffon's Needle Problem for a Rectangular Grid

If a needle of length l is dropped onto a floor constructed of evenly spaced wooden planks of width k , what is the probability that the needle crosses a crack between planks? This question is "Buffon's needle problem." Schroeder (1974), in an article in the *Mathematics Teacher*, considers this problem and shows that for the case $l \leq k$, the solution is

$$p = \frac{2l}{k\pi}.$$

He observes that since classrooms no longer have wooden floors, the needle problem could be brought up to date by considering what happens when a needle of length l is dropped on a floor covered with square tiles of side k , where $l \leq k$. He also asks what is the probability that the needle will cross a crack in the floor.

To solve this problem, it is necessary to represent the possible positions in which the needle could land, relative to the cracks in the floor, by a set of independent variables. How many such variables would be needed? Two variables—an x -coordinate and a y -coordinate—are required to specify the position of one point on the needle. But the needle could fall in infinitely many positions that have this point in the same position, relative to the cracks in the floor, but different "slopes." To determine a unique position of the needle, in addition to specifying the position of a particular point, we could specify the "slope" of the needle. This process would require a third variable. Thus we see that *three* independent variables are required to determine the result of a needle toss. In solving his variation of the needle problem, Schroeder mistakenly uses *two* variables to represent the set of all possible needle tosses and obtains an incorrect answer,

$$p = \frac{2l\sqrt{2}}{k\pi}.$$

Instead of the problem Schroeder proposed, we shall consider a more general problem, in which a needle of length l is tossed onto a rectangular grid composed of rectangles having dimensions $a \times b$. If a needle of length l is tossed onto a rectangular grid consisting of $a \times b$ rectangles, what is the probability that the needle will cross a grid line? We assume

that $l \leq a$ and $l \leq b$. (If

$$a < l < \sqrt{a^2 + b^2}$$

or

$$b < l < \sqrt{a^2 + b^2},$$

the solution of this problem becomes more difficult, and we leave this case for the interested reader to pursue. If

$$l \geq \sqrt{a^2 + b^2},$$

then the needle will always cross a line, and so the probability that the needle will cross a line will be 1.)

Schroeder's problem is the special case where $a = b = k$. The outcome of each toss of the needle is determined by x , y , and θ , where x is the distance from the left end of the needle to the nearest vertical line to the right, y is the distance from the lower end of the needle to the nearest horizontal line above it, and θ is the angle whose initial side is the ray extending from the left end of the needle horizontally to the right and whose terminal side is the ray extending from the left end of the needle through the needle (see **fig. 1**). We notice that $0 \leq x \leq a$, $0 \leq y \leq b$, and $-\pi/2 < \theta \leq \pi/2$. We can identify the set of all possible outcomes for our needle-tossing experiment with the points in the rectangular solid

$$S = \{(x, y, \theta) | 0 \leq x \leq a, 0 \leq y \leq b, -\pi/2 < \theta \leq \pi/2\}.$$

Let R be the subregion of S corresponding to the outcomes in which the needle crosses a line. Then the probability, p , of the needle's crossing a line will be


$$p = \frac{\text{volume of } R}{\text{volume of } S}.$$

If $0 \leq \theta \leq \pi/2$, the needle will cross a line if and only if $x \leq l \cos \theta$ or $y \leq l \sin \theta$. Let S^+ be the subregion of S consisting of those points in S for which

How do we solve the classic problem if we have tiles on the floor instead of floorboards?


Robert Fakler teaches at the University of Michigan—Dearborn, Dearborn, MI 48128. His interests include geometrical probability and its use to enhance mathematics teaching.

We can use
the
method
of
“volume by
slicing”



$0 \leq \theta \leq \pi/2$. In this case, the needle tosses that result in the needle's crossing a line can be identified with the points in the region R^+ consisting of those points in S^+ that are inside the union of the cylinders $x = l \cos \theta$ and $y = l \sin \theta$. The regions R^+ and S^+ are shown in figure 2.

The region R^+ is the union of two right cylinders of heights a and b with bases B_1 and B_2 , respective-



ly. These cylinders will have volumes

$$(\text{area of } B_1)a$$

and

$$(\text{area of } B_2)b,$$

respectively. Let I be the region formed by the intersection of these cylinders. Then the volume of R^+ is

$$(\text{area of } B_1)a + (\text{area of } B_2)b - \text{volume of } I.$$

So

$$\begin{aligned} \text{area of } B_1 &= \int_0^{\pi/2} l \sin \theta d\theta \\ &= -l \cos \theta \Big|_0^{\pi/2} \\ &= l, \end{aligned}$$

and

$$\begin{aligned} \text{area of } B_2 &= \int_0^{\pi/2} l \cos \theta d\theta \\ &= l \sin \theta \Big|_0^{\pi/2} \\ &= l. \end{aligned}$$

To find the volume of I , we can use the method of “volumes by slicing.” Notice that a typical cross section of I perpendicular to the θ -axis is a rectangle with dimensions $l \cos \theta$ and $l \sin \theta$. (See fig. 3.) Thus the area $A(\theta)$ of this cross section is given by

$$A(\theta) = (l \cos \theta)(l \sin \theta) = l^2 \cos \theta \sin \theta,$$


and so

$$\begin{aligned} \text{volume of } I &= \int_0^{\pi/2} A(\theta) d\theta \\ &= \int_0^{\pi/2} l^2 \sin \theta \cos \theta d\theta \\ &= l^2 \left(\frac{\sin^2 \theta}{2} \right) \Big|_0^{\pi/2} \\ &= \frac{l^2}{2}. \end{aligned}$$

Therefore,

$$\text{volume of } R^+ = la + lb - \frac{l^2}{2}.$$

Next suppose that $-\pi/2 < \theta < 0$. Then the needle will cross a line, that is, (x, y, θ) will belong to R , if and only if $x < l \cos \theta$ or $y < -l \sin \theta$. Let R^- be the subregion of R consisting of those points where $-\pi/2 < \theta < 0$. Then R^- will consist of all points (x, y, θ)



in R satisfying $-\pi/2 < \theta < 0$ and also $x < l \cos \theta$ or $y < -l \sin \theta$. We claim that the subregion of R^+ satisfying $0 < \theta < \pi/2$ is symmetric to R^- with respect to the xy -plane. To see this relationship, notice that $(x, y, -\theta)$ is the symmetric point to (x, y, θ) with respect to the xy -plane and that

$$\begin{aligned} (x, y, -\theta) \in R^- \\ \Leftrightarrow (x < l \cos(-\theta) \text{ or } y < -l \sin(-\theta)) \\ \text{and } -\frac{\pi}{2} < -\theta < 0 \\ \Leftrightarrow (x < l \cos \theta \text{ or } y < l \sin \theta) \\ \text{and } 0 < \theta < \frac{\pi}{2} \\ \Leftrightarrow (x, y, \theta) \in R^+ \text{ and } 0 < \theta < \frac{\pi}{2}. \end{aligned}$$

Therefore, the volume of the subregion of R^+ consisting of those points with $0 < \theta < \pi/2$ is the same as the volume of R^- . Since the volume of this subregion is also the same as the volume of R^+ , we observe that

$$\text{volume of } R = 2(\text{volume of } R^+).$$

Also, we easily see that

$$\text{volume of } S^+ = 2(\text{volume of } S).$$

Since the volume of S^+ is $(\pi ab)/2$, we obtain

$$\begin{aligned} p &= \frac{\text{volume of } R}{\text{volume of } S} \\ &= \frac{2(\text{volume of } R^+)}{2(\text{volume of } S^+)} \\ &= \frac{\text{volume of } R^+}{\text{volume of } S^+} \\ &= \frac{la + lb - \frac{l^2}{2}}{\frac{\pi ab}{2}} \\ &= \frac{2l(a + b) - l^2}{\pi ab}. \end{aligned}$$

When the $a \times b$ rectangles are squares with side k , we have $a = b = k$ and so

$$p = \frac{4lk - l^2}{\pi k^2},$$

which is the correct solution for the needle problem with a square grid considered by Schroeder.

An interesting observation is that if we let one of the dimensions, say a , of the rectangles in our rectangular grid approach infinity, our solution should approach that for the needle problem with parallel lines, where b is the distance between adjacent lines. We see that this outcome indeed occurs, since

$$\begin{aligned} \lim_{a \rightarrow \infty} \frac{2l(a + b) - l^2}{\pi ab} &= \lim_{a \rightarrow \infty} \frac{2l + \frac{2lb}{a} - \frac{l^2}{a}}{\pi b} \\ &= \frac{2l}{\pi b}. \end{aligned}$$

PROGRAM 1

```

10  REM VARIABLES
20  REM  PI . . . . THE CONSTANT PI
30  REM  N . . . . NUMBER OF TOSSES
40  REM  S . . . . NUMBER OF TIMES NEEDLE
                   CROSSES A LINE
50  REM  X . . . . DISTANCE BETWEEN
                   LEFTMOST END OF
                   NEEDLE AND NEAREST
                   VERTICAL LINE TO THE
                   RIGHT
60  REM  Y . . . . DISTANCE BETWEEN LOWER
                   END OF NEEDLE AND
                   NEAREST HORIZONTAL
                   LINE ABOVE IT
70  REM  THETA . ANGLE MADE BY NEEDLE
                   WITH A HORIZONTAL
                   LINE THROUGH ITS
                   LOWER END
80  REM  I . . . . INDEX VARIABLE FOR
                   FOR-NEXT LOOP
90  LET PI = 3.14159
100 PRINT "NO. OF TOSSES"; TAB (17);
        "PROBABILITY ESTIMATE"
110 FOR N = 1000 TO 10000 STEP 1000
120   RANDOMIZE (N)
130   LET S = 0
140   FOR I = 1 TO N
150     LET X = 2 * RND
160     LET Y = 3 * RND
170     LET THETA = (PI/2) * RND
180     IF X <= COS (THETA) OR
        Y <= SIN (ABS(THETA))
        THEN S = S + 1
190   NEXT I
200   PRINT TAB (5); N; TAB (23); S/N
210 NEXT N


```

NO. OF TOSSES	PROBABILITY ESTIMATE
1000	.465
2000	.4805
3000	.4743334
4000	.4995
5000	.4552
6000	.4805
7000	.475
8000	.474375
9000	.471
10000	.4735

**We can
avoid
calculus
by using a
computer
simulation**

Finally, we recall that our solution to the needle problem required the use of integration to obtain the volume of R . Another approach, which would avoid the use of integration, would be to write a computer program to simulate the needle-tossing experiment. We could have our program "toss" the needle a large number of times and obtain an empirical estimate of the probability of crossing a line by dividing the number of times the needle crosses a line by the number of times the needle is

TEXAS INSTRUMENTS
Calculators




CLASSROOM

TI 108
TI 12
TI 34
TI 30
TI 7

**CALL FOR
CURRENT
PRICING**

GRAPHING

TI 80
TI 81
TI 82
TI 85



CASIO
Calculators

GRAPHING CLASSROOM

FX6300G
FX7700GE
FX9700GE

SL450
AZ45F
FX55

**CALL FOR
CURRENT
PRICING**

**hp HEWLETT
PACKARD**
Graphing Calculators

HP 48G

Hundreds of built in equations. 32 KB Memory for complex problem solving. RS-232 lets you link to your computer through a serial port. Two way infrared I/O for easy data transfer. 3-D Graphics, calculus, and symbolic math functions

HP 48G Same as 48G with expandable port

**CALL FOR
CURRENT
PRICING**

SHARP
Calculators



FRACTIONS

ELE300

**CALL FOR
CURRENT
PRICING**

GRAPHING

EL9200
EL9300



Overhead Calculators and Class Sets
for these and other models are available!

Advantage
Marketing

Call for our New 1994/95 Catalog & Quantity Pricing Information!

800-937-9777

VISA OR MASTERCARD ACCEPTED

tossed. For example, suppose we take $l = 1$, $a = 2$, and $b = 3$. **Program 1** includes an IBM PC BASIC program that computes, in this case, empirical probability estimates starting with 1 000 tosses and continuing in steps of 1 000 up to 10 000 tosses, along with a sample run; the teacher should note that this program takes a long time to run. In this case, the mathematical probability of the needle's crossing a line is

$$p = \frac{2(1)(2+3) - 1^2}{\pi(2)(3)} = \frac{9}{6\pi} \approx 0.477.$$

REFERENCE

Schroeder, Lee L. "Buffon's Needle Problem: An Exciting Application of Many Mathematical Concepts." *Mathematics Teacher* 67 (February 1974):183-86.

73rd Annual Meeting

Boston • Massachusetts

6-9 April 1995

To register call 800-235-7566

The Power and Beauty of Mathematics

Instant Awards, Class Enrichment,

Bulletin Board Eyecatchers,

Correspondence with Colleagues,

Club Fund Raisers, etc.

Handsomely Printed Selections of

Striking Mathematical Formulas:

"Smart" Cards with Envelopes

Set of twelve, all different,

postpaid : . . . \$20.00

Send check to:

MATHEMATEX®

P. O. Box 181805

Dallas, TX 75218

*Please allow a minimum of
four to six weeks for delivery.*