

# Measure Theory

## Contents

|     |                                                                                              |    |
|-----|----------------------------------------------------------------------------------------------|----|
| 1   | Algebras of Sets .....                                                                       | 2  |
| 1.1 | Additive set-function on an algebra of sets .....                                            | 4  |
| 2   | Sigma Algebras .....                                                                         | 4  |
| 2.1 | Positive measure on a sigma algebra .....                                                    | 5  |
| 2.2 | A glimpse of the Lebesgue measure on $\mathbb{R}$ .....                                      | 7  |
| 3   | Measurable functions .....                                                                   | 8  |
| 3.1 | The space of functions $\text{Bor}(X, \mathbb{R})$ .....                                     | 8  |
| 4   | Convergence properties of $\text{Bor}(X, \mathbb{R})$ .....                                  | 10 |
| 4.1 | Approximation by simple functions .....                                                      | 12 |
| 5   | Integration of functions in $\text{Bor}^+(X, \mathbb{R})$ .....                              | 13 |
| 5.1 | The map $L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \rightarrow [0, \infty]$ .....                | 13 |
| 5.2 | The map $L^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty]$ .....                    | 17 |
| 6   | The Monotone Convergence Theorem .....                                                       | 18 |
| 6.1 | Proof of the Monotone Convergence Theorem .....                                              | 20 |
| 7   | The space of integrable functions $\mathcal{L}^1(\mu)$ .....                                 | 23 |
| 7.1 | Notion of integrable function, the map $L : \mathcal{L}^1(\mu) \rightarrow \mathbb{R}$ ..... | 23 |
| 7.2 | Integration of subsets .....                                                                 | 27 |
| 8   | Lebesgue's Dominated Convergence Theorem (LDCT) .....                                        | 30 |
| 8.1 | The "a.e.- $\mu$ " annoyance (and the "a.e.- $\mu$ " version of LDCT) .....                  | 35 |

**Problem**

Consider a random (infinite) string of 0's and 1's

$$(\ell_1, \dots, \ell_n, \dots) \quad \ell_n \in \{0, 1\} \forall n \in \mathbb{N}$$

What is the probability that somewhere along the string, I find 100 consecutive zeros?

**Comment**

The statement of the problem is unclear. Lebesgue's machinery will clarify this

Define

$$X = \{x = (\ell_1, \dots, \ell_n, \dots) \mid \ell_n \in \{0, 1\} \forall n \in \mathbb{N}\}$$

This is, in a natural way, a compact metric space. Use the "Hamming-style" distance:

For  $x = (\ell_1, \dots, \ell_n, \dots)$  and  $x' = (\ell'_1, \dots, \ell'_n, \dots)$

$$d(x, x') = \sum_{n \in \mathbb{N}} \frac{|\ell_n - \ell'_n|}{2^n}$$

Since this is a metric space, we can consider the topology:

$$\mathcal{T} = \{G \subseteq X \mid G \text{ is open}\} \subseteq 2^X = \mathcal{P}(X)$$

Can consider

$$\mathfrak{B} = \sigma\text{-Alg}(\mathcal{T}) \quad (\text{Borel sigma algebra})$$

Then a probability measure is

$$P : \mathcal{B} \longrightarrow [0, 1]$$

[Algebra of Sets \(Problem 1.1\)](#) says:

Let

$$S = \{x = (\ell_1, \dots, \ell_n, \dots) \in X \mid \exists m \in \mathbb{N} \text{ with } \ell_m = \ell_{m+1} = \dots = \ell_{m+99} = 0\}$$

check that  $S \in \mathcal{B}$ , and compute  $P(S)$

## 1 Algebras of Sets

Fix a non-empty set  $X$ .

**Definition 1.1****Algebra of Sets**

$\mathfrak{A} \subseteq 2^X$  is an *algebra of sets* if

1.  $\emptyset \in \mathfrak{A}$
2. If  $A \in \mathfrak{A}$ , then  $X \setminus A \in \mathfrak{A}$
3. If  $A, B \in \mathfrak{A}$ , then  $A \cap B \in \mathfrak{A}$

**Proposition 1.2****Properties of algebras of sets**

Let  $\mathfrak{A}$  be an algebra of sets on  $X$ . Then

1.  $X \in \mathfrak{A}$
2. If  $A, B \in \mathfrak{A}$ , then  $A \cup B \in \mathfrak{A}$
3. If  $A, B \in \mathfrak{A}$ , then  $A \setminus B \in \mathfrak{A}$
4. If  $\mathfrak{B} \subseteq \mathfrak{A}$  is finite, then  $\bigcap \mathfrak{B} \in \mathfrak{A}$
5. If  $\mathfrak{B} \subseteq \mathfrak{A}$  is finite, then  $\bigcup \mathfrak{B} \in \mathfrak{A}$

**Example 1.1**

Obviously,  $2^X$  and  $\{\emptyset, X\}$  are algebras of sets on  $X$

**Lemma 1.3****Intersections of algebras**

Let  $\mathfrak{F}$  be any collection of algebras of sets on  $X$ , then  $\bigcap \mathfrak{F}$  is an algebra of sets on  $X$ .

**Proof:** First note if  $\mathfrak{F} = \emptyset$ , then  $\bigcap \mathfrak{F} = 2^X$  which is an algebra of sets on  $X$ . We now proceed with checking the axioms of an algebra of sets on  $X$ .

1. For every  $\mathfrak{A} \in \mathfrak{F}$ , we have  $\emptyset \in \mathfrak{A}$ , thus  $\emptyset \in \bigcap \mathfrak{F}$ .
2. If  $A \in \bigcap \mathfrak{F}$ , then  $A \in \mathfrak{A}$  for every  $\mathfrak{A} \in \mathfrak{F}$ . Thus  $X \setminus A \in \mathfrak{A}$  for every  $\mathfrak{A} \in \mathfrak{F}$ , and thus  $X \setminus A \in \bigcap \mathfrak{F}$ .
3. If  $A, B \in \bigcap \mathfrak{F}$ , then  $A, B \in \mathfrak{A}$  for every  $\mathfrak{A} \in \mathfrak{F}$ . Thus  $A \cap B \in \mathfrak{A}$  for every  $\mathfrak{A} \in \mathfrak{F}$ , and thus  $A \cap B \in \bigcap \mathfrak{F}$ .

□

**Proposition 1.4****Algebra generated by a collection**

Let  $\mathfrak{B} \subseteq 2^X$  be any collection of subsets of  $X$ , then there exists a unique algebra of sets  $\mathfrak{A}$  on  $X$  with the following properties:

1.  $\mathfrak{B} \subseteq \mathfrak{A}$
2. Whenever  $\mathfrak{C}$  is an algebra of sets on  $X$  with  $\mathfrak{B} \subseteq \mathfrak{C}$ , then we have  $\mathfrak{A} \subseteq \mathfrak{C}$

**Proof:** Define  $\mathfrak{F}$  to be the collection of all algebras of sets on  $X$  that contain  $\mathfrak{B}$ . Then  $\bigcap \mathfrak{F}$  is an algebra of sets by [Intersections of algebras \(Lemma 1.3\)](#) and satisfies (1) and (2). Uniqueness follows from (2).

□

**Definition 1.5****Algebra of sets generated by a collection of subsets**

Let  $\mathfrak{B} \subseteq 2^X$  be any collection of subsets of  $X$ . We define the *algebra of sets generated by  $\mathfrak{B}$*  to be the unique algebra from [Algebra generated by a collection \(Proposition 1.4\)](#). We notate it as  $\text{Alg}(\mathfrak{B})$ .

## 1.1 Additive set-function on an algebra of sets

### Definition 1.6

### Additive set-function

Let  $\mathfrak{A} \subseteq 2^X$  be an algebra of sets on  $X$ . A function  $\mu : \mathfrak{A} \rightarrow [0, \infty]$  is called an *additive set-function on  $\mathfrak{A}$*  if

1. If  $A, B \in \mathfrak{A}$  with  $A \cap B = \emptyset$ , then  $\mu(A \cup B) = \mu(A) + \mu(B)$
2.  $\mu(\emptyset) = 0$

### Remark 1.7

The arithmetic of  $[0, \infty]$  is standard, except for  $0 \cdot \infty = \infty \cdot 0 = 0$ , which is a useful convention in measure theory.

### Remark 1.8

We almost don't need to require (2), it's just when  $\mu$  is identically  $\infty$  do things go wrong without (2)

### Proposition 1.9

### Properties of an additive set function

Let  $\mathfrak{A}$  be an algebra of sets on  $X$ , and let  $\mu$  be an additive set function on  $\mathfrak{A}$ . Then

1. If  $A, C \in \mathfrak{A}$  with  $A \subseteq C$ , then  $\mu(A) \leq \mu(C)$
2. If  $A, B \in \mathfrak{A}$ , then  $\mu(A \cup B) \leq \mu(A) + \mu(B)$  (sub additivity)
3. If  $A_1, \dots, A_n \in \mathfrak{A}$  are all pairwise disjoint, then  $\mu(A_1 \cup \dots \cup A_n) = \mu(A_1) + \dots + \mu(A_n)$
4. If  $A_1, \dots, A_n \in \mathfrak{A}$ , then  $\mu(A_1 \cup \dots \cup A_n) \leq \mu(A_1) + \dots + \mu(A_n)$

## 2 Sigma Algebras

### Definition 2.1

### Sigma Algebra

$\mathfrak{M} \subseteq 2^X$  is called a *sigma algebra* on  $X$  if

1.  $\emptyset \in \mathfrak{M}$
2. If  $A \in \mathfrak{M}$ , then  $X \setminus A \in \mathfrak{M}$
3. If  $(A_n)_{n=1}^\infty \subseteq \mathfrak{M}$ , then  $\bigcap_{n=1}^\infty A_n \in \mathfrak{M}$

### Remark 2.2

countable unions work too, and so do finite operations

### Lemma 2.3

### Intersections of sigma algebras

Let  $\mathfrak{F}$  be any collection of sigma algebras on  $X$ , then  $\bigcap \mathfrak{F}$  is a sigma algebra on  $X$ .

**Proof:** basically identical to the proof for algebras of sets □

**Proposition 2.4****Sigma algebra generated by a collection**

Let  $\mathfrak{B} \subseteq 2^X$  be any collection of subsets of  $X$ , then there exists a unique sigma algebra  $\mathfrak{A}$  on  $X$  with the following properties:

1.  $\mathfrak{B} \subseteq \mathfrak{A}$
2. Whenever  $\mathfrak{C}$  is a sigma algebra on  $X$  with  $\mathfrak{B} \subseteq \mathfrak{C}$ , then we have  $\mathfrak{A} \subseteq \mathfrak{C}$

**Proof:** basically identical to the proof for algebras of sets □

**Definition 2.5****Sigma algebra generated by a collection of subsets**

Let  $\mathfrak{B} \subseteq 2^X$  be any collection of subsets of  $X$ . We define the *sigma algebra generated by  $\mathfrak{B}$*  to be the unique sigma algebra from [Sigma algebra generated by a collection \(Proposition 2.4\)](#). We notate it as  $\sigma\text{-Alg}(\mathfrak{B})$ .

**Definition 2.6****Borel Sigma Algebra**

Given a topological space  $(X, \mathcal{T})$ , the sigma algebra generated by  $\mathcal{T}$  is called the *Borel sigma algebra on  $X$*  and written  $\mathcal{B}_X := \sigma\text{-Alg}(\mathcal{T})$

**2.1 Positive measure on a sigma algebra****Remark 2.7****Increasing limits and series in  $[0, \infty]$** 

1. Let  $(s_n)_{n=1}^\infty$  be an *increasing* sequence in  $[0, \infty]$  in the sense that

$$s_1 \leq s_2 \leq \dots$$

Such a sequence is sure to approach a limit  $L \in [0, \infty]$ . With a slight abuse of notation we can also write the formula  $L = \sup\{s_n \mid n \in \mathbb{N}\}$  meaning that  $L$  is the smallest upper bound for all the  $s_n$ 's. This is usually written in connection to a bounded increasing sequence of numbers in  $[0, \infty)$  – our “slight abuse of notation” is that we are also allowing it to cover the case  $L = \infty$ , which can occur when some of the  $s_n$ 's are equal to  $\infty$ , or when  $(s_n)_{n=1}^\infty$  is an unbounded sequence in  $[0, \infty)$ .

2. Let  $(t_n)_{n=1}^\infty$  be any sequence in  $[0, \infty]$ . We can consider the finite sums

$$s_n := \sum_{i=1}^n t_i$$

which forms an increasing sequence in  $[0, \infty]$ . We can consider  $L = \lim_{n \rightarrow \infty} s_n$  as in (1.), and use the familiar notation  $L = \sum_{n=1}^\infty t_n$

**Definition 2.8****Positive measure**

Let  $\mathfrak{M} \subseteq 2^X$  be sigma algebra of subsets of  $X$ . We say  $\mu : \mathfrak{M} \rightarrow [0, \infty]$  is a *positive measure* on  $\mathfrak{M}$  if it satisfies:

1.  $\mu(\emptyset) = 0$

2. 
$$\mu\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \mu(A_n)$$

Whenever  $(A_n)_{n=1}^{\infty}$  is a sequence of sets in  $\mathfrak{M}$  with the property that if  $i, j \in \mathbb{N}$  with  $i \neq j$  we have  $A_i \cap A_j = \emptyset$

**Remark 2.9**

A positive measure is, in particular, finitely additive. Thus we can use all the properties in [Properties of an additive set function \(Proposition 1.9\)](#)

**Definition 2.10****Nomenclature involving measures**

1. If  $\mathfrak{M}$  is a sigma algebra on  $X$ , then the pair  $(X, \mathfrak{M})$  is called a *measurable space*
2. If we also have a positive measure  $\mu$  on  $\mathfrak{M}$ , then the triple  $(X, \mathfrak{M}, \mu)$  is called a *measure space*
3. If  $\mu(X) < \infty$ , then we say  $\mu$  is a *finite measure*
4. If  $\mu(X) = 1$ , then we say  $\mu$  is a *probability measure* and that  $(X, \mathfrak{M}, \mu)$  is a *probability space*
5. We say that  $\mu$  is *sigma finite* if there exists a sequence of sets  $(X_n)_{n=1}^{\infty}$  in  $\mathfrak{M}$  with
  - a.  $X_1 \subseteq X_2 \subseteq \dots$  (The  $X_n$ 's are said to form an increasing chain)
  - b.  $\bigcup_{n=1}^{\infty} X_n = X$
  - c.  $\mu(X_n) < \infty$  for all  $n \in \mathbb{N}$

**Proposition 2.11****Sigma sub-additivity**

Let  $\mathfrak{M}$  be a sigma algebra on  $X$ , and let  $\mu$  be a positive measure on  $\mathfrak{M}$ . If  $(A_n)_{n=1}^{\infty}$  is a sequence of sets in  $\mathfrak{M}$ , then

$$\mu\left(\bigcup_{n=1}^{\infty} A_n\right) \leq \sum_{n=1}^{\infty} \mu(A_n)$$

**Proposition 2.12****Continuity along chains**

Let  $\mathfrak{M}$  be a sigma algebra on  $X$ , and let  $\mu$  be a positive measure on  $\mathfrak{M}$ .

1. If  $(A_n)_{n=1}^{\infty}$  is an increasing chain in  $\mathfrak{M}$ , then

$$\mu\left(\bigcup_{n=1}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \mu(A_n)$$

2. If  $(A_n)_{n=1}^{\infty}$  is a decreasing chain in  $\mathfrak{M}$  with  $\mu(A_1) < \infty$ , then

$$\mu\left(\bigcap_{n=1}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \mu(A_n)$$

**Example 2.1**

Example of “atomic”  $\mu$  based on weights of individual points  $x \in X$ . Suppose you have  $w : X \rightarrow [0, \infty]$ . Let  $\mathfrak{M} = 2^X$ , for every  $A \subseteq X$ , put

$$\mu(A) = \sup \left\{ \sum_{x \in F} w(x) \mid F \subseteq A \text{ finite} \right\}$$

Then  $(X, \mathfrak{M}, \mu)$  is a measure space. For the special case that  $w(x) = 1, \forall x \in X$  is called the *counting measure* on  $X$ . For the special case that  $w(x_0) = 1$  for some  $x_0 \in X$  and  $w(x) = 0, \forall x \neq x_0$  is called the *Dirac measure concentrated at  $x_0$*

**2.2 A glimpse of the Lebesgue measure on  $\mathbb{R}$** **Note**

1.  $\mathcal{B}_{\mathbb{R}}$  can be generated from the half open intervals
2. If  $\mu, \nu : \mathcal{B}_{\mathbb{R}} \rightarrow [0, \infty]$  are positive measures such that

$$\mu((a, b]) = \nu((a, b]) < \infty, \forall a < b \in \mathbb{R}$$

It follows that  $\mu = \nu$

Then the *Lebesgue measure*  $\lambda : \mathcal{B}_{\mathbb{R}} \rightarrow [0, \infty]$  is determined via with requirement that  $\lambda((a, b]) = b - a, \forall a < b \in \mathbb{R}$ .

Another approach:  $\lambda$  is uniquely determined by the requirement that  $\lambda(B + t) = \lambda(B)$  for every  $B \in \mathcal{B}_{\mathbb{R}}$  and  $t \in \mathbb{R}$  with the normalization that  $\lambda((0, 1]) = 1$ .

**Note**

For the *existence* of  $\lambda$ , will invoke an extension theorem of Caratheodory (to be done later)

**Note**

Interesting fact: there is no positive measure  $\mu : 2^{\mathbb{R}} \rightarrow [0, \infty]$  which is translation invariant and  $\mu((0, 1]) = 1$ . This shows, in particular that  $\mathcal{B}_{\mathbb{R}} \neq 2^{\mathbb{R}}$

### 3 Measurable functions

**Definition 3.1****Measurable function**

Let  $(X, \mathfrak{M}), (Y, \mathfrak{N})$  be measurable spaces and  $f : X \rightarrow Y$ . Then  $f$  is called  $\mathfrak{M}/\mathfrak{N}$ -measurable if  $f^{-1}(N) \in \mathfrak{M}$  for every  $N \in \mathfrak{N}$ .

**Remark 3.2**

Pre-images behave nicely with respect to the usual set operations.

**Proposition 3.3****Composition of measurable maps**

Compositions of measurable maps are measurable.

**Proof:** Obvious □

**Proposition 3.4****Measurability criterion**

Let  $(X, \mathfrak{M}), (Y, \mathfrak{N})$  be measurable spaces.  $f : X \rightarrow Y$ , and  $\mathfrak{C} \subseteq \mathfrak{N}$  such that  $\sigma\text{-Alg}(\mathfrak{C}) = \mathfrak{N}$  and such  $f^{-1}(C) \in \mathfrak{M}$  for every  $C \in \mathfrak{C}$ . Then  $f$  is measurable.

**Proof:** Let  $\mathfrak{Q} = \{Q \subseteq Y \mid f^{-1}(Q) \in \mathfrak{M}\} \subseteq 2^Y$

**Claim:**  $\mathfrak{Q}$  is a sigma algebra

Note by assumption  $\mathfrak{C} \subseteq \mathfrak{Q}$  thus  $\mathfrak{N} = \sigma\text{-Alg}(\mathfrak{C}) \subseteq \mathfrak{Q}$ . Thus  $f$  is measurable. □

**Corollary 3.5****Continuous maps are measurable**

Let  $X, Y$  be topological spaces,  $f : X \rightarrow Y$  continuous. Then  $f$  is  $\mathcal{B}_X/\mathcal{B}_Y$ -measurable.

#### 3.1 The space of functions $\text{Bor}(X, \mathbb{R})$

**Definition 3.6****Borel measurable real-valued functions**

Given a measurable space  $(X, \mathfrak{M})$ . Define

$$\text{Bor}(X, \mathbb{R}) := \{f : X \rightarrow \mathbb{R} \mid f \text{ is } \mathfrak{M}/\mathcal{B}_{\mathbb{R}} \text{ - measurable}\}$$



**Proposition 3.7****Algebra and lattice operations**

Let  $(X, \mathfrak{M})$  be a measurable space with  $f, g \in \text{Bor}(X, \mathbb{R})$ . Then

1.  $f + g \in \text{Bor}(X, \mathbb{R})$
2. For any  $c \in \mathbb{R}$ ,  $cf \in \text{Bor}(X, \mathbb{R})$
3.  $f \cdot g \in \text{Bor}(X, \mathbb{R})$
4.  $f \vee g, f \wedge g \in \text{Bor}(X, \mathbb{R})$

**Proof of 3:** Let  $f, g \in \text{Bor}(X, \mathbb{R})$ . Let  $h : X \rightarrow \mathbb{R}^2$  given by  $h(x) = (f(x), g(x))$ . Let  $p : \mathbb{R}^2 \rightarrow \mathbb{R}$  given by  $p(s, t) = s \cdot t$ .

**Claim:**  $h$  is measurable

**Claim:**  $p$  is measurable

**Claim:**  $f \cdot g = p \circ h$

Thus  $f \cdot g \in \text{Bor}(X, \mathbb{R})$  □

**Exercise 3.1**

Let  $\mathfrak{S}$  be the collection of open squares in  $\mathbb{R}^2$ . Then  $\sigma\text{-Alg}(\mathfrak{S}) = \mathcal{B}_{\mathbb{R}^2}$ .

**Remark 3.8**

Let  $(X, \mathfrak{M})$  be a measurable space. Then  $\text{Bor}(X, \mathbb{R})$  is an *algebra of functions* and a *lattice of functions*.

**Remark 3.9**

Operations with only one function. If  $f \in \text{Bor}(X, \mathbb{R})$ , then  $|f| \in \text{Bor}(X, \mathbb{R})$ . Use algebraic stability properties to prove this.

**Example 3.1****Problem 3.1**

**Claim:** Let  $\mathfrak{C} = \{(-\infty, t], \mid t \in \mathbb{R}\} \subseteq \mathcal{B}_{\mathbb{R}}$ . Then  $\sigma\text{-Alg}(\mathfrak{C}) = \mathcal{B}_{\mathbb{R}}$ .

**Proof:** Denote  $\sigma\text{-Alg}(\mathfrak{C}) = \mathfrak{P}$ . Clearly,  $\mathfrak{P} \subseteq \mathcal{B}_{\mathbb{R}}$ . If we can show that  $G \in \mathfrak{P}$  for all  $G \subseteq \mathbb{R}$  open, then we will be done because we will have that  $\mathfrak{P}$  is a sigma-algebra and that  $\mathfrak{P} \supseteq \mathcal{T}_{\mathbb{R}}$ .

Now, every  $\emptyset \neq G \subseteq \mathbb{R}$  open can be written as a countable union of open intervals. So then, it suffices to check that  $\mathfrak{P}$  contains all of the open intervals of  $\mathbb{R}$ . This is left as an exercise. □

Let  $(X, \mathfrak{M})$  be a measurable space, and suppose  $f : X \rightarrow \mathbb{R}$  satisfies  $f^{-1}((-\infty, t]) \in \mathfrak{M}$  for all  $t \in \mathbb{R}$ . That is,  $f^{-1}(C) \in \mathfrak{M}$  for all  $C \in \mathfrak{C}$ . Then by [Measurability criterion \(Proposition 3.4\)](#),  $f^{-1}(B) \in \mathfrak{M}$  for all  $B \in \mathcal{B}_{\mathbb{R}}$ , hence  $f \in \text{Bor}(X, \mathbb{R})$ .

## 4 Convergence properties of $\text{Bor}(X, \mathbb{R})$

### Proposition 4.1

### Closure under suprema and infima

Let  $(X, \mathfrak{M})$  be a measurable space.

1. Let  $f : X \rightarrow \mathbb{R}$  be a function. Let  $(f_n)_{n=1}^{\infty}$  be a sequence of functions in  $\text{Bor}(X, \mathbb{R})$  such that

$$\sup\{f_n(x) \mid n \in \mathbb{N}\} = f(x)$$

for all  $x \in X$ . Then it follows that  $f \in \text{Bor}(X, \mathbb{R})$ .

2. Let  $g : X \rightarrow \mathbb{R}$  be a function. Let  $(g_n)_{n=1}^{\infty}$  be a sequence of functions in  $\text{Bor}(X, \mathbb{R})$  such that

$$\inf\{g_n(x) \mid n \in \mathbb{N}\} = g(x)$$

for all  $x \in X$ . Then it follows that  $g \in \text{Bor}(X, \mathbb{R})$ .

### Proof:

1. It suffices to show that  $f^{-1}((-\infty, t]) \in \mathfrak{M}$  for all  $t \in \mathbb{R}$ . This is left as an exercise.
2. Exercise.

□

### Remark 4.2

Let  $(t_n)_{n=1}^{\infty}$  be a bounded sequence in  $\mathbb{R}$ . Then there exists  $\ell_+ \in \mathbb{R}$  such that

1. There exists a subsequence of  $(t_n)_{n=1}^{\infty}$  which converges to  $\ell_+$ .
2. There is no subsequence of  $(t_n)_{n=1}^{\infty}$  which converges to an  $\ell > \ell_+$ .

Define

$$\limsup_{n \rightarrow \infty} t_n := \ell_+$$

We can obtain  $\ell_+$  as

$$\ell_+ = \lim_{k \rightarrow \infty} \left( \sup_{n \geq k} t_n \right) = \inf_{k \geq 1} \left( \sup_{n \geq k} t_n \right)$$

We can define  $\liminf_{n \rightarrow \infty} t_n$  similarly.

Fix a measurable space  $(X, \mathfrak{M})$

**Proposition 4.3**

1. Let  $f : X \rightarrow \mathbb{R}$ . Suppose we could find  $f_1, f_2, \dots \in \text{Bor}(X, \mathbb{R})$  such that for every  $x \in X$ , the sequence  $(f_n(x))_{n=1}^\infty$  is bounded with

$$\limsup_{n \rightarrow \infty} f_n(x) = f(x)$$

Then it follows that  $f \in \text{Bor}(X, \mathbb{R})$

2. Let  $g : X \rightarrow \mathbb{R}$ . Suppose we could find  $g_1, g_2, \dots \in \text{Bor}(X, \mathbb{R})$  such that for every  $x \in X$ , the sequence  $(g_n(x))_{n=1}^\infty$  is bounded with

$$\liminf_{n \rightarrow \infty} g_n(x) = g(x)$$

Then it follows that  $g \in \text{Bor}(X, \mathbb{R})$

**Proof of 1:** For every  $k \in \mathbb{N}$ , define the helper function  $h_k : X \rightarrow \mathbb{R}$  by

$$h_k(x) = \sup \underbrace{\{f_n(x) \mid n \geq k\}}_{\text{bounded by hypothesis}}, \quad \forall x \in X$$

Observe  $f_k, f_{k+1}, \dots \in \text{Bor}(X, \mathbb{R})$ . We can invoke [Closure under suprema and infima \(Proposition 4.1\)](#), to conclude that  $h_k \in \text{Bor}(X, \mathbb{R})$ . Observe for every  $x \in X$ , we have

$$h_1(x) \geq h_2(x) \geq \dots$$

with

$$f(x) = \lim_{k \rightarrow \infty} h_k(x) = \inf\{h_k(x) \mid k \in \mathbb{N}\}$$

This is because  $f(x) = \limsup_{n \rightarrow \infty} f_n(x)$  by hypothesis. But then

$$h_k \in \text{Bor}(X, \mathbb{R}), \quad \forall k \in \mathbb{N}$$

and

$$f(x) = \inf\{h_k(x) \mid k \in \mathbb{N}\} \quad \forall x \in X$$

thus  $f \in \text{Bor}(X, \mathbb{R})$  by [Closure under suprema and infima \(Proposition 4.1\)](#). □

**Proof of 2:** We can reduce to (1), by introducing  $f : X \rightarrow \mathbb{R}$ ,  $f(x) = -g(x) \quad \forall x \in X$  and introducing  $f_n = -g_n \in \text{Bor}(X, \mathbb{R})$ ,  $\forall n \in \mathbb{N}$ , then apply (1) to  $f(x) = \limsup_{n \rightarrow \infty} f_n(x)$ , to get  $f \in \text{Bor}(X, \mathbb{R})$ , hence  $g \in \text{Bor}(X, \mathbb{R})$ . □

**Corollary 4.4**

Let  $f : X \rightarrow \mathbb{R}$ . Suppose we could find  $(f_n)_{n=1}^\infty$  in  $\text{Bor}(X, \mathbb{R})$  such that  $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ ,  $\forall x \in X$ . Then it follows that  $f \in \text{Bor}(X, \mathbb{R})$ .

**Proof:** Use [Proposition 4.3](#). □

**Exercise 4.1**

Suppose that  $(X, \mathfrak{M}) = (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ . Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be differentiable and consider the function  $f' : \mathbb{R} \rightarrow \mathbb{R}$  ( $f'$  may not be continuous).

**Solution:**

Idea is to produce  $(f_n)_{n=1}^{\infty}$  in  $\text{Bor}(\mathbb{R}, \mathbb{R})$  such that  $\lim_{n \rightarrow \infty} f_n(x) = f'(x)$ . If we find the  $f_n$ 's then  $f' \in \text{Bor}(\mathbb{R}, \mathbb{R})$  due to [Corollary 4.4](#). How do we find the  $f_n$ 's? For  $x \in \mathbb{R}$

$$f'(x) = \lim_{n \rightarrow \infty} \frac{f(x + \frac{1}{n}) - f(x)}{\frac{1}{n}} = \lim_{n \rightarrow \infty} n(u_n(x) - f(x))$$

With  $u_n(x) = f(x + \frac{1}{n})$ . So then, for every  $n \in \mathbb{N}$ , let  $u_n : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $u_n(x) = f(x + \frac{1}{n})$ ,  $\forall x \in \mathbb{R}$  and let  $f_n = n(u_n - f)$ . Every  $f_n$  is continuous, hence in  $\text{Bor}(\mathbb{R}, \mathbb{R})$  by [Continuous maps are measurable \(Corollary 3.5\)](#) and  $\lim_{n \rightarrow \infty} f_n(x) = f'(x)$  as we wanted.

**4.1 Approximation by simple functions****Definition 4.5****Simple functions**

A function  $f \in \text{Bor}(X, \mathbb{R})$  is said to be *simple* when it takes finitely many values, that is  $f(X)$  is a finite subset of  $\mathbb{R}$ . We denote

$$\text{Bor}_s(X, \mathbb{R}) := \{f \in \text{Bor}(X, \mathbb{R}) \mid f \text{ is simple}\}$$

We also introduce the spaces

$$\text{Bor}^+(X, \mathbb{R}) = \{f \in \text{Bor}(X, \mathbb{R}) \mid f(x) \geq 0 \forall x \in X\}$$

and

$$\text{Bor}_s^+(X, \mathbb{R}) := \text{Bor}_s(X, \mathbb{R}) \cap \text{Bor}^+(X, \mathbb{R})$$

**Note**

For  $A \in \mathfrak{M}$ , we have  $\chi_A \in \text{Bor}_s(X, \mathbb{R})$ , where  $\chi_A : X \rightarrow \mathbb{R}$  is defined by

$$\chi_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \in X \setminus A \end{cases}$$

called the *indicator function* of  $A$ . Also note that  $\chi_A$  is  $\mathfrak{M}/\mathcal{B}_{\mathbb{R}}$ -measurable since for every  $S \in \mathcal{B}_{\mathbb{R}}$  we have that

$$\chi_A^{-1}(S) \in \{\emptyset, A, X \setminus A, X\} \subseteq \mathfrak{M}$$

**Note**

It can be shown that

$$\text{Bor}_s(X, \mathbb{R}) = \underbrace{\text{span}\{\chi_A \mid A \in \mathfrak{M}\}}_{\substack{\text{set of all linear combinations} \\ \text{of indicator functions} \\ \chi_A \text{ with } A \in \mathfrak{M}}}$$

**Remark 4.6****Binning values of a non-negative Borel function**

Let  $f \in \text{Bor}^+(X, \mathbb{R})$ . For every  $n \in \mathbb{N}$ , let  $f_n : X \rightarrow \mathbb{R}$  be the  $n$ -th binning of the values of  $f$ , defined as follows. For each  $x \in X$ :

1. If  $f(x) \geq n$ , define  $f_n(x) := n$ .
2. If  $f(x) < n$ , pick the unique  $k \in \{1, \dots, n \cdot 2^n\}$  such that

$$\frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n}$$

and define  $f_n(x) := \frac{k-1}{2^n}$ .

**Proposition 4.7****Binning approximation**

Let  $f \in \text{Bor}^+(X, \mathbb{R})$ , and for every  $n \in \mathbb{N}$ , let  $f_n$  be the  $n$ -th binning of the values of  $f$ . Then

1.  $f_n \in \text{Bor}_s^+(X, \mathbb{R})$  for every  $n \in \mathbb{N}$
2.  $f_n \leq f_{n+1}$  for every  $n \in \mathbb{N}$
3. For every  $x \in X$ ,

$$\lim_{n \rightarrow \infty} f_n(x) = f(x)$$

**5 Integration of functions in  $\text{Bor}^+(X, \mathbb{R})$** **5.1 The map  $L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \rightarrow [0, \infty]$** 

Fix a measure space  $(X, \mathfrak{M}, \mu)$

**Note**

For didactical reasons, it is convenient to first define  $\int_X f(x) d\mu(x)$  for  $f \in \text{Bor}^+(X, \mathbb{R})$ . In this case, the integral is always defined but may be  $\infty$ . For general  $f \in \text{Bor}(X, \mathbb{R})$ , the integral may not be defined; to get a finite integral, we need  $f \in \mathcal{L}^1(\mu)$ .

Moreover, it will be convenient to introduce the integral on  $\text{Bor}^+(X, \mathbb{R})$  as a map

$$L^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty].$$

We will get  $L^+$  in two parts:

1. Define  $L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \rightarrow [0, \infty]$
2. Extend  $L_s^+$  to a map  $L^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty]$

The map  $L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \rightarrow [0, \infty]$

**Remark 5.1****Easily verified properties of  $\text{Bor}_s^+(X, \mathbb{R})$** 

1. If  $f, g \in \text{Bor}_s^+(X, \mathbb{R})$ , then  $f + g \in \text{Bor}_s^+(X, \mathbb{R})$
2. If  $f \in \text{Bor}_s^+(X, \mathbb{R})$  and  $\alpha \in [0, \infty)$ , then  $\alpha f \in \text{Bor}_s^+(X, \mathbb{R})$
3. If  $A \in \mathfrak{M}$ , then  $\chi_A \in \text{Bor}_s^+(X, \mathbb{R})$

**Note**

As a consequence of (3), taking  $A = \emptyset$  and  $A = X$ , we get  $\underline{1}, \underline{0} : X \rightarrow \mathbb{R} \in \text{Bor}_s^+(X, \mathbb{R})$ .

**Theorem 5.2****Integral of non-negative simple functions**

There exists a map

$$L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \longrightarrow [0, \infty],$$

uniquely determined, such that the following conditions are satisfied:

(i) Additivity:

$$L_s^+(f + g) = L_s^+(f) + L_s^+(g)$$

for all  $f, g \in \text{Bor}_s^+(X, \mathbb{R})$

(ii) Positive homogeneity:

$$L_s^+(\alpha f) = \alpha L_s^+(f)$$

for all  $f \in \text{Bor}_s^+(X, \mathbb{R})$  and  $\alpha \in [0, \infty)$

(iii) Relation to the measure:

$$L_s^+(\chi_A) = \mu(A)$$

for all  $A \in \mathfrak{M}$

In order to prove the existence of the map  $L_s^+$ , we will take advantage of the fact that one can describe the functions from  $\text{Bor}_s^+(X, \mathbb{R})$  in terms of measurable divisions of  $X$ .

**Definition 5.3****Measurable division and intersections of measurable divisions**

1. A *measurable division* of  $X$  is a set  $\{A_1, \dots, A_n\} \subseteq \mathfrak{M}$  of pairwise disjoint, non-empty sets such that  $\bigcup_{i=1}^n A_i = X$
2. Given two measurable divisions  $\Delta = \{A_1, \dots, A_n\}$  and  $\Gamma = \{B_1, \dots, B_m\}$  of  $X$ , one can “intersect” them, to form a new measurable division of  $X$ , denoted  $\Delta \wedge \Gamma$ , defined by

$$\Delta \wedge \Gamma := \{A_i \cap B_j \mid 1 \leq i \leq n, 1 \leq j \leq m, \text{ and } A_i \cap B_j \neq \emptyset\}$$

**Lemma 5.4****Simple functions and measurable divisions**

Let  $f \in \text{Bor}_s^+(X, \mathbb{R})$ .

1. There exists a measurable division  $\Delta = \{A_1, \dots, A_n\}$  of  $X$  and numbers  $\alpha_1, \dots, \alpha_n \in [0, \infty)$  such that

$$f = \alpha_1 \chi_{A_1} + \dots + \alpha_n \chi_{A_n}$$

2. Suppose that  $\Delta = \{A_1, \dots, A_n\}$  and  $\alpha_1, \dots, \alpha_n \in [0, \infty)$  are as in (1.), and suppose that  $\Gamma = \{B_1, \dots, B_m\}$  is a second measurable division of  $X$ . Then

$$f = \sum_{\substack{1 \leq i \leq n, \\ 1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \alpha_i \chi_{A_i \cap B_j}$$

**Proof of 2:** For every  $1 \leq i \leq n$ , write

$$\chi_{A_i} = \chi_{A_i} \cdot 1 = \chi_{A_i} (\chi_{B_1} + \dots + \chi_{B_m}) = \sum_{j=1}^m \chi_{A_i \cap B_j} = \sum_{\substack{1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \chi_{A_i \cap B_j}$$

Then

$$f = \sum_{i=1}^n \alpha_i \chi_{A_i} = \sum_{i=1}^n \alpha_i \left( \sum_{\substack{1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \chi_{A_i \cap B_j} \right) = \sum_{\substack{1 \leq i \leq n, \\ 1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \alpha_i \chi_{A_i \cap B_j}$$

as desired. □

**Lemma 5.5****Well-definedness of the simple integral formula**

Let  $f \in \text{Bor}_s^+(X, \mathbb{R})$ . Suppose we have two ways of writing  $f$  as a linear combination:

$$f = \alpha_1 \chi_{A_1} + \dots + \alpha_n \chi_{A_n}$$

and

$$f = \beta_1 \chi_{B_1} + \dots + \beta_m \chi_{B_m},$$

where  $\Delta = \{A_1, \dots, A_n\}$  and  $\Gamma = \{B_1, \dots, B_m\}$  are measurable divisions of  $X$ , and  $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m \in [0, \infty)$ . Then

$$\sum_{i=1}^n \alpha_i \mu(A_i) = \sum_{j=1}^m \beta_j \mu(B_j) \in [0, \infty].$$

**Proof:** For every  $1 \leq i \leq n$ , let us write

$$A_i = A_i \cap X = A_i \cap (B_1 \cup \dots \cup B_m) = \bigcup_{j=1}^m (A_i \cap B_j) = \bigcup_{\substack{1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} (A_i \cap B_j)$$

Since the union is disjoint,

$$\mu(A_i) = \sum_{\substack{1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \mu(A_i \cap B_j)$$

so

$$\sum_{i=1}^n \alpha_i \mu(A_i) = \sum_{i=1}^n \alpha_i \sum_{\substack{1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \mu(A_i \cap B_j) = \sum_{\substack{1 \leq i \leq n, \\ 1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \alpha_i \mu(A_i \cap B_j)$$

Similarly,

$$\sum_{j=1}^m \beta_j \mu(B_j) = \sum_{j=1}^m \beta_j \sum_{\substack{1 \leq i \leq n, \\ B_j \cap A_i \neq \emptyset}} \mu(B_j \cap A_i) = \sum_{\substack{1 \leq i \leq n, \\ 1 \leq j \leq m, \\ A_i \cap B_j \neq \emptyset}} \beta_j \mu(A_i \cap B_j)$$

**Claim:** If  $A_i \cap B_j \neq \emptyset$ , then  $\alpha_i = \beta_j$

Observe that this claim will finish the proof. To verify the claim, suppose  $A_i \cap B_j \neq \emptyset$ . Pick  $x \in A_i \cap B_j$ , and evaluate  $f(x)$  in two ways:

$$f(x) = \alpha_1 \chi_{A_1}(x) + \dots + \alpha_n \chi_{A_n}(x) = \alpha_i$$

similarly

$$f(x) = \beta_1 \chi_{B_1}(x) + \dots + \beta_m \chi_{B_m}(x) = \beta_j,$$

so  $\alpha_i = f(x) = \beta_j$

□

**Proof of existence of  $L_s^+$  from *Integral of non-negative simple functions (Theorem 5.2)*:** For every  $f \in \text{Bor}_s^+(X, \mathbb{R})$ , we define the quantity  $L_s^+(f) \in [0, \infty]$  by the following procedure:

Consider a writing

$$f = \alpha_1 \chi_{A_1} + \dots + \alpha_n \chi_{A_n}$$

with  $\Delta = \{A_1, \dots, A_n\}$  a measurable division of  $X$  and  $\alpha_1, \dots, \alpha_n \in [0, \infty)$ ,  
and put

$$L_s^+(f) := \sum_{i=1}^n \alpha_i \mu(A_i) \in [0, \infty].$$

The map  $L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \longrightarrow [0, \infty]$



**Well-definedness of the simple integral formula (Lemma 5.5)** assures us that the procedure  $(*)$  is coherent, in the sense that the proposed quantity  $L_s^+(f)$  only depends on  $f$  and not on the choice of how we write  $f$  as a linear combination of indicators.

With  $(*)$ , we have thus defined a map

$$L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \longrightarrow [0, \infty].$$

We must now check that this map has the properties (i), (ii), (iii) indicated in **Integral of non-negative simple functions (Theorem 5.2)**. □

## 5.2 The map $L^+ : \text{Bor}^+(X, \mathbb{R}) \longrightarrow [0, \infty]$

### Definition 5.6

### Integral of non-negative Borel functions

For every function  $u \in \text{Bor}^+(X, \mathbb{R})$ , we define

$$L^+(u) := \sup\{L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\} \in [0, \infty].$$

### Remark 5.7

### Comments on the formula for $L^+$

- We use letters such as  $u, v, \dots$  for general functions in  $\text{Bor}^+(X, \mathbb{R})$ , reserving  $f, g, \dots$  for functions in  $\text{Bor}_s^+(X, \mathbb{R})$ .
- In the formula above,  $f \leq u$  means  $f(x) \leq u(x)$  for every  $x \in X$ .
- The set over which we take the supremum is non-empty because it contains  $\underline{0}$ .
- The supremum is taken in  $[0, \infty]$ , so it is allowed to be  $\infty$ .

### Proposition 5.8

### $L^+$ extends $L_s^+$

$L^+$  is an extension of  $L_s^+$ . That is, if  $u \in \text{Bor}_s^+(X, \mathbb{R})$ , then the newly defined quantity  $L^+(u)$  is the same as  $L_s^+(u)$ .

**Proof:** Must show

$$\sup\{L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\} = L_s^+(u)$$

“ $\geq$ ” holds because  $L_s^+(f)$  is an element of  $\{L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\}$ . For “ $\leq$ ”, we must show  $L_s^+(u)$  is an upper bound for  $\{L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\}$ . Indeed, if  $f \in \text{Bor}_s^+(X, \mathbb{R})$  with  $f \leq u$ , then, using **Integral of non-negative simple functions (Theorem 5.2)**,

$$L_s^+(u) = L_s^+(f + (u - f)) = L_s^+(f) + L_s^+(u - f) \geq L_s^+(f)$$

□

### Proposition 5.9

### Monotonicity and homogeneity of $L^+$

1.  $L^+$  is an increasing map: if  $u, v \in \text{Bor}^+(X, \mathbb{R})$  with  $u \leq v$ , then  $L^+(u) \leq L^+(v)$
2.  $L^+$  is homogeneous: if  $u \in \text{Bor}^+(X, \mathbb{R})$ ,  $\alpha \in [0, \infty)$ , then  $L^+(\alpha u) = \alpha L^+(u)$

**Proof of 1:** Since  $u \leq v$ , we have

The map  $L^+ : \text{Bor}^+(X, \mathbb{R}) \longrightarrow [0, \infty]$

$$\{f \in \text{Bor}_s^+(X, \mathbb{R}) \mid f \leq u\} \subseteq \{f \in \text{Bor}_s^+(X, \mathbb{R}) \mid f \leq v\}.$$

Thus

$$\begin{aligned} L^+(u) &= \sup\{L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\} \\ &\leq \sup\{L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq v\} \\ &= L^+(v). \end{aligned}$$

□

**Proof of 2:** If  $\alpha = 0$ , then  $L^+(\alpha u) = 0 = \alpha L^+(u)$ , where we use the convention  $0 \cdot \infty = 0$ .

Suppose  $\alpha > 0$ . Then

$$\{g \in \text{Bor}_s^+(X, \mathbb{R}) \mid g \leq \alpha u\} = \{\alpha f \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\}.$$

Thus,

$$\begin{aligned} L^+(\alpha u) &= \sup\{L_s^+(\alpha f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\} \\ &= \sup\{\alpha L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\} \\ &= \alpha \sup\{L_s^+(f) \mid f \in \text{Bor}_s^+(X, \mathbb{R}), f \leq u\} \\ &= \alpha L^+(u), \end{aligned}$$

where the second equality uses [Integral of non-negative simple functions \(Theorem 5.2\)](#).

□

#### Remark 5.10

#### Additivity and MCT

$L^+$  is additive:  $L^+(u + v) = L^+(u) + L^+(v)$  for every  $u, v \in \text{Bor}^+(X, \mathbb{R})$ . We can show  $L^+(u + v) \geq L^+(u) + L^+(v)$  by an easy argument using the definition, but the other inequality is not so easy. Instead we move to the Monotone Convergence Theorem (MCT), which will prove it. Note we have to make sure not to use additivity in the proof of MCT.

## 6 The Monotone Convergence Theorem

Fix a measure space  $(X, \mathfrak{M}, \mu)$  from chapter 5. We will work with the map  $L^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty]$ .

### Notation 6.1

### Increasing convergence

For  $u$  and  $u_1, u_2, \dots \in \text{Bor}^+(X, \mathbb{R})$ , we will use the shorthand  $u_n \nearrow u$  to mean that  $u_1 \leq u_2 \leq \dots$  and  $u_n \rightarrow u$  pointwise.

### Theorem 6.2

### Monotone Convergence Theorem

Let  $u, u_1, u_2, \dots \in \text{Bor}^+(X, \mathbb{R})$  with  $u_n \nearrow u$ . Then

$$\lim_{n \rightarrow \infty} L^+(u_n) = L^+(u)$$

(an increasing limit in  $[0, \infty]$ )

**Proposition 6.3****The missing inequality for additivity of  $L^+$** 

Let  $u, v \in \text{Bor}^+(X, \mathbb{R})$ . Then

$$L^+(u + v) \leq L^+(u) + L^+(v).$$

**Proof:** By [Binning approximation \(Proposition 4.7\)](#), choose sequences  $(f_n)_{n=1}^\infty$  and  $(g_n)_{n=1}^\infty$  in  $\text{Bor}_s^+(X, \mathbb{R})$  such that  $f_n \nearrow u$  and  $g_n \nearrow v$ . It follows that  $(f_n + g_n)_{n=1}^\infty$  is a sequence in  $\text{Bor}_s^+(X, \mathbb{R})$  such that  $f_n + g_n \nearrow u + v$ .

Thus,

$$\begin{aligned} L^+(u + v) &= \lim_{n \rightarrow \infty} L^+(f_n + g_n) \\ &= \lim_{n \rightarrow \infty} L_s^+(f_n + g_n) \\ &= \lim_{n \rightarrow \infty} (L_s^+(f_n) + L_s^+(g_n)) \\ &= \lim_{n \rightarrow \infty} (L^+(f_n) + L^+(g_n)). \end{aligned}$$

The first equality uses [Monotone Convergence Theorem \(Theorem 6.2\)](#), the second and fourth use  [\$L^+\$  extends  \$L\_s^+\$  \(Proposition 5.8\)](#), and the third uses [Integral of non-negative simple functions \(Theorem 5.2\)](#).

Now for every  $n \in \mathbb{N}$ , since  $f_n \leq u$  and  $g_n \leq v$ , [Monotonicity and homogeneity of  \$L^+\$  \(Proposition 5.9\)](#) gives

$$L^+(f_n) + L^+(g_n) \leq L^+(u) + L^+(v).$$

Hence

$$L^+(u + v) \leq L^+(u) + L^+(v),$$

as desired. □

## 6.1 Proof of the Monotone Convergence Theorem

### Remark 6.4

If  $u_n \nearrow u$  in  $\text{Bor}^+(X, \mathbb{R})$ , then

$$\Lambda := \lim_{n \rightarrow \infty} L^+(u_n) = \sup\{L^+(u_n) \mid n \in \mathbb{N}\} \quad (\text{point-to-prove})$$

will satisfy the inequality  $\Lambda \geq L^+(u)$ .

To see that this is enough to prove [Monotone Convergence Theorem \(Theorem 6.2\)](#), suppose  $u_n \nearrow u$  in  $\text{Bor}^+(X, \mathbb{R})$ . Since  $u_n \leq u$  for every  $n \in \mathbb{N}$ , [Monotonicity and homogeneity of  \$L^+\$  \(Proposition 5.9\)](#) gives

$$L^+(u_n) \leq L^+(u)$$

for every  $n \in \mathbb{N}$ . Hence  $L^+(u)$  is an upper bound for  $\{L^+(u_n) \mid n \in \mathbb{N}\}$ , so

$$\Lambda = \sup\{L^+(u_n) \mid n \in \mathbb{N}\} \leq L^+(u).$$

Thus, if we prove (point-to-prove), then  $\Lambda = L^+(u)$ , which is exactly the conclusion of [Monotone Convergence Theorem \(Theorem 6.2\)](#).

We will obtain (point-to-prove) in [Proposition 6.7](#) below. Towards that end, we will prove two special cases of (point-to-prove), which are covered by [Lemma 6.5](#) and [Lemma 6.6](#).

### Lemma 6.5

Let  $u_1 \leq u_2 \leq \dots \leq u_n \leq \dots$  be in  $\text{Bor}^+(X, \mathbb{R})$  and let  $A \in \mathfrak{M}$  be such that  $u_n \nearrow \chi_A$ . Consider the increasing limit

$$\Lambda := \lim_{n \rightarrow \infty} L^+(u_n) \in [0, \infty].$$

Then  $\Lambda \geq \mu(A)$ , where we note that  $\mu(A) = L^+(\chi_A)$ .

**Proof:** Assume  $A \neq \emptyset$ , since if  $A = \emptyset$ , then  $\mu(A) = 0$  and there is nothing to prove. Fix for the moment a  $\theta \in (0, 1)$ . For every  $n \in \mathbb{N}$ , define

$$A_n := \{x \in A \mid u_n(x) > \theta\}$$

Observe:

1.  $A_n \in \mathfrak{M}$  for all  $n \in \mathbb{N}$ . This is because  $A_n = A \cap u_n^{-1}((\theta, \infty))$  with  $A, u_n^{-1}((\theta, \infty)) \in \mathfrak{M}$ .
2.  $A_n \subseteq A_{n+1}$  for all  $n \in \mathbb{N}$ . Indeed, if  $x \in A_n$ , then  $u_n(x) > \theta$ , and since  $u_n \leq u_{n+1}$ , we get  $u_{n+1}(x) > \theta$ , so  $x \in A_{n+1}$ .
3.  $\bigcup_{n=1}^{\infty} A_n = A$ . Indeed,  $A_n \subseteq A$  for all  $n \in \mathbb{N}$ , hence  $\bigcup_{n=1}^{\infty} A_n \subseteq A$ . For “ $\supseteq$ ”, pick  $x \in A$ . Since  $u_n \nearrow \chi_A$ , we have  $u_n(x) \rightarrow \chi_A(x) = 1$ . Since  $\theta < 1$ , there exists  $n \in \mathbb{N}$  such that  $u_n(x) > \theta$ , hence  $x \in A_n$ .

Thus  $(A_n)_{n=1}^{\infty}$  is an increasing chain of sets in  $\mathfrak{M}$  with  $\bigcup_{n=1}^{\infty} A_n = A$ . Using [Continuity along chains \(Proposition 2.12\)](#), we have

$$\lim_{n \rightarrow \infty} \mu(A_n) = \mu(A) \quad (*)$$

Now observe that for every  $n \in \mathbb{N}$  we have  $u_n \geq \theta \chi_{A_n}$ . Indeed, if  $x \notin A_n$ , then  $\theta \chi_{A_n}(x) = 0 \leq u_n(x)$ , while if  $x \in A_n$ , then  $\theta \chi_{A_n}(x) = \theta < u_n(x)$ .

Hence by [Monotonicity and homogeneity of  \$L^+\$](#)  (Proposition 5.9),  $L^+$  extends  $L_s^+$  (Proposition 5.8), and [Integral of non-negative simple functions](#) (Theorem 5.2),

$$L^+(u_n) \geq L^+(\theta \chi_{A_n}) = \theta L^+(\chi_{A_n}) = \theta L_s^+(\chi_{A_n}) = \theta \mu(A_n).$$

So we have

$$\text{For every } n \in \mathbb{N}, L^+(u_n) \geq \theta \mu(A_n) \quad (**)$$

Letting  $n \rightarrow \infty$  in  $(**)$ , we get

$$\Lambda \geq \theta \mu(A)$$

where we use  $(*)$  for the right-hand side. Finally, unfix  $\theta$ . We have  $\Lambda \geq \theta \mu(A)$  for every  $\theta \in (0, 1)$ , and hence  $\Lambda \geq \mu(A)$ , as desired.  $\square$

### Lemma 6.6

Let  $u_1 \leq u_2 \leq \dots \leq u_n \leq \dots$  be in  $\text{Bor}^+(X, \mathbb{R})$  and let  $f_0 \in \text{Bor}_s^+(X, \mathbb{R})$  be such that  $u_n \nearrow f_0$ . Consider the increasing limit

$$\Lambda := \lim_{n \rightarrow \infty} L^+(u_n) \in [0, \infty].$$

Then  $\Lambda \geq L_s^+(f_0)$ , where we note that  $L_s^+(f_0) = L^+(f_0)$ .

### Proposition 6.7

Let  $u_n \nearrow u$  in  $\text{Bor}^+(X, \mathbb{R})$ , and let us consider the increasing limit

$$\Lambda := \lim_{n \rightarrow \infty} L^+(u_n) \in [0, \infty].$$

Then  $\Lambda \geq L^+(u)$ .

### Theorem 6.8

### Characterization of $L^+$

The map  $L^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty]$  has the following properties, which determine it uniquely:

- (i)  $L^+(\chi_A) = \mu(A)$  for every  $A \in \mathfrak{M}$ .
- (ii)  $L^+(u + v) = L^+(u) + L^+(v)$  for every  $u, v \in \text{Bor}^+(X, \mathbb{R})$ .
- (iii)  $L^+(\alpha u) = \alpha \cdot L^+(u)$  for every  $u \in \text{Bor}^+(X, \mathbb{R})$  and  $\alpha \in [0, \infty)$ .
- (iv)  $\lim_{n \rightarrow \infty} L^+(u_n) = L^+(u)$  whenever  $u$  and  $(u_n)_{n=1}^\infty$  are in  $\text{Bor}^+(X, \mathbb{R})$  and  $u_n \nearrow u$ .

**Note**

For the existence part in [Characterization of  \$L^+\$  \(Theorem 6.8\)](#), property (iv) is [Monotone Convergence Theorem \(Theorem 6.2\)](#), while properties (i), (ii), (iii) follow from [Integral of non-negative simple functions \(Theorem 5.2\)](#), combined with the fact that  $L^+$  is an extension of the map  $L_s^+ : \text{Bor}_s^+(X, \mathbb{R}) \rightarrow [0, \infty]$ .

For the uniqueness part in [Characterization of  \$L^+\$  \(Theorem 6.8\)](#), it is convenient to record the following lemma, which will also turn out to be useful in subsequent lectures.

**Lemma 6.9****Method of the friendly functions**

Let  $(X, \mathfrak{M})$  be a measurable space and consider a set of functions  $\mathcal{G} \subseteq \text{Bor}^+(X, \mathbb{R})$  which has the properties listed below.

- (j)  $\chi_A \in \mathcal{G}$  for every  $A \in \mathfrak{M}$ .
- (jj) If  $g_1, g_2 \in \mathcal{G}$ , then  $g_1 + g_2 \in \mathcal{G}$ .
- (jjj) If  $g \in \mathcal{G}$  and  $\alpha \in [0, \infty)$ , then  $\alpha g \in \mathcal{G}$ .
- (jv) If  $g \in \text{Bor}^+(X, \mathbb{R})$ ,  $g_1 \leq g_2 \leq \dots \leq g_n \leq \dots$  in  $\mathcal{G}$ , and  $g_n \nearrow g$ , then  $g \in \mathcal{G}$ .

Under these hypotheses, it follows that  $\mathcal{G} = \text{Bor}^+(X, \mathbb{R})$ .

**Proof:**

- First, every  $f \in \text{Bor}_s^+(X, \mathbb{R})$  belongs to  $\mathcal{G}$ . Indeed, by [Simple functions and measurable divisions \(Lemma 5.4\)](#), we can write

$$f = \alpha_1 \chi_{A_1} + \dots + \alpha_n \chi_{A_n}$$

with  $A_1, \dots, A_n \in \mathfrak{M}$  and  $\alpha_1, \dots, \alpha_n \in [0, \infty)$ . By properties (j), (jj), and (jjj), it follows that  $f \in \mathcal{G}$ .

- Now let  $u \in \text{Bor}^+(X, \mathbb{R})$ . By [Binning approximation \(Proposition 4.7\)](#), there exists a sequence  $(f_n)_{n=1}^\infty$  in  $\text{Bor}_s^+(X, \mathbb{R})$  such that  $f_n \nearrow u$ . By the previous bullet,  $f_n \in \mathcal{G}$  for every  $n \in \mathbb{N}$ , and hence property (jv) gives  $u \in \mathcal{G}$ .

□

**Proof of uniqueness in [Characterization of  \$L^+\$  \(Theorem 6.8\)](#):** Suppose we are given a map

$$M^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty]$$

which also satisfies the conditions (i)-(iv) in [Characterization of  \$L^+\$  \(Theorem 6.8\)](#). Let

$$\mathcal{G} = \{g \in \text{Bor}^+(X, \mathbb{R}) \mid L^+(g) = M^+(g)\} \subseteq \text{Bor}^+(X, \mathbb{R})$$

We will show that  $\mathcal{G} = \text{Bor}^+(X, \mathbb{R})$  by checking the hypotheses of [Method of the friendly functions \(Lemma 6.9\)](#).

- If  $A \in \mathfrak{M}$ , then  $\chi_A \in \mathcal{G}$  because

$$L^+(\chi_A) = \mu(A) = M^+(\chi_A)$$

- If  $g_1, g_2 \in \mathcal{G}$ , then  $g_1 + g_2 \in \mathcal{G}$  because

$$L^+(g_1 + g_2) = L^+(g_1) + L^+(g_2) = M^+(g_1) + M^+(g_2) = M^+(g_1 + g_2).$$

- If  $g \in \mathcal{G}$  and  $\alpha \in [0, \infty)$ , then  $\alpha g \in \mathcal{G}$  because

$$L^+(\alpha g) = \alpha L^+(g) = \alpha M^+(g) = M^+(\alpha g).$$

- Suppose  $g \in \text{Bor}^+(X, \mathbb{R})$ ,  $g_1 \leq g_2 \leq \dots \leq g_n \leq \dots$  in  $\mathcal{G}$ , and  $g_n \nearrow g$ . Then

$$L^+(g) = \lim_{n \rightarrow \infty} L^+(g_n) = \lim_{n \rightarrow \infty} M^+(g_n) = M^+(g),$$

so  $g \in \mathcal{G}$ .

Thus  $\mathcal{G} = \text{Bor}^+(X, \mathbb{R})$  by [Method of the friendly functions \(Lemma 6.9\)](#). Hence  $L^+(g) = M^+(g)$  for every  $g \in \text{Bor}^+(X, \mathbb{R})$ , so  $L^+$  is uniquely determined.  $\square$

## 7 The space of integrable functions $\mathcal{L}^1(\mu)$

### 7.1 Notion of integrable function, the map $L : \mathcal{L}^1(\mu) \rightarrow \mathbb{R}$

#### Remark 7.1

#### Framework

In this part, we continue to work with the fixed measure space  $(X, \mathfrak{M}, \mu)$  from the previous chapter. There we considered

$$\text{Bor}^+(X, \mathbb{R}) = \{f \in \text{Bor}(X, \mathbb{R}) \mid f(x) \geq 0 \ \forall x \in X\}$$

and defined a map  $L^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty]$ , which is trying to become the integral with respect to  $\mu$ .

We now want to work with the full space  $\text{Bor}(X, \mathbb{R})$ . The idea is to reduce a function  $f \in \text{Bor}(X, \mathbb{R})$  to a pair of functions  $f_+, f_- \in \text{Bor}^+(X, \mathbb{R})$ , called the *positive part* and *negative part* of  $f$ , defined by

$$f_+ := f \vee \underline{0} \quad \text{and} \quad f_- := (-f) \vee \underline{0}.$$

Here  $\vee$  denotes the pointwise maximum of two functions. Equivalently, for every  $x \in X$ ,

$$f_+(x) = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ 0 & \text{if } f(x) < 0 \end{cases} \quad \text{and} \quad f_-(x) = \begin{cases} 0 & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}.$$

From these formulas we have

$$f_+ + f_- = |f| \quad \text{and} \quad f_+ - f_- = f.$$

We also have the useful expressions

$$f_+ = \frac{f + |f|}{2} \quad \text{and} \quad f_- = \frac{|f| - f}{2}.$$

#### Lemma 7.2

For  $f \in \text{Bor}(X, \mathbb{R})$ , we have the equivalence

$$L^+(|f|) < \infty \iff (L^+(f_+) < \infty \text{ and } L^+(f_-) < \infty).$$

**Proof:** This is immediate from the fact that

$$L^+(|f|) = L^+(f_+) + L^+(f_-),$$

which follows from  $|f| = f_+ + f_-$  and the additivity of  $L^+$  from [Characterization of  \$L^+\$  \(Theorem 6.8\)](#).  
□

### Definition 7.3

### Integrable functions and the map $L$

A function  $f : X \rightarrow \mathbb{R}$  is said to be *integrable with respect to  $\mu$*  if  $f \in \text{Bor}(X, \mathbb{R})$  and the equivalent conditions from [Lemma 7.2](#) hold. We denote

$$\mathcal{L}^1(\mu) := \{f : X \rightarrow \mathbb{R} \mid f \text{ is integrable with respect to } \mu\}.$$

We define a map  $L : \mathcal{L}^1(\mu) \rightarrow \mathbb{R}$  by putting, for every  $f \in \mathcal{L}^1(\mu)$ ,

$$L(f) := L^+(f_+) - L^+(f_-).$$

### Note

We will prove in [Linearity of  \$L\$  \(Theorem 7.5\)](#) that the function  $L$  is linear. Towards that end, we first prove a lemma which gives some leeway in how  $L(f)$  is calculated: instead of having to stick with  $f = f_+ - f_-$ , the lemma allows us to use a writing  $f = h_1 - h_2$ .

### Lemma 7.4

Let  $f \in \mathcal{L}^1(\mu)$  and suppose that we can write  $f = h_1 - h_2$  with  $h_1, h_2 \in \text{Bor}^+(X, \mathbb{R})$  such that  $L^+(h_1), L^+(h_2) < \infty$ . Then

$$L(f) = L^+(h_1) - L^+(h_2).$$

**Proof:** Since  $f = h_1 - h_2$  and  $f = f_+ - f_-$ , we have

$$h_1 + f_- = h_2 + f_+.$$

Applying  $L^+$  to both sides, and using additivity from [Characterization of  \$L^+\$  \(Theorem 6.8\)](#), gives

$$L^+(h_1) + L^+(f_-) = L^+(h_2) + L^+(f_+).$$

All four quantities appearing here are finite, so we can rearrange this equality to obtain

$$L^+(f_+) - L^+(f_-) = L^+(h_1) - L^+(h_2).$$

The left-hand side is  $L(f)$  by definition, and the result follows. □

### Theorem 7.5

### Linearity of $L$

- 1  $\mathcal{L}^1(\mu)$  is stable under linear combinations, hence is a linear subspace of  $\text{Bor}(X, \mathbb{R})$ .
- 2 The map  $L : \mathcal{L}^1(\mu) \rightarrow \mathbb{R}$  is linear.

**Proof of 1:** Pick some  $f, g \in \mathcal{L}^1(\mu)$  and  $\alpha, \beta \in \mathbb{R}$ , and form the linear combination



$$h := \alpha f + \beta g \in \text{Bor}(X, \mathbb{R}).$$

We want to prove that  $h \in \mathcal{L}^1(\mu)$ . Observe that

$$|h| = |\alpha f + \beta g| \leq |\alpha| \cdot |f| + |\beta| \cdot |g|,$$

which implies that

$$\begin{aligned} L^+(|h|) &\leq L^+(|\alpha| \cdot |f| + |\beta| \cdot |g|) \\ &= |\alpha| \cdot L^+(|f|) + |\beta| \cdot L^+(|g|) \\ &< \infty. \end{aligned}$$

The first inequality follows from the fact that  $L^+$  is increasing, and the second equality follows from the additivity and homogeneity properties in [Characterization of  \$L^+\$  \(Theorem 6.8\)](#). Since  $f, g \in \mathcal{L}^1(\mu)$ , the last expression is finite. It follows that  $L^+(|h|) < \infty$ , so  $h \in \mathcal{L}^1(\mu)$  by [Lemma 7.2](#).  $\square$

**Proof of 2:** We will verify separately the two statements which, together, give linearity:

- (a)  $L(f + g) = L(f) + L(g)$  for every  $f, g \in \mathcal{L}^1(\mu)$ .
- (b)  $L(\alpha f) = \alpha L(f)$  for every  $f \in \mathcal{L}^1(\mu)$  and  $\alpha \in \mathbb{R}$ .

*Verification of (a):*

Pick  $f, g \in \mathcal{L}^1(\mu)$ . By 1,  $f + g \in \mathcal{L}^1(\mu)$ . We can write

$$f + g = (f_+ + g_+) - (f_- + g_-),$$

where  $f_+ + g_+$  and  $f_- + g_-$  are in  $\text{Bor}^+(X, \mathbb{R})$ . Moreover,

$$L^+(f_+ + g_+) = L^+(f_+) + L^+(g_+) < \infty$$

and

$$L^+(f_- + g_-) = L^+(f_-) + L^+(g_-) < \infty,$$

since  $f, g \in \mathcal{L}^1(\mu)$ . Thus [Lemma 7.4](#) gives

$$\begin{aligned} L(f + g) &= L^+(f_+ + g_+) - L^+(f_- + g_-) \\ &= L^+(f_+) + L^+(g_+) - L^+(f_-) - L^+(g_-) \\ &= L(f) + L(g). \end{aligned}$$

*Verification of (b):*

Pick  $f \in \mathcal{L}^1(\mu)$  and  $\alpha \in \mathbb{R}$ . By 1,  $\alpha f \in \mathcal{L}^1(\mu)$ .

If  $\alpha \geq 0$ , then

$$\alpha f = \alpha f_+ - \alpha f_-,$$

and  $L^+(\alpha f_+) = \alpha L^+(f_+) < \infty$ ,  $L^+(\alpha f_-) = \alpha L^+(f_-) < \infty$ . Hence [Lemma 7.4](#) gives

$$L(\alpha f) = \alpha L^+(f_+) - \alpha L^+(f_-) = \alpha L(f).$$

If  $\alpha < 0$ , then

$$\alpha f = (-\alpha)f_- - (-\alpha)f_+,$$

and  $L^+((- \alpha)f_-) = (- \alpha)L^+(f_-) < \infty$ ,  $L^+((- \alpha)f_+) = (- \alpha)L^+(f_+) < \infty$ . Hence [Lemma 7.4](#) gives

$$L(\alpha f) = (- \alpha)L^+(f_-) - (- \alpha)L^+(f_+) = \alpha L(f).$$

□

### Proposition 7.6

1. If  $f \in \mathcal{L}^1(\mu)$  and also  $f \in \text{Bor}^+(X, \mathbb{R})$ , then  $L(f) = L^+(f)$ .
2. If  $f, g \in \mathcal{L}^1(\mu)$  are such that  $f \leq g$ , then  $L(f) \leq L(g)$ .
3. If  $f \in \mathcal{L}^1(\mu)$ , then  $|f| \in \mathcal{L}^1(\mu)$  as well, and

$$|L(f)| \leq L(|f|).$$

**Proof of 1:** Since  $f \in \text{Bor}^+(X, \mathbb{R})$ , we have  $f_+ = f$  and  $f_- = \underline{0}$ . Hence

$$L(f) = L^+(f_+) - L^+(f_-) = L^+(f) - L^+(\underline{0}) = L^+(f).$$

□

**Proof of 2:** If  $f \leq g$ , then  $g - f \in \mathcal{L}^1(\mu)$  by [Linearity of  \$L\$  \(Theorem 7.5\)](#), and moreover  $g - f \in \text{Bor}^+(X, \mathbb{R})$ . By part 1,

$$L(g - f) = L^+(g - f) \geq 0.$$

Using the linearity of  $L$ , this says that  $L(g) - L(f) \geq 0$ , hence  $L(f) \leq L(g)$ .

□

**Proof of 3:** Since  $f \in \mathcal{L}^1(\mu)$ , [Lemma 7.2](#) gives  $L^+(|f|) < \infty$ , so  $|f| \in \mathcal{L}^1(\mu)$ .

As functions, we have

$$-|f| \leq f \leq |f|.$$

Applying part 2 gives

$$-L(|f|) \leq L(f) \leq L(|f|),$$

where we also used the linearity of  $L$  on the left. Thus

$$|L(f)| \leq L(|f|).$$

□

**Remark 7.7****Switching to integral notation**

For our measure space  $(X, \mathfrak{M}, \mu)$ , we have now constructed two maps

$$L^+ : \text{Bor}^+(X, \mathbb{R}) \rightarrow [0, \infty] \quad \text{and} \quad L : \mathcal{L}^1(\mu) \rightarrow \mathbb{R},$$

which are trying to become “the integral with respect to  $\mu$ ”. If  $f \in \text{Bor}(X, \mathbb{R})$  is such that both values  $L^+(f)$  and  $L(f)$  are defined, then [Proposition 7.6](#) says that  $L(f) = L^+(f)$ .

It is therefore safe to switch to the customary integral notation:

if at least one of  $L^+(f), L(f)$  is defined, we denote that value by

$$\int_X f(x) \, d\mu(x).$$

This is often abbreviated to just  $\int f \, d\mu$ .

While we now have uniform notation for all our integrals, it is useful to remember that the integral notation can have two meanings:

- an  $L^+$  meaning, when it has values in  $[0, \infty]$  and we use the properties of  $L^+$ ;
- an  $L$  meaning, when it has values in  $\mathbb{R}$  and we use the properties of  $L$ .

This usually causes no harm. For instance, the inequality from [Proposition 7.6](#) becomes

$$\left| \int_X f(x) \, d\mu(x) \right| \leq \int_X |f(x)| \, d\mu(x), \quad \forall f \in \mathcal{L}^1(\mu),$$

where the integral on the right-hand side can be viewed either in the  $L^+$  sense or in the  $L$  sense.

## 7.2 Integration of subsets

### Notation 7.8

Let  $(X, \mathfrak{M}, \mu)$  be a measure space.

1. For any  $f \in \text{Bor}^+(X, \mathbb{R})$  and  $A \in \mathfrak{M}$ , we denote

$$\int_A f \, d\mu := \int_X f \cdot \chi_A \, d\mu \in [0, \infty],$$

where, as usual,  $\chi_A$  denotes the indicator function of  $A$ .

2. For any  $f \in \mathcal{L}^1(\mu)$  and  $A \in \mathfrak{M}$ , observe that  $f \cdot \chi_A \in \mathcal{L}^1(\mu)$ , since  $|f \cdot \chi_A| \leq |f|$ . We denote

$$\int_A f \, d\mu := \int_X f \cdot \chi_A \, d\mu \in \mathbb{R}.$$

**Proposition 7.9**

Let  $(X, \mathfrak{M}, \mu)$  be a measure space and let  $h \in \text{Bor}^+(X, \mathbb{R})$ . We define a set-function  $\nu : \mathfrak{M} \rightarrow [0, \infty]$  by putting

$$\nu(A) := \int_A h \, d\mu, \quad A \in \mathfrak{M}.$$

Then  $\nu$  is a positive measure on  $\mathfrak{M}$ .

**Proof:**

1.  $\nu(\emptyset) = \int_X h \cdot \chi_{\emptyset} \, d\mu = \int_X 0 \, d\mu = 0.$
2. Fix  $(A_n)_{n=1}^{\infty}$  in  $\mathfrak{M}$  such that  $A_i \cap A_j = \emptyset$  for  $i \neq j$ , and let  $U = \bigcup_{n=1}^{\infty} A_n$ . We must prove that

$$\nu(U) = \sum_{n=1}^{\infty} \nu(A_n).$$

For every  $N \in \mathbb{N}$ , let

$$s_N = \sum_{n=1}^N h \cdot \chi_{A_n}.$$

Then  $(s_N)_{N=1}^{\infty}$  is an increasing sequence in  $\text{Bor}^+(X, \mathbb{R})$ , and

$$s_N \rightarrow h \cdot \chi_U$$

pointwise. Hence, by [Monotone Convergence Theorem \(Theorem 6.2\)](#),

$$\begin{aligned} \nu(U) &= \int_X h \cdot \chi_U \, d\mu \\ &= \lim_{N \rightarrow \infty} \int_X s_N \, d\mu \\ &= \lim_{N \rightarrow \infty} \sum_{n=1}^N \int_X h \chi_{A_n} \, d\mu \\ &= \sum_{n=1}^{\infty} \nu(A_n). \end{aligned}$$

□

**Proposition 7.10**

Let  $(X, \mathfrak{M}, \mu)$ ,  $h \in \text{Bor}^+(X, \mathbb{R})$ , and the positive measure  $\nu : \mathfrak{M} \rightarrow [0, \infty]$  be as in [Proposition 7.9](#). Then, for every  $f \in \text{Bor}^+(X, \mathbb{R})$ ,

$$\int_X f \, d\nu = \int_X f \cdot h \, d\mu \in [0, \infty].$$

This is an equality of integrals considered in the  $L^+$  sense.

**Proof:** We will use the method of the friendly functions ([Method of the friendly functions \(Lemma 6.9\)](#)). Let

$$\mathcal{G} = \left\{ g \in \text{Bor}^+(X, \mathbb{R}) \mid \int_X g \, d\nu = \int_X g \cdot h \, d\mu \right\}$$

We verify that  $\mathcal{G}$  satisfies the conditions of [Method of the friendly functions \(Lemma 6.9\)](#).

- For  $A \in \mathfrak{M}$ , we have

$$\int_X \chi_A \, d\nu = \nu(A) = \int_A h \, d\mu = \int_X \chi_A \cdot h \, d\mu,$$

so  $\chi_A \in \mathcal{G}$ .

- If  $g_1, g_2 \in \mathcal{G}$ , then

$$\begin{aligned} \int_X (g_1 + g_2) \, d\nu &= \int_X g_1 \, d\nu + \int_X g_2 \, d\nu \\ &= \int_X g_1 \cdot h \, d\mu + \int_X g_2 \cdot h \, d\mu \\ &= \int_X (g_1 + g_2) \cdot h \, d\mu, \end{aligned}$$

so  $g_1 + g_2 \in \mathcal{G}$ .

- If  $g \in \mathcal{G}$  and  $\alpha \in [0, \infty)$ , then

$$\begin{aligned} \int_X \alpha g \, d\nu &= \alpha \int_X g \, d\nu \\ &= \alpha \int_X g \cdot h \, d\mu \\ &= \int_X (\alpha g) \cdot h \, d\mu, \end{aligned}$$

so  $\alpha g \in \mathcal{G}$ .

- Suppose that  $g_1 \leq g_2 \leq \dots \leq g_n \leq \dots$  are in  $\mathcal{G}$  and  $g_n \rightarrow g$  for some  $g \in \text{Bor}^+(X, \mathbb{R})$ . Since  $h \geq 0$ , we also have  $g_n \cdot h \rightarrow g \cdot h$  increasingly. By [Monotone Convergence Theorem \(Theorem 6.2\)](#), applied once with respect to  $\nu$  and once with respect to  $\mu$ ,

$$\begin{aligned}
\int_X g \, d\nu &= \lim_{n \rightarrow \infty} \int_X g_n \, d\nu \\
&= \lim_{n \rightarrow \infty} \int_X g_n \cdot h \, d\mu \\
&= \int_X g \cdot h \, d\mu.
\end{aligned}$$

Thus  $g \in \mathcal{G}$ .

Then [Method of the friendly functions \(Lemma 6.9\)](#) gives  $\mathcal{G} = \text{Bor}^+(X, \mathbb{R})$ , which means that

$$\int_X g \, d\nu = \int_X g \cdot h \, d\mu$$

for all  $g \in \text{Bor}^+(X, \mathbb{R})$ , as required. □

### Note

In connection with [Proposition 7.10](#), it is customary to write the connection between the positive measures  $\mu$  and  $\nu$  in the form

$$d\nu = h \, d\mu.$$

This can be thought of as cancelling the common integral expression in

$$\int_X f \, d\nu = \int_X f \cdot h \, d\mu.$$

Some further abuse of notation rewrites this relation as

$$h = \frac{d\nu}{d\mu},$$

where one thinks of  $h$  as a kind of “derivative” of  $\nu$  with respect to  $\mu$ .

## 8 Lebesgue’s Dominated Convergence Theorem (LDCT)

Fix a measure space  $(X, \mathfrak{M}, \mu)$ .

**Theorem 8.1****LDCT**

Let  $f$  and  $(f_n)_{n=1}^{\infty}$  be functions in  $\text{Bor}(X, \mathbb{R})$  such that

$$\lim_{n \rightarrow \infty} f_n(x) = f(x), \quad \forall x \in X.$$

Suppose moreover that there exists a function  $h \in \mathcal{L}^1(\mu) \cap \text{Bor}^+(X, \mathbb{R})$  which dominates all the  $f_n$ , in the sense that

$$|f_n| \leq h, \quad \forall n \in \mathbb{N}.$$

Then  $f$  and  $f_1, f_2, \dots, f_n, \dots$  are all in  $\mathcal{L}^1(\mu)$ , and

$$\lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \int_X f \, d\mu.$$

**Note**

The last equation can be written as

$$\lim_{n \rightarrow \infty} \left( \int_X f_n \, d\mu \right) = \int_X \left( \lim_{n \rightarrow \infty} f_n \right) d\mu,$$

so this is a formula where one interchanges integration with a limit. In principle, this interchange can have issues. The point of LDCT is that the existence of the dominating function  $h$  is a useful sufficient condition which makes the issues go away.

It is convenient to think of LDCT as going in parallel with the MCT from [Monotone Convergence Theorem \(Theorem 6.2\)](#). In connection to that, we will observe in [Reverse-MCT \(Proposition 8.2\)](#) that a special case of LDCT gives a “Reverse-MCT” statement.

**Proposition 8.2****Reverse-MCT**

Let  $u$  and  $(u_n)_{n=1}^{\infty}$  be functions in  $\text{Bor}^+(X, \mathbb{R})$  such that

$$u_1 \geq u_2 \geq \dots \geq u_n \geq \dots$$

and such that

$$\lim_{n \rightarrow \infty} u_n(x) = u(x), \quad \forall x \in X.$$

(compactly written  $u_n \searrow u$ ). We assume, in addition, that  $\int_X u_1 \, d\mu < \infty$ . Then

$$\lim_{n \rightarrow \infty} \int_X u_n \, d\mu = \int_X u \, d\mu.$$

**Note**

Reverse-MCT is a special case of LDCT because we can use  $h = u_1$  as the dominating function. Then [LDCT \(Theorem 8.1\)](#) gives the required conclusion.

For the plan of how we prove LDCT, it is convenient to also consider its special case  $\text{LDCT}_0$ , which considers a dominated sequence in  $\text{Bor}^+(X, \mathbb{R})$  converging pointwise to 0. This is [LDCT<sub>0</sub> \(Proposition 8.3\)](#).

**Proposition 8.3** $\text{LDCT}_0$ 

Let  $(g_n)_{n=1}^\infty$  be a sequence of functions in  $\text{Bor}^+(X, \mathbb{R})$  such that

$$\lim_{n \rightarrow \infty} g_n(x) = 0, \quad \forall x \in X.$$

Suppose moreover that there exists a function  $h \in \mathcal{L}^1(\mu) \cap \text{Bor}^+(X, \mathbb{R})$  such that

$$g_n \leq h, \quad \forall n \in \mathbb{N}.$$

Then

$$\lim_{n \rightarrow \infty} \int_X g_n \, d\mu = 0.$$

**Remark 8.4**

Scheme of proof of LDCT

We will go in the following way:

$$\text{MCT} \implies \text{Reverse-MCT} \implies \text{LDCT}_0 \implies \text{LDCT}.$$

The first implication is proved by subtracting everything from the top. The second implication uses the trick of Fatou. The third implication is an immediate reduction, by putting

$$g_n := |f_n - f|.$$

**8.5. Proof of the implication (1).** Define

$$v_n := u_1 - u_n, \quad n \in \mathbb{N}, \quad \text{and} \quad v := u_1 - u.$$

Then  $v_n \in \text{Bor}^+(X, \mathbb{R})$  for every  $n$ , and  $v_n \nearrow v$ . By [Monotone Convergence Theorem \(Theorem 6.2\)](#),

$$\lim_{n \rightarrow \infty} \int_X v_n \, d\mu = \int_X v \, d\mu.$$

Since  $u_1 = v_n + u_n$  and  $u_1 = v + u$ , and since  $\int_X u_1 \, d\mu < \infty$ , all the quantities below are finite and we can write



$$\begin{aligned}
\int_X u_n \, d\mu &= \int_X u_1 \, d\mu - \int_X v_n \, d\mu \\
&\rightarrow \int_X u_1 \, d\mu - \int_X v \, d\mu \\
&= \int_X u \, d\mu.
\end{aligned}$$

□

**8.6. Proof of the implication (2).** The idea is to make a helper sequence  $(u_n)_{n=1}^\infty$  to which we can apply [Reverse-MCT \(Proposition 8.2\)](#). We break the proof into steps:

1. For every  $n \in \mathbb{N}$ , define  $u_n : X \rightarrow [0, \infty)$  by

$$u_n(x) := \sup\{g_k(x) \mid k \geq n\}.$$

Since each  $g_k \in \text{Bor}^+(X, \mathbb{R})$ , [Closure under suprema and infima \(Proposition 4.1\)](#) gives  $u_n \in \text{Bor}(X, \mathbb{R})$ . Also  $0 \leq u_n \leq h$ , so  $u_n \in \text{Bor}^+(X, \mathbb{R})$ . In particular,

$$\int_X u_1 \, d\mu \leq \int_X h \, d\mu < \infty,$$

where the first inequality uses the monotonicity of  $L^+$  from [Characterization of  \$L^+\$  \(Theorem 6.8\)](#), and the finiteness of the last term follows from [Proposition 7.6](#). Finally, from the definition of the tail suprema we have

$$u_1 \geq u_2 \geq \cdots \geq u_n \geq \cdots.$$

2. We claim that  $u_n \searrow 0$ . Fix  $x \in X$  and let  $\varepsilon > 0$  be given. Since  $g_n(x) \rightarrow 0$ , there is  $n_0 \in \mathbb{N}$  such that  $g_k(x) < \varepsilon$  for all  $k \geq n_0$ . For this  $n_0$ , we have

$$u_{n_0}(x) = \sup\{g_k(x) \mid k \geq n_0\} \leq \varepsilon.$$

Since the sequence  $(u_n(x))_{n=1}^\infty$  is decreasing, it follows that

$$0 \leq u_n(x) \leq u_{n_0}(x) \leq \varepsilon, \quad \forall n \geq n_0.$$

Thus  $u_n(x) \rightarrow 0$  for every  $x \in X$ .

3. Now [Reverse-MCT \(Proposition 8.2\)](#) applied to  $(u_n)_{n=1}^\infty$  gives

$$\lim_{n \rightarrow \infty} \int_X u_n \, d\mu = 0.$$

Since  $g_n \leq u_n$  for every  $n \in \mathbb{N}$ , the monotonicity of  $L^+$  gives

$$0 \leq \int_X g_n \, d\mu \leq \int_X u_n \, d\mu, \quad \forall n \in \mathbb{N}.$$

Hence  $\int_X g_n \, d\mu \rightarrow 0$  as required.

□

**8.7. Proof of the implication (3).** Have  $f$  and  $(f_n)_{n=1}^\infty$  in  $\text{Bor}(X, \mathbb{R})$  such that  $f_n(x) \xrightarrow{n \rightarrow \infty} f(x)$  for all  $x \in X$ . Also, we have  $h \in \text{Bor}^+(X, \mathbb{R}) \cap \mathcal{L}^1(\mu)$  such that  $|f_n| \leq h$  for all  $n \in \mathbb{N}$ . We want  $f_n \in \mathcal{L}^1(\mu)$  for all  $n \in \mathbb{N}$ ,  $f \in \mathcal{L}^1(\mu)$  and

$$\int_X f_n \, d\mu \xrightarrow{n \rightarrow \infty} \int_X f \, d\mu$$

• For every  $n \in \mathbb{N}$ , we have

$$\begin{aligned} |f_n| \leq h &\implies \int_X |f_n| \, d\mu \leq \int_X h \, d\mu < \infty \\ &\implies f_n \in \mathcal{L}^1(\mu) \end{aligned}$$

How about  $f$ ? Observe that  $|f| \leq h$ , which holds because

$$|f(x)| = \lim_{n \rightarrow \infty} |f_n(x)| \leq h(x) \quad \forall x \in X$$

Thus

$$\int_X |f| \, d\mu \leq \int_X h \, d\mu < \infty$$

hence  $f \in \mathcal{L}^1(\mu)$  as well. It remains to check that

$$\int_X f_n \, d\mu \xrightarrow{n \rightarrow \infty} \int_X f \, d\mu$$

Towards that end, consider

$$g_n := |f_n - f| \in \text{Bor}^+(X, \mathbb{R}) \quad \forall n \in \mathbb{N}$$

**Claim:** The  $g_n$ 's satisfy the hypothesis of [LDCT<sub>0</sub> \(Proposition 8.3\)](#). That is, we have

$$(i) \quad g_n(x) \xrightarrow{n \rightarrow \infty} 0 \quad \forall x \in X$$

(ii) There is  $\tilde{h} \in \text{Bor}^+(X, \mathbb{R}) \cap \mathcal{L}^1(\mu)$  such that  $g_n \leq \tilde{h}$  for all  $n \in \mathbb{N}$

**Verification of Claim:**

(i) For every  $x \in X$  we have

$$g_n(x) = |f_n(x) - f(x)| \xrightarrow{n \rightarrow \infty} 0$$

due to the hypothesis that  $f_n(x) \xrightarrow{n \rightarrow \infty} f(x)$ .

(ii) For every  $n \in \mathbb{N}$  and  $x \in X$  we have

$$g_n(x) = |f_n(x) - f(x)| \leq |f_n(x)| + |f(x)| \leq 2h(x)$$

Hence  $g_n \leq 2h$  for all  $n \in \mathbb{N}$ , where

$$\int_X 2h \, d\mu = 2 \int_X h \, d\mu < \infty$$

so we can use  $\tilde{h} = 2h$

Due to the claim, we can apply **LDCT<sub>0</sub>** (**Proposition 8.3**) to  $(g_n)_{n=1}^\infty$  to get that

$$\int_X g_n \, d\mu \xrightarrow{n \rightarrow \infty} 0 \quad (*)$$

This implies the fact that

$$\int_X f_n \, d\mu \xrightarrow{n \rightarrow \infty} \int_X f \, d\mu$$

because we can write

$$0 \leq \left| \int_X f_n \, d\mu - \int_X f \, d\mu \right| = \left| \int_X f_n - f \, d\mu \right| \leq \int_X |f_n - f| \, d\mu = \int_X g_n \, d\mu \xrightarrow{n \rightarrow \infty} 0$$

which completes the proof. □

## 8.1 The “a.e.- $\mu$ ” annoyance (and the “a.e.- $\mu$ ” version of LDCT)

### Note

A rule of thumb: changes made on a set of measure 0 will not affect integrability or the value of the integral.

### Definition 8.8

We say that two functions  $f, g : X \rightarrow \mathbb{R}$  are *equal almost everywhere with respect to  $\mu$* , denoted  $f = g$  a.e.- $\mu$ , to mean there exists a set  $Z \in \mathfrak{M}$  such that

- (i)  $\mu(Z) = 0$ .
- (ii)  $f(x) = g(x)$  for all  $x \in X \setminus Z$ .

The preceding definition makes sense even if  $f$  and  $g$  are not Borel functions. In the case of interest when  $f, g \in \text{Bor}(X, \mathbb{R})$ , we can use the following equivalent description of equality almost everywhere.

### Proposition 8.9

Let  $f, g \in \text{Bor}(X, \mathbb{R})$ , and denote

$$M := \{x \in X \mid f(x) \neq g(x)\}.$$

1. We have  $M \in \mathfrak{M}$ , hence the quantity  $\mu(M)$  is well-defined.
2. We have

$$(f = g \text{ a.e.-}\mu) \iff (\mu(M) = 0).$$

**Proof of 1:** We can write

$$M = (f - g)^{-1}(\mathbb{R} \setminus \{0\}).$$

Since  $f - g \in \text{Bor}(X, \mathbb{R})$  and  $\mathbb{R} \setminus \{0\} \in \mathcal{B}_{\mathbb{R}}$ , it follows that  $M \in \mathfrak{M}$ .  $\square$

**Proof of 2:** “ $\implies$ ” Suppose  $f = g$  a.e.- $\mu$ . Then there exists  $Z \in \mathfrak{M}$  such that  $\mu(Z) = 0$  and  $f(x) = g(x)$  for every  $x \in X \setminus Z$ . It follows that  $M \subseteq Z$ , so  $0 \leq \mu(M) \leq \mu(Z) = 0$ , hence  $\mu(M) = 0$ .

“ $\impliedby$ ” Suppose  $\mu(M) = 0$ . Taking  $Z := M$ , we have  $Z \in \mathfrak{M}$  by part 1,  $\mu(Z) = 0$ , and  $f(x) = g(x)$  for every  $x \in X \setminus Z$ . Hence  $f = g$  a.e.- $\mu$ .  $\square$

### Lemma 8.10

Let  $f \in \text{Bor}^+(X, \mathbb{R})$  be such that  $f = \underline{0}$  a.e.- $\mu$ . Then

$$\int_X f \, d\mu = 0.$$

In particular,  $f \in \mathcal{L}^1(\mu)$ .

**Proof:** For every  $n \in \mathbb{N}$ , let  $f_n = f \wedge \underline{n}$ , where  $\underline{n} : X \rightarrow \mathbb{R}$  is identically equal to  $n$ . That is,

$$f_n(x) = \begin{cases} f(x) & \text{if } f(x) \leq n \\ n & \text{if } f(x) > n \end{cases}$$

We have  $f_n \in \text{Bor}(X, \mathbb{R})$  because  $\text{Bor}(X, \mathbb{R})$  is a lattice of functions. It is also easy to see that  $f_n \nearrow f$ . By [Monotone Convergence Theorem \(Theorem 6.2\)](#) we have

$$\int_X f \, d\mu = \lim_{n \rightarrow \infty} \int_X f_n \, d\mu.$$

So in order to get  $\int_X f \, d\mu = 0$ , it suffices to check that  $\int_X f_n \, d\mu = 0$  for every  $n \in \mathbb{N}$ . Let  $Z = \{x \in X \mid f(x) \neq 0\}$ . By [Proposition 8.9](#), we have  $\mu(Z) = 0$ . Now observe that  $f_n \leq n\chi_Z$  for all  $n \in \mathbb{N}$ : for  $x \in Z$  use that  $f_n(x) \leq n = n\chi_Z(x)$ , while for  $x \in X \setminus Z$  we have  $f_n(x) = 0 = n\chi_Z(x)$ . Thus

$$\int_X f_n \, d\mu \leq \int_X n\chi_Z \, d\mu = n\mu(Z) = 0,$$

hence  $\int_X f_n \, d\mu = 0$  as desired.  $\square$

### Proposition 8.11

Let  $f, g \in \text{Bor}(X, \mathbb{R})$  be such that  $f = g$  a.e.- $\mu$ . Suppose that one of these functions, say  $f$ , is in  $\mathcal{L}^1(\mu)$ . Then  $g \in \mathcal{L}^1(\mu)$  as well, and

$$\int_X f \, d\mu = \int_X g \, d\mu.$$