

Complex Analysis

Contents

1	Complex Number System	3
1.1	Algebraic Structure	3
1.2	Conjugation and Polar Form	3
1.3	Topology and Argument	4
2	Differentiation	5
2.1	Complex Derivatives	5
2.2	Cauchy-Riemann Equations	5
2.3	Holomorphic Functions and Exponential	6
2.4	Logarithms and Branches	6
2.5	Further Holomorphicity Criteria	7
3	Series	7
3.1	Numerical Series	7
3.2	Function Series	8
3.3	Power Series	8
3.4	Consequences of Power Series	10
4	Complex Integration	10
4.1	Complex Integrals on Intervals	10
4.2	Path Integrals	10
4.3	Estimates and Uniform Convergence	14
4.4	Cauchy's Theorem	15
4.5	Cauchy's Theorem for Convex Regions	17
4.6	Cauchy's Integral Formula	18
4.7	Consequences of Cauchy's Formula	19
4.8	Cauchy Integrals and Analyticity	20
4.9	Derivative Formula	22
4.10	Operations on Power Series	23
4.11	Converse Results	23
4.12	Liouville's Theorem and Algebraic Consequences	24
5	Zeros and the Identity Theorem	25
5.1	Orders of Zeros	25
5.2	Identity Theorem	26
6	Convergence and Maximum Principles	26
6.1	Identity Theorem Applications	26
6.2	Locally Uniform Limits	27
6.3	Maximum Modulus Principle	27
6.4	Schwarz's Lemma	28
6.5	Minimum Modulus Principle	29
7	Laurent Series	29
7.1	Examples	29
7.2	Definition and Convergence	30

7.3	Annuli	33
7.4	Cauchy's Formula on an Annulus	34
8	Isolated Singularities	35
8.1	Laurent Series Representation	35
8.2	Classification	36
8.3	Singularity at Infinity	39
8.4	Removable Singularities	39

1 Complex Number System

1.1 Algebraic Structure

Definition 1.1

Complex Numbers

The set of complex numbers $\mathbb{C}$ is $\mathbb{R}^2$, together with the operations

$$(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

and

$$(x_1, y_1) \cdot (x_2, y_2) = (x_1x_2 - y_1y_2, x_1y_2 + x_2y_1)$$

Theorem 1.1

Field Structure

$\mathbb{C}$ is a field. With $0 = (0, 0)$, $-(x, y) = (-x, -y)$, $1 = (1, 0)$ and

$$(x, y)^{-1} = \left(\frac{x}{x^2 + y^2}, -\frac{y}{x^2 + y^2} \right) \text{ for } (x, y) \neq (0, 0)$$

Notation 1.1

If $z = (x, y)$ we call x the “real part” of z and write $x = \operatorname{Re}(z)$.

We call y the “imaginary part” of z and write $y = \operatorname{Im}(z)$.

Define $i = (0, 1)$.

Remark 1.2

There is an embedding

$$\begin{aligned} \mathbb{R} &\hookrightarrow \mathbb{C} \\ x &\mapsto (x, 0) \end{aligned}$$

so we view $\mathbb{R} \subseteq \mathbb{C}$.

Notation 1.3

Under the above embedding, any $z \in \mathbb{C}$ can be written as $z = \operatorname{Re}(z) + \operatorname{Im}(z) \cdot i$ and $i^2 = -1$

1.2 Conjugation and Polar Form

Definition 1.2

Conjugation

Define *conjugation* to be the map

$$\begin{aligned} - : \mathbb{C} &\longrightarrow \mathbb{C} \\ z = x + yi &\longmapsto x - yi = \bar{z} \end{aligned}$$

Note

$$\operatorname{Re}(z) = \frac{z + \bar{z}}{2}, \quad \operatorname{Im}(z) = \frac{z - \bar{z}}{2i}$$

Note

If $z = x + yi \in \mathbb{C}$ it has a polar representation $(x, y) \mapsto (r, \theta)$ via $z = r \cos \theta + ir \sin \theta$ where $r = \sqrt{x^2 + y^2} \geq 0$, $x = r \cos \theta$, $y = r \sin \theta$. The angle θ is defined modulo 2π when $z \neq 0$, and is arbitrary when $z = 0$.

Remark 1.4

If z_1, z_2 have polar representations

$$z_1 : (r_1, \theta_1), \quad z_2 : (r_2, \theta_2)$$

Then

$$\begin{aligned} z_1 \cdot z_2 &= (r_1 \cos \theta_1 + ir_1 \sin \theta_1) \cdot (r_2 \cos \theta_2 + ir_2 \sin \theta_2) \\ &= r_1 r_2 ((\cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2) + i(\cos \theta_1 \sin \theta_2 + \sin \theta_1 \cos \theta_2)) \\ &= r_1 r_2 (\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)) \end{aligned}$$

1.3 Topology and Argument**Note**

If $z = x + iy$ and $r = \sqrt{x^2 + y^2}$, then $r^2 = z\bar{z}$.

Remark 1.5

$\mathbb{C}$ is a metric space and thus the triangle inequality holds.

Remark 1.6

$\mathbb{C}$ is a topological space, so topological notions make sense.

Definition 1.3**Region**

A *region* is a non-empty subset of $\mathbb{C}$ that is open and connected.

Definition 1.4**Argument**

If $z \in \mathbb{C} \setminus \{0\}$ has polar representation (r, θ) , we call θ the *argument* of z and write $\theta = \arg(z)$. We'll think about $\arg$ as a multivalued function. We write $\operatorname{Arg}(z) = \theta$ if $\theta \in (-\pi, \pi]$, the *principal value*.

Example 1.1

If $z = r(\cos \theta + i \sin \theta)$ and $z \neq 0$ then we can find an n^{th} root of z (actually n of them).

2 Differentiation

2.1 Complex Derivatives

Definition 2.1**Complex Derivative**

Suppose $f : U \rightarrow \mathbb{C}$ where $U \subseteq \mathbb{C}$ is open and contains z_0 . We'll say that f has derivative $f'(z_0)$ at z_0 if

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

Theorem 2.1**Product Rule**

If f and g are differentiable at z_0 then so is fg and

$$(f \cdot g)'(z_0) = f'(z_0)g(z_0) + f(z_0)g'(z_0)$$

2.2 Cauchy-Riemann Equations

Remark 2.1

If $f : \mathbb{C} \rightarrow \mathbb{C}$, then we may write $f(z) = u(z) + iv(z)$ where $u, v : \mathbb{C} \rightarrow \mathbb{R}$. If f is differentiable at $z_0 = x_0 + iy_0$ (say $f'(z_0) = a + ib$), then

$$\begin{aligned} R(z) &= f(z) - f(z_0) - f'(z_0)(z - z_0) \\ &= u(z) - u(z_0) - a(x - x_0) + b(y - y_0) + i(v(z) - v(z_0) - b(x - x_0) - a(y - y_0)) \\ &= R_1(z) + iR_2(z) \end{aligned}$$

Then $\lim_{z \rightarrow z_0} R_1 \frac{z}{|z - z_0|} = \lim_{z \rightarrow z_0} R_2 \frac{z}{|z - z_0|} = 0$. Thus both u and v are differentiable and $\frac{\partial u}{\partial x}(z_0) = a$, $\frac{\partial u}{\partial y}(z_0) = -b$, $\frac{\partial v}{\partial x}(z_0) = b$, $\frac{\partial v}{\partial y}(z_0) = a$.

Theorem 2.2**Cauchy-Riemann Equations**

If $f : \mathbb{C} \rightarrow \mathbb{C}$, $f(z) = u(z) + iv(z)$, then f is differentiable at z_0 (as a complex function) iff both u and v are differentiable at z_0 and

$$\begin{aligned} \frac{\partial u}{\partial x}(z_0) &= \frac{\partial v}{\partial y}(z_0) \quad \text{and} \\ \frac{\partial v}{\partial x}(z_0) &= -\frac{\partial u}{\partial y}(z_0) \end{aligned}$$

These are called the Cauchy-Riemann equations.

Note

If $u : \mathbb{R}^2 \rightarrow \mathbb{R}$, then u is differentiable at (x_0, y_0) if $\frac{\partial u}{\partial x}$ and $\frac{\partial u}{\partial y}$ exist in a neighborhood of (x_0, y_0) and are continuous at (x_0, y_0) .

2.3 Holomorphic Functions and Exponential**Definition 2.2****Exponential function**

The exponential function is defined to be the function

$$\exp(x + iy) = e^x(\cos y + i \sin y)$$

we write $e^z := \exp(z)$.

Remark 2.2

For $y \in \mathbb{R}$, we have $e^{iy} = \cos(y) + i \sin(y)$.

Definition 2.3**Holomorphic**

If $f : U \rightarrow \mathbb{C}$ with $U \subseteq \mathbb{C}$ open, such that f is differentiable at each point in U , then f is *holomorphic*.

Definition 2.4**Entire**

$f : \mathbb{C} \rightarrow \mathbb{C}$ is called *entire* if it is holomorphic on all of $\mathbb{C}$.

2.4 Logarithms and Branches**Definition 2.5****Logarithm**

A complex number w is a *logarithm* of z if $e^w = z$.

Remark 2.3

The range of the exponential is $\mathbb{C} \setminus \{0\}$. So there is no logarithm of 0.

Definition 2.6**Branch of Logarithm**

If $G \subseteq \mathbb{C} \setminus \{0\}$ is a non-empty open set, then a *branch of log* is a continuous function $\ell : G \rightarrow \mathbb{C}$ such that for all $z \in G$, $\ell(z)$ is a logarithm of z , i.e. $e^{\ell(z)} = z$

Note

A branch of log may or may not exist depending on the shape of G .

Example 2.1

$G := \mathbb{C} \setminus (-\infty, 0]$. $\ell(z) = \ln|z| + i \operatorname{Arg}(z)$. Then

$$\begin{aligned} e^{\ell(z)} &= e^{\ln|z|}(\cos(\operatorname{Arg}(z)) + i \sin(\operatorname{Arg}(z))) \\ &= |z|(\cos(\operatorname{Arg}(z)) + i \sin(\operatorname{Arg}(z))) \\ &= z \end{aligned}$$

This is called the *principal branch of log* and is denoted by $\operatorname{Log}(z)$

Note

If we define $\ell(z) = \operatorname{Log}(z) + i2\pi k$ on $\mathbb{C} \setminus (-\infty, 0]$ for fixed $k \in \mathbb{Z}$, then we get a branch of log.

Remark 2.4

It is not possible to find a branch of log on $\mathbb{C} \setminus \{0\}$.

2.5 Further Holomorphicity Criteria**Exercise 2.5**

In polar form, the Cauchy-Riemann equations become, if $f(z) = u(z) + iv(z)$

$$\begin{aligned} ru_r &= v_\theta \\ \text{and } u_\theta &= -rv_r \end{aligned}$$

Theorem 2.3**Derivative of Logarithm**

If $\ell : G \rightarrow \mathbb{C}$ is a branch of log, then ℓ is holomorphic and $\ell'(z) = \frac{1}{z}$.

Theorem 2.4**Polynomial Holomorphicity Criterion**

Suppose f is a polynomial in variables x and y (with complex coefficients), then f is holomorphic iff f can be written as polynomial in $z = x + iy$.

3 Series**3.1 Numerical Series****Definition 3.1****Series**

A *series* is a formal sum $\sum_{n=0}^{\infty} c_n$ where the terms $c_n \in \mathbb{C}$. We say a series *converges* if the sequence of partial sums converge otherwise we say it *diverges*.

Lemma 3.1**Term Test**

If $\sum_{n=0}^{\infty} c_n$ converges, then $c_n \rightarrow 0$.

Definition 3.2**Absolute Convergence**

A series $\sum_{n=0}^{\infty} c_n$ converges absolutely if $\sum_{n=0}^{\infty} |c_n| < \infty$

Lemma 3.2**Absolute Convergence Test**

If $\sum_{n=0}^{\infty} c_n$ converges absolutely, then $\sum_{n=0}^{\infty} c_n$ converges and $|\sum_{n=0}^{\infty} c_n| \leq \sum_{n=0}^{\infty} |c_n|$

Example 3.1**Geometric Series**

$\sum_{n=0}^{\infty} c^n$ converges iff $|c| < 1$. If $|c| < 1$ then $\sum_{n=0}^{\infty} c^n = \frac{1}{1-c}$

3.2 Function Series**Definition 3.3****Convergence of Functions**

Suppose $(f_n)_{n=1}^{\infty}$ is a sequence of $\mathbb{C}$ -valued functions on $G \subseteq \mathbb{C}$.

- $(f_n)_{n=1}^{\infty}$ converges pointwise to f on $S \subseteq G$ if $\lim_{n \rightarrow \infty} f_n(z) = f(z)$ for every $z \in S$.
- $(f_n)_{n=1}^{\infty}$ converges uniformly to f on S if $\sup_{z \in S} |f_n(z) - f(z)| \rightarrow 0$
- $(f_n)_{n=1}^{\infty}$ converges locally uniformly to f on G if for each $z \in G$, there is some open ball $B \subseteq G$ with centre z such that $(f_n)_{n=1}^{\infty}$ converges uniformly to f on B .

Definition 3.4**Series of Functions**

If $(f_n)_{n=0}^{\infty}$ is a sequence of functions, $\sum_{n=0}^{\infty} f_n$ is the associated series of functions. The series converges to f if the partial sums do.

3.3 Power Series**Notation 3.1**

For $z_0 \in \mathbb{C}$ and $r > 0$, write

$$D(z_0, r) = \{z \in \mathbb{C} \mid |z - z_0| < r\}$$

for the open disk centered at z_0 with radius r . We write

$$\mathbb{D} = D(0, 1)$$

for the open unit disk.

Example 3.2

$z_0 \in \mathbb{C}$ is fixed. $(f_n)_{n=0}^{\infty}$, where $f_n(z) = a_n(z - z_0)^n$ we get the series $\sum_{n=0}^{\infty} a_n(z - z_0)^n$. The power series centred at z_0 .

Proposition 3.3**Region of Convergence**

The region of convergence for a power series $\sum_{n=0}^{\infty} a_n(z - z_0)^n$ is of the form either

1. $\{z_0\}$
2. $\mathbb{C}$
3. The points in some disk centred at z_0 and possibly points on the boundary.

Example 3.3

$\sum_{n=0}^{\infty} z^n$ has radius of convergence 1. This represents the function $\frac{1}{1-z}$ on $\mathbb{D}$. This converges locally uniformly.

Theorem 3.4**Comparison Test for Power Series**

Suppose $\sum_{n=0}^{\infty} a_n(z - z_0)^n$ and $\sum_{n=0}^{\infty} b_n(z - z_0)^n$ are power series with radius of convergence R_1, R_2 respectively. If $|b_n| \leq M|a_n|$ for some $M > 0$, then $R_1 \leq R_2$

Proof: This follows from the [Region of Convergence](#) because if $|z - z_0| < R$, where R is the radius of convergence, then $\sum a_n(z - z_0)^n$ converges absolutely. $\square$

Theorem 3.5**Cauchy-Hadamard Theorem**

The radius of convergence for a power series $\sum a_n(z - z_0)^n$ is

$$\frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}}$$

Theorem 3.6**Differentiation of Power Series**

Suppose $\sum_{n=0}^{\infty} a_n(z - z_0)^n$ has radius of convergence $R > 0$. Let f be the function it represents on $D(z_0, R)$. Then the radius of convergence for $\sum_{n=1}^{\infty} n a_n(z - z_0)^{n-1}$ is also R and if g is the function it represents, then f is differentiable and $f'(z) = g(z)$ for $z \in D(z_0, R)$.

Proof: Note

$$\begin{aligned} & \frac{1}{\limsup_{n \rightarrow \infty} |n a_n|^{\frac{1}{n}}} \\ &= \frac{1}{\lim_{n \rightarrow \infty} n^{\frac{1}{n}}} \cdot \frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}} \\ &= \frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}} \\ &= R \end{aligned}$$

$\square$

Example 3.4

$f(z) = \sum_{n=0}^{\infty} \frac{1}{n!} z^n$ then f is entire, $f'(z) = f(z)$ and $f(0) = 1$.

3.4 Consequences of Power Series**Corollary 3.7****Smoothness of Power Series**

If f is represented by a power series on its disk of convergence, then derivatives of all orders exist on that disk.

4 Complex Integration**4.1 Complex Integrals on Intervals****Definition 4.1****Complex Integral**

Suppose $\varphi : [a, b] \rightarrow \mathbb{C}$ is piecewise continuous (continuous except at finitely many points where one-sided limits exist). Then $\operatorname{Re} \varphi$ and $\operatorname{Im} \varphi$ are also piecewise continuous. Set

$$\int_a^b \varphi := \int_a^b \operatorname{Re} \varphi + i \int_a^b \operatorname{Im} \varphi$$

Note this is complex linear in φ

Theorem 4.1**Fundamental Theorem of Calculus**

If φ is continuous and piecewise $\mathcal{C}^1$ on $[a, b]$, then

$$\int_a^b \varphi'(t) dt = \varphi(b) - \varphi(a)$$

Theorem 4.2**Integral Triangle Inequality**

$$\left| \int_a^b \varphi(t) dt \right| \leq \int_a^b |\varphi(t)| dt$$

4.2 Path Integrals**Definition 4.2****Path**

A path is a function $[a, b] \rightarrow \mathbb{C}$.

Definition 4.3**Piecewise $\mathcal{C}^1$ Path**

A path $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise $\mathcal{C}^1$ if γ is continuous and there is a partition of $[a, b]$ such that γ is $\mathcal{C}^1$ on each subinterval, with one-sided limits for γ' at the breakpoints.

Definition 4.4**Path Integral**

Suppose $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$ and $f : G \rightarrow \mathbb{C}$ is continuous on a set G containing the image of γ . Then define

$$\int_{\gamma} f(z) dz := \int_a^b f(\gamma(t))\gamma'(t) dt$$

Example 4.1

$f(z) = ie^x$ where $z = x + iy$, $\gamma : [a, b] \rightarrow \mathbb{C}$, $\gamma(t) = t$ then

$$\int_{\gamma} f(z) dz = \int_a^b ie^t dt = i(e^b - e^a)$$

Theorem 4.3**Fundamental Theorem for Path Integrals**

If $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$ and if f is holomorphic on an open set that contains the range of γ then

$$\int_{\gamma} f'(z) dz = f(\gamma(b)) - f(\gamma(a))$$

Proof:

$$\int_{\gamma} f'(z) dz = \int_a^b f'(\gamma(t))\gamma'(t) dt = \int_a^b (f \circ \gamma)'(t) dt = (f \circ \gamma)(b) - (f \circ \gamma)(a)$$

□

Proposition 4.4**Reparametrization Invariance**

If $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$ and $\beta : [a_1, b_1] \rightarrow [a, b]$ is increasing and piecewise- $\mathcal{C}^1$ with $\beta(a_1) = a$ and $\beta(b_1) = b$. Set $\gamma_1 = \gamma \circ \beta$. Suppose f is continuous on the image of γ . Then

$$\int_{\gamma_1} f(z) dz = \int_{\gamma} f(z) dz.$$

Proof: By definition of the path integral and the chain rule,

$$\begin{aligned} \int_{\gamma_1} f(z) dz &= \int_{a_1}^{b_1} f(\gamma_1(s))\gamma_1'(s) ds \\ &= \int_{a_1}^{b_1} f(\gamma(\beta(s)))\gamma'(\beta(s))\beta'(s) ds \end{aligned}$$

By the substitution theorem with $t = \beta(s)$,

$$\begin{aligned}
\int_{a_1}^{b_1} f(\gamma(\beta(s)))\gamma'(\beta(s))\beta'(s) \, ds &= \int_a^b f(\gamma(t))\gamma'(t) \, dt \\
&= \int_{\gamma} f(z) \, dz
\end{aligned}$$

□

Proposition 4.5**Properties of Path Integrals**

Suppose f is continuous on the image of the paths below.

1. If $a < c < b$ and $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$, let $\gamma_1 = \gamma|_{[a, c]}$ and $\gamma_2 = \gamma|_{[c, b]}$. Then

$$\int_{\gamma} f(z) \, dz = \int_{\gamma_1} f(z) \, dz + \int_{\gamma_2} f(z) \, dz.$$

2. If $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$, define the reverse path $-\gamma : [a, b] \rightarrow \mathbb{C}$ by $(-\gamma)(t) = \gamma(a + b - t)$. Then

$$\int_{-\gamma} f(z) \, dz = - \int_{\gamma} f(z) \, dz.$$

Proof: For the first property, by the definition of path integral,

$$\begin{aligned}
\int_{\gamma} f(z) \, dz &= \int_a^b f(\gamma(t))\gamma'(t) \, dt \\
&= \int_a^c f(\gamma(t))\gamma'(t) \, dt + \int_c^b f(\gamma(t))\gamma'(t) \, dt \\
&= \int_{\gamma_1} f(z) \, dz + \int_{\gamma_2} f(z) \, dz.
\end{aligned}$$

For the second property, by the chain rule,

$$(-\gamma)'(t) = -\gamma'(a + b - t).$$

Hence

$$\int_{-\gamma} f(z) \, dz = \int_a^b f(\gamma(a + b - t)) \cdot (-\gamma'(a + b - t)) \, dt.$$

Using the substitution $s = a + b - t$, we get

$$\begin{aligned}
\int_{-\gamma} f(z) \, dz &= \int_b^a f(\gamma(s)) \gamma'(s) \, ds \\
&= - \int_a^b f(\gamma(s)) \gamma'(s) \, ds \\
&= - \int_{\gamma} f(z) \, dz.
\end{aligned}$$

□

Example 4.2

$z_0 \in \mathbb{C}$, $R > 0$, $n \in \mathbb{Z}$, $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$, $\gamma(t) = z_0 + Re^{it}$. If $n \neq -1$, then

$$\int_{\gamma} (z - z_0)^n \, dz = 0$$

If $n = -1$, then

$$\int_{\gamma} (z - z_0)^n \, dz = 2\pi i$$

Indeed, if $n \neq -1$, then $F(z) = \frac{1}{n+1}(z - z_0)^{n+1}$ is an antiderivative of $(z - z_0)^n$, so

$$\begin{aligned}
\int_{\gamma} (z - z_0)^n \, dz &= F(\gamma(2\pi)) - F(\gamma(0)) \\
&= \frac{1}{n+1}(\gamma(2\pi) - z_0)^{n+1} - \frac{1}{n+1}(\gamma(0) - z_0)^{n+1} \\
&= 0.
\end{aligned}$$

If $n = -1$, then

$$\begin{aligned}
\int_{\gamma} \frac{1}{z - z_0} \, dz &= \int_0^{2\pi} \frac{1}{\gamma(t) - z_0} \gamma'(t) \, dt \\
&= \int_0^{2\pi} \frac{1}{Re^{it}} iRe^{it} \, dt \\
&= \int_0^{2\pi} i \, dt \\
&= 2\pi i.
\end{aligned}$$

4.3 Estimates and Uniform Convergence

Definition 4.5

Length of a Path

If $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$ its length is

$$L(\gamma) := \int_a^b |\gamma'(t)| \, dt$$

Theorem 4.6

ML Inequality

If $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$ and f is continuous on the image of γ , then

$$\left| \int_{\gamma} f(z) \, dz \right| \leq L(\gamma) M_f$$

where M_f is the maximum value of $|(f \circ \gamma)(t)|$ for $t \in [a, b]$.

Proof:

$$\begin{aligned} \left| \int_{\gamma} f(z) \, dz \right| &= \left| \int_a^b f(\gamma(t)) \gamma'(t) \, dt \right| \\ &\leq \int_a^b |f(\gamma(t)) \gamma'(t)| \, dt \\ &= \int_a^b |f(\gamma(t))| |\gamma'(t)| \, dt \\ &\leq \int_a^b M_f |\gamma'(t)| \, dt \\ &= M_f L(\gamma) \end{aligned}$$

□

Theorem 4.7

Uniform Convergence for Path Integrals

Suppose $\gamma : [a, b] \rightarrow \mathbb{C}$ is piecewise- $\mathcal{C}^1$ and $(f_n)_{n=1}^{\infty}$ are continuous on the image of γ and (f_n) converges uniformly to f on the range of γ . Then

$$\int_{\gamma} f = \lim_{n \rightarrow \infty} \int_{\gamma} f_n$$

Proof: Since $f_n \rightarrow f$ uniformly and each f_n is continuous on the image of γ , f is continuous on the image of γ . By the [ML Inequality](#),

$$\begin{aligned}
\left| \int_{\gamma} f_n - \int_{\gamma} f \right| &= \left| \int_{\gamma} (f_n - f) \right| \\
&\leq L(\gamma) \sup_{t \in [a, b]} |f_n(\gamma(t)) - f(\gamma(t))| \\
&= L(\gamma) \sup_{z \in \gamma([a, b])} |f_n(z) - f(z)|.
\end{aligned}$$

Since $f_n \rightarrow f$ uniformly on the range of γ , the last expression tends to 0. Hence $\int_{\gamma} f_n \rightarrow \int_{\gamma} f$. $\square$

4.4 Cauchy's Theorem

Example 4.3

Fourier Transform of the Gaussian

Let $a, b > 0$ and let $R_{a,b}$ be the positively oriented boundary of the rectangle with vertices $-a$, a , $a + ib$, and $-a + ib$. Consider

$$\int_{R_{a,b}} e^{-z^2} dz.$$

Since

$$e^{-z^2} = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n}}{n!},$$

define

$$g(z) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)n!}.$$

The power series for g has infinite radius of convergence, and termwise differentiation gives $g'(z) = e^{-z^2}$. Hence

$$\int_{R_{a,b}} e^{-z^2} dz = \int_{R_{a,b}} g'(z) dz = 0.$$

Writing the four sides of $R_{a,b}$ separately gives

$$0 = \int_{-a}^a e^{-x^2} dx + \int_0^b e^{-(a+iy)^2} i dy - \int_{-a}^a e^{-(x+ib)^2} dx - \int_0^b e^{-(-a+iy)^2} i dy.$$

Thus

$$\int_{-a}^a e^{-(x+ib)^2} dx = \int_{-a}^a e^{-x^2} dx + i \int_0^b e^{-(a+iy)^2} dy - i \int_0^b e^{-(-a+iy)^2} dy.$$

Since

$$e^{-(x+ib)^2} = e^{-x^2+b^2-2ibx},$$

we get

$$\int_{-a}^a e^{-x^2} e^{-2ibx} dx = e^{-b^2} \int_{-a}^a e^{-x^2} dx + ie^{-b^2} \int_0^b e^{-(a+iy)^2} dy - ie^{-b^2} \int_0^b e^{-(-a+iy)^2} dy.$$

For $0 \leq y \leq b$,

$$\left| e^{-(a+iy)^2} \right| = e^{-a^2+y^2} \leq e^{-a^2+b^2}$$

and the same bound holds for $e^{-(-a+iy)^2}$. Hence the two vertical integrals tend to 0 as $a \rightarrow \infty$. Therefore

$$\int_{-\infty}^{\infty} e^{-x^2} e^{-2ibx} dx = e^{-b^2} \int_{-\infty}^{\infty} e^{-x^2} dx.$$

Since $\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$,

$$\int_{-\infty}^{\infty} e^{-x^2} e^{-2ibx} dx = \sqrt{\pi} e^{-b^2}.$$

Theorem 4.8

Cauchy's Theorem for a Triangle (Goursat's Lemma)

If T is the positively oriented boundary of a triangle in $\mathbb{C}$ and if f is holomorphic in a region that contains T and its interior, then

$$\int_T f(z) dz = 0.$$

Proof: Let Δ be the closed filled triangle with boundary T . Let

$$I = \int_T f(z) dz$$

Divide Δ into four congruent closed subtriangles by joining the midpoints of the sides. The boundary integrals over the four subtriangles add to I , because the integrals over the shared sides cancel. Hence one subtriangle, call it Δ_1 , with positively oriented boundary $T_1 = \partial\Delta_1$, satisfies

$$\left| \int_{T_1} f(z) dz \right| \geq \frac{|I|}{4}.$$

Repeating this construction gives nested closed filled triangles

$$\Delta \supseteq \Delta_1 \supseteq \Delta_2 \supseteq \cdots$$

with positively oriented boundaries $T_n = \partial\Delta_n$ such that

$$\left| \int_{T_n} f(z) dz \right| \geq \frac{|I|}{4^n}.$$

If P is the perimeter of T and D is the diameter of Δ , then the perimeter of T_n and the diameter of Δ_n are $\frac{P}{2^n}$ and $\frac{D}{2^n}$ respectively.

Since the Δ_n are nested compact sets and their diameters tend to 0, there is a unique point

$$z_0 \in \bigcap_{n=1}^{\infty} \Delta_n.$$

Since f is holomorphic at z_0 ,

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + \eta(z)(z - z_0),$$

where $\eta(z) \rightarrow 0$ as $z \rightarrow z_0$. The first two terms have primitives, so their integrals around T_n are 0. Thus

$$\int_{T_n} f(z) dz = \int_{T_n} \eta(z)(z - z_0) dz.$$

Let $\varepsilon > 0$. Since $\eta(z) \rightarrow 0$, for all sufficiently large n we have $|\eta(z)| < \varepsilon$ for all $z \in \Delta_n$. Hence, by the [ML Inequality](#),

$$\left| \int_{T_n} f(z) dz \right| \leq \varepsilon \cdot \left(\frac{D}{2^n} \right) \cdot \left(\frac{P}{2^n} \right) = \frac{\varepsilon PD}{4^n}.$$

Combining this with the lower bound gives, for all sufficiently large n ,

$$\frac{|I|}{4^n} \leq \left| \int_{T_n} f(z) dz \right| \leq \frac{\varepsilon PD}{4^n}.$$

Multiplying by 4^n , we get $|I| \leq \varepsilon PD$. Since $\varepsilon > 0$ was arbitrary, $I = 0$. □

4.5 Cauchy's Theorem for Convex Regions

Theorem 4.9

Cauchy's Theorem for Convex Regions

Suppose G is a convex region and f is holomorphic in G . Then for any piecewise- $\mathcal{C}^1$ closed curve γ in G we have

$$\int_{\gamma} f(z) dz = 0$$

Proof: Fix $z_0 \in G$. For $z \in G$, define

$$F(z) = \int_{[z_0, z]} f(w) dw.$$

This is well-defined since G is convex. We claim that $F' = f$.

Let $z \in G$. Since G is open, for h sufficiently small we have $z + h \in G$. By convexity, the triangle with vertices $z_0, z, z + h$ is contained in G , so by [Cauchy's Theorem for a Triangle \(Goursat's Lemma\)](#),

$$\int_{[z_0, z]} f(w) dw + \int_{[z, z+h]} f(w) dw - \int_{[z_0, z+h]} f(w) dw = 0.$$

Hence

$$F(z+h) - F(z) = \int_{[z, z+h]} f(w) dw.$$

Parametrizing the line segment $[z, z+h]$ by $w = z + th$, $0 \leq t \leq 1$, gives

$$\frac{F(z+h) - F(z)}{h} = \int_0^1 f(z+th) dt.$$

Therefore

$$\frac{F(z+h) - F(z)}{h} - f(z) = \int_0^1 (f(z+th) - f(z)) dt.$$

Since f is continuous at z , the right side tends to 0 as $h \rightarrow 0$. Thus $F'(z) = f(z)$.

Now if $\gamma : [a, b] \rightarrow G$ is a piecewise- $\mathcal{C}^1$ closed curve, then by the [Fundamental Theorem for Path Integrals](#),

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)) = 0,$$

since $\gamma(a) = \gamma(b)$. □

4.6 Cauchy's Integral Formula

Theorem 4.10

Cauchy's Formula for the Circle

Suppose C is a circle oriented counter-clockwise and f is holomorphic in a region that contains C and its interior, then if z is in the interior of C

$$f(z) = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} d\zeta$$

Proof: Let

$$g(\zeta) = \begin{cases} \frac{f(\zeta) - f(z)}{\zeta - z} & \text{if } \zeta \neq z \\ f'(z) & \text{if } \zeta = z. \end{cases}$$

Then g is holomorphic in the interior of C , so by [Cauchy's Theorem for Convex Regions](#),

$$\int_C g(\zeta) d\zeta = 0.$$

Hence

$$\int_C \frac{f(\zeta)}{\zeta - z} d\zeta = f(z) \int_C \frac{1}{\zeta - z} d\zeta.$$

It remains to compute the last integral. Write $C(t) = z_0 + Re^{it}$, $0 \leq t \leq 2\pi$, and set $a = \frac{z - z_0}{R}$. Since z is in the interior of C , $|a| < 1$. Then

$$\begin{aligned} \int_C \frac{1}{\zeta - z} d\zeta &= \int_0^{2\pi} \frac{iRe^{it}}{z_0 + Re^{it} - z} dt \\ &= \int_0^{2\pi} \frac{i}{1 - ae^{-it}} dt \\ &= \int_0^{2\pi} i \sum_{n=0}^{\infty} a^n e^{-int} dt \\ &= 2\pi i. \end{aligned}$$

The third equality uses the geometric series expansion, valid uniformly since $|a| < 1$. The last equality follows by integrating the uniformly convergent series term-by-term: all terms with $n \geq 1$ integrate to 0, and the $n = 0$ term gives $2\pi i$. Therefore

$$\int_C \frac{f(\zeta)}{\zeta - z} d\zeta = 2\pi i f(z),$$

which gives the result. □

Note

The integral is 0 for z outside the circle

4.7 Consequences of Cauchy's Formula

Corollary 4.11

Mean Value Property

Suppose f is holomorphic on a region containing $D(z_0, R)$. If $0 < r < R$, then

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{it}) dt.$$

Proof: Let C_r be the circle of radius r centered at z_0 , oriented counter-clockwise. By [Cauchy's Formula for the Circle](#),

$$f(z_0) = \frac{1}{2\pi i} \int_{C_r} \frac{f(\zeta)}{\zeta - z_0} d\zeta.$$

Parametrize C_r by $\gamma(t) = z_0 + re^{it}$ for $0 \leq t \leq 2\pi$. Then $\gamma'(t) = ire^{it}$, so

$$\begin{aligned}
 f(z_0) &= \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z_0 + re^{it})}{re^{it}} ire^{it} dt \\
 &= \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{it}) dt.
 \end{aligned}$$

□

Note

This is called the *mean value property* because $f(z_0)$ is equal to the average of the values of f on the circle centered at z_0 .

4.8 Cauchy Integrals and Analyticity**Definition 4.6**

If φ is a continuous function on the range of some piecewise- $\mathcal{C}^1$ path γ . The *Cauchy Integral* over γ is the function

$$z \mapsto \int_{\gamma} \frac{\varphi(\zeta)}{\zeta - z} d\zeta$$

This is defined for z not in the range of γ

Definition 4.7**Distance to a Set**

If $A \subseteq \mathbb{C}$ and $z_0 \in \mathbb{C}$, define

$$\text{dist}(z_0, A) = \inf_{z \in A} |z - z_0|.$$

Proposition 4.12**Power Series of a Cauchy Integral**

Let $\gamma : [a, b] \rightarrow \mathbb{C}$ be a piecewise- $\mathcal{C}^1$ path, let $\Gamma = \gamma([a, b])$, and suppose φ is continuous on Γ . Define $f : \mathbb{C} \setminus \Gamma \rightarrow \mathbb{C}$ by

$$f(z) = \int_{\gamma} \frac{\varphi(\zeta)}{\zeta - z} d\zeta$$

If $z_0 \notin \Gamma$ and

$$R = \text{dist}(z_0, \Gamma),$$

then f has a power series representation on $D(z_0, R)$.

Proof: Let $z \in D(z_0, R)$. Then for $\zeta \in \Gamma$,

$$\left| \frac{z - z_0}{\zeta - z_0} \right| < 1.$$

Thus

$$\begin{aligned}\frac{1}{\zeta - z} &= \frac{1}{\zeta - z_0 - (z - z_0)} \\ &= \frac{1}{\zeta - z_0} \cdot \frac{1}{1 - \frac{z - z_0}{\zeta - z_0}} \\ &= \sum_{n=0}^{\infty} \frac{(z - z_0)^n}{(\zeta - z_0)^{n+1}}.\end{aligned}$$

Therefore

$$f(z) = \int_{\gamma} \varphi(\zeta) \sum_{n=0}^{\infty} \frac{(z - z_0)^n}{(\zeta - z_0)^{n+1}} d\zeta.$$

If $0 < \rho < R$ and $|z - z_0| \leq \rho$, then

$$\left| \frac{z - z_0}{\zeta - z_0} \right| \leq \frac{\rho}{R} < 1$$

for all $\zeta \in \Gamma$. Hence the geometric series converges uniformly for $z \in \overline{D(z_0, \rho)}$ and $\zeta \in \Gamma$, so we may integrate term by term. Thus

$$f(z) = \sum_{n=0}^{\infty} \left(\int_{\gamma} \frac{\varphi(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \right) (z - z_0)^n.$$

This is a power series centered at z_0 . □

Theorem 4.13

Analyticity of Holomorphic Functions

Suppose f is holomorphic on a region containing $D(z_0, R)$. Then f has a power series representation centered at z_0 with radius of convergence at least R . More precisely, for $0 < r < R$ and $|z - z_0| < r$,

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n,$$

where

$$a_n = \frac{1}{2\pi i} \int_{C_r} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta,$$

and C_r is the positively oriented circle of radius r centered at z_0 .

Proof: Fix $0 < r < R$. By [Cauchy's Formula for the Circle](#), for $|z - z_0| < r$,

$$f(z) = \frac{1}{2\pi i} \int_{C_r} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

This is a Cauchy integral with $\varphi(\zeta) = \frac{f(\zeta)}{2\pi i}$. By the [Power Series of a Cauchy Integral](#),

$$f(z) = \sum_{n=0}^{\infty} \left(\int_{C_r} \frac{f(\zeta)}{2\pi i (\zeta - z_0)^{n+1}} d\zeta \right) (z - z_0)^n.$$

Hence

$$a_n = \frac{1}{2\pi i} \int_{C_r} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta.$$

Since $r < R$ was arbitrary, the radius of convergence is at least R . □

4.9 Derivative Formula

Corollary 4.14

Cauchy's Integral Formula for Derivatives

Suppose f is holomorphic on a region containing $D(z_0, R)$. If $0 < r < R$ and C_r is the positively oriented circle of radius r centered at z_0 , then for every $n \geq 0$,

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_{C_r} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta.$$

Proof: By the [Analyticity of Holomorphic Functions](#), for $|z - z_0| < r$,

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n,$$

where

$$a_n = \frac{1}{2\pi i} \int_{C_r} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta.$$

Since power series can be differentiated term-by-term inside their radius of convergence,

$$f^{(n)}(z_0) = n! a_n.$$

Substituting the formula for a_n gives

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_{C_r} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta.$$

□

Corollary 4.15

Holomorphicity of Derivatives

If f is holomorphic on a region G , then $f^{(n)}$ is holomorphic on G for every $n \geq 0$.

Proof: Let $z_0 \in G$. Since G is open, there is $R > 0$ such that $D(z_0, R) \subseteq G$. By the [Analyticity of Holomorphic Functions](#), f has a power series representation on this disk. Power series can be differentiated term-by-term inside their radius of convergence, so f' is holomorphic on the disk. Repeating this argument gives that $f^{(n)}$ is holomorphic near z_0 for every $n \geq 0$. Since $z_0 \in G$ was arbitrary, $f^{(n)}$ is holomorphic on G . □

4.10 Operations on Power Series

Proposition 4.16

Cauchy Product of Power Series

Suppose $\sum_{n=0}^{\infty} a_n(z - z_0)^n$ has radius of convergence R_1 and $\sum_{n=0}^{\infty} b_n(z - z_0)^n$ has radius of convergence R_2 . Consider

$$\sum_{n=0}^{\infty} \left(\sum_{k=0}^n a_k b_{n-k} \right) (z - z_0)^n$$

Then the radius of convergence for this power series is at least $\min\{R_1, R_2\}$. If f and g are the functions defined by the first two power series, then for $|z - z_0| < \min\{R_1, R_2\}$,

$$f(z)g(z) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n a_k b_{n-k} \right) (z - z_0)^n.$$

Proof: Let $R = \min\{R_1, R_2\}$ and define $h(z) = f(z)g(z)$ on $D(z_0, R)$. Since f and g are holomorphic on this disk, so is h . Thus h has a power series representation

$$h(z) = \sum_{n=0}^{\infty} c_n (z - z_0)^n$$

on $D(z_0, R)$, where $c_n = h^{(n)}(z_0)/n!$. By Leibniz's rule,

$$\begin{aligned} h^{(n)}(z_0) &= \sum_{k=0}^n \binom{n}{k} f^{(k)}(z_0) g^{(n-k)}(z_0) \\ &= \sum_{k=0}^n \frac{n!}{k!(n-k)!} \cdot k! a_k \cdot (n-k)! b_{n-k} \\ &= n! \sum_{k=0}^n a_k b_{n-k}. \end{aligned}$$

Hence

$$c_n = \sum_{k=0}^n a_k b_{n-k}.$$

Therefore the Cauchy product represents $h = fg$ on $D(z_0, R)$, so its radius of convergence is at least R .

□

4.11 Converse Results

Proposition 4.17

Converse of Goursat's Lemma

If G is a region, $f : G \rightarrow \mathbb{C}$ is continuous, and $\int_T f = 0$ for every triangle T whose edges and interior are contained in G , then f is holomorphic.

Proof: It is enough to show that f is holomorphic in a neighbourhood of each point. Fix $z_0 \in G$ and choose a disk D centered at z_0 with $D \subseteq G$.

For $z \in D$, define

$$g(z) = \int_{[z_0, z]} f(\zeta) d\zeta.$$

The triangle hypothesis implies that this integral is path independent in D : if $z_1, z_2, z_3 \in D$, then the integral around the triangle with these vertices is 0.

By the same line-segment argument as in the proof of [Cauchy's Theorem for Convex Regions](#), $g' = f$ on D . Therefore g is holomorphic on D . Since derivatives of holomorphic functions are holomorphic by [Holomorphicity of Derivatives](#), $f = g'$ is holomorphic on D . Since z_0 was arbitrary, f is holomorphic on G . $\square$

4.12 Liouville's Theorem and Algebraic Consequences

Theorem 4.18

Liouville's Theorem

Suppose $f : \mathbb{C} \rightarrow \mathbb{C}$ is entire and bounded. Then f is constant.

Proof: Suppose $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Fix $z_0 \in \mathbb{C}$ and $r > 0$. Let C_r be the circle centered at z_0 with radius r , oriented counter-clockwise. By [Cauchy's Integral Formula for Derivatives](#),

$$f'(z_0) = \frac{1}{2\pi i} \int_{C_r} \frac{f(\zeta)}{(\zeta - z_0)^2} d\zeta.$$

By the [ML Inequality](#),

$$|f'(z_0)| \leq \frac{1}{2\pi} \cdot \left(\frac{M}{r^2} \right) \cdot 2\pi r = \frac{M}{r}.$$

Since this holds for every $r > 0$, letting $r \rightarrow \infty$ gives $f'(z_0) = 0$. Since z_0 was arbitrary, $f' = 0$ on $\mathbb{C}$, so f is constant. $\square$

Theorem 4.19

The Fundamental Theorem of Algebra

If $p(z) = \sum_{k=0}^n a_k z^k$ is a nonconstant polynomial with complex coefficients and $a_n \neq 0$, then there exist $z_1, \dots, z_n \in \mathbb{C}$ such that

$$p(z) = a_n \prod_{k=1}^n (z - z_k).$$

Proof: It is enough to show that p has a zero, since after finding a zero z_1 , we may factor $p(z) = (z - z_1)q(z)$ and repeat by induction on the degree.

Suppose, for contradiction, that p has no zero. Then $\frac{1}{p}$ is entire. We claim that $\frac{1}{p}$ is bounded. Since $a_n \neq 0$,

$$p(z) = z^n \left(a_n + \frac{a_{n-1}}{z} + \dots + \frac{a_0}{z^n} \right).$$

For $|z|$ sufficiently large, the term in parentheses has absolute value at least $\frac{|a_n|}{2}$, so

$$|p(z)| \geq \frac{|a_n|}{2} \cdot |z|^n.$$

Hence $\left|\frac{1}{p(z)}\right| \rightarrow 0$ as $|z| \rightarrow \infty$, so $\frac{1}{p}$ is bounded outside a large disk. On the closed disk, $\frac{1}{p}$ is continuous, hence bounded. Thus $\frac{1}{p}$ is bounded on $\mathbb{C}$.

By [Liouville's Theorem](#), $\frac{1}{p}$ is constant, so p is constant, contradicting that p is nonconstant. Therefore p has a zero, and the factorization follows by induction. $\square$

5 Zeros and the Identity Theorem

5.1 Orders of Zeros

Definition 5.1

Zero of Order n

Suppose f is holomorphic on G and $f(z_0) = 0$. We say z_0 is a *zero of order n* if

$$0 = f(z_0) = f'(z_0) = \dots = f^{(n-1)}(z_0)$$

but $f^{(n)}(z_0) \neq 0$. A zero of order 1 is called a *simple zero*.

Proposition 5.1

Infinite-Order Zero

Suppose G is open, f is holomorphic on G , and f has a zero of infinite order at $z_0 \in G$. Then $f = 0$ on the connected component of G containing z_0 .

Proof: Let H be the connected component of G containing z_0 . Define

$$A = \{z \in H \mid f \text{ is identically zero on some disk centered at } z\}.$$

Since f has a zero of infinite order at z_0 , every coefficient in the power series for f centered at z_0 is 0. Thus f is identically zero on some disk centered at z_0 , so $z_0 \in A$ and A is non-empty.

The set A is open in H by definition. We show that A is closed in H . Suppose $z_k \in A$ and $z_k \rightarrow z \in H$. For each $m \geq 0$, since f is identically zero near each z_k , we have

$$f^{(m)}(z_k) = 0.$$

Since $f^{(m)}$ is continuous, $f^{(m)}(z) = 0$. Hence all derivatives of f vanish at z , so the power series for f centered at z is identically zero on some disk centered at z . Thus $z \in A$.

Therefore A is non-empty, open, and closed in H . Since H is connected, $A = H$, so $f = 0$ on H . $\square$

Proposition 5.2

Finite-Order Zeros are Isolated

Suppose f is holomorphic on an open set G and z_0 is a zero of finite order n . Then there is a disk centered at z_0 on which z_0 is the only zero of f .

Proof: Since z_0 is a zero of order n , the power series for f centered at z_0 has the form

$$f(z) = \sum_{k=n}^{\infty} a_k (z - z_0)^k$$

with $a_n \neq 0$. Thus

$$f(z) = (z - z_0)^n g(z),$$

where

$$g(z) = \sum_{k=0}^{\infty} a_{n+k} (z - z_0)^k.$$

Then g is holomorphic near z_0 and $g(z_0) = a_n \neq 0$. By continuity, $g(z) \neq 0$ on some disk centered at z_0 . On this disk, the only zero of $f(z) = (z - z_0)^n g(z)$ is z_0 . $\square$

5.2 Identity Theorem

Corollary 5.3

Identity Theorem

Suppose f and g are holomorphic on an open set G . If f and g agree on a subset $S \subseteq G$ that has a limit point $z_0 \in G$, then $f = g$ on the connected component of G containing z_0 .

Proof: Let $h = f - g$. Then h is holomorphic on G and $h = 0$ on S . Since z_0 is a limit point of S , there is a sequence (z_n) in S with $z_n \rightarrow z_0$ and $z_n \neq z_0$. Since h is continuous, $h(z_0) = 0$.

If h had a zero of finite order at z_0 , then z_0 would be an isolated zero of h , contradicting that $z_n \rightarrow z_0$ with $z_n \neq z_0$ and $h(z_n) = 0$. Hence h has a zero of infinite order at z_0 . By the [Infinite-Order Zero](#), $h = 0$ on the connected component of G containing z_0 . Thus $f = g$ there. $\square$

6 Convergence and Maximum Principles

6.1 Identity Theorem Applications

Example 6.1

Suppose $f : \mathbb{D} \rightarrow \mathbb{C}$ is holomorphic and $f(\frac{1}{n}) = \frac{1}{n^2}$ for all $n \in \mathbb{N}$. What is $f'(\frac{i}{2})$?

Proof: Let $g(z) = z^2$. Then $f(\frac{1}{n}) = g(\frac{1}{n})$ for all $n \in \mathbb{N}$, and $\frac{1}{n} \rightarrow 0 \in \mathbb{D}$. By the [Identity Theorem](#), $f = g$ on $\mathbb{D}$. Hence

$$f'\left(\frac{i}{2}\right) = g'\left(\frac{i}{2}\right) = 2 \cdot \frac{i}{2} = i.$$

$\square$

6.2 Locally Uniform Limits

Proposition 6.1

Weierstrass Theorem

Suppose $G \subseteq \mathbb{C}$ is open and $f_k : G \rightarrow \mathbb{C}$ is holomorphic for each $k \in \mathbb{N}$. If $f_k \rightarrow f$ locally uniformly on G , then f is holomorphic. Moreover, for each $n \geq 0$, $f_k^{(n)} \rightarrow f^{(n)}$ locally uniformly on G .

Proof: Fix $z_0 \in G$ and choose $R > 0$ such that the closed disk $\overline{B(z_0, R)} \subseteq G$. Let C be the positively oriented circle centered at z_0 with radius R .

If $z \in B(z_0, R)$, then by [Cauchy's Formula for the Circle](#),

$$f_k(z) = \frac{1}{2\pi i} \int_C \frac{f_k(\zeta)}{\zeta - z} d\zeta.$$

Since $f_k \rightarrow f$ uniformly on C , the [Uniform Convergence for Path Integrals](#) gives

$$f(z) = \lim_{k \rightarrow \infty} f_k(z) = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Thus f is a Cauchy integral on $B(z_0, R)$, so f is holomorphic there. Since z_0 was arbitrary, f is holomorphic on G .

If $0 < r < R$ and $z \in B(z_0, r)$, then by [Cauchy's Integral Formula for Derivatives](#),

$$f_k^{(n)}(z) - f^{(n)}(z) = \frac{n!}{2\pi i} \int_C \frac{f_k(\zeta) - f(\zeta)}{(\zeta - z)^{n+1}} d\zeta.$$

Since $|\zeta - z| \geq R - r$ for $\zeta \in C$ and $z \in B(z_0, r)$, the [ML Inequality](#) gives

$$\left| f_k^{(n)}(z) - f^{(n)}(z) \right| \leq n! \frac{R}{(R - r)^{n+1}} \sup_{\zeta \in C} |f_k(\zeta) - f(\zeta)|.$$

The supremum tends to 0 by local uniform convergence, so $f_k^{(n)} \rightarrow f^{(n)}$ uniformly on $B(z_0, r)$. Since z_0 was arbitrary, the convergence is locally uniform on G . $\square$

6.3 Maximum Modulus Principle

Theorem 6.2

Maximum Modulus Principle

If G is a region and $f : G \rightarrow \mathbb{C}$ is holomorphic and nonconstant, then $|f|$ has no local maximum in G .

Proof: Suppose $|f|$ has a local maximum at $z_0 \in G$. Choose $R > 0$ such that $D(z_0, R) \subseteq G$ and

$$|f(z)| \leq |f(z_0)|$$

for all $z \in D(z_0, R)$.

If $f(z_0) = 0$, then $|f(z)| = 0$ on this disk, so f is locally constant. By the [Identity Theorem](#), f is constant on G , a contradiction. Thus $f(z_0) \neq 0$.

For $0 < r < R$, the [Mean Value Property](#) gives

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{it}) dt.$$

Taking absolute values,

$$|f(z_0)| \leq \frac{1}{2\pi} \int_0^{2\pi} |f(z_0 + re^{it})| dt \leq |f(z_0)|.$$

Hence equality holds throughout. Since the continuous function

$$t \mapsto |f(z_0)| - |f(z_0 + re^{it})|$$

is nonnegative and has integral 0, it is identically 0. Thus $|f(z_0 + re^{it})| = |f(z_0)|$ for all t .

Equality in the triangle inequality also forces all values $f(z_0 + re^{it})$ to have the same argument as $f(z_0)$. Therefore

$$f(z_0 + re^{it}) = f(z_0)$$

for all t . Since this holds for every $0 < r < R$, $f = f(z_0)$ on the disk centered at z_0 with radius R . By the [Identity Theorem](#), f is constant on G , contradicting the hypothesis. Therefore $|f|$ has no local maximum in G . □

Corollary 6.3

If G is a region, $f : G \rightarrow \mathbb{C}$ is holomorphic and nonconstant, and $K \subseteq G$ is compact and nonempty, then $|f|$ attains its maximum on ∂K .

6.4 Schwarz's Lemma

Theorem 6.4

Schwarz's Lemma

Suppose $f : \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$. Then $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$. Moreover, if $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D}$, $z_0 \neq 0$, then there exists $\lambda \in \mathbb{C}$ with $|\lambda| = 1$ such that $f(z) = \lambda z$ for all $z \in \mathbb{D}$.

Proof: Define

$$g(z) = \begin{cases} \frac{f(z)}{z} & \text{if } z \neq 0 \\ f'(0) & \text{if } z = 0. \end{cases}$$

Since $f(0) = 0$, the singularity at 0 is removable, so g is holomorphic on $\mathbb{D}$.

Fix $0 < r < 1$. On the circle $|z| = r$,

$$|g(z)| = \frac{|f(z)|}{|z|} < \frac{1}{r}.$$

By the [Maximum Modulus Principle](#), for $|z| \leq r$ we have $|g(z)| \leq \frac{1}{r}$. Letting $r \rightarrow 1$ gives $|g(z)| \leq 1$ for all $z \in \mathbb{D}$. Hence, for $z \neq 0$,

$$|f(z)| = |z||g(z)| \leq |z|,$$

and the inequality is also true at $z = 0$.

Now suppose $|f(z_0)| = |z_0|$ for some $z_0 \neq 0$. Then $|g(z_0)| = 1$, so $|g|$ has a local maximum in $\mathbb{D}$. By the [Maximum Modulus Principle](#), g is constant. Thus $g(z) = \lambda$ for some $\lambda \in \mathbb{C}$ with $|\lambda| = 1$, and so $f(z) = \lambda z$ for all $z \in \mathbb{D}$. $\square$

6.5 Minimum Modulus Principle

Proposition 6.5

Minimum Modulus Principle

Suppose G is a region and $f : G \rightarrow \mathbb{C}$ is holomorphic and nonconstant. If $|f|$ has a local minimum at $z_0 \in G$, then $f(z_0) = 0$.

Proof: Suppose $|f|$ has a local minimum at z_0 and $f(z_0) \neq 0$. Then f is nonzero on some disk centered at z_0 , so $\frac{1}{f}$ is holomorphic on that disk. Since $|f|$ has a local minimum at z_0 , $\left|\frac{1}{f}\right|$ has a local maximum at z_0 . By the [Maximum Modulus Principle](#), $\frac{1}{f}$ is constant on a disk centered at z_0 , so f is constant on that disk. By the [Identity Theorem](#), f is constant on G , a contradiction. Thus $f(z_0) = 0$. $\square$

Example 6.2

The conclusion is possible: if $f(z) = z$ and $K = \overline{\mathbb{D}}$, then $|f|$ attains its minimum at 0, and $f(0) = 0$.

7 Laurent Series

7.1 Examples

Example 7.1

For $z \in \mathbb{D}$,

$$\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n.$$

For $|z| > 1$,

$$\frac{1}{1-z} = -\frac{1}{z} \cdot \frac{1}{1-\frac{1}{z}} = -\sum_{n=0}^{\infty} z^{-n-1} = -\sum_{n=1}^{\infty} z^{-n}.$$

Example 7.2

Consider

$$\frac{1}{(z-1)(z-2)} = \frac{1}{z-2} - \frac{1}{z-1}.$$

The Laurent expansion depends on the annulus.

If $z \in \mathbb{D}$, then

$$\begin{aligned}\frac{1}{z-2} &= -\frac{1}{2} \cdot \frac{1}{1-\frac{z}{2}} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \\ -\frac{1}{z-1} &= \frac{1}{1-z} = \sum_{n=0}^{\infty} z^n.\end{aligned}$$

Hence

$$\frac{1}{(z-1)(z-2)} = \sum_{n=0}^{\infty} \left(1 - \frac{1}{2^{n+1}}\right) z^n.$$

If $1 < |z| < 2$, then

$$\begin{aligned}\frac{1}{z-2} &= -\frac{1}{2} \cdot \frac{1}{1-\frac{z}{2}} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \\ -\frac{1}{z-1} &= -\frac{1}{z} \cdot \frac{1}{1-\frac{1}{z}} = -\sum_{n=0}^{\infty} z^{-n-1}.\end{aligned}$$

Hence

$$\frac{1}{(z-1)(z-2)} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} - \sum_{n=1}^{\infty} z^{-n}.$$

If $|z| > 2$, then

$$\begin{aligned}\frac{1}{z-2} &= \frac{1}{z} \cdot \frac{1}{1-\frac{2}{z}} = \sum_{n=0}^{\infty} 2^n z^{-n-1}, \\ -\frac{1}{z-1} &= -\frac{1}{z} \cdot \frac{1}{1-\frac{1}{z}} = -\sum_{n=0}^{\infty} z^{-n-1}.\end{aligned}$$

Hence

$$\frac{1}{(z-1)(z-2)} = \sum_{n=0}^{\infty} (2^n - 1) z^{-n-1}.$$

7.2 Definition and Convergence

Notation 7.1

If $z_0 \in \mathbb{C}$ and $0 \leq R_1 < R_2 \leq \infty$, write

$$A(z_0; R_1, R_2) = \{z \in \mathbb{C} \mid R_1 < |z - z_0| < R_2\}$$

for the annulus centered at z_0 with inner radius R_1 and outer radius R_2 .

Definition 7.1**Laurent Series**

A *Laurent Series* centered at z_0 is a series of the form

$$\sum_{n=-\infty}^{\infty} a_n (z - z_0)^n$$

where $a_n \in \mathbb{C}$. This converges at z if both

$$\sum_{n=0}^{\infty} a_n (z - z_0)^n \quad \text{and} \quad \sum_{n=1}^{\infty} a_{-n} (z - z_0)^{-n}$$

converge. Equivalently, the symmetric partial sums

$$\sum_{n=-N}^N a_n (z - z_0)^n$$

converge as $N \rightarrow \infty$. In that case,

$$\lim_{N \rightarrow \infty} \sum_{n=-N}^N a_n (z - z_0)^n$$

is defined to be the sum of the Laurent series.

Theorem 7.1**Convergence of Laurent Series**

Let

$$\sum_{n=-\infty}^{\infty} a_n (z - z_0)^n$$

be a Laurent series. Define

$$R_2 = \frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{\frac{1}{n}}}$$

and

$$R_1 = \limsup_{n \rightarrow \infty} |a_{-n}|^{\frac{1}{n}}.$$

Then the positive part converges absolutely and locally uniformly on

$$D(z_0, R_2),$$

and the negative part converges absolutely and locally uniformly on

$$|z - z_0| > R_1.$$

Hence, if $R_1 < R_2$, the Laurent series converges absolutely and locally uniformly on the annulus

$$A(z_0; R_1, R_2),$$

and the symmetric partial sums

$$\sum_{n=-N}^N a_n (z - z_0)^n$$

converge locally uniformly there. This annulus is called the *annulus of convergence* of the Laurent series.

Proof: By the [Cauchy-Hadamard Theorem](#), the positive part is an ordinary power series with radius of convergence R_2 , so it converges absolutely and locally uniformly on $D(z_0, R_2)$.

For the negative part, the root test gives

$$\limsup_{n \rightarrow \infty} |a_{-n} (z - z_0)^{-n}|^{\frac{1}{n}} = \frac{R_1}{|z - z_0|}.$$

Thus the negative part converges absolutely when $\frac{R_1}{|z - z_0|} < 1$, i.e. when $|z - z_0| > R_1$. The same estimate gives locally uniform convergence on compact subsets of $|z - z_0| > R_1$.

On every compact subannulus

$$R_1 < r \leq |z - z_0| \leq R < R_2,$$

both the positive and negative parts converge uniformly. Hence, if $R_1 < R_2$, the Laurent series converges absolutely and the symmetric partial sums converge locally uniformly on $A(z_0; R_1, R_2)$. $\square$

Theorem 7.2

Differentiating Laurent Series

Suppose the Laurent series

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n$$

converges on the annulus $A(z_0; R_1, R_2)$. Then f is holomorphic on this annulus, and

$$f'(z) = \sum_{n=-\infty}^{\infty} n a_n (z - z_0)^{n-1}.$$

Proof: By the [Convergence of Laurent Series](#), the symmetric partial sums converge locally uniformly on the annulus. Each symmetric partial sum is holomorphic on the annulus, so by the [Weierstrass Theorem](#) the limit f is holomorphic.

Again by the [Weierstrass Theorem](#), the derivatives of the symmetric partial sums converge locally uniformly to f' . Since

$$\frac{d}{dz} \sum_{n=-N}^N a_n (z - z_0)^n = \sum_{n=-N}^N n a_n (z - z_0)^{n-1},$$

it follows that

$$f'(z) = \sum_{n=-\infty}^{\infty} n a_n (z - z_0)^{n-1}.$$

$\square$

7.3 Annuli

Theorem 7.3

Cauchy's Theorem for an Annulus

Suppose $0 \leq R_1 < R_2 \leq \infty$ and f is holomorphic on the annulus $A(z_0; R_1, R_2)$. For $R_1 < r < R_2$, let C_r be the positively oriented circle centered at z_0 with radius r . Then $\int_{C_r} f(z) dz$ is independent of r . In other words, if $R_1 < r < s < R_2$, then

$$\int_{C_r} f(z) dz = \int_{C_s} f(z) dz.$$

Proof: Let

$$I(r) = \int_{C_r} f(z) dz.$$

Using the parametrization $z = z_0 + re^{it}$, $0 \leq t \leq 2\pi$, we have

$$I(r) = \int_0^{2\pi} f(z_0 + re^{it}) ire^{it} dt.$$

The integrand has continuous partial derivatives in r and t , so we may differentiate under the integral sign:

$$I'(r) = \int_0^{2\pi} \frac{\partial}{\partial r} (f(z_0 + re^{it}) ire^{it}) dt.$$

Now

$$\begin{aligned} \frac{\partial}{\partial r} (f(z_0 + re^{it}) ire^{it}) &= f'(z_0 + re^{it}) ire^{it} + f(z_0 + re^{it}) ie^{it} \\ &= \frac{\partial}{\partial t} (f(z_0 + re^{it}) e^{it}). \end{aligned}$$

Therefore

$$\begin{aligned} I'(r) &= \int_0^{2\pi} \frac{\partial}{\partial t} (f(z_0 + re^{it}) e^{it}) dt \\ &= f(z_0 + re^{2\pi i}) e^{2\pi i} - f(z_0 + r) \\ &= f(z_0 + r) - f(z_0 + r) \\ &= 0. \end{aligned}$$

Hence I is constant on (R_1, R_2) , so $\int_{C_r} f(z) dz$ is independent of r . □

7.4 Cauchy's Formula on an Annulus

Theorem 7.4

Cauchy's Formula for an Annulus

Suppose $0 \leq R_1 < R_2 \leq \infty$ and f is holomorphic on the annulus $A(z_0; R_1, R_2)$. For $R_1 < r < R_2$, let C_r be the positively oriented circle centered at z_0 with radius r . If

$$R_1 < r_1 < |w - z_0| < r_2 < R_2,$$

then

$$f(w) = \frac{1}{2\pi i} \int_{C_{r_2}} \frac{f(z)}{z - w} dz - \frac{1}{2\pi i} \int_{C_{r_1}} \frac{f(z)}{z - w} dz.$$

Proof: Fix w as above and define

$$g(z) = \begin{cases} \frac{f(z) - f(w)}{z - w} & \text{if } z \neq w \\ f'(w) & \text{if } z = w. \end{cases}$$

Then g is holomorphic on the annulus. By [Cauchy's Theorem for an Annulus](#),

$$\int_{C_{r_1}} g(z) dz = \int_{C_{r_2}} g(z) dz.$$

Hence

$$\int_{C_{r_2}} \frac{f(z)}{z-w} dz - \int_{C_{r_1}} \frac{f(z)}{z-w} dz = f(w) \int_{C_{r_2}} \frac{1}{z-w} dz - f(w) \int_{C_{r_1}} \frac{1}{z-w} dz.$$

The first integral on the right is $2\pi i$ by [Cauchy's Formula for the Circle](#), since w lies inside C_{r_2} . The second integral is 0 by [Cauchy's Theorem for Convex Regions](#), since w lies outside C_{r_1} and $z \mapsto \frac{1}{z-w}$ is holomorphic on the disk bounded by C_{r_1} . Therefore

$$\int_{C_{r_2}} \frac{f(z)}{z-w} dz - \int_{C_{r_1}} \frac{f(z)}{z-w} dz = 2\pi i f(w),$$

which gives the formula. □

8 Isolated Singularities

8.1 Laurent Series Representation

Corollary 8.1

Laurent Series Representation

Suppose $0 \leq R_1 < R_2 \leq \infty$ and write $A = A(z_0; R_1, R_2)$. If f is holomorphic on A , then f has a Laurent series representation on A :

$$f(w) = \sum_{n=-\infty}^{\infty} a_n (w - z_0)^n.$$

If $R_1 < r < R_2$ and C_r is the positively oriented circle centered at z_0 with radius r , then

$$a_n = \frac{1}{2\pi i} \int_{C_r} \frac{f(z)}{(z - z_0)^{n+1}} dz$$

for every $n \in \mathbb{Z}$.

Proof: Fix $w \in A$, and choose r_1, r_2 such that

$$R_1 < r_1 < |w - z_0| < r_2 < R_2.$$

By [Cauchy's Formula for an Annulus](#),

$$f(w) = \frac{1}{2\pi i} \int_{C_{r_2}} \frac{f(z)}{z-w} dz - \frac{1}{2\pi i} \int_{C_{r_1}} \frac{f(z)}{z-w} dz.$$

On C_{r_2} we have $|w - z_0| < |z - z_0|$, so

$$\frac{1}{z-w} = \frac{1}{z-z_0} \cdot \frac{1}{1 - \frac{w-z_0}{z-z_0}} = \sum_{n=0}^{\infty} \frac{(w-z_0)^n}{(z-z_0)^{n+1}},$$

locally uniformly in w . Hence the outer integral contributes

$$\sum_{n=0}^{\infty} \left(\frac{1}{2\pi i} \int_{C_{r_2}} \frac{f(z)}{(z - z_0)^{n+1}} dz \right) (w - z_0)^n.$$

On C_{r_1} we have $|z - z_0| < |w - z_0|$, so

$$\frac{1}{z - w} = -\frac{1}{w - z_0} \cdot \frac{1}{1 - \frac{z - z_0}{w - z_0}} = -\sum_{n=0}^{\infty} \frac{(z - z_0)^n}{(w - z_0)^{n+1}},$$

locally uniformly in w . Because the inner integral is subtracted, this contributes

$$\sum_{n=0}^{\infty} \left(\frac{1}{2\pi i} \int_{C_{r_1}} f(z)(z - z_0)^n dz \right) (w - z_0)^{-n-1}.$$

These are exactly the Laurent coefficients above. The coefficient formula is independent of the chosen radius r by [Cauchy's Theorem for an Annulus](#) applied to $z \mapsto \frac{f(z)}{(z - z_0)^{n+1}}$. $\square$

8.2 Classification

Notation 8.1

For $z_0 \in \mathbb{C}$ and $r > 0$, write

$$D^*(z_0, r) = \{z \in \mathbb{C} \mid 0 < |z - z_0| < r\}$$

for the punctured disk centered at z_0 with radius r .

Definition 8.1

Isolated Singularity

Suppose G is a region and $f : G \rightarrow \mathbb{C}$ is holomorphic. A point z_0 is an *isolated singularity* of f if $z_0 \notin G$ and there is some $R > 0$ such that

$$D^*(z_0, R) \subseteq G.$$

Note

If z_0 is an isolated singularity of f , then for sufficiently small $R > 0$, f is holomorphic on the punctured disk $D^*(z_0, R)$. Thus f has a Laurent series

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$$

near z_0 .

Definition 8.2

Removable Singularity

An isolated singularity z_0 of $f : G \rightarrow \mathbb{C}$ is *removable* if there is a holomorphic function $F : G \cup \{z_0\} \rightarrow \mathbb{C}$ such that $F|_G = f$.

Proposition 8.2**Removable Singularity Criterion**

Let z_0 be an isolated singularity of f , and write the Laurent series of f near z_0 as

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n.$$

Then z_0 is removable if and only if $a_n = 0$ for all $n < 0$.

Proof: If z_0 is removable, then the holomorphic extension has a power series at z_0 , so the Laurent series has no negative-power terms. Conversely, if all negative-power terms vanish, then the Laurent series is a power series in a disk centered at z_0 , so defining $F(z_0) = a_0$ gives a holomorphic extension. $\square$

Example 8.1

The function $f(z) = z$ on $\mathbb{C} \setminus \{0\}$ has a removable singularity at 0.

Example 8.2

The function

$$f(z) = \frac{z}{e^z - 1}$$

on $\mathbb{C} \setminus \{0\}$ has a removable singularity at 0, since $e^z - 1 = z + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots$

Definition 8.3**Pole**

An isolated singularity z_0 of f is a *pole of order m* if, in the Laurent series of f near z_0 ,

$$f(z) = \sum_{n=-m}^{\infty} a_n (z - z_0)^n$$

with $a_{-m} \neq 0$. A pole of order 1 is called a *simple pole*.

Proposition 8.3**Characterization of Poles**

Let z_0 be an isolated singularity of f . Then f has a pole of order m at z_0 if and only if $(z - z_0)^m f(z)$ has a removable singularity at z_0 and its removable extension is nonzero at z_0 .

Proof: If f has a pole of order m , then

$$f(z) = \sum_{n=-m}^{\infty} a_n (z - z_0)^n$$

with $a_{-m} \neq 0$. Hence

$$(z - z_0)^m f(z) = \sum_{n=-m}^{\infty} a_n (z - z_0)^{n+m} = \sum_{k=0}^{\infty} a_{k-m} (z - z_0)^k,$$

so $(z - z_0)^m f(z)$ has a removable singularity at z_0 , and the value of its extension at z_0 is $a_{-m} \neq 0$.

Conversely, if $(z - z_0)^m f(z)$ has a removable singularity at z_0 with nonzero value, write its holomorphic extension as

$$g(z) = \sum_{k=0}^{\infty} b_k (z - z_0)^k$$

with $b_0 \neq 0$. Then

$$f(z) = (z - z_0)^{-m} g(z) = \sum_{k=0}^{\infty} b_k (z - z_0)^{k-m},$$

so the first nonzero term is the $(z - z_0)^{-m}$ term. Thus f has a pole of order m at z_0 . $\square$

Example 8.3

The function $f(z) = \frac{1}{z}$ has a simple pole at 0.

Definition 8.4

Essential Singularity

An isolated singularity z_0 of f is an *essential singularity* if the Laurent series of f near z_0 has infinitely many nonzero negative-power terms.

Proposition 8.4

Classification of Isolated Singularities

Every isolated singularity is exactly one of the following: removable, a pole, or essential.

Proof: Let

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n$$

be the Laurent series near the isolated singularity z_0 . If $a_n = 0$ for all $n < 0$, then z_0 is removable. If there are finitely many nonzero negative-power terms, let $-m$ be the smallest index with $a_{-m} \neq 0$. Then z_0 is a pole of order m . If there are infinitely many nonzero negative-power terms, then z_0 is essential. $\square$

Example 8.4

The function

$$f(z) = e^{\frac{1}{z}} = \sum_{n=0}^{\infty} \frac{1}{n!} z^{-n}$$

has an essential singularity at 0.

8.3 Singularity at Infinity

Definition 8.5

Isolated Singularity at Infinity

Suppose G is a region and $f : G \rightarrow \mathbb{C}$ is holomorphic. The point ∞ is an isolated singularity of f if there exists some $R > 0$ such that

$$\{z \in \mathbb{C} \mid |z| > R\} \subseteq G.$$

Definition 8.6

Classification at Infinity

Suppose ∞ is an isolated singularity of f . Define

$$g(w) = f\left(\frac{1}{w}\right)$$

for $0 < |w| < \frac{1}{R}$. We say that ∞ is a removable singularity, a pole of order m , or an essential singularity of f if 0 is respectively a removable singularity, a pole of order m , or an essential singularity of g .

Example 8.5

The function $f(z) = \frac{1}{z}$ has a removable singularity at ∞ , since $f\left(\frac{1}{w}\right) = w$ has a removable singularity at 0.

8.4 Removable Singularities

Theorem 8.5

Removable Singularity Theorem

Let z_0 be an isolated singularity of $f : G \rightarrow \mathbb{C}$. Then z_0 is removable if and only if f is bounded on some punctured disk around z_0 . That is, there exist $r > 0$ and $M > 0$ such that

$$D^*(z_0, r) \subseteq G$$

and $|f(z)| \leq M$ for all $z \in D^*(z_0, r)$.

Proof: First suppose z_0 is removable. Let $F : G \cup \{z_0\} \rightarrow \mathbb{C}$ be a holomorphic extension of f . Since F is continuous at z_0 , there is some $r > 0$ such that F is bounded on

$$D(z_0, r).$$

Hence f is bounded on the punctured disk $D^*(z_0, r)$.

Conversely, suppose $|f(z)| \leq M$ for all $z \in D^*(z_0, r)$. Write the Laurent series of f near z_0 as

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n.$$

We show that all negative coefficients vanish. Fix $n \geq 1$. For $0 < \rho < r$, let C_ρ be the positively oriented circle centered at z_0 with radius ρ . By the [Laurent Series Representation](#),

$$a_{-n} = \frac{1}{2\pi i} \int_{C_\rho} f(z)(z - z_0)^{n-1} dz.$$

Therefore, by the [ML Inequality](#),

$$|a_{-n}| \leq \frac{1}{2\pi} \cdot M\rho^{n-1} \cdot 2\pi\rho = M\rho^n.$$

Since this holds for every $0 < \rho < r$, letting $\rho \rightarrow 0$ gives $a_{-n} = 0$. Thus $a_k = 0$ for all $k < 0$, so by the [Removable Singularity Criterion](#), z_0 is removable. $\square$

Theorem 8.6

Pole Characterization by Growth

Suppose z_0 is an isolated singularity of f . Then z_0 is a pole if and only if

$$\lim_{z \rightarrow z_0} |f(z)| = \infty.$$

Proof: First suppose z_0 is a pole of order m . By the [Characterization of Poles](#), the function

$$g(z) = (z - z_0)^m f(z)$$

has a removable singularity at z_0 , and its removable extension satisfies $g(z_0) \neq 0$. Hence there are $r > 0$ and $c > 0$ such that $|g(z)| \geq c$ for all $z \in D(z_0, r)$. Thus, for $z \in D^*(z_0, r)$,

$$|f(z)| = \frac{|g(z)|}{|z - z_0|^m} \geq \frac{c}{|z - z_0|^m}.$$

Letting $z \rightarrow z_0$ gives $|f(z)| \rightarrow \infty$.

Conversely, suppose

$$\lim_{z \rightarrow z_0} |f(z)| = \infty.$$

Then there is $r > 0$ such that $f(z) \neq 0$ for all $z \in D^*(z_0, r)$. Define $h(z) = \frac{1}{f(z)}$ on this punctured disk. Since $|h(z)| \rightarrow 0$ as $z \rightarrow z_0$, the function h is bounded near z_0 , so by the [Removable Singularity Theorem](#), h has a removable singularity at z_0 . Define its extension by $h(z_0) = 0$.

The zero of h at z_0 has finite order. Indeed, h is not identically zero on any punctured disk around z_0 , since f is finite there, so h cannot have a zero of infinite order at z_0 . Thus

$$h(z) = (z - z_0)^m k(z)$$

for some $m \geq 1$, where k is holomorphic near z_0 and $k(z_0) \neq 0$. Therefore

$$f(z) = \frac{1}{h(z)} = (z - z_0)^{-m} \frac{1}{k(z)},$$

where $\frac{1}{k}$ is holomorphic near z_0 . By the [Characterization of Poles](#), z_0 is a pole of order m . $\square$

Theorem 8.7**Picard's Theorem**

Let z_0 be an essential singularity of $f : G \rightarrow \mathbb{C}$. For every $\varepsilon > 0$ such that

$$D^*(z_0, \varepsilon) \subseteq G,$$

the image

$$f(D^*(z_0, \varepsilon))$$

omits at most one complex number.

Note

The proof of Picard's theorem is out of scope.

Theorem 8.8**Casorati-Weierstrass Theorem**

Let z_0 be an essential singularity of $f : G \rightarrow \mathbb{C}$. For every $\varepsilon > 0$ such that

$$D^*(z_0, \varepsilon) \subseteq G,$$

the image

$$f(D^*(z_0, \varepsilon))$$

is dense in $\mathbb{C}$.

Proof: Suppose not. Then there exist $\varepsilon > 0$, $w \in \mathbb{C}$, and $\delta > 0$ such that

$$|f(z) - w| \geq \delta$$

for all $z \in D^*(z_0, \varepsilon)$. In particular, $f(z) \neq w$ on this punctured disk, so

$$g(z) = \frac{1}{f(z) - w}$$

is holomorphic there. Moreover, $|g(z)| \leq \frac{1}{\delta}$, so by the [Removable Singularity Theorem](#), g has a removable singularity at z_0 .

Let G_0 be the holomorphic extension of g to a disk centered at z_0 . If $G_0(z_0) \neq 0$, then $\frac{1}{G_0}$ is holomorphic in a smaller disk centered at z_0 , so

$$f(z) = w + \frac{1}{G_0}(z)$$

has a removable singularity at z_0 , a contradiction.

If $G_0(z_0) = 0$, then $|g(z)| \rightarrow 0$ as $z \rightarrow z_0$, so

$$|f(z) - w| = \frac{1}{|g(z)|} \rightarrow \infty.$$

By the [Pole Characterization by Growth](#), $f - w$ has a pole at z_0 , and hence f has a pole at z_0 , again a contradiction. Therefore the image of every punctured disk around z_0 is dense in $\mathbb{C}$. $\square$