

Homework 1 Solution

PubH 7440: Introduction to Bayesian Analysis

Spring 2020

1. (Carlin and Louis, Chapter 2, Problem 8)

- a.

The Fisher information for θ being:

$$\mathbb{I}(\theta) = -\mathbb{E}\left[\frac{d^2}{d\theta^2} \log\left(\frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\theta)^2/2\sigma^2}\right)\right] = \frac{1}{\sigma^2}$$

Then $p(\theta) \propto \sqrt{\mathbb{I}(\theta)} = \sqrt{\frac{1}{\sigma^2}} \propto 1$

- b.

The Fisher information for σ being:

$$\mathbb{I}(\sigma) = -\mathbb{E}\left[\frac{d^2}{d\sigma^2} \log\left(\frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\theta)^2/2\sigma^2}\right)\right] = -\mathbb{E}\left[\frac{1}{\sigma^2} - \frac{3(x-\theta)^2}{\sigma^4}\right] = \frac{2}{\sigma^2}$$

Then $p(\sigma) \propto \sqrt{\mathbb{I}(\sigma)} = \sqrt{\frac{2}{\sigma^2}} \propto \frac{1}{\sigma}$

2. (Gelman, Chapter 2, Problem 5)

- a.

$$\begin{aligned}\Pr(y = k) &= \int_0^1 \Pr(y = k|\theta) d\theta \\ &= \int_0^1 \mathbf{C}_n^k \theta^k (1-\theta)^{n-k} d\theta \\ &= \mathbf{C}_n^k \frac{\Gamma(k+1)\Gamma(n-k+1)}{\Gamma(n+2)} \\ &= \frac{1}{n+1}\end{aligned}$$

Where the third equal sign uses the property of Beta distribution and the fourth equal sign comes from the fact that $\Gamma(x) = (x-1)!$

- b.

The posterior distribution $\Pr(\theta|y) \propto \theta^y (1-\theta)^{(n-y)} \theta^{\alpha-1} (1-\theta)^{\beta-1} = \theta^{(y+\alpha-1)} (1-\theta)^{n+\beta-y-1}$, hence $\theta|y \sim \mathbf{Beta}(y+\alpha, n+\beta-y)$, and the posterior mean is $\frac{y+\alpha}{\alpha+n-\beta}$

Suppose there exists λ such that $\frac{y+\alpha}{\alpha+n-\beta} = \lambda \frac{\alpha}{\alpha+\beta} + (1-\lambda) \frac{y}{n}$, solve it for λ , $\lambda = \frac{\alpha+\beta}{\alpha+\beta+n}$. Since α, β, n are all positive, $\lambda \in (0, 1)$, i.e. the posterior mean is a weighted average of the prior mean and the observed relative frequency of heads, this yields the required inequality.

- c.

Through the discussion in a., when the prior is uniform, the posterior distribution is **Beta**($y+1, n-y+1$), hence the posterior variance of θ is $\frac{(y+1)(n-y+1)}{(n+2)^2(n+3)}$, while the prior variance is $\frac{1}{12}$.

$$\begin{aligned} \frac{(y+1)(n-y+1)}{(n+2)^2(n+3)} &\leq \frac{\frac{1}{4}(y+1+n-y+1)^2}{(n+2)^2(n+3)} \\ &= \frac{1}{4(n+3)} \\ &\leq \frac{1}{12} \end{aligned}$$

Where the first inequality comes from the fact: $(a+b)^2 \geq 4ab$, and the second inequality comes from the condition that $n \geq 1$

3. (Gelman, Chapter 2, Problem 8)

- a. From the lecture that for n i.i.d. observations $\mathbf{y} = (y_1, \dots, y_n)$ from normal distribution $N(\theta, \sigma^2)$, and the prior $\pi(\theta) = N(\mu, \tau^2)$, we have:

$$p(\theta|\mathbf{y}) = N\left(\frac{\frac{\sigma^2}{n}\mu + \tau^2\bar{y}}{\frac{\sigma^2}{n} + \tau^2}, \frac{\sigma^2\tau^2}{\sigma^2 + n\tau^2}\right)$$

Hence the posterior distribution for θ is:

$$N\left(\frac{\frac{20^2}{n}180 + 40^2 \times 150}{\frac{20^2}{n} + 40^2}, \frac{20^2 \times 40^2}{20^2 + n \times 40^2}\right)$$

- b.

A normal likelihood with normal prior will result in a normal predictive distribution. The mean and variance can be found by iterated expectation:

$$E[\tilde{y}|\mathbf{y}] = E[E[\tilde{y}|\theta, \mathbf{y}]|\mathbf{y}] = E[\theta|\mathbf{y}] = \frac{\frac{20^2}{n}180 + 40^2 \times 150}{\frac{20^2}{n} + 40^2}$$

$$Var[\tilde{y}|\mathbf{y}] = E[Var[\tilde{y}|\theta, \mathbf{y}]] + Var[E[\tilde{y}|\theta, \mathbf{y}]] = E[20^2|\mathbf{y}] + Var[\theta|\mathbf{y}] = 20^2 + \frac{20^2 \times 40^2}{20^2 + 40^2 n}$$

- c.

Example R code:

```
post_mean <- function(n){
  #function calculating posterior mean
  #same for posterior predictive mean
  (20^2 * 180/n + 40^2 * 150)/(20^2/n + 40^2)
}
```

```

post_var <- function(n){
  #calculating posterior variance
  (20^2 * 40^2 / n) / (20^2 / n + 40^2)
}

pred_var <- function(n){
  #calculating posterior predictive variance
  20^2 + (40^2 * 20^2 / n) / (20^2 / n + 40^2)
}

#### credible interval ####
round(qnorm(c(0.025, .975), mean = post_mean(10), sd = sqrt(post_var(10))),1)

## [1] 138.5 163.0

#### prediction interval ####
round(qnorm(c(0.025, .975), mean = post_mean(10), sd = sqrt(pred_var(10))),1)

## [1] 109.7 191.8

• d.

####credible interval###
round(qnorm(c(0.025, .975), mean = post_mean(100), sd = sqrt(post_var(100))),1)

## [1] 146.2 154.0

####prediction interval
round(qnorm(c(0.025, .975), mean = post_mean(100), sd = sqrt(pred_var(100))),1)

## [1] 110.7 189.5

```

4. (Gelman, Chapter 2, Problem 9)

- a.

From the property of Beta distribution:

$$E(\theta) = \frac{\alpha}{\alpha + \beta} = 0.6$$

$$Var(\theta) = \frac{\alpha\beta}{(\alpha + \beta)^2(\alpha + \beta + 1)} = 0.3^2$$

solve it for α, β : $\alpha = 1$ and $\beta = \frac{2}{3}$

- b.

$$p(\theta|\mathbf{y}) \propto \theta^{650}(1 - \theta)^{350}\theta^{\alpha-1}(1 - \theta)^{\beta-1}$$

which means $\theta|\mathbf{y} \sim \text{Beta}(650 + \alpha, 350 + \beta)$. Hence $E(\theta|\mathbf{y}) = \frac{650+\alpha}{1000+\alpha+\beta} = 0.650$, $Var(\theta|\mathbf{y}) = \frac{(650+\alpha)(350+\beta)}{(1000+\alpha+\beta)^2(1000+\alpha+\beta+1)} = 0.000227$

Example R code for plots:

```

p = seq(0,1, length=1000)
plot(p, dbeta(p, 651, 350.667), ylab="density", type="l")

```

- c.

Example R code for plots:

```
p = seq(0.5,0.7, length=100)
plot(p, dbeta(p, 650, 350), ylab="density", type="l", col = 1)
lines(p, dbeta(p, 651, 350.667), col = 2)
lines(p, dbeta(p, 680, 370), col = 3)
legend(0.5, 25, legend = c("Non-informative", "Beta(1,2/3)", "Beta(30,20)"), lty=c(1,1,1), col=c(1,2,3))
```

5. (Gelman, Chapter 2, Problem 14)

- b.

For $k \leq n$, denote $\mathbf{y}_k = (y_1, \dots, y_k)$, $\theta_k = \frac{\frac{\sigma^2}{k}\mu + \tau^2\bar{\mathbf{y}}_k}{\frac{\sigma^2}{k} + \tau^2}$, and $\sigma_k^2 = \frac{\sigma^2\tau^2}{\sigma^2 + k\tau^2}$

For the case of \mathbf{y}_1 , it is already shown in the lecture notes. Using mathematical induction, suppose the formula holds for all \mathbf{y}_k , $k \leq m-1$, then

$$\begin{aligned} p(\theta|\mathbf{y}_m) &\propto p(y_m|\theta)p(\theta|\mathbf{y}_{m-1}) \\ &\propto \exp\left(-\frac{(y_m - \theta)^2}{2\sigma^2}\right)\exp\left(-\frac{(\theta - \theta_{m-1})^2}{2\sigma_{m-1}^2}\right) \\ &\propto \exp\left(-\frac{\theta^2 - 2\frac{\sigma_{m-1}^2 y_m + \sigma^2 \theta_{m-1}}{\sigma_{m-1}^2 + \sigma^2}\theta}{2\frac{\sigma^2 \sigma_{m-1}^2}{\sigma_{m-1}^2 + \sigma^2}}\right) \\ &\propto \exp\left(-\frac{(\theta - \theta_m)^2}{2\sigma_m^2}\right) \end{aligned}$$

Hence $p(\theta|\mathbf{y}_n) \sim N(\theta_n, \sigma_n^2)$