strengthOfSchedule — how hard was each team's schedule?

Theory: some teams face tougher opponents than others, depending on who they play and when.

Regular season only (byes excluded):

1. **Score distributions:** points against, points for and point differential per game, one panel per team.
2. **Points against by week:** when each team ran into big scores.
3. **Opponent strength vs. points faced:** separates *facing good teams* from *facing teams on a hot week*.
4. **Points scored each week:** every team's score against that week's median.
5. **Regressions:** how strength-of-schedule metrics relate to wins, this season.
6. **Pooled regressions:** the same models on full 2022–2025 seasons (40 team-seasons).

Points against measures how well your opponent played *that week*. Opponent strength (their average in their other games) measures how good they are in general. The gap between them is luck.

Chart style

Load the season

```
The Corrupt League 2026: 7 regular-season weeks, league average 119.3
```

Strength of schedule table

- `avg_pa` : average points scored against the team.
- `opp_strength` : how good its opponents are. For each game, the opponent's average score in *their other games*, so the game against this team doesn't count. Averaged over the team's schedule.
- `luck` : `avg_pa − opp_strength` . Positive means opponents played *above* their usual level against this team (unlucky). Negative means they played below it (lucky).


team	games	wins	losses	avg_pf	avg_pa	med_pa	sd_pa	opp_strength	luck
Will Power	7	1	6	117.4	138.3	136.4	20.6	121.6	16.7
Chubland	7	3	4	128.8	137.4	143.1	35.0	113.9	23.5
Benevolent Commish	7	3	4	127.9	122.5	130.5	21.1	117.7	4.8
Chase You to the Playoffs	7	2	5	117.5	122.0	115.0	16.5	119.7	2.3
On a Train Back to Boston	7	3	4	100.8	118.1	117.0	7.6	123.6	-5.6
I didnt hear no bell	7	2	5	103.4	116.1	120.1	17.6	127.0	-10.9
The Williams Family	7	5	2	124.8	111.6	107.6	13.9	112.3	-0.7
The Germiphobes	7	5	2	123.5	111.2	119.5	31.0	116.5	-5.3
To Infinity and Bijan	7	5	2	112.9	108.6	103.4	29.8	120.0	-11.3
#EATITFROMTHEBACK	7	6	1	135.8	106.9	107.4	14.3	117.5	-10.7

1. Score distributions per team

One panel per team, for three per-game measures: points against, points for, and point differential (points for minus points against). Bars count games in 10-point bins. The solid line is the team's average and the shaded band is ±1 standard deviation around it. The dashed line is the league reference.


Points against

Hardest schedule first.




Points for

Highest-scoring team first. The league average is the same number as for points against: every team's score is its opponent's points against.




Point differential

Points for minus points against, per game. Positive means the team won that game, so blue bars are wins and red bars are losses. Best average margin first. A wide spread means a mix of blowouts and close games.



2. Points against by week

Same team order. Darker means the opponent scored more that week. This shows whether a tough schedule was spread out or concentrated in a few weeks. The table below the chart has the exact values.




week	1	2	3	4	5	6	7
team							
Will Power	166.4	127.6	106.4	148.3	157.2	136.4	125.4
Chubland	159.8	143.1	151.9	188.6	125.9	80.7	111.9
Benevolent Commish	136.2	128.3	130.5	81.3	107.6	142.1	131.8
Chase You to the Playoffs	136.6	115.0	104.8	107.2	137.0	109.9	143.8
On a Train Back to Boston	125.3	108.3	110.0	116.1	129.1	120.7	117.0
I didnt hear no bell	120.1	96.9	122.4	131.1	118.6	87.5	135.9
The Williams Family	113.3	136.9	95.7	121.9	104.3	101.8	107.6
The Germiphobes	128.9	86.4	129.8	119.5	151.7	103.3	58.9
To Infinity and Bijan	95.5	116.5	82.1	103.4	167.9	80.3	114.7
#EATITFROMTHEBACK	110.6	99.7	81.0	123.6	121.1	104.7	107.4

3. Opponent strength vs. points actually faced


Light dot: how good the team's opponents are on average (`opp_strength`). Dark dot: what they actually scored against this team (`avg_pa`). Rows run from hardest opponents down.

- Dark dot right of the light dot: opponents played above their usual level (unlucky).
- Dark dot left of it: opponents played below their usual level (lucky).



4. Points scored each week, against the median

One panel per week. Each dot is a team's score, highest at the top. The vertical line is that week's median. Blue means the team scored above the median, red means below. Beating the median is roughly "would have beaten half the league that week".



Each team's score minus that week's median (positive = above), plus how many weeks it finished above the median. Sorted by weeks above the median.

team	wk_1	wk_2	wk_3	wk_4	wk_5	wk_6	wk_7	weeks_above	avg_vs_median
#EATITFROMTHEBACK	9.5	0.8	14.1	67.9	-8.9	32.4	16.0	6	18.8
Chubland	-13.8	11.9	-3.4	2.9	40.4	16.7	28.0	5	11.8
The Williams Family	32.7	-7.4	-1.8	10.4	24.2	-23.7	20.1	4	7.8
The Germiphobes	-1.8	-18.8	22.3	27.7	-23.1	38.1	1.1	4	6.5
Benevolent Commish	39.3	-0.8	21.6	-4.6	29.8	-0.7	-8.5	3	10.9
Chase You to the Playoffs	-16.5	12.6	43.6	-17.3	1.6	-16.5	-3.9	3	0.5
Will Power	9.1	27.4	-12.5	-1.2	-19.9	0.7	-1.1	3	0.3
On a Train Back to Boston	1.8	21.1	-26.1	-39.4	9.5	-23.3	-56.9	3	-16.2
To Infinity and Bijan	-7.0	-16.0	1.8	-13.5	-1.6	-2.2	9.6	2	-4.1
I didnt hear no bell	-31.6	-29.3	-27.3	1.2	-6.3	5.9	-8.2	2	-13.7

5. Do strength-of-schedule metrics predict wins?

One data point per team. Each model regresses **wins** on one or more metrics:


Model	Question
`wins ~ avg_pa`	Do teams that face more points win less?
`wins ~ opp_strength`	Do teams that face better opponents win less?
`wins ~ luck`	Do teams whose opponents overperform against them win less?
`wins ~ avg_pf`	Baseline: how much does a team's *own* scoring explain?
`wins ~ avg_pf + avg_pa`	How much does schedule add once scoring is accounted for?

How to read the output: **coef** is the change in wins per 1 extra point per game. **P>|t|** below 0.05 is the usual bar for "unlikely to be chance". **R-squared** is the share of the variation in wins the model explains. With only one season of teams (n = 10), expect wide confidence intervals. A slope can look large and still not be distinguishable from zero.

model	formula	term	coef	ci_low	ci_high	p_value	r_squared	adj_r_squared	n
Points against	wins ~ avg_pa	avg_pa	-0.115	-0.192	-0.038	0.009	0.597	0.547	10
Opponent strength	wins ~ opp_strength	opp_strength	-0.200	-0.459	0.060	0.114	0.283	0.193	10
Luck	wins ~ luck	luck	-0.073	-0.169	0.023	0.119	0.275	0.185	10
Points for (baseline)	wins ~ avg_pf	avg_pf	0.069	-0.037	0.175	0.173	0.219	0.121	10
Points for + points against	wins ~ avg_pf + avg_pa	avg_pa	-0.116	-0.171	-0.061	0.002	0.829	0.780	10

Regression plots

One panel per single-metric model. The line is the fitted regression and the shaded band is its 95% confidence interval. Points against, opponent strength and luck are the schedule metrics. Points for is the baseline for comparison.



Full regression output

```
                            Points against
==============================================================================
Dep. Variable:                   wins   R-squared:                       0.597
Model:                            OLS   Adj. R-squared:                  0.547
Method:                 Least Squares   F-statistic:                     11.86
Date:                Sat, 31 Oct 2026   Prob (F-statistic):            0.00877
Time:                        09:30:15   Log-Likelihood:                -14.123
No. Observations:                  10   AIC:                             32.25
Df Residuals:                       8   BIC:                             32.85
Df Model:                           1
Covariance Type:            nonrobust
==============================================================================
                 coef    std err          t      P>|t|      [0.025      0.975]
------------------------------------------------------------------------------
Intercept     17.1939      3.992      4.307      0.003       7.989      26.399
avg_pa        -0.1148      0.033     -3.444      0.009      -0.192      -0.038
==============================================================================
Omnibus:                        0.269   Durbin-Watson:                   1.660
Prob(Omnibus):                  0.874   Jarque-Bera (JB):                0.356
Skew:                          -0.289   Prob(JB):                        0.837
Kurtosis:                       2.278   Cond. No.                     1.36e+03
==============================================================================

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 1.36e+03. This might indicate that there are
strong multicollinearity or other numerical problems.
```

Opponent strength

Dep. Variable:	wins	R-squared:	0.283
Model:	OLS	Adj. R-squared:	0.193
Method:	Least Squares	F-statistic:	3.150
Date:	Sat, 31 Oct 2026	Prob (F-statistic):	0.114
Time:	09:30:15	Log-Likelihood:	-17.010
No. Observations:	10	AIC:	38.02
Df Residuals:	8	BIC:	38.62
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P>|t|	[0.025	0.975]
Intercept	27.2690	13.401	2.035	0.076	-3.633	58.171
opp_strength	-0.1998	0.113	-1.775	0.114	-0.459	0.060

Omnibus:	0.715	Durbin-Watson:	0.336
Prob(Omnibus):	0.699	Jarque-Bera (JB):	0.569
Skew:	0.116	Prob(JB):	0.753
Kurtosis:	1.855	Cond. No.	3.40e+03

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 3.4e+03. This might indicate that there are
strong multicollinearity or other numerical problems.

Luck

Dep. Variable:	wins	R-squared:	0.275
Model:	OLS	Adj. R-squared:	0.185
Method:	Least Squares	F-statistic:	3.039
Date:	Sat, 31 Oct 2026	Prob (F-statistic):	0.119
Time:	09:30:15	Log-Likelihood:	-17.060
No. Observations:	10	AIC:	38.12
Df Residuals:	8	BIC:	38.72
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P>|t|	[0.025	0.975]
Intercept	3.5209	0.471	7.471	0.000	2.434	4.608
luck	-0.0727	0.042	-1.743	0.119	-0.169	0.023

Omnibus:	1.935	Durbin-Watson:	1.520
Prob(Omnibus):	0.380	Jarque-Bera (JB):	0.902
Skew:	-0.285	Prob(JB):	0.637
Kurtosis:	1.644	Cond. No.	11.3

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Points for (baseline)

Dep. Variable:	wins	R-squared:	0.219
Model:	OLS	Adj. R-squared:	0.121
Method:	Least Squares	F-statistic:	2.241
Date:	Sat, 31 Oct 2026	Prob (F-statistic):	0.173
Time:	09:30:15	Log-Likelihood:	-17.435
No. Observations:	10	AIC:	38.87
Df Residuals:	8	BIC:	39.47
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P>|t|	[0.025	0.975]
Intercept	-4.7181	5.511	-0.856	0.417	-17.426	7.990
avg_pf	0.0689	0.046	1.497	0.173	-0.037	0.175

Omnibus:	1.842	Durbin-Watson:	0.562
Prob(Omnibus):	0.398	Jarque-Bera (JB):	0.841
Skew:	-0.202	Prob(JB):	0.657
Kurtosis:	1.638	Cond. No.	1.35e+03

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 1.35e+03. This might indicate that there are
strong multicollinearity or other numerical problems.

```
                        Points for + points against
==============================================================================
Dep. Variable:                   wins   R-squared:                       0.829
Model:                            OLS   Adj. R-squared:                  0.780
Method:                 Least Squares   F-statistic:                     16.99
Date:                Sat, 31 Oct 2026   Prob (F-statistic):           0.00206
Time:                        09:30:15   Log-Likelihood:                -9.8346
No. Observations:                  10   AIC:                             25.67
Df Residuals:                       7   BIC:                             26.58
Df Model:                           2
Covariance Type:            nonrobust
==============================================================================
                 coef    std err          t      P>|t|      [0.025      0.975]
------------------------------------------------------------------------------
Intercept      8.8835      3.872      2.295      0.055      -0.271      18.038
avg_pf         0.0709      0.023      3.083      0.018       0.017       0.125
avg_pa        -0.1161      0.023     -5.001      0.002      -0.171      -0.061
==============================================================================
Omnibus:                        1.251   Durbin-Watson:                   2.590
Prob(Omnibus):                  0.535   Jarque-Bera (JB):                0.716
Skew:                          -0.179   Prob(JB):                        0.699
Kurtosis:                       1.739   Cond. No.                     2.68e+03
==============================================================================

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 2.68e+03. This might indicate that there are
strong multicollinearity or other numerical problems.
```

Reading the combined model

`wins ~ avg_pf + avg_pa` asks: for two teams that score the same, how many wins does facing an extra point per game cost? If `avg_pa` still has a clear negative coefficient here, schedule matters beyond a team's own scoring. If its p-value is large, any schedule effect is too small to see in this sample.

"The condition number is large" note in the outputs comes from the predictors sitting around 100–140 rather than near zero, which inflates the intercept's scale. It isn't a sign that `avg_pf` and `avg_pa` are too correlated to separate. The coefficients and p-values are fine.

More data helps most. Section 6 pools four full seasons, giving 40 team-seasons instead of 10.

6. Pooled seasons: 2022–2025

The same analysis on four complete seasons. Each **team-season** is one data point: 10 teams × 4 seasons = 40 rows, each with a full 14-game regular season. Strength-of-schedule metrics are computed within each season, so `opp_strength` compares a team only with that season's opponents.

The league format is the same in every season: 10 teams, 14 regular-season weeks, the same starting lineup. The only change is the bench, which went from 7 spots to 6 in 2025. That doesn't affect scoring.

Loading takes a little while (four full seasons of box scores).

```
        teams  games_each  total_wins  avg_pf
season
2022       10          14          70   115.6
2023       10          14          70   117.7
2024       10          14          70   121.0
2025       10          14          70   118.7
```

	team	games	wins	losses	avg_pf	avg_pa	med_pa	sd_pa	opp_strength	luck	season	owner
0	Chubb Chubb	14	7	7	119.4	120.9	116.1	28.4	113.3	7.6	2022	Samuel Stein
1	Team Handsome	14	8	6	120.1	119.8	117.3	17.9	115.4	4.3	2022	Ryan O'Connor
2	Corrupt league	14	3	11	106.2	119.5	119.7	18.3	117.6	1.9	2022	ben englander
3	GRUDENS GONE!!!!!!	14	3	11	104.1	119.2	121.4	20.7	117.4	1.8	2022	Richard Kinches
4	Team Gesin	14	7	7	113.0	117.6	120.9	22.9	115.2	2.5	2022	Mo Gesin
5	We Need a Minyan Man	14	10	4	125.6	115.0	110.9	18.8	112.8	2.2	2022	Sam Levitt
6	Bishop Sycamore	14	8	6	111.3	112.0	119.4	25.4	117.0	-5.0	2022	Aaron Fandel
7	Team Birnbach	14	8	6	113.8	111.7	113.2	20.8	116.7	-5.0	2022	Jake Birnbach
8	I Like Ike	14	9	5	130.8	110.5	108.4	21.5	114.5	-4.0	2022	Isaac Leibowitz
9	Team Warren	14	7	7	112.2	110.2	105.2	26.5	116.1	-5.9	2022	Gavriel Warren
10	Mossterbation	14	7	7	124.3	132.5	129.8	22.5	114.8	17.8	2023	Ryan O'Connor
11	Tuck Em Diggers	14	6	8	119.3	124.9	116.8	25.6	116.6	8.3	2023	Jake Birnbach
12	Etienne Word	14	7	7	118.3	122.8	121.1	24.1	117.1	5.7	2023	Isaac Leibowitz
13	BH	14	7	7	115.7	119.1	123.3	23.7	118.3	0.8	2023	Richard Kinches
14	I Have a Team	14	9	5	122.5	116.8	115.0	20.9	117.6	-0.7	2023	Mo Gesin
15	Team Warren	14	5	9	109.5	114.9	116.4	15.4	118.6	-3.7	2023	Gavriel Warren
16	🤡🤡🤡	14	7	7	121.9	113.2	114.8	22.7	117.5	-4.4	2023	ben englander
17	Bishop Sycamore	14	6	8	105.2	112.7	111.7	20.3	119.4	-6.7	2023	Aaron Fandel
18	We Need a Minyan Man	14	7	7	119.3	110.3	110.9	23.1	118.9	-8.6	2023	Sam Levitt
19	Stroud Boys	14	9	5	120.8	109.6	106.7	14.6	118.0	-8.3	2023	Samuel Stein
20	Bishop Sycamore	14	3	11	113.2	129.0	124.6	14.6	120.1	9.0	2024	Aaron Fandel
21	🤡🤡🤡	14	4	10	110.2	125.1	124.9	24.7	121.8	3.3	2024	ben englander
22	BH	14	8	6	123.6	122.4	117.2	26.9	122.4	0.0	2024	Richard Kinches
23	Team Warren	14	7	7	119.3	121.5	118.6	17.3	121.9	-0.4	2024	Gavriel Warren
24	I Have a Team	14	6	8	117.7	120.7	122.6	26.1	121.2	-0.5	2024	Moshe Gesin
25	I Love Saquon Barkley	14	7	7	124.8	120.5	119.8	21.6	120.5	-0.0	2024	Isaac Leibowitz
26	Skibidi rizz	14	9	5	130.7	120.4	122.7	20.2	118.8	1.6	2024	Ryan O'Connor
27	Big Black Machers	14	6	8	119.2	119.1	118.2	13.8	122.1	-3.0	2024	Jake Birnbach
28	We Need a Minyan Man	14	10	4	130.2	116.0	115.3	16.8	120.8	-4.8	2024	Sam Levitt
29	Big Josh Fan Club	14	10	4	120.6	114.8	119.2	22.7	120.9	-6.2	2024	Samuel Stein
30	Start Diggsin in ur Kupp twin	14	6	8	115.9	125.6	134.2	30.2	116.9	8.7	2025	Ryan O'Connor
31	They Hit the Second Bower	14	7	7	123.8	124.8	117.7	24.5	116.6	8.2	2025	Aaron Fandel
32	Seeing Is Believing	14	5	9	116.2	123.8	124.1	31.8	120.0	3.8	2025	Samuel Stein
33	Give me 25 Jaylen Warren	14	7	7	116.6	123.3	123.3	19.3	118.8	4.5	2025	Isaac Leibowitz
34	It Hurts When I'm Tuten	14	7	7	114.8	120.1	116.3	25.0	118.8	1.2	2025	Moshe Gesin
35	Corrupt Commish	14	7	7	114.1	119.0	122.1	24.4	118.5	0.6	2025	ben englander
36	I didn't reach for Lamar	14	7	7	118.6	117.8	117.2	17.4	120.2	-2.4	2025	Jake Birnbach
37	Injury Buck	14	6	8	111.1	112.4	112.5	20.4	121.6	-9.2	2025	Richard Kinches
38	We sleep in Maye	14	10	4	137.9	110.4	112.2	22.5	117.5	-7.0	2025	Gavriel Warren
39	CMC You in the Playoffs	14	8	6	118.4	110.3	104.0	26.4	120.1	-9.9	2025	Sam Levitt


Pooled models

The same five models, plus one more. Scoring levels differ from season to season (league average points move around), so the last model adds a **season fixed effect**, `C(season)`, which compares each team only with other teams from the same season.

model	formula	term	coef	ci_low	ci_high	p_value	r_squared	adj_r_squared	n
Points against	wins ~ avg_pa	avg_pa	-0.144	-0.238	-0.050	0.004	0.202	0.181	40
Opponent strength	wins ~ opp_strength	opp_strength	-0.152	-0.391	0.088	0.207	0.042	0.016	40
Luck	wins ~ luck	luck	-0.102	-0.195	-0.009	0.032	0.115	0.092	40
Points for (baseline)	wins ~ avg_pf	avg_pf	0.188	0.132	0.243	0.000	0.550	0.538	40
Points for + points against	wins ~ avg_pf + avg_pa	avg_pa	-0.132	-0.189	-0.076	0.000	0.720	0.704	40
Points for + points against + season	wins ~ avg_pf + avg_pa + C(season)	avg_pa	-0.125	-0.188	-0.061	0.000	0.723	0.683	40

Pooled regression plots

40 points per panel, so points aren't labeled. The table above has every team-season.



Full pooled regression output

Points against (pooled 2022–2025)

Dep. Variable:	wins	R-squared:	0.202
Model:	OLS	Adj. R-squared:	0.181
Method:	Least Squares	F-statistic:	9.633
Date:	Sat, 31 Oct 2026	Prob (F-statistic):	0.00360
Time:	09:30:57	Log-Likelihood:	−75.502
No. Observations:	40	AIC:	155.0
Df Residuals:	38	BIC:	158.4
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P>|t|	[0.025	0.975]
Intercept	24.0487	5.499	4.373	0.000	12.917	35.181
avg_pa	−0.1442	0.046	−3.104	0.004	−0.238	−0.050

Omnibus:	1.915	Durbin–Watson:	1.890
Prob(Omnibus):	0.384	Jarque–Bera (JB):	1.477
Skew:	−0.470	Prob(JB):	0.478
Kurtosis:	2.936	Cond. No.	2.51e+03

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 2.51e+03. This might indicate that there are
strong multicollinearity or other numerical problems.

Opponent strength (pooled 2022–2025)

Dep. Variable:	wins	R-squared:	0.042
Model:	OLS	Adj. R-squared:	0.016
Method:	Least Squares	F-statistic:	1.646
Date:	Sat, 31 Oct 2026	Prob (F-statistic):	0.207
Time:	09:30:57	Log-Likelihood:	−79.173
No. Observations:	40	AIC:	162.3
Df Residuals:	38	BIC:	165.7
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P>|t|	[0.025	0.975]
Intercept	24.9554	13.999	1.783	0.083	−3.385	53.296
opp_strength	−0.1518	0.118	−1.283	0.207	−0.391	0.088

Omnibus:	1.990	Durbin–Watson:	1.610
Prob(Omnibus):	0.370	Jarque–Bera (JB):	1.118
Skew:	−0.374	Prob(JB):	0.572
Kurtosis:	3.333	Cond. No.	5.83e+03

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 5.83e+03. This might indicate that there are
strong multicollinearity or other numerical problems.

Luck (pooled 2022–2025)

Dep. Variable:	wins	R-squared:	0.115
Model:	OLS	Adj. R-squared:	0.092
Method:	Least Squares	F-statistic:	4.951
Date:	Sat, 31 Oct 2026	Prob (F-statistic):	0.0321
Time:	09:30:57	Log-Likelihood:	−77.571
No. Observations:	40	AIC:	159.1
Df Residuals:	38	BIC:	162.5
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P>|t|	[0.025	0.975]
Intercept	6.9952	0.273	25.627	0.000	6.443	7.548
luck	−0.1021	0.046	−2.225	0.032	−0.195	−0.009

Omnibus:	1.531	Durbin–Watson:	1.933
Prob(Omnibus):	0.465	Jarque–Bera (JB):	1.317
Skew:	−0.431	Prob(JB):	0.518
Kurtosis:	2.785	Cond. No.	5.95

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.

Points for (baseline) (pooled 2022–2025)

Dep. Variable:	wins	R-squared:	0.550
Model:	OLS	Adj. R-squared:	0.538
Method:	Least Squares	F-statistic:	46.46
Date:	Sat, 31 Oct 2026	Prob (F-statistic):	4.37e−08
Time:	09:30:57	Log-Likelihood:	−64.048

```
No. Observations:                  40   AIC:                              132.1
Df Residuals:                      38   BIC:                              135.5
Df Model:                           1
Covariance Type:            nonrobust
==============================================================================
                 coef    std err          t      P>|t|      [0.025      0.975]
------------------------------------------------------------------------------
Intercept    -15.1881      3.261     -4.657      0.000     -21.790      -8.586
avg_pf         0.1876      0.028      6.816      0.000       0.132       0.243
==============================================================================
Omnibus:                        0.000   Durbin-Watson:                   1.626
Prob(Omnibus):                  1.000   Jarque-Bera (JB):                0.121
Skew:                          -0.004   Prob(JB):                        0.941
Kurtosis:                       2.730   Cond. No.                     1.98e+03
==============================================================================

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 1.98e+03. This might indicate that there are
strong multicollinearity or other numerical problems.
```

Points for + points against (pooled 2022–2025)

```
==============================================================================
Dep. Variable:                   wins   R-squared:                       0.720
Model:                            OLS   Adj. R-squared:                  0.704
Method:                 Least Squares   F-statistic:                     47.49
Date:                Sat, 31 Oct 2026   Prob (F-statistic):           6.06e-11
Time:                        09:30:57   Log-Likelihood:                -54.587
No. Observations:                  40   AIC:                             115.2
Df Residuals:                      37   BIC:                             120.2
Df Model:                           2
Covariance Type:            nonrobust
==============================================================================
                 coef    std err          t      P>|t|      [0.025      0.975]
------------------------------------------------------------------------------
Intercept      1.0844      4.317      0.251      0.803      -7.663       9.832
avg_pf         0.1822      0.022      8.263      0.000       0.138       0.227
avg_pa        -0.1322      0.028     -4.731      0.000      -0.189      -0.076
==============================================================================
Omnibus:                        2.357   Durbin-Watson:                   1.735
Prob(Omnibus):                  0.308   Jarque-Bera (JB):                1.381
Skew:                          -0.128   Prob(JB):                        0.501
Kurtosis:                       2.126   Cond. No.                     4.64e+03
==============================================================================

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 4.64e+03. This might indicate that there are
strong multicollinearity or other numerical problems.
```

Points for + points against + season (pooled 2022–2025)

```
==============================================================================
Dep. Variable:                   wins   R-squared:                       0.723
Model:                            OLS   Adj. R-squared:                  0.683
Method:                 Least Squares   F-statistic:                     17.77
Date:                Sat, 31 Oct 2026   Prob (F-statistic):           1.22e-08
Time:                        09:30:57   Log-Likelihood:                -54.329
No. Observations:                  40   AIC:                             120.7
Df Residuals:                      34   BIC:                             130.8
Df Model:                           5
Covariance Type:            nonrobust
===============================================================================================
                              coef    std err          t      P>|t|      [0.025      0.975]
-----------------------------------------------------------------------------------------------
Intercept                  -0.2411      4.903     -0.049      0.961     -10.205       9.723
C(season)[T.2023]          -0.1275      0.465     -0.274      0.785      -1.072       0.817
C(season)[T.2024]          -0.3322      0.509     -0.653      0.518      -1.366       0.702
C(season)[T.2025]          -0.1943      0.475     -0.409      0.685      -1.160       0.771
avg_pf                      0.1871      0.024      7.790      0.000       0.138       0.236
avg_pa                     -0.1245      0.031     -3.991      0.000      -0.188      -0.061
==============================================================================
Omnibus:                        0.595   Durbin-Watson:                   1.783
Prob(Omnibus):                  0.743   Jarque-Bera (JB):                0.690
Skew:                          -0.126   Prob(JB):                        0.708
Kurtosis:                       2.408   Cond. No.                     5.09e+03
==============================================================================

Notes:
[1] Standard Errors assume that the covariance matrix of the errors is correctly specified.
[2] The condition number is large, 5.09e+03. This might indicate that there are
strong multicollinearity or other numerical problems.
```

This season vs. pooled

Each model's key coefficient, R^2 and p-value, from {SEASON} alone (n = 10) and from the pooled seasons (n = 40). Agreement in sign and size suggests a real pattern rather than one season's noise. A p-value that drops sharply with more data means the effect was real but the single season was too small to show it.

		2026			2022–2025		
	term	coef	p_value	r_squared	coef	p_value	r_squared
model							
Points against	avg_pa	-0.115	0.009	0.597	-0.144	0.004	0.202
Opponent strength	opp_strength	-0.200	0.114	0.283	-0.152	0.207	0.042
Luck	luck	-0.073	0.119	0.275	-0.102	0.032	0.115
Points for (baseline)	avg_pf	0.069	0.173	0.219	0.188	0.000	0.550
Points for + points against	avg_pa	-0.116	0.002	0.829	-0.132	0.000	0.720
Points for + points against + season	NaN	NaN	NaN	NaN	-0.125	0.000	0.723

Caveats

- **Small samples.** Mid-season there are only a handful of games per team, so one blowout week moves a team's average a lot. Re-run as the season goes on, or set `SEASON` to a finished year (2019 onward).
- **`opp_strength` is an average, not a ranking of matchups.** A team can face strong opponents and still catch them on off weeks. That's what the luck column captures.
- **Correlation with record.** To test whether schedule actually decided standings, compare `luck` (or `avg_pa`) with wins in the table above.