

## PRETALK 1: CHOW GROUPS

**Exercise 1.** Suppose  $X$  is a finite type separated  $k$ -scheme, with closed subscheme  $\iota : Z \hookrightarrow X$ . Show that there is an exact sequence

$$\mathrm{CH}_i(Z) \xrightarrow{\iota_*} \mathrm{CH}_i(X) \xrightarrow{j^*} \mathrm{CH}_i(X \setminus Z) \rightarrow 0,$$

where  $j : X \setminus Z \hookrightarrow X$ .

**Exercise 2.** Consider  $X = V(x_1, x_2) \cup V(x_3, x_4) \subseteq \mathbb{P}^4$ . Show that  $[X] \cdot h^2$  is not equal to  $[X \cap V(x_1 - x_3, x_2 - x_4)]$  (despite the fact that the latter has the appropriate dimension).

**Exercise 3.** Say a finite type separated  $k$ -scheme  $X$  has the Chow–Künneth generation property (CKgP) if for all  $Y$ , the Künneth map

$$\mathrm{CH}(X) \otimes \mathrm{CH}(Y) \rightarrow \mathrm{CH}(X \times Y)$$

is an isomorphism. Show that  $\mathbb{P}^n$  has the CKgP.

**Exercise 4.** Let  $E$  be a genus one curve. Show that the image of  $\mathrm{CH}(E) \otimes \mathrm{CH}(E) \rightarrow \mathrm{CH}(E \times E)$  does not contain the class of the diagonal (hence  $E$  does not have the CKgP). Hint: Use the adjunction formula.

**Exercise 5.** Let  $X \subseteq \mathbb{P}^n$  be a subvariety. Show that  $[X] \neq 0$ . Hint: First show it for a point  $p \in \mathbb{P}^n$ .

**Exercise 6.** Let  $X$  be an equidimensional finite type separated  $k$ -scheme of positive dimension. Show that given  $\alpha, \beta \in Z^0(X)$ , there exists  $\alpha' \in Z^0(X)$  such that  $\alpha' \sim \alpha$  and  $|\alpha'| \cap |\beta| = \emptyset$ .