



## Lessons from a Decade of IPS and LLC

Joakim Westerlund & Jörg Breitung

To cite this article: Joakim Westerlund & Jörg Breitung (2013) Lessons from a Decade of IPS and LLC, *Econometric Reviews*, 32:5-6, 547-591, DOI: [10.1080/07474938.2013.741023](https://doi.org/10.1080/07474938.2013.741023)

To link to this article: <http://dx.doi.org/10.1080/07474938.2013.741023>



Accepted author version posted online: 25 Oct 2012.  
Published online: 25 Oct 2012.



Submit your article to this journal [↗](#)



Article views: 373



View related articles [↗](#)



Citing articles: 36 View citing articles [↗](#)

## LESSONS FROM A DECADE OF IPS AND LLC

Joakim Westerlund<sup>1</sup> and Jörg Breitung<sup>2</sup>

<sup>1</sup>*Department of Economics, University of Gothenburg, Gothenburg, Sweden*

<sup>2</sup>*Department of Economics, University of Bonn, Bonn, Germany*

□ *This paper points to some of the facts that have emerged from 20 years of research into the analysis of unit roots in panel data, an area that has been heavily influenced by two studies, IPS (Im, Pesaran, and Shin, 2003) and LLC (Levin et al., 2002). Some of these facts are known, others are not. But they all have in common that, if ignored, the effects can be very serious. This is demonstrated using both simulations and theoretical arguments.*

**Keywords** Cross-section dependence; Multidimensional limits; Non-stationary panel data; Unit root tests.

**JEL Classification** C13; C33.

### 1. INTRODUCTION

Starting with the working paper versions of Quah (1994) and Breitung and Meyer (1994) that were available already in 1989, the literature concerned with the analysis of unit roots in panel data covers more than 20 years. While during the first decade the topic was rather peripheral, it has by now become a very active research area, see Breitung and Pesaran (2008) for a recent survey of the literature. Today panel unit root tests are standard econometric tools within most fields of empirical economics, especially in macroeconomics and financial economics, and some are now available in commercial software packages such as EViews and STATA.

In the beginning when the panel unit root literature was still in its infancy econometricians tended to view the extension of the conventional unit root analysis to panel data as a rather straightforward and less exciting exercise. However, it has since then become clear that this is not the

Address correspondence to Joakim Westerlund, Department of Economics, University of Gothenburg, P. O. Box 640, SE 405 30 Gothenburg, Sweden; E-mail: joakim.westerlund@economics.gu.se

case. Indeed, subsequent work has revealed a number of fundamental differences in the way statistical inference with non-stationary data is performed and it seems fair to say that panel unit root analysis has established itself as a legitimate branch of literature.

Two of the most influential contributions to this field of research are those of Levin, Lin, and Chu (2002) (LLC) and Im, Peseran, and Shin, (2003) (IPS), who were among the first to develop so-called first-generation tests that are appropriate when the cross-sectional units are independent of each other. These works have not only been important for applied researchers, to which vast amounts of panel data are now available, but also for econometricians concerned with the development of new tests and methods for non-stationary panels. In fact, among the large number of studies that has recently been published within this field there are only a few, if any, that do not build, either directly or indirectly, on the works of LLC and IPS.

To the best of our knowledge, however, there is still no study that investigates basic principles, or facts, upon which these tests, and therefore also the whole panel unit root literature, are built. There are of course literature reviews such as those of Baltagi and Kao (2000), Bowman (1999), Choi (2006), Bai and Kao (2006), and Breitung and Pesaran (2008), but the purpose of these is more to review, and not so much to study in detail the underlying facts. Consequently, they do not either emphasize enough the consequences of overlooking these facts. This paper goes back to the basics and shows how they make a difference. It is an in-dept review and theoretical companion to specific research papers.

In general in econometrics and statistics most methods are based on certain established facts that are assumed to be well-known. For example, time series unit root tests, like the one suggested by Dickey and Fuller (1979), henceforth denoted DF, can suffer from poor power if the autoregressive root is local-to-unity, especially in the presence of deterministic constant and trend terms. It also a well-known fact that one way to improve the power of such tests is to perform the detrending regression in a way that is efficient under the local alternative hypothesis, an idea that was first suggested by Elliott et al. (1996).

Panel unit root analysis also has its fair share of facts like this. But in contrast to the conventional time series analysis, which has been around for almost half a century now, these are not very well-understood, and some are just being developed. The purpose of this paper is in part to point to these facts, in part to illustrate, both analytically and numerically, their importance, and in so doing we will make use of the LLC and IPS tests as leading examples. Some of the facts are known, others are not. But they all share the feature that if ignored the effects upon analysis can be dramatic. Proper understanding of these facts, and the lessons that can be drawn from them, is therefore key in any research with non-stationary panel data.

To place our results in the context of the literature on dynamic panel data models in general, note that in this paper we are only concerned with those aspects that are in some sense unique to the unit root testing problem. The facts we analyze can be divided roughly into the following three broad categories: (i) asymptotic properties under the unit root null and stationary alternative hypotheses in absence of nuisance parameters, (ii) the treatment of deterministic components, and (iii) the effect of initial condition, serial correlation and cross-section dependence. These categories have of course been treated before but for different reasons than they are here.

The plan of the paper is the following. Section 2 focuses on the simplest case without any deterministic terms, short-run dynamics or cross-sectional dependence. Although admittedly very restrictive, this setup allows us to focus on some of the most primitive facts of panel unit root testing. In Section 3 we generalize the setup of Section 2 to allow for deterministic constant and trend terms. The analysis reveal that this small change has major implications for the resulting tests. Models with short-run dynamics are considered in Section 4, and in Section 5 we address the problems that arise when the cross-sectional units are no longer independent. Section 6 offers some concluding remarks.

## 2. THE SIMPLEST CASE

Consider the double indexed variable  $y_{it}$ , observable for  $t = 1, \dots, T$  time periods, and  $i = 1, \dots, N$  cross-sectional units. Initially we will assume that  $y_{it}$  has no deterministic part, so that

$$y_{it} = y_{it}^s, \quad (1)$$

where  $y_{it}^s$  is the stochastic part of  $y_{it}$ , which is assumed to evolve according to the following first-order autoregressive (AR) process:

$$y_{it}^s = \rho_i y_{it-1}^s + \varepsilon_{it}, \quad (2)$$

or, equivalently,

$$\Delta y_{it} = (\rho_i - 1)y_{it-1} + \varepsilon_{it} = \alpha_i y_{it-1} + \varepsilon_{it}. \quad (3)$$

In this section we assume that the error  $\varepsilon_{it}$  is mean zero and independent across both  $i$  and  $t$ . To make life even simpler, we assume that the errors are homoscedastic so that  $E(\varepsilon_{it}^2) = \sigma^2$  for all  $i$  and  $t$ . Note that while unduely restrictive for most practical purposes, this data generating process has the advantage of being simple and illustrative.

The null hypothesis of interest is

$$H_0 : \alpha_i = 0 \quad \text{for all } i,$$

which corresponds to a fully non-stationary panel. As for the alternative hypothesis, we will consider two candidates,  $H_{1a}$  and  $H_{1b}$ . The first is specified as

$$H_{1a} : \alpha_i = \alpha < 0 \quad \text{for all } i,$$

and corresponds to a fully stationary panel with the same degree of mean reversion for all units. It is therefore quite restrictive. The second alternative is more relaxed. It reads

$$H_{1b} : \alpha_i < 0 \quad \text{for } i = 1, \dots, N_1 \quad \text{with } \frac{N_1}{N} \rightarrow \delta_1 > 0 \quad \text{as } N_1, N \rightarrow \infty,$$

which corresponds to a mixed panel with  $\delta_1$  being the limiting fraction of stationary units. Note that in this formulation, there are no homogeneity restrictions with regards to the degree of mean reversion. Note also that at this point we make no assumptions concerning the remaining  $N - N_1$  slopes,  $\alpha_{N_1+1}, \dots, \alpha_N$ , which may all be zero, negative or a mixture of both. However, we do require that  $\delta_1 > 0$ , as otherwise the panel would escape stationarity as  $N_1, N \rightarrow \infty$ . The two alternative hypotheses  $H_{1a}$  and  $H_{1b}$  are chosen to match the LLC and IPS tests, respectively.

Before considering these tests, however, it is useful to introduce some notation. In particular, we define  $M = \sum_{i=1}^N M_i$ , where

$$M_i = \begin{pmatrix} M_{11i} & M_{12i} \\ M_{12i} & M_{22i} \end{pmatrix} = \sum_{t=2}^T \begin{pmatrix} (\Delta y_{it})^2 & y_{it-1} \Delta y_{it} \\ y_{it-1} \Delta y_{it} & y_{it-1}^2 \end{pmatrix}$$

is the non-normalized moment matrix of the variables contained in the regression in (3). As  $T \rightarrow \infty$  we have under the null hypothesis

$$\begin{pmatrix} \frac{1}{T} M_{11i} & \frac{1}{T} M_{12i} \\ \frac{1}{T} M_{12i} & \frac{1}{T^2} M_{22i} \end{pmatrix} \Rightarrow \sigma^2 M_i^\circ,$$

where the symbol  $\Rightarrow$  signifies weak convergence and

$$M_i^\circ = \begin{pmatrix} M_{11i}^\circ & M_{12i}^\circ \\ M_{12i}^\circ & M_{22i}^\circ \end{pmatrix} = \begin{pmatrix} 1 & \int_0^1 W_i(s) dW_i(s) \\ \int_0^1 W_i(s) dW_i(s) & \int_0^1 W_i(s)^2 ds \end{pmatrix}.$$

The results reported in this paper are derived using either the joint limit method wherein  $N, T \rightarrow \infty$  simultaneously, or the sequential limit method wherein one of the indices is passed to infinity before the other, see Phillips and Moon (1999). In any case, since the purpose here is more to illustrate rather than to prove, details that are not essential for the understanding of the main point will be omitted. The derivations will therefore not be complete, and readers are referred to the relevant original works for a more detailed treatment.

Having introduced the main notation, we now go on to discuss the IPS and LLC tests. With no serial correlation or heteroskedasticity, and no deterministic constant or trend terms, the Levin and Lin (1992) statistic is given by

$$\tau_{LLC} = \frac{M_{12}}{\hat{\sigma} \sqrt{M_{22}}} = \hat{\alpha} \frac{\sqrt{M_{22}}}{\hat{\sigma}},$$

where  $\hat{\sigma}^2 = \frac{1}{NT}(M_{11} - \hat{\alpha} M_{12})$  with  $\hat{\alpha} = M_{12}/M_{22}$  being the least squares (LS) estimator of  $\alpha$ , whose standard error is given by  $\hat{\sigma}/\sqrt{M_{22}}$ . Note that although in this setting the Levin and Lin (1992) statistic is the same as the LLC statistic that assumes no deterministic component and no short-run dynamics, at times it will be important to keep the distinction, as this similarity is not always going to hold when we go on to discuss more general models.

The IPS test is given by

$$\tau_{IPS} = \frac{\sqrt{N}(\bar{\tau} - E(\tau))}{\sqrt{\text{var}(\tau)}},$$

where  $\bar{\tau} = \frac{1}{N} \sum_{i=1}^N \tau_i$  and  $\tau_i$  is the usual DF test statistic,

$$\tau_i = \frac{M_{12i}}{\hat{\sigma}_i \sqrt{M_{22i}}} = \hat{\alpha}_i \frac{\sqrt{M_{22i}}}{\hat{\sigma}_i}$$

with an obvious definition of  $\hat{\sigma}_i^2$  and  $\hat{\alpha}_i$ . It is well-known that

$$\tau_i \Rightarrow \frac{M_{12i}^\circ}{\sqrt{M_{22i}^\circ}}$$

as  $T \rightarrow \infty$ . The constants  $E(\tau)$  and  $\text{var}(\tau)$  are simply the mean and variance of this limiting distribution. Note that since  $M_i^\circ$  is identically distributed,  $E(\tau)$  and  $\text{var}(\tau)$  do not need to carry an  $i$  index.

**Fact 1: The IPS and LLC Statistics are Standard Normally Distributed as  $N \rightarrow \infty$  even if  $T$  is Fixed**

In order to establish the asymptotic normality of  $\tau_{LLC}$  and  $\tau_{IPS}$  as  $N \rightarrow \infty$  with  $T$  fixed we invoke two of the most important tools of the analysis of non-stationary panel data, the weak law of large numbers and the Lindeberg–Levy central limit theorem in the cross-sectional dimension.

Consider first the LLC statistic, which can be written as

$$\tau_{LLC} = \frac{\frac{1}{T\sqrt{N}}M_{12}}{\hat{\sigma}\sqrt{\frac{1}{NT^2}M_{22}}}.$$

We begin by analyzing the denominator under  $H_0$ , which by the law of large numbers as  $N \rightarrow \infty$  becomes

$$\frac{1}{NT^2}M_{22} \xrightarrow{p} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \frac{1}{T^2} \sum_{t=2}^T E(y_{it-1}^2) = \sigma^2 \frac{1}{T^2} \sum_{t=1}^{T-1} t = \sigma^2 \frac{T-1}{2T},$$

where  $\xrightarrow{p}$  signifies convergence in probability. Similarly,  $\frac{1}{NT}M_{12} \xrightarrow{p} 0$  and  $\frac{1}{NT}M_{11} \xrightarrow{p} \sigma^2$  as  $N \rightarrow \infty$ , from which we deduce that  $\hat{\alpha} \xrightarrow{p} 0$  and  $\hat{\sigma}^2 \xrightarrow{p} \sigma^2$ .

Moreover,

$$\text{var}\left(\frac{1}{T}M_{12i}\right) = \frac{1}{T^2} \sum_{t=2}^T \text{var}(y_{it-1}\Delta y_{it}) = \sigma^2 \frac{1}{T^2} E(M_{22i}) = \sigma^4 \frac{T-1}{2T}.$$

Note that for a fixed  $T$ ,  $\sigma^4(T-1)/2T < \infty$ . In view of this and the assumed independence across  $i$ , we have that by the Lindeberg–Levy central limit theorem,

$$\frac{1}{T\sqrt{N}}M_{12} = \frac{1}{T\sqrt{N}} \sum_{i=1}^N M_{12i} \xrightarrow{d} \sigma^2 \sqrt{\frac{T-1}{2T}} \mathcal{N}(0, 1)$$

as  $N \rightarrow \infty$ , where  $\xrightarrow{d}$  denotes convergence in distribution. Thus, by putting everything together we get

$$\tau_{LLC} = \frac{\frac{1}{T\sqrt{N}}M_{12}}{\hat{\sigma}\sqrt{\frac{1}{T^2N}M_{22}}} \xrightarrow{d} \mathcal{N}(0, 1).$$

Note that this result holds for any  $T$ . Hence, the asymptotic normality of the LLC statistic does not require  $T \rightarrow \infty$ . However, if individual specific parameters relating to for example deterministic terms or short-run dynamics are introduced, then this is no longer true. The reason is

that consistent estimation of these parameters requires  $T \rightarrow \infty$ , see for example Harris and Tzavalis (1999) and LLC.

In a similar manner it can be shown that  $\tau_{IPS}$  is also standard normally distributed as  $N \rightarrow \infty$  with  $T$  held fixed.<sup>1</sup> This is done in IPS. The idea behind the proof is that since  $\tau_i$  is cross-sectionally independent by assumption, the application of the Lindeberg–Levy central limit theorem requires that the variance is finite. Although this is generally far from trivial to show, if we assume that  $\varepsilon_{it}$  is normally distributed, then  $\tau_i^2 = (T - 1)R^2$ , where  $R^2$  is the usual  $R^2$  statistic of the regression in (3). Clearly, since  $R^2 \in [0, 1]$ , the variance exists for finite  $T$  and therefore

$$\tau_{IPS} \xrightarrow{d} \mathcal{N}(0, 1)$$

as  $N \rightarrow \infty$  by the Lindeberg–Levy central limit theorem.<sup>2</sup> Thus, as long as  $N \rightarrow \infty$  normality of these statistics does not require passing  $T \rightarrow \infty$ , a fact that is oftentimes not considered in theoretical work. As a practical matter, this means that the tests can be applied even in situations in which  $N \gg T$ .

The performance under the stationary alternative is the topic of the next section.

## Fact 2: The LLC Test Can Be More Powerful than the IPS Test

It has become standard to treat  $\tau_{LLC}$  as a test against  $H_{1a}$  and  $\tau_{IPS}$  as a test against  $H_{1b}$ . Therefore, since  $H_{1b}$  is less restrictive than  $H_{1a}$ , one might be led to believe that  $\tau_{IPS}$  should dominate  $\tau_{LLC}$  in terms of power, at least under the heterogeneous alternative. However, this is not true.

Consider first the case when the slope coefficient  $\alpha_i$  is fixed under the alternative. If  $H_{1a}$  holds, then we write

$$\frac{1}{\sqrt{NT}} \tau_{LLC} = \alpha \frac{\sqrt{\frac{1}{NT} M_{22}}}{\hat{\sigma}} + (\hat{\alpha} - \alpha) \frac{\sqrt{\frac{1}{NT} M_{22}}}{\hat{\sigma}} = O_p(1) + O_p\left(\frac{1}{\sqrt{NT}}\right) O_p(1),$$

which implies that  $\tau_{LLC} = O_p(\sqrt{NT})$ . Similarly, if  $H_{1b}$  holds, and assuming for simplicity that the last  $N - N_1$  units are non-stationary,

$$\sqrt{\text{var}(\tau)} \tau_{IPS} = \frac{1}{\sqrt{N}} \sum_{i=1}^N (\tau_i - E(\tau))$$

<sup>1</sup>Instead of the Lindeberg condition the Liapounov condition is usually invoked, in which case one has to assume that the individual test statistics  $\tau_i$  have more than two moments.

<sup>2</sup>Strictly speaking, for this result to hold, we require that  $E(\tau)$  and  $\text{var}(\tau)$  are evaluated for a finite  $T$ .



$$\begin{aligned}
&= \sqrt{\frac{N_1}{N}} \frac{1}{\sqrt{N_1}} \sum_{i=1}^{N_1} (\tau_i - E(\tau)) + \sqrt{1 - \frac{N_1}{N}} \frac{1}{\sqrt{N - N_1}} \sum_{i=N_1+1}^N (\tau_i - E(\tau)) \\
&= \sqrt{\delta_1} O_p(\sqrt{NT}) + \sqrt{1 - \delta_1} O_p(1),
\end{aligned}$$

where we have used that under a fixed alternative

$$\frac{1}{\sqrt{T}} E(\tau_i) = \frac{\alpha_i}{\sigma} E\left(\sqrt{\frac{1}{T}} M_{22i}\right) + O_p\left(\frac{1}{\sqrt{T}}\right) \rightarrow \frac{\alpha_i}{\sqrt{1 - \rho_i^2}} \neq E(\tau)$$

as  $T \rightarrow \infty$ , implying

$$\frac{1}{N_1} \sum_{i=1}^{N_1} (\tau_i - E(\tau)) \xrightarrow{p} E(\tau_i) - E(\tau) = O_p(\sqrt{T})$$

so that  $\frac{1}{\sqrt{N_1}} \sum_{i=1}^{N_1} (\tau_i - E(\tau)) = O_p(\sqrt{N_1 T})$ , which is  $O_p(\sqrt{NT})$  provided that  $\delta_1 > 0$ . It follows that  $\tau_{IPS} = O_p(\sqrt{NT})$ .

The rate of divergence is therefore the same for both tests. Note also that the rate of divergence of  $\tau_{IPS}$  is independent of the value taken by  $\delta_1$ , as long as  $\delta_1 > 0$ . The divergence rate of this test in a panel where for example only half of the units are stationary is therefore the same as that in a panel where all units are stationary.

In other words, under a fixed alternative hypothesis and the conditions considered here little can be said about the power of the LLC and IPS tests. A more detailed analysis requires more assumptions. This is done in Bowman (1999), who studies the power of some panel unit root tests in the absence of nuisance parameters. He characterizes the class of admissible panel unit root tests, and shows that while the LLC test is admissible, and in fact uniformly most powerful against the homogenous alternative  $H_{1b}$ , the IPS test is inadmissible. The LLC test is therefore expected to be more powerful than the IPS test.

Consider next the case when  $\alpha_i$  is local-to-unity,

$$H_{1c} : \alpha_i = \frac{c_i}{T\sqrt{N}}, \quad (4)$$

where  $c_i < 0$  is a constant such that  $\frac{1}{N} \sum_{i=1}^N c_i \rightarrow \bar{c}$  as  $N \rightarrow \infty$ . This corresponds to an autoregressive coefficient that approaches one with increasing values of  $N$  and  $T$ . Since  $c_i < 0$  this means that  $\alpha_i$  approaches one from below and so  $y_{it}$  is locally stationary. Note also that since  $c_i$  is not restricted to be equal across  $i$ , the mean reversion is essentially heterogenous. Alternatives of this kind has been extensively used in the time series literature to study the power of unit root test statistics. The rate

of shrinking of the local alternative is faster in our case, though, because of the faster rate of convergence of the pooled LS estimator of  $\alpha$  under  $H_0$ .

Let us assume that  $y_{i0} = 0$ .<sup>3</sup> Then, by Taylor expansion,

$$\begin{aligned} \frac{1}{\sigma\sqrt{T}} y_{it} &= \frac{1}{\sigma\sqrt{T}} \sum_{j=0}^t \rho_i^j \varepsilon_{it-j} \simeq \frac{1}{\sigma\sqrt{T}} \sum_{j=0}^t \varepsilon_{it-j} + \frac{c_i}{\sigma\sqrt{NT}} \sum_{j=1}^t \frac{j}{T} \varepsilon_{it-j} \\ &\Rightarrow W_i(s) + \frac{c_i}{\sqrt{N}} U_i(s) \end{aligned}$$

as  $T \rightarrow \infty$ , where  $U_i(s) = \int_0^s W_i(r) dr$ . Thus, by subsequently passing  $N \rightarrow \infty$ ,

$$\begin{aligned} \frac{1}{\sigma^2 T \sqrt{N}} \sum_{i=1}^N \sum_{t=2}^T y_{it-1} \varepsilon_{it} &\xrightarrow{d} \mathcal{N}\left(0, \frac{1}{2}\right) + \bar{c} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N E\left(\int_0^1 U_i(s) dW_i(s)\right) \\ &\sim \mathcal{N}\left(0, \frac{1}{2}\right), \end{aligned}$$

which uses the fact that  $E\left(\int_0^1 U_i(s) dW_i(s)\right) = 0$ . But we also have  $\frac{1}{T^2 N} M_{22} \xrightarrow{p} \frac{\sigma^2}{2}$  as  $N, T \rightarrow \infty$ , and therefore

$$\tau_{LLC} = \frac{1}{\hat{\sigma} \sqrt{\frac{1}{NT^2} M_{22}}} \frac{1}{T \sqrt{N}} \sum_{i=1}^N \sum_{t=2}^T \left( \frac{c_i}{T \sqrt{N}} y_{it-1}^2 + y_{it-1} \varepsilon_{it} \right) \xrightarrow{d} \mathcal{N}\left(\frac{\bar{c}}{\sqrt{2}}, 1\right).$$

Thus, in spite of the heterogeneity of this alternative, we find that the LLC test has nontrivial local power. The basic reason for this is that the power only depends on  $\bar{c}$ , the mean of  $c_i$ , which means that it does not matter whether  $c_i$  is equal across  $i$ . It is also interesting to note that the denominator of the LLC statistic does not contribute to the local power of the test, which stands in sharp contrast to the DF statistic, whose local power depends on both the numerator and the denominator.

Let us now consider the local power of the IPS statistic. Using Taylor expansion and then inserting

$$\begin{aligned} \frac{1}{\sigma^2 T} \sum_{t=2}^T y_{it-1} \Delta y_{it} &\Rightarrow \int_0^1 \left( W_i(r) + \frac{c_i}{\sqrt{N}} U_i(r) \right) \left( dW_i(r) + \frac{c_i}{\sqrt{N}} W_i(r) \right) dr \\ &= \int_0^1 W_i(r) dW_i(r) + \frac{c_i}{\sqrt{N}} \left( \int_0^1 U_i(r) dW_i(r) + \int_0^1 W_i(r)^2 dr \right) \end{aligned}$$

<sup>3</sup>The effect of a non-zero initial value is considered in Fact 5.

$$\begin{aligned}
& + O_p\left(\frac{1}{N}\right) \\
& = M_{12i}^\circ + \frac{c_i}{\sqrt{N}}(R_{1i} + M_{22i}^\circ) + O_p\left(\frac{1}{N}\right), \\
\frac{1}{\sigma^2 T^2} \sum_{t=2}^T y_{it-1}^2 & \Rightarrow \int_0^1 \left( W_i(r) + \frac{c_i}{\sqrt{N}} U_i(r) \right)^2 \\
& = \int_0^1 W_i(r)^2 dr + \frac{2c_i}{\sqrt{N}} \int_0^1 W_i(r) U_i(r) dr + O_p\left(\frac{1}{N}\right) \\
& = M_{22i}^\circ + \frac{2c_i}{\sqrt{N}} R_{2i} + O_p\left(\frac{1}{N}\right),
\end{aligned}$$

we obtain

$$\tau_i \Rightarrow \frac{M_{12i}^\circ}{\sqrt{M_{22i}^\circ}} + \frac{c_i}{\sqrt{N}} \left( \sqrt{M_{22i}^\circ} + \frac{R_{1i}}{\sqrt{M_{22i}^\circ}} - \frac{M_{12i}^\circ R_{2i}}{(M_{22i}^\circ)^{3/2}} \right) + O_p\left(\frac{1}{N}\right).$$

It follows that as  $N, T \rightarrow \infty$ ,

$$\tau_{IPS} \xrightarrow{d} \mathcal{N}(0, 1) + \frac{\bar{c}}{\sqrt{\text{var}(\tau)}} E \left( \sqrt{M_{22i}^\circ} + \frac{R_{1i}}{\sqrt{M_{22i}^\circ}} - \frac{M_{12i}^\circ R_{2i}}{(M_{22i}^\circ)^{3/2}} \right),$$

see Harris et al. (2010).

Using simulations where the Brownian motion  $W_i(r)$  is approximated by a random walk of length  $T = 1,000$  we find

$$E \left( \sqrt{M_{22i}^\circ} + \frac{R_{1i}}{\sqrt{M_{22i}^\circ}} - \frac{M_{12i}^\circ R_{2i}}{(M_{22i}^\circ)^{3/2}} \right) = 0.6221 - 0.0794 + 0.0382 = 0.581.$$

Since  $0.581/\sqrt{\text{var}(\tau)} = 0.581/0.985 \cong 0.6 < 1/\sqrt{2} = 0.707$  it follows that the local power of the IPS test is always smaller than that of the LLC test, which corroborates the results of Bowman (1999).

To illustrate these findings a small simulation experiment was conducted using (1), (2) and (4) with  $\varepsilon_{it} \sim \mathcal{N}(0, 1)$  and  $y_{i0} = 0$  to generate the data. Two specifications are considered. In the first,  $c_i = c$  for all  $i$ , suggesting a completely homogenous AR parameter, while in the second,  $c_i \sim U(2c, 0)$ . Hence,  $\text{var}(c_i) = c^2/3 > 0$  whenever  $c < 0$  and so the individual AR coefficients are no longer restricted to be equal. However, the mean is still  $c$ , just as in the first specification. The empirical rejection

**TABLE 1** Power against different local alternatives

$T, N$	$c$	$c_i = c$		$c_i \sim U(2c, 0)$	
		LLC	IPS	LLC	IPS
20	−1	15.4	12.5	15.3	12.4
	−2	33.5	24.0	31.3	23.3
	−5	90.4	73.8	76.6	67.9
50	−1	16.6	13.1	16.2	12.8
	−2	36.6	26.8	33.9	26.1
	−5	93.7	80.4	85.0	75.9
100	−1	16.8	13.3	16.5	13.3
	−2	38.7	26.8	36.7	26.1
	−5	94.8	83.5	89.8	79.9
Asymptotic	−1	17.4	14.8	17.4	14.8
	−2	40.9	32.8	40.9	32.8
	−5	97.1	91.23	97.1	91.2

The table reports the 5% rejection frequencies when the AR parameter is set to  $\alpha_i = c_i/T\sqrt{N}$ .

frequencies are based on 5,000 replications and the 5% critical value.<sup>4</sup> The results are summarized in Table 1. We see that in agreement with the theoretical results,  $\tau_{LLC}$  is uniformly more powerful than  $\tau_{IPS}$ . We also see that the actual power corresponds roughly to the asymptotic power, at least for large samples and small values of  $c$ .

These results suggest that if it is power that we are concerned about, then the LLC test is the one to use.

### 3. MODELS WITH DETERMINISTIC TERMS

#### **Fact 3: Deterministic Components Need not Be Treated as in the DF Approach**

In the presence of deterministic constant and trend terms, LLC and IPS suggest following the DF proposal of using LS demeaning. One might therefore think that this is also the simplest way to handle such terms. But this is not true.<sup>5</sup>

Consider the model

$$y_{it} = \mu_i + y_{it}^s, \quad (5)$$

<sup>4</sup>From now on all simulations will be conducted at the 5% level using 5,000 replications. Also, in order to reduce the effect of the initial condition, the first 100 observations of each cross-sectional unit will henceforth be disregarded.

<sup>5</sup>As an alternative, the local-to-unity demeaning procedure of Elliott et al. (1996) can be used, which is expected to lead to better small sample properties, see for example Choi (2001) and Lopez (2009).

where the constant  $\mu_i$  now represents the deterministic part of  $y_{it}$ , while  $y_{it}^s$  again represents the stochastic part. As usual, the allowance for deterministic terms of this kind makes it necessary to appropriately augment the regression in (3). Let us therefore introduce  $x_{it}$  to denote a generic vector containing all regressors other than  $y_{it-1}$  with  $\gamma_i$  being the associated vector of slope coefficients. In the current case with a constant this yields

$$\Delta y_{it} = \alpha_i y_{it-1} - \alpha_i \mu_i + \varepsilon_{it} = \alpha_i y_{it-1} + \gamma_i x_{it} + \varepsilon_{it}, \quad (6)$$

where  $\gamma_i = -\alpha_i \mu_i$  and  $x_{it} = 1$  for all  $i$  and  $t$ . The matrix of sample moments is augmented accordingly as

$$M_i = \begin{pmatrix} M_{11i} & M_{12i} & M_{13i} \\ M_{12i} & M_{22i} & M_{23i} \\ M'_{13i} & M'_{23i} & M_{33i} \end{pmatrix} = \sum_{t=2}^T \begin{pmatrix} (\Delta y_{it})^2 & y_{it-1} \Delta y_{it} & \Delta y_{it} x'_{it} \\ y_{it-1} \Delta y_{it} & y_{it-1}^2 & y_{it-1} x'_{it} \\ x_{it} \Delta y_{it} & x_{it} y_{it-1} & x_{it} x'_{it} \end{pmatrix}$$

with  $x_{it}$  ordered last. Moreover, since the focus here is on  $\alpha_i$  and not on  $\gamma_i$ , the analysis will be carried out in two steps, where the first involves projecting  $\Delta y_{it}$  and  $y_{it-1}$  upon  $x_{it}$ . The second step is then to test for a unit root in the resulting projection errors, whose sum of squares can be written in terms of the partitions of  $M_i$  as

$$M_{abi}^p = M_{abi} - M_{a3i} M_{33i}^{-1} M_{3bi}.$$

The corresponding limiting sum of squared projection errors is defined as

$$M_{abi}^{\circ p} = M_{abi}^{\circ} - M_{a3i}^{\circ} (M_{33i}^{\circ})^{-1} M_{3bi}^{\circ}$$

with an obvious definition of  $M_{abi}^{\circ}$ .

Also, except for  $M^p$ , to simplify the notation let us from now on suppress any dependence upon  $p$ . For example, we write  $\hat{\sigma}^2 = \frac{1}{NT} (M_{11}^p - \hat{\alpha} M_{12}^p)$  and  $\hat{\alpha} = M_{12}^p / M_{22}^p$ , which are the same definitions as in Section 2 but with the elements of  $M^p$  in place of the corresponding elements of  $M$ .

Consider now the DF approach of using LS demeaning, in which case

$$M_{abi}^p = M_{abi} - \frac{1}{T} M_{a3i} M_{3bi},$$

so that for example  $M_{12i}^p = \sum_{t=2}^T (y_{it-1} - \bar{y}_{i,-1}) \Delta y_{it}$ , where  $\bar{y}_{i,-1} = \frac{1}{T} \sum_{t=2}^T y_{it-1}$ . The limiting version of this quantity is given by  $M_{12i}^{\circ p} = \int_0^1 (W_i(s) - \bar{W}_i) dW_i(s)$ , where  $\bar{W}_i = \int_0^1 W_i(s) ds$ . Thus, since  $E(M_{12i}^{\circ p}) = -1/2$  under

$H_0$ , we have that in the sequential limit, where  $T \rightarrow \infty$  is followed by  $N \rightarrow \infty$

$$\frac{1}{TN} M_{12}^p \xrightarrow{p} \sigma^2 E(M_{12i}^{op}) = -\frac{\sigma^2}{2}.$$

Since  $\hat{\sigma}^2 \xrightarrow{p} \sigma^2$  and  $\frac{1}{T^2} E(M_{22i}^p) \rightarrow \frac{\sigma^2}{6}$  we have that as  $N, T \rightarrow \infty$

$$\frac{1}{\sqrt{N}} \tau_{LLC} = \frac{\frac{1}{TN} M_{12}^p}{\hat{\sigma} \sqrt{\frac{1}{T^2 N} M_{22}^p}} \xrightarrow{p} -\frac{\sqrt{6}}{2}$$

and by using  $\text{var}(M_{12i}^p)/T^2 \rightarrow \sigma^4/12$ ,

$$\frac{\sqrt{12N}}{\sigma^2} \left( \frac{1}{TN} M_{12}^p + \frac{\sigma^2}{2} \right) \xrightarrow{d} \mathcal{N}(0, 1).$$

It follows that

$$\tau_{LLC}^c = \frac{\sqrt{2N} \left( \frac{1}{TN} M_{12}^p + \frac{\hat{\sigma}^2}{2} \right)}{\hat{\sigma} \sqrt{\frac{1}{NT^2} M_{22}^p}} \xrightarrow{d} \mathcal{N}(0, 1).$$

This is the bias-adjusted LLC statistic, which has been superscripted by  $c$  to indicate that it is robust to the presence of the constant in the model. The point here is that LS demeaning is not enough to get rid off the effect of  $\mu_i$ . There is also a bias that needs to be accounted for, which complicates the testing considerably. This is the so-called Nickell bias first analyzed in the context of a fixed  $T$  by Nickell (1981), which has been further analyzed by Hahn and Kuersteiner (2002) in cases when  $N, T \rightarrow \infty$  such that  $\frac{N}{T}$  converges to a constant.

As mentioned in Section 2, as soon as one moves away from the simplest case with no deterministic components and no short-run dynamics, the statistic proposed in Levin and Lin (1992) need not be the same as the one in LLC. The main difference between the two is that while the LLC test is adjusted for the nonzero mean in the numerator, the Levin and Lin (1992) test is adjusted for the nonzero mean in the entire statistic (see Moon and Perron, 2008). In the current setting, Levin and Lin (1992) statistic can be written as

$$\tau_{LL}^c = \frac{\sqrt{5}}{2} \tau_{LLC} + \sqrt{\frac{15N}{8}}.$$

To appreciate the effect of the two approaches let us begin by expanding  $\tau_{LL}^c$  as

$$\begin{aligned} \frac{2}{\sqrt{5}} \tau_{LL}^c &= \tau_{LLC} + \sqrt{\frac{3N}{2}} = \sqrt{N} \frac{\frac{1}{NT} M_{12}^p}{\hat{\sigma} \sqrt{\frac{1}{NT^2} M_{22}^p}} + \sqrt{N} \frac{\frac{1}{2} \sigma^2}{\sigma \sqrt{\frac{\sigma^2}{6}}} \\ &= \frac{\sqrt{N} \left( \frac{1}{NT} M_{12}^p + \frac{1}{2} \sigma^2 \right)}{\hat{\sigma} \sqrt{\frac{1}{NT^2} M_{22}^p}} - \frac{1}{2} \sigma^2 \sqrt{N} \left( \frac{1}{\hat{\sigma} \sqrt{\frac{1}{NT^2} M_{22}^p}} - \frac{1}{\sigma \sqrt{\frac{\sigma^2}{6}}} \right), \end{aligned}$$

which, by Taylor expansion of the second term, yields

$$\begin{aligned} \frac{2}{\sqrt{5}} \tau_{LL}^c &\simeq \frac{\sqrt{N} \left( \frac{1}{NT} M_{12}^p + \frac{1}{2} \sigma^2 \right)}{\sqrt{\hat{\sigma}^2 \frac{1}{NT^2} M_{22}^p}} + \hat{\sigma}^2 \sqrt{\frac{27}{2\sigma^8}} \sqrt{N} \left( \frac{1}{NT^2} M_{22}^p - \frac{\sigma^2}{6} \right) \\ &\quad + \frac{1}{NT^2} M_{22}^p \sqrt{\frac{27}{72\sigma^8}} \sqrt{N} (\hat{\sigma}^2 - \sigma^2) \xrightarrow{d} \frac{\frac{\sigma^2}{\sqrt{12}} Z_1}{\sigma \sqrt{\frac{\sigma^2}{6}}} + \sigma^2 \sqrt{\frac{27}{2\sigma^8}} \frac{\sigma^2}{\sqrt{45}} Z_2, \end{aligned}$$

where  $Z_1$  and  $Z_2$  are two independent standard normal random variables. In order to obtain this results, we have used that  $\frac{1}{T^2} \text{var}(M_{22i}^p) \rightarrow \frac{\sigma^4}{45}$  and  $\sqrt{N}(\hat{\sigma}^2 - \sigma^2) = o_p(1)$ , see Lemma 2 of Moon and Phillips (2004). It follows that

$$\frac{2}{\sqrt{5}} \tau_{LL}^c \xrightarrow{d} \left( \frac{1}{\sqrt{2}} + \sqrt{\frac{3}{10}} \right) \mathcal{N}(0, 1) \sim \frac{2}{\sqrt{5}} \mathcal{N}(0, 1),$$

or  $\tau_{LL}^c \xrightarrow{d} \mathcal{N}(0, 1)$ .

Thus, although the end result is the same as for  $\tau_{LLC}^c$ , the route to normality is more complicated, and involves additional approximations, which is suggestive of poor small-sample properties. On the other hand, the bias-adjustment of LLC requires estimation of  $\sigma^2$ , which obviously increases the variability of their test.

The relationship between  $\tau_{LL}^c$  and  $\tau_{LLC}^c$  is easily seen by noting that

$$\begin{aligned} \frac{2}{\sqrt{5}} \tau_{LL}^c &= \tau_{LLC} + \sqrt{\frac{3N}{2}} = \tau_{LLC} + \frac{1}{2} \sqrt{N} \frac{\sigma}{\sqrt{\frac{\sigma^2}{6}}} \\ &= \tau_{LLC} + \frac{1}{2} \sqrt{N} \text{plim}_{N, N \rightarrow \infty} \frac{\hat{\sigma}}{\sqrt{\frac{1}{NT^2} M_{22}^p}}, \end{aligned}$$

which is asymptotically equivalent to

$$\frac{1}{\sqrt{2}}\tau_{LLC}^c = \tau_{LLC} + \frac{\sqrt{N}}{2} \frac{\hat{\sigma}}{\sqrt{\frac{1}{NT^2}M_{22}^p}}.$$

The demeaning not only complicates the route to normality but also impact the local power of the tests. Consider  $\tau_{LLC}^c$ . From Moon and Perron (2008) we have that under  $H_{1c}$ ,

$$\frac{1}{\sigma\sqrt{T}}(y_{it} - \bar{y}_i) \Rightarrow W_i(s) - \bar{W}_i + \frac{c_i}{\sqrt{N}}(U_i(s) - \bar{U}_i) + O_p\left(\frac{1}{N^{3/4}}\right)$$

as  $T \rightarrow \infty$ , which implies

$$\frac{1}{\sigma^2 T}M_{12i}^p \Rightarrow M_{12i}^{op} + \frac{c_i}{\sqrt{N}}\left(M_{22i}^{op} + \int_0^1 (U_i(r) - \bar{U}_i)dW_i(r)\right) + O_p\left(\frac{1}{N^{3/4}}\right).$$

Using  $E(M_{12i}^{op}) = \sigma^2/2$ ,  $E(M_{22i}^{op}) = \sigma^2/6$ ,  $\text{var}(M_{12i}^{op}) = \sigma^2/12$ , and

$$E\left(\int_0^1 (U_i(r) - \bar{U}_i)dW_i(r)\right) = -E(W_i(1)\bar{U}_i) = -\frac{1}{6}$$

it is possible to show that as  $N, T \rightarrow \infty$

$$\begin{aligned} \frac{\sqrt{12N}}{\sigma^2} \left( \frac{1}{NT}M_{12}^p + \frac{\sigma^2}{2} \right) &\xrightarrow{d} \mathcal{N}(0, 1) + \sqrt{12} \bar{c} E\left(M_{22i}^{op} + \int_0^1 (U_i(r) - \bar{U}_i)dW_i(r)\right) \\ &\sim \mathcal{N}(0, 1). \end{aligned}$$

Hence, under the typical local alternative given by (4) the limiting distribution of the numerator of  $\tau_{LLC}^c$  does not depend on  $c_i$ . For the denominator we have as  $T \rightarrow \infty$

$$\frac{1}{\sigma^2 T^2}M_{22i}^p \Rightarrow M_{22i}^{op} + \frac{2c_i}{\sqrt{N}} \int_0^1 (W_i(r) - \bar{W}_i)(U_i(r) - \bar{U}_i)dr + O_p\left(\frac{1}{N^{3/4}}\right),$$


suggesting that

$$\frac{1}{NT^2}M_{22}^p \xrightarrow{p} \sigma^2 E(M_{22i}^{op}) = \frac{\sigma^2}{6}$$

as  $N, T \rightarrow \infty$ , from which it follows that

$$\tau_{LLC}^c = \frac{\sqrt{2N} \left( \frac{1}{TN}M_{12}^p + \frac{\hat{\sigma}^2}{2} \right)}{\hat{\sigma} \sqrt{\frac{1}{NT^2}M_{22}^p}} \xrightarrow{d} \mathcal{N}(0, 1).$$





**FIGURE 1** Local power of  $\tau_{LL}^c$  and  $\tau_{LLC}^c$  as a function of  $c$ . (Figure available in color online.)

In other words,  $\tau_{LLC}^c$  does not have any power against  $H_{1c}$ . This is illustrated in Figure 1, which plots the local power as a function of  $c$  when the data are generated from (1), (2), and (4) with  $c_i \sim U(2c, 0)$ . As in Table 1, the results are based on 5,000 replications and the 5% critical value. Note in particular that the power function of  $\tau_{LLC}^c$  is flat, thereby corroborating the theoretical results. We also see that the power function of  $\tau_{LL}^c$  is strictly increasing in  $c$ , suggesting that unlike  $\tau_{LLC}^c$ ,  $\tau_{LL}^c$  does have power against  $H_{1c}$ .

This loss of power is rather unexpected because while it is fairly well-known that the presence of individual specific trends can be problematic in this regard, an issue that will be discussed in detail in the next section, here we only consider the case with individual specific intercepts. The reason is the absence of serial correlation, or rather the absence of correction thereof, which turns out to be rather important for a number of reasons, see Section 4 for a more detailed analysis.

The point here is that these complications are all due to the fact that the constant is removed by LS demeaning. Thus, in order to avoid bias, one needs to consider alternatives to LS demeaning. For example, Breitung and Meyer (1994) suggest using the initial value  $y_{i0}$  as an estimator of the intercept. This suggests running a regression of  $\Delta y_{it}$  on  $y_{it-1}^* = y_{it-1} - y_{i0}$ . To analyze the effect of this modification, note that under

the null hypothesis

$$E(\Delta y_{it} y_{it-1}^*) = E(\Delta y_{it} (y_{it-1} - y_{i0})) = E\left(\varepsilon_{it} \sum_{s=1}^{t-1} \varepsilon_{is}\right) = 0.$$

In other words, using  $y_{i0}$  as an estimator of the intercept removes the bias. In fact, it is not difficult to show that as  $N, T \rightarrow \infty$

$$\tau_{BM}^c = \frac{\sum_{i=1}^N \sum_{t=2}^T y_{it-1}^* \Delta y_{it}}{\hat{\sigma} \sqrt{\sum_{i=1}^N \sum_{t=2}^T (y_{it-1}^*)^2}} \xrightarrow{d} \mathcal{N}(0, 1),$$

where  $\tau_{BM}^c$  is the Breitung and Meyer (1994) statistic.

Interestingly, as pointed out by Schmidt and Phillips (1992),  $y_{i0}$  is also the maximum likelihood (ML) estimator of the intercept under  $H_0$ , which has been shown to lead to significant power gains when compared to LS demeaning, see Madsen (2010). In fact, it is not difficult to see that under  $H_{1c}$ ,

$$\tau_{BM}^c \xrightarrow{d} \mathcal{N}\left(\frac{\bar{c}}{\sqrt{2}}, 1\right)$$

as  $N, T \rightarrow \infty$ , which is the same results we obtained earlier for the LLC statistic in the model without any deterministic terms.

To examine the extent of these gains in small samples, Table 2 reports some results based on data generated from (1), (2), and (4) with  $c_i \sim U(2c, 0)$ . Consistent with the results of Madsen (2010) we see that the tests based on removing the initial condition are almost uniformly more powerful than those based on LS demeaning. We also see that this increase in power comes at no cost in terms of size accuracy. Note that if the demeaning is accomplished by subtracting the initial observation the LLC and Levin and Lin (1992) tests are the same and hence we only report the results of the latter test.<sup>6</sup>

Another possibility is to demean  $y_{it-1}$  recursively as  $y_{it-1} - \frac{1}{t-1} \sum_{s=1}^{t-1} y_{is}$ , which can be used instead of  $y_{it-1}^*$  to produce yet another unbiased and standard normally distributed test statistic.

These alternative demeaning approaches can be seen as special cases of a more general class of test statistics. In particular, letting  $\Delta y_i = (\Delta y_{i2}, \dots, \Delta y_{iT})'$  and  $y_{i,-1} = (y_{i1}, \dots, y_{iT-1})'$ , these statistics can be written as

$$\frac{\sum_{i=1}^N (\Delta y_i)' C y_{i,-1}}{\hat{\sigma} \sqrt{\sum_{i=1}^N y_{i,-1}' C' C y_{i,-1}}}.$$

<sup>6</sup>The performance of the LLC test under LS demeaning can be inferred from Fig. 1.

**TABLE 2** Size and local power for different demeaning procedures

$c$	$N$	$T$	LL		IPS	
			LS	ML	LS	ML
0	10	50	7.1	7.1	7.3	5.8
	20	50	6.8	6.9	7.4	6.3
	10	100	6.4	7.5	4.8	5.4
	20	100	6.6	7.0	5.3	5.8
-1	10	50	10.1	11.9	9.4	8.5
	20	50	9.3	12.9	9.7	10.3
	10	100	8.6	13.5	6.7	10.0
	20	100	9.6	13.2	8.2	11.2
-2	10	50	13.2	18.5	10.5	12.5
	20	50	12.5	20.1	11.6	14.4
	10	100	10.9	18.0	7.0	14.4
	20	100	11.7	19.9	10.0	16.2
-5	10	50	25.9	41.0	19.8	31.2
	20	50	23.7	46.1	20.9	37.3
	10	100	21.6	39.8	15.3	33.3
	20	100	24.0	43.3	17.3	36.8

The table reports the 5% rejection frequencies when the AR parameter is set to  $\alpha_i = c_i/T\sqrt{N}$ , where  $c_i \sim U(2c, 0)$ . LS and ML refers to demeaning by LS and maximum likelihood, respectively, where the latter is based on removing the first observation from  $y_{it}$ . LL refers to the Levin and Lin (1992) test.

The matrix  $C$  has the property that  $C \iota_{T-1} = 0$ , where  $\iota_{T-1}$  is a  $(T-1) \times 1$  vector of ones. Therefore, pre-multiplying  $y_{i,-1}$  by  $C$  eliminates the individual specific constant. The statistic has expectation zero if

$$E((\Delta y_i)' C y_{i,-1}) = \sigma^2 \text{tr}(CD) = 0,$$

where  $D$  is a matrix with elements  $d_{jk} = 1$  if  $j < k$  and  $d_{jk} = 0$  for  $j \geq k$ . Note that in the case of LS demeaning,  $C = I_{T-1} - \frac{1}{T-1} \iota_{T-1} \iota_{T-1}'$ , where  $I_{T-1}$  is the identity matrix. Since in this case  $\text{tr}(CD) \neq 0$ , bias correction is needed. The same principle can be used to construct bias-corrected statistics in models with trends, an issue to be discussed in the next section.

#### **Fact 4: Incidental Trends Reduce the Local Power of the LLC Test**

Suppose now that instead of (5), we have

$$y_{it} = \mu_i + \beta_i t + y_{it}^s, \quad (7)$$

where  $\beta_i t$  is a unit specific trend term, giving

$$\begin{aligned}\Delta y_{it} &= -\alpha_i \mu_i + (\alpha_i + 1)\beta_i - \alpha_i \beta_i t + \alpha_i y_{it-1} + \varepsilon_{it} \\ &= \alpha_i y_{it-1} + \gamma'_i x_{it} + \varepsilon_{it}\end{aligned}\quad (8)$$

with  $x_{it} = (1, t)'$ .

The incidental trends problem refers to the need of having to estimate an increasing number of trend coefficients  $\beta_1, \dots, \beta_N$  as  $N \rightarrow \infty$ , which reduces the discriminatory power against  $H_0$ , see Moon and Phillips (1999).<sup>7</sup> In particular, as we will now demonstrate the presence of trends even has an order effect on the neighborhoods around the unit root null for which asymptotic power is non-negligible.

As Moon et al. (2007) show in the case with incidental trends the LLC statistic is asymptotically equivalent to

$$\tau_{LLC}^t = \frac{193}{112} \tau_{LLC} + \sqrt{\frac{252}{772}} \frac{10}{T} \frac{\sqrt{M_{22}^p}}{\hat{\sigma}},$$

where the superscript  $t$  indicates invariance with respect to the trend, while  $\tau_{LLC}$  is now the LLC statistic based on the detrended data. Moon and Phillips (2004) consider another statistic, which in the present setting may be written as

$$\tau_{MP}^t = \tau_{LLC} + \frac{NT}{2} \frac{\hat{\sigma}}{\sqrt{M_{22}^p}}.$$

It follows that

$$\sqrt{\frac{193}{112}} \tau_{LLC}^t = \tau_{MP}^t + \frac{15}{2T} \frac{(M_{22}^p - \frac{1}{15} \hat{\sigma}^2)}{\hat{\sigma} \sqrt{M_{22}^p}},$$

suggesting that  $\tau_{LLC}^t$  will inherit some of the asymptotic properties of  $\tau_{MP}^t$ . In particular, from Theorem 4 of Moon and Phillips (2004) we know that  $\tau_{MP}^t$  has power within  $\frac{1}{N^{1/6}T}$  neighborhoods of  $H_0$ , but not for any higher powers of  $N$  and  $T$ . In particular,  $\tau_{MP}^t$  has no power against  $H_{1c}$  when the neighborhood is of order  $\frac{1}{T\sqrt{N}}$ . The above relationship suggests that  $\tau_{LLC}^t$  should have the same property. Thus, just as in the case of an intercept, we see that the presence of the trend leads to a loss of power. This is illustrated in Table 3, which plots the local power of the LLC and IPS tests

<sup>7</sup>Moon and Phillips (1999) show that the ML estimator of the local-to-unity parameter is inconsistent. They call this phenomenon an incidental trend problem because it is analogous to the well-known incidental parameter problem in dynamic panels when  $N \rightarrow \infty$  and  $T$  is fixed.

**TABLE 3** Local power in the presence of incidental trends

$c$	$N$	$T$	LLC	IPS
-4	20	50	10.6	12.2
	10	100	6.9	7.7
	20	100	8.2	8.4
	10	50	13.4	12.4
	20	50	12.6	14.3
	10	100	10.2	9.4
-8	20	100	10.0	10.4
	10	50	22.8	22.1
	20	50	20.0	20.5
	10	100	16.9	14.6
	20	100	14.9	14.9

The table reports the 5% rejection frequencies when the AR parameter is set to  $\alpha_i = c_i/T\sqrt{N}$ , where  $c_i \sim U(2c, 0)$ .

for some different values of  $c$  when  $\alpha_i$  is generated according to (4) with  $c_i \sim U(2c, 0)$ . In accordance with the theoretical results we see that the power can be very low and practically nonexistent in many cases if there is a trend in the model.

In view of the previous fact one might think that this loss of power is due to the fact that the detrending is carried out using LS. However, this is not true. Take as an example the study of Breitung (2000), who proposes a generalized version of the demeaning by initial value procedure discussed in the previous section. Specifically, using  $y_{i0}$  and  $\frac{1}{T} \sum_{t=2}^T \Delta y_{it} = \frac{1}{T}(y_{iT} - y_{i0})$  as estimators of the constant and trend, respectively, Breitung (2000) proposes replacing (8) with a regression of  $\Delta y_{it}^*$  on  $y_{it-1}^*$ , where  $y_{it}^* = y_{it} - y_{i0} - \frac{1}{T}(y_{iT} - y_{i0})t$  and

$$\Delta y_{it}^* = s_t \left( \Delta y_{it} - \frac{1}{T-t}(y_{iT} - y_{it}) \right)$$

with  $s_t^2 = (T-t)/(T-t+1)$ . The effect of this is easily seen by noting that

$$\begin{aligned}
 E(\Delta y_{it}^* y_{it-1}^*) &= s_t E \left( \left( \Delta y_{it} - \frac{1}{T-t}(y_{iT} - y_{it}) \right) \left( y_{it} - y_{i0} - \frac{1}{T}(y_{iT} - y_{i0}) \right) \right) \\
 &= s_t E \left( \left( \Delta y_{it}^s - \frac{1}{T-t}(y_{iT}^s - y_{it}^s) \right) \left( y_{it}^s - y_{i0}^s - \frac{1}{T}(y_{iT}^s - y_{i0}^s) \right) \right) \\
 &= s_t E \left( \left( \varepsilon_{it} - \frac{1}{T-t}(y_{iT}^s - y_{it}^s) \right) \left( y_{it-1}^s - \frac{t-1}{T} y_{iT}^s \right) \right) \\
 &= s_t \left( \frac{t-1}{T} \sigma^2 - \frac{(t-1)(T-t)}{(T-t)T} \sigma^2 \right) = 0,
 \end{aligned}$$

showing that the bias has been successfully eliminated.

However, as Moon et al. (2006) show, just as with  $\tau_{MP}^t$  and  $\tau_{LLC}^t$ , the Breitung (2000) test has no power in neighborhoods that shrinks to zero at a faster rate than  $\frac{1}{N^{1/4}T}$ . The reduced power effect in the presence of trends is therefore not specific to  $\tau_{MP}^t$  and  $\tau_{LLC}^t$  but is a general property of this type of tests. In fact, as Ploberger and Phillips (2002) show, the panel unit root test that maximizes the average local power has nontrivial power in local neighborhoods that shrink at the same rate,  $\frac{1}{N^{1/4}T}$ .

### Fact 5: The Initial Condition May Affect the Asymptotic Properties of the Tests

The power of panel unit root tests is usually evaluated while assuming that all  $N$  units are initiated at zero. Although this is a convenient assumption that simplifies the theoretical considerations, as Moon et al. (2007) points out it is very unrealistic and, as we will see, by no means innocuous. Suppose for example that  $y_{it}$  is generated according to (5) with a constant and where  $y_{it}^s$  is as in (2). But suppose now that instead of setting  $y_{i0}^s$  to zero, we follow the convention in the time series context and consider the initial condition as a draw from the unconditional distribution under the stationary alternative, see for example Phillips and Sul (2003). That is, we set

$$y_{i0}^s = \frac{\sigma}{\sqrt{1 - \rho_i^2}} \eta_i,$$

where  $\eta_i$  is independent and identically distributed with mean  $\bar{\eta}$  and variance  $\sigma_\eta^2$ . Note that for  $\rho_i < 1$  and  $\eta_i \sim \mathcal{N}(0, 1)$  this initial condition implies that  $y_{it}^s$  is stationary.

Similar to what we had before when  $y_{i0}^s = 0$ , Harris et al. (2010) show that under  $H_{1c}$ , as  $T \rightarrow \infty$

$$\begin{aligned} \frac{1}{\sigma\sqrt{T}}(y_{it} - \bar{y}_i) &\Rightarrow \frac{\eta_i}{N^{1/4}}\sqrt{\frac{-c_i}{2}}\left(r - \frac{1}{2}\right) + W_i(s) - \bar{W}_i + \frac{c_i}{\sqrt{N}}(U_i(s) - \bar{U}_i) \\ &\quad + O_p\left(\frac{1}{N^{3/4}}\right), \end{aligned}$$

where  $U_i(s)$  is again the integral of  $W_i(r)$  over  $[0, s]$ , implying

$$\begin{aligned} \frac{1}{\sigma^2 T} M_{12i}^p &\Rightarrow M_{12i}^{op} + \frac{c_i}{\sqrt{N}} \left( M_{22i}^{op} + \int_0^1 (U_i(r) - \bar{U}_i) dW_i(r) \right) \\ &\quad - \frac{\eta_i}{N^{1/4}} \sqrt{\frac{-c_i}{2}} \int_0^1 \left( r - \frac{1}{2} \right) dW_i(r) + O_p\left(\frac{1}{N^{3/4}}\right). \end{aligned}$$

It follows that as  $N, T \rightarrow \infty$

$$\frac{\sqrt{12N}}{\sigma^2} \left( \frac{1}{NT} M_{12}^p + \frac{\sigma^2}{2} \right) \xrightarrow{d} \mathcal{N}(0, 1) + \sqrt{12} \bar{c} E \left( M_{22i}^{op} + \int_0^1 (U_i(r) - \bar{U}_i) dW_i(r) \right) \\ - \text{plim}_{N \rightarrow \infty} \frac{1}{N^{3/4}} \sum_{i=1}^N \eta_i \sqrt{\frac{-c_i}{2}} \int_0^1 \left( r - \frac{1}{2} \right) dW_i(r).$$

But  $E(dW_i(r)) = 0$ , suggesting that the last term on the right-hand side is  $O_p(1/N^{1/4})$ . Thus, since the second term is zero,

$$\frac{\sqrt{12N}}{\sigma^2} \left( \frac{1}{NT} M_{12}^p + \frac{\sigma^2}{2} \right) \xrightarrow{d} \mathcal{N}(0, 1),$$

which is the same result as the one we obtained when  $y_{i0}^s = 0$ . By further using  $\frac{1}{T^2 N} M_{22}^p \xrightarrow{p} \sigma^2/6$  it follows that  $\tau_{LLC}^c \xrightarrow{d} \mathcal{N}(0, 1)$ . Thus, just as before the asymptotic distribution of  $\tau_{LLC}^c$  is independent of both  $\bar{c}$  and  $y_{i0}^s$ .

The analysis for  $\tau_{LL}^c$  is more tedious, but the qualitative effect is easily appreciated by focusing on  $\sqrt{N} \left( \frac{1}{NT^2} M_{22}^p - \frac{\sigma^2}{6} \right)$ . Remember that from before that

$$\frac{2}{\sqrt{5}} \tau_{LL}^c = \frac{\sqrt{N} \left( \frac{1}{NT} M_{12}^p + \frac{1}{2} \sigma^2 \right)}{\sqrt{\hat{\sigma}^2 \frac{1}{NT^2} M_{22}^p}} + \hat{\sigma}^2 \sqrt{\frac{27}{2\sigma^8}} \sqrt{N} \left( \frac{1}{NT^2} M_{22}^p - \frac{\sigma^2}{6} \right) + O_p(1),$$

where the  $O_p(1)$  term is a remainder comprised of  $\sqrt{N}$  times the error of a Taylor expansion of the inverse square root of  $\frac{1}{NT^2} M_{22}^p$  around  $\sigma^2/6$  plus another term that is asymptotically negligible. We already know that the first term on the right-hand side is independent of  $\bar{c}$  and  $y_{i0}^s$ . Any dependence must therefore come through the second term, which then also determines the  $O_p(1)$  term. By using the same steps as before,

$$\frac{1}{\sigma^2 T^2} M_{22i}^p \Rightarrow M_{22i}^{op} + \frac{2c_i}{\sqrt{N}} \int_0^1 (U_i(r) - \bar{U}_i)(W_i(r) - \bar{W}_i) dr \\ + \frac{\eta_i^2 c_i}{2\sqrt{N}} \int_0^1 \left( r - \frac{1}{2} \right)^2 dr \\ + \frac{\eta_i \sqrt{-2c_i}}{N^{1/4}} \int_0^1 \left( r - \frac{1}{2} \right) (W_i(r) - \bar{W}_i) dr + O_p \left( \frac{1}{N^{3/4}} \right)$$

as  $T \rightarrow \infty$ , and therefore,

$$\sqrt{N} \left( \frac{1}{NT^2} M_{22}^p - \frac{\sigma^2}{6} \right)$$


$$\begin{aligned} &\xrightarrow{d} \frac{\sigma^2}{\sqrt{45}} \mathcal{N}(0, 1) + 2\bar{c}E \left( \int_0^1 (U_i(r) - \bar{U}_i)(W_i(r) - \bar{W}_i) dr \right) \\ &\quad + \frac{E(\eta_i^2)\bar{c}}{2} \int_0^1 \left( r - \frac{1}{2} \right)^2 dr \\ &\quad + \text{plim}_{N \rightarrow \infty} \frac{1}{N^{3/4}} \sum_{i=1}^N \eta_i \sqrt{-2c_i} \int_0^1 \left( r - \frac{1}{2} \right) (W_i(r) - \bar{W}_i) dr \end{aligned}$$

as  $T \rightarrow \infty$  and then  $N \rightarrow \infty$ . But  $E(W_i(r) - \bar{W}_i) = 0$  and  $\int_0^1 (r - 1/2)^2 dr = 1/12$ , which means that while the fourth term is eliminated, the third term is not, and therefore the dependence upon  $\eta_i$  remains. Thus, unlike  $\tau_{LLC}^c$ , the asymptotic distribution of  $\tau_{LL}^c$  does depend on the initial condition.

The same is true for the IPS test. In fact, as Harris et al. (2010) show,


$$\tau_{IPS}^c \xrightarrow{d} \mathcal{N}(0, 1) + \bar{c} \left( 0.282 - 0.135(\bar{\eta}^2 + \sigma_\eta^2) \right),$$

which shows that the local power of the IPS test is decreasing in  $\bar{\eta}^2$  and  $\sigma_\eta^2$ . In particular, note that if the initial condition is large enough so that  $0.282 > 0.135(\bar{\eta}^2 + \sigma_\eta^2)$ , then this test is no longer unbiased. This is illustrated in Figs. 2 and 3, which plot the local power of the IPS test for different combinations of  $N$  and  $\bar{\eta}$  when  $\sigma_\eta^2 = 0$  and  $\alpha_i$  is generated as in (4) with  $c_i = -10$  for all  $i$ . We see that if there is a constant present then the power is decreasing in both  $N$  and  $\bar{\eta}$ , while if there is no deterministic component then the power is very close to one.



**FIGURE 2** Local power of  $\tau_{IPS}^c$  for different initial values. (Figure available in color online.)





**FIGURE 3** Local power of  $\tau_{PS}$  for different initial values. (Figure available in color online.)

### Fact 6: The GMM Approach Can Be Used also in the Unit Root Case

As an alternative to using bias-adjusted LS or removing the initial condition, one may use generalized LS (GLS) estimation, which is especially appealing for panels with  $N \gg T$ . The problem is that most generalized method of moments (GMM) estimators of dynamic panel data models are rendered invalid in case of a unit root, see Breitung (1997), Kruiniger (2008), and Madsen (2010). Consider for example the well-known GMM estimator of Arellano and Bond (1991), which, in model (6) with no short-run dynamics, is based on the following moment condition:

$$E(y_{it}(\Delta y_{it} - \rho \Delta y_{it-1})) = 0 \quad \text{for } s = 1, 2, \dots, t-2.$$

It is important to note that under the unit root hypothesis the instruments  $y_{it-2}, y_{it-3}, \dots, y_{i1}$  fail to identify the common AR parameter  $\rho$ , and therefore the resulting GMM estimator is inconsistent and has a nonstandard limiting distribution. The instrumental variables estimator of Anderson and Hsiao (1982) suffers from the same problem, and in fact has a limiting Cauchy distribution as  $N \rightarrow \infty$  with  $T$  held fixed, see Breitung (1997) and Kruiniger (2009).

However, there are moment conditions that can be used to identify  $\rho$ , even under the unit root hypothesis. Breitung (1997), Bond et al. (2005), and Kruiniger (2009) consider the level equations of the system GMM estimator of Arellano and Bover (1995), and Blundell and Bond (1998).

The associated moment condition is given by

$$E(\Delta y_{it-1}(y_{it} - \rho y_{it-1})) = 0.$$

It is not difficult to see that if  $\rho \neq 1$ ,  $E(\Delta y_{it-1}(y_{it} - \rho y_{it-1})) \neq 0$ , and therefore GMM based on this moment condition is consistent and asymptotically normal. Another moment condition that can also be used is

$$E(\Delta y_{it-1}(2\Delta y_{it} + (1 - \rho)\Delta y_{it-1})) = 0,$$

which is particularly convenient because it implies that a consistent and asymptotically normal estimator of  $\rho$  can be obtained by applying LS to a regression of  $2\Delta y_{it} + \Delta y_{it-1}$  onto  $\Delta y_{it-1}$ . In fact, as Han and Phillips (2010) show, this estimator has a limiting normal distribution regardless of whether there is a unit root or not. The rate of consistency is  $\sqrt{NT}$ , which is slower than the usual unit root rate of  $T\sqrt{N}$ .

It is also possible to construct test statistics that are robust against the weak instruments problem, see Arellano and Hahn (2007). Breitung (1997) uses the Sargan statistic to test the unit root hypothesis based on the Arellano and Bond (1991) instruments. In the special case considered here, the test statistic can easily be performed by forming the  $(T - 2)(T - 1)/2$  dimensional vector

$$m_{Ti} = ((y_{i1}\Delta y_{i3}), (y_{i2}\Delta y_{i4}), \dots, (y_{iT-2}\Delta y_{iT}))'.$$

If  $T$  is fixed, the test statistic

$$J_N = \left( \sum_{i=1}^N m_{Ti} \right)' \left( \sum_{i=1}^N m_{Ti} m_{Ti}' \right)^{-1} \left( \sum_{i=1}^N m_{Ti} \right)$$

has a  $\chi^2$  limiting distribution with  $(T - 2)(T - 1)/2$  degrees of freedom. In practice, this means that  $N \gg (T - 2)(T - 1)/2$ .

#### 4. MODELS WITH SHORT-RUN DYNAMICS

##### **Fact 7: Lag Augmentation Does not Remove the Effects of Serial Correlation**

Suppose that (1) holds so that  $y_{it}$  is purely stochastic, but that the error  $\varepsilon_{it}$  in (2) is no longer independent across  $t$ . In particular, suppose that  $\varepsilon_{it}$  follows a stationary and invertible AR process of order  $\ell$ ,

$$\phi_i(L)\varepsilon_{it} = \left( 1 - \sum_{j=1}^{\ell} \phi_{ij}L^j \right) \varepsilon_{it} = \varepsilon_{it} - \sum_{j=1}^{\ell} \phi_{ij}\varepsilon_{it-j} = e_{it}, \quad (9)$$

where  $L$  is the lag operator and  $e_{it}$  is a mean zero error that has variance  $\sigma^2$  for all  $i$  but is otherwise independent across both  $i$  and  $t$ . Note that the long-run variance of  $\varepsilon_{it}$ , henceforth denoted  $\omega_i^2$ , can be expressed in terms of the lag polynomial  $\phi_i(L)$  as

$$\omega_i^2 = \frac{\sigma^2}{\phi_i(1)^2}.$$

Under  $H_0$ , (1), (2) and (9) can be combined to obtain the following augmented DF (ADF) regression:

$$\Delta y_{it} = \alpha_i y_{it-1} + \sum_{j=1}^{\ell} \phi_{ij} \Delta y_{it-j} + e_{it} = \alpha_i y_{it-1} + \gamma_i' x_{it} + e_{it}, \quad (10)$$

where  $x_{it} = (\Delta y_{it-1}, \dots, \Delta y_{it-\ell})'$  is now the vector of lagged differences with  $\gamma_i = (\phi_{i1}, \dots, \phi_{i\ell})'$  being the associated vector of lag coefficients. This gives rise to the ADF test statistic,

$$\tau_i = \frac{M_{12i}^p}{\hat{\sigma}_i \sqrt{M_{22i}^p}} = \hat{\alpha}_i \frac{\sqrt{M_{22i}^p}}{\hat{\sigma}_i},$$

where  $\hat{\sigma}_i^2 = \frac{1}{T}(M_{11i}^p - \hat{\alpha}_i M_{12i}^p)$  and  $\hat{\alpha}_i = M_{12i}^p / M_{22i}^p$ , which again suppresses the dependence upon  $\ell$ . Note also that in this setup  $M_{abi}^p$  takes the projection onto the lags of  $\Delta y_{it}$  rather than onto a vector of deterministic components as in Section 3.

Under  $H_0$ , using  $M_{12i}^e$  and  $M_{13i}^e$  to denote  $M_{12i}$  and  $M_{13i}$  with  $e_{it}$  in place of  $\Delta y_{it}$ ,

$$\begin{aligned} \frac{1}{T} M_{12i}^{ep} &= \frac{1}{T} M_{12i}^e - \frac{1}{T} M_{13i}^e M_{33i}^{-1} M_{32i} = \frac{1}{T} M_{12i}^e + O_p\left(\frac{1}{\sqrt{T}}\right), \\ \frac{1}{T^2} M_{22i}^p &= \frac{1}{T^2} M_{22i}^e - \frac{1}{T^2} M_{23i}^e M_{33i}^{-1} M_{32i} = \frac{1}{T^2} M_{22i}^e + O_p\left(\frac{1}{T}\right). \end{aligned}$$

These results, together with  $\frac{1}{T} M_{11i}^p \xrightarrow{p} \sigma^2$  and

$$\begin{pmatrix} \frac{1}{T} M_{12i}^e \\ \frac{1}{T^2} M_{22i}^e \end{pmatrix} \Rightarrow \begin{pmatrix} \sigma \omega_i M_{12i}^o \\ \omega_i^2 M_{22i}^o \end{pmatrix}$$

imply that

$$\tau_i = \frac{M_{12i}^p}{\hat{\sigma}_i \sqrt{M_{22i}^p}} \Rightarrow \frac{M_{12i}^o}{\sqrt{M_{22i}^o}}.$$

Thus, the asymptotic distribution of  $\tau_i$  is not affected by the presence of short-run dynamics, suggesting that the distribution of the IPS statistic should be unaffected too, see Section 4 of IPS. In other words, with this test lag augmentation successfully removes the short-run dynamics of the panel. However, this is not true for the LLC statistic. Note first that if there are no deterministic components, then as  $N, T \rightarrow \infty$

$$\frac{1}{T\sqrt{N}} \sum_{i=1}^N M_{12i}^p \xrightarrow{d} \frac{\sigma \sqrt{\bar{\omega}^2}}{\sqrt{2}} \mathcal{N}(0, 1),$$

where  $\bar{\omega}^2 = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \omega_i^2$ . Furthermore,

$$\frac{1}{NT^2} M_{22}^p \xrightarrow{p} \frac{\bar{\omega}^2}{2},$$

and therefore,

$$\tau_{LLC} = \frac{M_{12}^p}{\hat{\sigma} \sqrt{M_{22}^p}} \xrightarrow{d} \mathcal{N}(0, 1).$$

Thus, in this case lag augmentation removes the effect of the short-run dynamics also for the LLC statistic. However, the situation changes dramatically if the model includes a constant or a linear time trend. Consider for example the case with short-run dynamics and a constant, in which we let  $x_{it} = (1, \Delta y_{it-1}, \dots, \Delta y_{it-\ell})'$  and re-define  $M_{12i}^p$  and  $M_{22i}^p$  accordingly. This yields

$$\lim_{N, T \rightarrow \infty} E \left( \frac{1}{TN} M_{12}^p \right) = \sigma \bar{\omega} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N E(M_{12i}^o) \rightarrow -\frac{\sigma \bar{\omega}}{2},$$

where  $\bar{\omega} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \omega_i$ . It follows that in this case lag augmentation does not remove short-run parameters from the mean of the statistic. To cope with this problem LLC propose a bias and serial correlation corrected version of  $\tau_{LLC}$ , which we again denote by  $\tau_{LLC}^c$ .<sup>8</sup> The problem is that since the mean of  $\tau_{LLC}$  depends on both  $\sigma^2$  and  $\omega_i^2$ , for the bias-correction to work these parameters have to be estimated consistently, an issue that will be considered in more detail below. It follows that in the presence of deterministic terms lag augmentation alone is not enough to remove the short-run parameters from the asymptotic distribution of the LLC statistic.

<sup>8</sup>Note that because  $\sigma^2$  is assumed to be equal across  $i$ , there is no need for any correction for heteroskedasticity as in LLC.

For the estimation of  $\omega_i^2$  LLC propose using

$$\hat{\omega}_i^2 = \frac{1}{T} \sum_{t=2}^T (\Delta y_{it})^2 + \frac{2}{T} \sum_{j=1}^{q-1} \left(1 - \frac{j}{q}\right) \sum_{t=j+1}^T \Delta y_{it} \Delta y_{it-j},$$

which is the conventional Newey and West (1987) long-run variance estimator. It is important to note that by using  $\Delta y_{it}$  this estimator is in fact imposing  $H_0$ . Thus, if  $H_0$  holds then we have from Andrews (1991) that  $\hat{\omega}_i^2 \xrightarrow{p} \omega_i^2$  as  $T \rightarrow \infty$  with  $q \rightarrow \infty$  and  $\frac{q}{T} \rightarrow 0$ . This indicates that the following bias-corrected statistic can be used:

$$\tau_{LLC}^c = \sqrt{2} \tau_{LLC} + \frac{NT}{\sqrt{2}} \frac{\hat{\omega}}{\sqrt{M_{22}^p}}$$

with  $\hat{\omega} = \frac{1}{N} \sum_{i=1}^N \hat{\omega}_i$ , whose asymptotic distribution is given by

$$\tau_{LLC}^c = \frac{\sqrt{2N} \left( \frac{1}{TN} M_{12}^p + \frac{1}{2} \hat{\sigma} \hat{\omega} \right)}{\hat{\sigma} \sqrt{\frac{1}{NT^2} M_{22}^p}} \xrightarrow{d} \frac{\frac{\sigma \sqrt{\omega^2}}{\sqrt{6}} \mathcal{N}(0, 1)}{\sigma \sqrt{\frac{\omega^2}{6}}} \sim \mathcal{N}(0, 1),$$

see Theorem 5 of LLC. However, although theoretically not an issue as long as  $q \rightarrow \infty$  and  $\frac{q}{T} \rightarrow 0$ , in practice the optimal truncation lag  $q$  unknown. There is also the problem that  $\hat{\omega}_i^2$  tends to zero under  $H_{1a}$ , an issue that we will discuss in more detail in the next section.

To sidestep these difficulties, Breitung and Das (2005) propose to pre-whiten the variables. Their idea is as follows. Under  $H_0$  we have

$$\Delta y_{it} = \sum_{j=1}^{\ell} \phi_{ij} \Delta y_{it-j} + e_{it}, \quad (11)$$

or, in terms of levels,

$$y_{it} = \sum_{s=1}^t \Delta y_{is} = \sum_{j=1}^{\ell} \phi_{ij} \sum_{s=1}^t \Delta y_{is-j} + \sum_{s=1}^t e_{is} = \sum_{j=1}^{\ell} \phi_{ij} y_{it-j} + y_{it}^s,$$

where  $y_{it}^s$  is as in (2) but with  $H_0$  imposed. It is a random walk with serially uncorrelated increments  $e_{it}$ . We also assume that  $y_{i0} = 0$ . Thus, in contrast to  $\frac{1}{\sqrt{T}} y_{it}$ , whose long-run variance is given by  $\omega_i^2$ , the long-run variance of  $\frac{1}{\sqrt{T}} y_{it}^s$  is just  $\sigma^2$ . For the estimation of the lag coefficients  $\phi_{ij}$ , Breitung and

Das (2005) recommend using the above regression in first differences with  $H_0$  imposed. For a fixed  $\ell$  this yields

$$\frac{1}{\sqrt{T}} y_{it}^* = \frac{1}{\sqrt{T}} y_{it}^s - \sum_{j=1}^{\ell} (\hat{\phi}_{ij} - \phi_{ij}) \frac{1}{\sqrt{T}} y_{it-j} = \frac{1}{\sqrt{T}} y_{it}^s + o_p\left(\frac{1}{\sqrt{T}}\right) o_p(1),$$

where  $\hat{\phi}_{ij}$  is the LS estimate of  $\phi_{ij}$  in (11), which means that  $\sqrt{T}(\hat{\phi}_{ij} - \phi_{ij}) = O_p(1)$ . Similarly,

$$\Delta y_{it}^* = e_{it} - \sum_{j=1}^{\ell} (\hat{\phi}_{ij} - \phi_{ij}) \Delta y_{it-j} = e_{it} + o_p\left(\frac{1}{\sqrt{T}}\right).$$

Thus, replacing  $\Delta y_{it}$  and  $y_{it-1}$  by  $\Delta y_{it}^*$  and  $y_{it-1}^*$ , respectively, eliminates the effects of the serial correlation without requiring any estimation of  $\omega_i^2$ .

In Table 4 we compare the size accuracy of the IPS and LLC tests using both LS lag augmentation and pre-whitening. The data are generated from

**TABLE 4** Size for different corrections for short-run dynamics

		$\phi_1 = 0.5$				$\phi_1 = -0.5$			
		LLC		IPS		LLC		IPS	
$N$	$T$	Aug	Pre	Aug	Pre	Aug	Pre	Aug	Pre
No lags									
10	20	7.5	7.5	3.8	3.8	13.4	13.4	21.7	21.7
20	20	5.2	5.2	3.1	3.1	22.4	22.4	31.7	31.7
40	20	2.7	2.7	0.6	0.6	45.1	45.1	54.4	54.4
10	50	4.8	4.8	1.7	1.7	29.2	29.2	42.0	42.0
20	50	2.8	2.8	0.4	0.4	50.0	50.0	67.5	67.5
40	50	0.7	0.7	0.1	0.1	82.9	82.9	90.6	90.6
The true number of lags									
10	20	8.9	7.1	5.6	3.5	8.4	7.6	4.5	4.0
20	20	8.9	5.8	3.5	2.7	7.6	7.2	3.2	3.0
40	20	8.2	6.1	2.0	0.9	6.0	5.7	1.7	1.3
10	50	7.5	6.7	3.6	2.6	6.5	6.0	3.5	3.4
20	50	6.9	5.5	3.5	2.8	6.2	5.8	3.6	3.5
40	50	6.1	5.3	2.5	2.0	5.2	5.2	2.8	2.7
The Schwarz Bayesian information criterion									
10	20	10.0	7.3	6.0	3.2	8.1	8.0	4.8	4.2
20	20	9.3	6.8	4.3	2.8	7.9	7.1	3.5	3.0
40	20	8.0	5.9	2.1	0.9	6.0	5.3	2.1	1.5
10	50	7.5	6.7	3.6	2.6	6.4	6.1	3.4	3.4
20	50	6.9	5.6	3.6	2.8	6.3	5.8	3.6	3.4
40	50	5.9	5.2	2.5	1.7	5.2	5.3	2.9	2.7

The table reports the 5% rejection frequencies under  $H_0$ .  $\phi_1$  refers to the first-order AR serial correlation parameter. Aug and Pre refer to LS augmentation and pre-whitening, respectively.

(1), (2) and (9) with  $\ell = 1$ , which makes  $\phi_1$ , the assumed homogenous first-order AR coefficient, an interesting nuisance parameter to study. For the choice of lag length we consider three alternatives. The first is to ignore the serial correlation and to set  $\ell = 0$ , while the second is to set  $\ell$  equal to its true value. The third alternative is to choose  $\ell$  in a data-dependent fashion by using the Schwarz Bayesian information criterion with a maximum of five lags.

The first thing to notice is the size distortions that result from ignoring the serial correlation, especially when  $\phi_1 = -0.5$ , and the effectiveness by which they are removed in the two correction procedures. Note also that there are basically no differences in the results depending on whether  $\ell$  is treated as known or not. Similarly, unreported simulation results suggest that there is no difference in power. The point here is that since the performance is so similar, why not use pre-whitening, which leads to a much simpler test?

### Fact 8: The Consistency of the LLC Test Depends on the Long-Run Variance Estimator

As long as that  $q \rightarrow \infty$  such that  $\frac{q}{T} \rightarrow 0$ , we have under  $H_0$ ,  $\hat{\omega}_i^2 - \omega_i^2 = o_p(1)$ . Thus, provided that  $H_0$  holds,  $\hat{\omega}_i$  is consistent for  $\omega_i$ , which as we have seen is a requirement for  $\tau_{LLC}^c$  to be asymptotically normal. The problem is under the fixed alternative hypothesis  $H_{1a}$   $y_{it}$  is stationary, and therefore  $\Delta y_{it}$  is over-differenced with a zero long-run variance. Thus, in contrast to what happens under  $H_0$ , in this case  $\hat{\omega}_i^2$  does not converge to  $\omega_i^2$  but in fact goes to zero suggesting that  $\hat{\omega}_i$  should go to zero too. In fact, as Westerlund (2010) shows, if  $q \rightarrow \infty$  with  $N, T \rightarrow \infty$ , and  $\frac{q}{T} \rightarrow 0$ ,

$$\hat{\omega} = \frac{1}{N} \sum_{i=1}^N \hat{\omega}_i = O_p\left(\frac{1}{\sqrt{q}}\right).$$

From Section 2 we know that  $\tau_{LLC} = O_p(\sqrt{NT})$  under  $H_{1a}$ , suggesting that  $\tau_{LLC}^c$  is of the same order. Therefore, to determine the effect of the inconsistency of  $\hat{\omega}$  on

$$\tau_{LLC}^c = \sqrt{2} \tau_{LLC} + \frac{NT}{\sqrt{2}} \frac{\hat{\omega}}{\sqrt{M_{22}^p}},$$

we only need to consider the second term, the bias, which under  $H_{1a}$  can be written as

$$\frac{NT}{\sqrt{2}} \frac{\hat{\omega}}{\sqrt{M_{22}^p}} = \frac{\sqrt{NT}}{\sqrt{2}} \hat{\omega} \frac{1}{\sqrt{\frac{1}{NT} M_{22}^p}} = \sqrt{NT} O_p\left(\frac{1}{\sqrt{q}}\right) O_p(1).$$

In order to appreciate the implications of this last result, suppose that  $q$  is set independent of  $T$ , so that the bias term is  $O_p(\sqrt{NT})$ . The problem here is that while  $\tau_{LLC} \rightarrow -\infty$ , the bias term is diverging at the same rate but in the opposite direction, which means that  $\tau_{LLC}^c$  need not be a consistent test. The only way to prevent this from happening is therefore to set  $q$  as a function of  $T$ , ensuring that the order of the bias term is lower than  $O_p(\sqrt{NT})$ , and hence that  $\tau_{LLC}^c \rightarrow -\infty$ . In practice, however, the fact that  $q$  should be allowed to increase with  $T$  offers little guidance, because there are many choices that are consistent with a particular expansion rate. There are so-called pug-in rules that facilitate a data-dependent choice of  $q$ , but these are subject to estimation error. Hence, plug-in rules may not work, in the sense that estimated value of  $q$  may not be large enough for the test to have power. A similar problem arises when trying to use the residual  $\hat{\varepsilon}_{it} = \Delta y_{it} - \hat{\alpha}_i y_{it-1}$  instead of  $\Delta y_{it}$ , which would otherwise seem like a natural thing to do in view of the inconsistency of  $\hat{\omega}_i^2$  under the stationary alternative. However, this means that under the alternative  $\hat{\omega}_i^2 \xrightarrow{p} \omega_i^2 > 0$ , which again leads to consistency problems.

To illustrate these results Table 5 reports the size-adjusted power of  $\tau_{LLC}^c$  for three different bandwidth selection rules. The first is the automatic rule of Newey and West (1994), while the other two are deterministic, and involve setting  $q$  either equal to  $4(T/100)^{2/9}$  as suggested by Newey and West (1987) or equal to  $3.21T^{1/3}$  as in LLC. The data are

**TABLE 5** Size-adjusted power of the LLC test for different bandwidths

$c$	$N$	$T$	Bandwidth selection rule		
			NW	$4(T/100)^{2/9}$	$3.21T^{1/3}$
-5	10	100	3.9	3.9	6.2
	20	100	3.6	3.5	5.0
	10	200	1.9	1.6	3.5
	20	200	2.5	2.5	2.8
-10	10	100	3.9	3.6	10.8
	20	100	3.7	3.7	7.9
	10	200	2.1	1.3	5.6
	20	200	2.7	1.8	3.9
-20	10	100	7.2	6.0	38.5
	20	100	6.8	6.8	29.7
	10	200	5.2	1.6	20.6
	20	200	3.4	2.0	11.6
-40	10	100	40.4	39.7	86.0
	20	100	27.3	28.3	85.2
	10	200	31.4	8.9	73.3
	20	200	18.5	4.2	65.2

The table reports the 5% size-adjusted rejection frequencies when the AR parameter is set to  $\alpha_i = c_i/T\sqrt{N}$ , where  $c_i \sim U(2c, 0)$ . NW refers to the automatic bandwidth selection rule of Newey and West (1994).



generated from (2), (4) and (5) with  $c_i \sim U(2c, 0)$ . In agreement with the results of Westerlund (2010) we see that the power can be very low and practically nonexistent unless  $q = 3.21T^{1/3}$ , which is also the most generous rule considered.<sup>9</sup> For example, if  $T = 100$ , then  $4(T/100)^{2/9} = 4$  while  $3.21T^{1/3} \simeq 15$ , an increase by almost a factor of four. For some simulation results of the size of the test we make reference to Westerlund (2010).

## 5. CROSS-SECTION DEPENDENCE

### Fact 9: Cross-Section Dependence Leads to Deceptive Inference

We consider two types of dependence, weak and strong. The first type refers to a situation in which all the eigenvalues of the covariance matrix of  $y_{it}$  are bounded as  $N \rightarrow \infty$ , which rules out the presence of unobserved common factors, but allows the cross-sectional units to be for example spatially correlated. This is the case considered by O'Connell (1998) in his pioneering work on the effects of cross-section dependence on panel unit root tests. The second type of dependence refers to a situation when at least one eigenvalue diverges with  $N$ , which arises when cross-section dependence is due to common factors.

Suppose as in the previous section that (3) holds so that

$$\begin{pmatrix} \Delta y_{1t} \\ \vdots \\ \Delta y_{Nt} \end{pmatrix} = \begin{pmatrix} \alpha_1 & & 0 \\ & \ddots & \\ 0 & & \alpha_N \end{pmatrix} \begin{pmatrix} y_{1t-1} \\ \vdots \\ y_{Nt-1} \end{pmatrix} + \begin{pmatrix} \varepsilon_{1t} \\ \vdots \\ \varepsilon_{Nt} \end{pmatrix}$$

or, in matrix format,

$$\Delta y_t = \Pi y_{t-1} + \varepsilon_t.$$

However, instead of looking at the case when  $\varepsilon_{it}$  is dependent across  $t$ , we now consider the case when it is dependent across  $i$ .

In particular, let us begin by assuming that all eigenvalues of the covariance matrix

$$\Omega = \text{cov}(\varepsilon_t) = E(\varepsilon_t \varepsilon_t')$$

are bounded as  $N \rightarrow \infty$ , which means that the dependence is of the weak form. By the spectral decomposition,  $\Omega = \Omega^{1/2}(\Omega^{1/2})' = V\Lambda V'$ , where  $\Lambda$  is the diagonal matrix with the ordered eigenvalues  $\lambda_1 \geq \dots \geq \lambda_N$  along

<sup>9</sup>Interestingly, even with the most generous choice of bandwidth the power is relatively poor, see Moon et al. (2007) for some simulation evidence of existing tests.

the diagonal, while  $V$  is the matrix of orthonormal eigenvectors. Let  $y_t^* = (y_{1t}^*, \dots, y_{Nt}^*)' = \Omega^{-1/2} y_t$ , which under  $H_0$  is nothing but a vector of uncorrelated random walks.

The above assumptions imply that  $M_{12}$  can be written as

$$M_{12} = \sum_{t=2}^T y'_{t-1} \Delta y_t = \sum_{t=2}^T (y_{t-1}^*)' \Lambda \Delta y_t^* = \sum_{i=1}^N \lambda_i M_{12i}^*,$$

where  $M_{12i}^* = \sum_{t=2}^T y_{it-1}^* \Delta y_{it}^*$ . Let  $\bar{\lambda} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \lambda_i$  with a similar definition of  $\bar{\lambda}^2$ . By using the results of the previous sections,

$$\frac{1}{NT} M_{12} \xrightarrow{p} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \lambda_i \frac{1}{T} E(M_{12i}^*) = \bar{\lambda} \frac{1}{T} E(M_{12i}^*) = 0$$

such that  $\frac{1}{T\sqrt{N}} M_{12} \xrightarrow{d} \frac{\sqrt{\lambda^2}}{\sqrt{2}} \mathcal{N}(0, 1)$  as  $N, T \rightarrow \infty$ . Similarly,

$$\frac{1}{NT^2} M_{22} = \frac{1}{NT^2} \sum_{t=2}^T y'_{t-1} y_{t-1} = \frac{1}{NT^2} \sum_{i=1}^N \lambda_i M_{22i}^* \xrightarrow{p} \bar{\lambda} \frac{1}{T^2} E(M_{22i}^*) \rightarrow \frac{\bar{\lambda}}{2},$$

where the limit is taken as  $N \rightarrow \infty$  followed by  $T \rightarrow \infty$ . This result, together with  $\hat{\sigma}^2 \xrightarrow{p} \bar{\lambda}$ , suggest that as  $N, T \rightarrow \infty$

$$\tau_{LLC} = \frac{\frac{1}{T\sqrt{N}} M_{12}}{\hat{\sigma} \sqrt{\frac{1}{NT^2} M_{22}}} \xrightarrow{d} \frac{\frac{\sqrt{\lambda^2}}{\sqrt{2}} \mathcal{N}(0, 1)}{\sqrt{\frac{\bar{\lambda}}{2}}} \sim \frac{\sqrt{\lambda^2}}{\bar{\lambda}} \mathcal{N}(0, 1),$$

which summarizes Theorem 1 of Breitung and Das (2005). In other words, if the dependence is weak then  $\tau_{LLC}$  is still asymptotically normal. However, as long as  $\lambda_i \neq \lambda_j$  for at least some  $i \neq j$ ,  $\frac{\sqrt{\lambda^2}}{\bar{\lambda}} > 1$  and so the variance will tend to increase with deceptive inference as a result. A similar result applies to  $\tau_{IPS}$ . That is, the IPS test will also tend to be misleading in the presence of weak cross-section dependence.

In order to analyze the effects of strong dependence suppose that the error  $\varepsilon_{it}$  in (3) has the following common factor representation:

$$\varepsilon_{it} = \theta_i g_t + \epsilon_{it}. \quad (12)$$

This representation is used by Phillips and Sul (2003), Moon and Phillips (2004), Moon et al. (2007), and Pesaran (2007).

Assuming that  $\epsilon_{it}$  and  $g_t$  are white noise processes with unit variance, the covariance matrix of  $\varepsilon_{it}$  can be written as

$$\Omega = \Theta\Theta' + I_N,$$

where  $\Theta = (\theta_1, \dots, \theta_N)'$ . The main difference here in comparison to the case with weak dependence is the presence of  $g_t$ , which suggests that the largest eigenvalue of  $\Omega$  is no longer bounded but is in fact  $O(N)$ , while the remaining eigenvalues are  $O(1)$ . Intuitively, the information regarding the common component  $\Theta g_t$  accumulates as we sum up the observations across  $i$  and therefore the largest eigenvalue will increase with  $N$ . It follows that

$$\begin{aligned} \frac{1}{NT} M_{12} &= \frac{\lambda_1}{NT} \sum_{t=2}^T y_{1t-1}^* \Delta y_{1t}^* + o_p\left(\frac{1}{\sqrt{N}}\right), \\ \frac{1}{NT^2} M_{22} &= \frac{\lambda_1}{NT^2} \sum_{t=2}^T (y_{1t-1}^*)^2 + o_p\left(\frac{1}{\sqrt{N}}\right) \end{aligned}$$

and therefore,

$$\frac{1}{\sqrt{N}} \tau_{LLC} \Rightarrow \frac{\int_0^1 W(s) dW(s)}{\sqrt{\int_0^1 W(s)^2 ds}},$$

where  $W(s)$  is a standard Brownian motion that does not depend on  $i$ . This is the limiting distribution of the DF statistic, see Breitung and Das (2008). Thus,  $\tau_{LLC} = O_p(\sqrt{N})$ , and moreover,

$$\begin{aligned} \lim_{N, T \rightarrow \infty} P(\tau_{LLC} < -1.645) &= \lim_{N, T \rightarrow \infty} P\left(\frac{1}{\sqrt{N}} \tau_{LLC} < -\frac{1.645}{\sqrt{N}}\right) \\ &= P\left(\int_0^1 W(s) dW(s) < 0\right) \simeq 0.7, \end{aligned}$$

showing that the LLC test is upwards biased in the case of strong cross-section dependence.

Let us now consider the case when there is a stationary factor that enters the AR process in the following fashion:

$$y_{it} = \theta_i f_t + y_{it}^s, \quad (13)$$

where  $y_{it}^s$  is as in (2), while the common factor  $f_t$  is assumed to satisfy  $f_t = \delta f_{t-1} + u_t$  with  $|\delta| < 1$  and  $u_t$  is white noise. Assuming  $\rho_i = \rho$  for all  $i$ , (12) can be written in terms of (13) as follows:

$$y_{it} = \theta_i(\rho f_{t-1} + g_t) + (\rho y_{it-1}^s + \epsilon_{it}) = \theta_i f_t + y_{it}^s.$$

It follows that if  $\rho = 1$ , then both components are non-stationary. Conversely, if  $|\rho| < 1$ , then both components are stationary. Thus, by using (12) the common and idiosyncratic components of the above model are restricted to have the same order of integration. Note also that if  $\rho_i$  is heterogeneous the above model cannot be expressed in terms of (13).

Under  $H_0$ ,

$$\Delta y_t = \Theta \Delta f_t + \Delta y_t^s = \Theta((1 - \delta)f_t + u_t) + \varepsilon_t,$$

suggesting that

$$\text{cov}(\Delta y_t) = \frac{(1 - \delta)^2}{1 - \delta^2} \Theta \Theta' + \sigma^2 I_N.$$

To see how this is going to affect the limiting distribution of the LLC test write

$$\frac{1}{NT} M_{12} = \frac{1}{NT} \sum_{t=2}^T (f_{t-1} \Theta' \Theta \Delta f_t + (y_{t-1}^s)' \varepsilon_t + f_{t-1}' \Theta' \varepsilon_t + (y_{t-1}^s)' \Theta \Delta f_t),$$

where, since  $\varepsilon_{it}$  is assumed to be independent of  $\varepsilon_{jt}$  and  $f_t$ ,

$$\frac{1}{\sqrt{NT}} \sum_{t=2}^T f_{t-1}' \Theta' \varepsilon_t = \frac{1}{N} \sum_{i=1}^N \theta_i \frac{1}{\sqrt{T}} \sum_{t=2}^T f_{t-1} \varepsilon_{it} = O_p(1).$$

Moreover, since

$$\frac{1}{T\sqrt{N}} \sum_{t=2}^T (y_{t-1}^s)' \varepsilon_t = O_p(1),$$

where  $\frac{1}{T\sqrt{N}} \sum_{t=2}^T (y_{t-1}^s)' \Theta \Delta f_t$  is of the same order, we obtain

$$\frac{1}{NT} M_{12} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=2}^T \theta_i^2 f_{t-1} \Delta f_t + O_p\left(\frac{1}{\sqrt{N}}\right) \xrightarrow{p} -\frac{1 - \delta}{1 - \delta^2} \bar{\theta}^2$$

as  $N, T \rightarrow \infty$ , where  $\bar{\theta}^2 = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \theta_i^2$  (Breitung and Das, 2008).

Also,

$$\begin{aligned} \frac{1}{NT^2} M_{22} &= \frac{1}{NT^2} \sum_{t=2}^T (\Theta f_{t-1} + y_{t-1}^s)' (\Theta f_{t-1} + y_{t-1}^s) \\ &= \frac{1}{NT^2} \sum_{t=2}^T (y_{t-1}^s)' y_{t-1}^s + O_p\left(\frac{1}{T}\right) + O_p\left(\frac{1}{T\sqrt{N}}\right) \xrightarrow{p} \frac{\sigma^2}{2}. \end{aligned}$$

By collecting these results we obtain

$$\frac{1}{\sqrt{N}} \tau_{LLC} \xrightarrow{p} -\frac{\frac{1-\delta}{1-\delta^2} \overline{\theta^2}}{\sigma^2/\sqrt{2}},$$

or  $\tau_{LLC} = O_p(\sqrt{N})$ , suggesting that the size of the LLC test will tend to one as  $N \rightarrow \infty$ .

As for the IPS test, note that

$$\begin{aligned} \frac{1}{T} M_{12i} &= \frac{1}{T} \sum_{t=2}^T y_{it-1} \Delta y_{it} = \frac{1}{T} \sum_{t=2}^T (\theta_i f_{t-1} + y_{it-1}^s) (\theta_i \Delta f_t + \varepsilon_{it}) \\ &= \frac{1}{T} \sum_{t=2}^T y_{it-1}^s \varepsilon_{it} + \theta_i \frac{1}{T} \sum_{t=2}^T y_{it-1}^s \Delta f_t + \theta_i^2 \frac{1}{T} \sum_{t=2}^T f_{t-1} \Delta f_t + O_p\left(\frac{1}{\sqrt{T}}\right), \end{aligned}$$

while

$$\hat{\sigma}_i^2 \frac{1}{T^2} M_{22i} = \hat{\sigma}_i^2 \frac{1}{T^2} \sum_{t=2}^T y_{it-1}^2 = \hat{\sigma}_i^2 \frac{1}{T^2} \sum_{t=2}^T (\theta_i f_{t-1} + y_{it-1}^s)^2 = Q_i + O_p\left(\frac{1}{T}\right),$$

where  $Q_i = \sigma^2 \frac{1}{T^2} \sum_{t=2}^T (y_{it-1}^s)^2$ . Hence,

$$\tau_i = \frac{M_{12i}}{\hat{\sigma}_i \sqrt{M_{22i}}} = \tau_i^s + \frac{\theta_i}{\sqrt{Q_i}} \frac{1}{T} \sum_{t=2}^T y_{it-1}^s \Delta f_t + \frac{\theta_i^2}{\sqrt{Q_i}} \frac{1}{T} \sum_{t=2}^T f_{t-1} \Delta f_t + O_p\left(\frac{1}{\sqrt{T}}\right),$$

where  $\tau_i^s$  is the DF test based on  $y_{it}^s$ . Using the law of iterated expectations with respect to the sigma field generated by  $f_t$ ,  $\mathcal{F}$  say, and passing  $T \rightarrow \infty$ , we get

$$\begin{aligned} E(\tau_i | \mathcal{F}) &= E(\tau_i^s) + \frac{1}{T} \sum_{t=2}^T E\left(\frac{\theta_i}{\sqrt{Q_i}} y_{it-1}^s | \mathcal{F}\right) \Delta f_t \\ &\quad + E\left(\frac{\theta_i^2}{\sqrt{Q_i}} | \mathcal{F}\right) \frac{1}{T} \sum_{t=2}^T f_{t-1} \Delta f_t + O_p\left(\frac{1}{\sqrt{T}}\right) \end{aligned}$$

and therefore,  $E(\tau_i) \neq E(\tau_i^s)$ . In other words, the usual correction terms for the mean of  $\tau_i$  are incorrect as they ignore the correlation between  $f_{t-1}$  and  $\Delta f_t$ . It follows that under  $H_0$  the IPS statistic is  $O_p(\sqrt{N})$ .

Summarizing the results reported in this section we find that the presence of cross-section dependence is likely to lead to misleading inference. The extreme case being when the dependence is of the strong form, in which the test statistics actually become divergent. This last

**TABLE 6** Size in the presence of a stationary common factor.

$\theta$	$N$	$T$	LLC	IPS
0	10	50	7.4	4.3
	20	50	6.1	3.3
	10	100	6.7	4.0
	20	100	6.1	4.6
1	10	50	29.7	41.4
	20	50	53.2	64.1
	10	100	47.3	59.9
	20	100	73.3	84.5
2	10	50	92.0	95.0
	20	50	99.4	99.8
	10	100	99.1	99.7
	20	100	100.0	100.0

The table reports the 5% rejection frequencies under  $H_0$ .  $\theta$  refers to the factor loading which is identical for all  $i$ .

result is particularly interesting since in our setup  $f_t$  is stationary, and the presence of unit roots usually eliminates the effects of such variables. This is illustrated in Table 6, which depicts the size of the LLC and IPS tests in the presence of a single common factor. For simplicity the data are generated from (2) and (13) with no deterministic components or serial correlation. The factor is generated with AR coefficient  $\delta = 0$  and loading  $\theta_i = \theta$  for all  $i$ . The results show that the distortions are increasing in  $N$ , which is clearly in agreement with the theoretical predictions.

The lesson we learn from there results is that since cross-section dependence is more of a rule rather than the exception, some form of correction is usually needed, see Breitung and Pesaran (2008) and the references therein.

### **Fact 10: The IPS and LLC Tests Fail Under Cross-Unit Cointegration**

Suppose that (13) holds, and that  $\rho_i = \rho$  for all  $i$  with  $|\rho| < 1$  such that  $y_{it}^s$  is stationary. Suppose also that  $\delta = 1$ , so that  $f_t$  is unit root non-stationary. Under these conditions,  $y_{it}$  and  $y_{jt}$  are cointegrated, a situation commonly referred to as cross-unit cointegration.

If we assume that  $\theta_i \neq 0$  for all  $i$ , the vector  $y_t$  has a unit root and therefore a panel unit root test should accept the null hypothesis that all  $N$  units have a unit root. However, it is not difficult to show that in this case the LLC and IPS tests are going to be oversized, leading to the wrong conclusion.

Consider first the LLC test. By using

$$\begin{aligned}\frac{1}{NT} M_{12} &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=2}^T \theta_i^2 f_{t-1} \Delta f_t + \frac{1}{NT} \sum_{i=1}^N \sum_{t=2}^T y_{it-1}^s \Delta y_{it} + o_p\left(\frac{1}{\sqrt{N}}\right) \\ &\Rightarrow \overline{\theta^2} \int_0^1 W(s) dW(s) - \frac{1-\rho}{1-\rho^2} \sigma^2, \\ \frac{1}{NT^2} M_{22} &= \frac{1}{NT^2} \sum_{t=2}^T (\Theta f_{t-1} + y_{t-1}^s)' (\Theta f_{t-1} + y_{t-1}^s) \\ &= \frac{1}{NT^2} \sum_{i=1}^N \sum_{t=2}^T \theta_i^2 f_{t-1}^2 + o_p\left(\frac{1}{\sqrt{N}}\right) \Rightarrow \overline{\theta^2} \int_0^1 W(s)^2 ds,\end{aligned}$$

it follows that

$$\frac{1}{\sqrt{N}} \tau_{LLC} \Rightarrow \frac{\overline{\theta^2} \int_0^1 W(s) dW(s) - \frac{1-\rho}{1-\rho^2} \sigma^2}{\sqrt{\overline{\theta^2} \int_0^1 W(s)^2 ds}}$$

or  $\tau_{LLC} = O_p(\sqrt{N})$ , showing that the size of the LLC test is tending to one as  $N \rightarrow \infty$ .

The analysis of the IPS test is very similar to the case when  $f_t$  what white noise. We begin by noting that

$$\begin{aligned}\frac{1}{T} M_{12i} &= \frac{1}{T} \sum_{t=2}^T (\theta_i f_{t-1} + y_{it-1}^s)(\theta_i u_t + \Delta y_{it}^s) \\ &= \theta_i^2 \frac{1}{T} \sum_{t=2}^T f_{t-1} u_t + \theta_i \frac{1}{T} \sum_{t=2}^T f_{t-1} \Delta y_{it}^s + \frac{1}{T} \sum_{t=2}^T y_{it-1}^s \Delta y_{it}^s + o_p\left(\frac{1}{\sqrt{T}}\right), \\ \hat{\sigma}_i^2 \frac{1}{T^2} M_{22i} &= \hat{\sigma}_i^2 \frac{1}{T^2} \sum_{t=2}^T (\theta_i f_{t-1} + y_{it-1}^s)^2 = V_i + o_p\left(\frac{1}{T}\right),\end{aligned}$$

where  $V_i = \sigma_i^2 \frac{1}{T^2} \sum_{t=2}^T \theta_i^2 f_{t-1}^2$ . Thus, using  $\tau^f$  to denote the DF test based on  $f_t$ ,

$$\tau_i = \tau^f + \frac{\theta_i}{\sqrt{V_i}} \frac{1}{T} \sum_{t=2}^T f_{t-1} \Delta y_{it}^s + \frac{1}{\sqrt{V_i}} \frac{1}{T} \sum_{t=2}^T y_{it-1}^s \Delta y_{it}^s + o_p\left(\frac{1}{\sqrt{T}}\right),$$

whose asymptotic distribution as  $T \rightarrow \infty$  is clearly different from the usual DF distribution, as the second and third terms on the right-hand side are  $O_p(1)$ . The IPS statistic is therefore  $O_p(\sqrt{N})$ .

Consequently, the presence of cross-unit cointegrating relationships causes the IPS and LLC statistics to become divergent, which is in agreement with the simulation results of Banerjee et al. (2004), showing that the presence of such relationships can lead to severe size distortions. One way to alleviate this problem in practice is to use the PANIC approach of Bai and Ng (2004), where separate tests are employed for the common and idiosyncratic components.

### Fact 11: Sequential Limits Need not Imply Joint Limits

As pointed out by Phillips and Moon (1999), the sequential limit theory wherein  $T \rightarrow \infty$  is followed by  $N \rightarrow \infty$ , is very straightforward to apply and generally leads to quick asymptotic results in a variety of settings. The main reason for this is that passing  $T \rightarrow \infty$  while holding  $N$  fixed in the first step allows one to focus only on the first-order terms, as the higher order terms are eliminated prior to averaging over  $N$ . However, this feature can also be deceptive in its simplicity because it hides the need to control the relative expansion rate of the two dimensions. Indeed, as Phillips and Moon (1999) show, sequential convergence does not necessarily imply convergence in the joint limit as  $N, T \rightarrow \infty$  simultaneously. In some situations the sequential limit theory may therefore break down. The problem is that the connection between the two methods is not well understood, and many researchers view breakdowns as extreme events.

Consider as an example the generalized LS test of Breitung and Das (2005) and Harvey and Bates (2005),

$$\tau_{GLS} = \frac{\sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} \Delta y_t}{\sqrt{\sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} y_{t-1}}},$$

which is suitable for testing  $H_0$  in the presence of weak cross-section dependence. Here  $y_t$  is again the vector stacking  $y_{it}$ , while  $\hat{\Omega}$  is such that  $\sqrt{T}(\hat{\Omega} - \Omega) = O_p(1)$ .

We begin by deriving the sequential limit distribution of  $\tau_{GLS}$ , and then we show that this distribution need not be the same as the one obtained when using joint limits.

Applying a Taylor expansion yields

$$\hat{\Omega}^{-1} = \Omega^{-1} + O_p\left(\frac{1}{\sqrt{T}}\right).$$

Moreover, by a functional central limit theorem,  $\frac{1}{\sqrt{T}} y_t \Rightarrow B(s)$  as  $T \rightarrow \infty$ , where  $B(s) = \Omega^{1/2} W(s)$  and  $W(s)$  is now a  $N$ -dimensional vector standard



Brownian motion. It follows that as  $T \rightarrow \infty$  with  $N$  kept fixed,

$$\tau_{GLS} = \frac{\frac{1}{T} \sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} \Delta y_t}{\sqrt{\frac{1}{T^2} \sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} y_{t-1}}} \Rightarrow \frac{\int_0^1 B(s)' \Omega^{-1} dB(s)}{\sqrt{\int_0^1 B(s)' \Omega^{-1} B(s) ds}} = \frac{\int_0^1 W(s)' dW(s)}{\sqrt{\int_0^1 W(s)' W(s) ds}}.$$

But the elements of  $W(s)$  are independent, suggesting that as  $N \rightarrow \infty$

$$\frac{\int_0^1 W(s)' dW(s)}{\sqrt{\int_0^1 W(s)' W(s) ds}} = \frac{\frac{1}{\sqrt{N}} \int_0^1 W(s)' dW(s)}{\sqrt{\frac{1}{N} \int_0^1 W(s)' W(s) ds}} \xrightarrow{d} \mathcal{N}(0, 1).$$

Thus, in the sequential limit as  $T \rightarrow \infty$  and then  $N \rightarrow \infty$

$$\tau_{GLS} \xrightarrow{d} \mathcal{N}(0, 1).$$

Consider next the joint limit of the same test statistic. By using the results of Breitung and Das (2005),

$$\begin{aligned} \frac{1}{T\sqrt{N}} \sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} \Delta y_t &= \frac{1}{T\sqrt{N}} \sum_{t=2}^T y'_{t-1} \Omega^{-1} \Delta y_t + o_p\left(\frac{N}{\sqrt{T}}\right), \\ \frac{1}{NT^2} \sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} y_{t-1} &= \frac{1}{NT^2} \sum_{t=2}^T y'_{t-1} \Omega^{-1} y_{t-1} + o_p\left(\frac{\sqrt{N}}{\sqrt{T}}\right) \end{aligned}$$

as  $N, T \rightarrow \infty$ , from which it follows that

$$\tau_{GLS} = \frac{\frac{1}{T\sqrt{N}} \sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} \Delta y_t}{\sqrt{\frac{1}{NT^2} \sum_{t=2}^T y'_{t-1} \hat{\Omega}^{-1} y_{t-1}}} = \frac{\frac{1}{T\sqrt{N}} \sum_{t=2}^T y'_{t-1} \Omega^{-1} \Delta y_t}{\sqrt{\frac{1}{NT^2} \sum_{t=2}^T y'_{t-1} \Omega^{-1} y_{t-1}}} + o_p\left(\frac{N}{\sqrt{T}}\right)$$

Using the previous results we obtain

$$\frac{\frac{1}{T\sqrt{N}} \sum_{t=2}^T (y_{t-1}^*)' \Delta y_t^*}{\sqrt{\frac{1}{NT^2} \sum_{t=2}^T (y_{t-1}^*)' y_{t-1}^*}} \xrightarrow{d} \mathcal{N}(0, 1).$$

Only if we assume that  $\frac{N}{\sqrt{T}} \rightarrow 0$  as  $N, T \rightarrow \infty$  we end up with the same asymptotic distribution resulting from a sequential limit theory. In other words,  $T$  must be large relative to  $N^2$  to achieve a reliable approximation by the standard normal distribution. This makes sense even from an empirical point of view, as  $\hat{\Omega}$  is singular unless  $T \geq N$ , a fact not accounted for when using the sequential limit theory. It also explains the poor small sample properties of the test if  $T$  is small relative to  $N^2$ , as documented by Breitung and Das (2005).

Of course, this example is still quite specific, which makes it difficult to appreciate the generality of the critique. Let us therefore consider another example. In particular, let us reconsider the limiting null distribution of  $\tau_{IPS}$  under the assumption that  $\varepsilon_{it}$  is normal, independent and identically distributed, in which case we know from Phillips (1987) that

$$\frac{1}{\sqrt{T}} y_{it} = \sigma W_i(s) + O_p\left(\frac{1}{T}\right).$$

Using a Taylor series expansion and  $\sqrt{T}(\hat{\sigma}_i^2 - \sigma^2) = O_p(1)$  we obtain


$$\tau_i = \frac{M_{12i}}{\hat{\sigma}_i \sqrt{M_{22i}}} = \frac{M_{12i}^\circ}{\sqrt{M_{22i}^\circ}} + O_p\left(\frac{1}{T}\right),$$

implying that


$$\begin{aligned} \tau_{IPS} &= \frac{\sqrt{N}(\bar{\tau} - E(\tau))}{\sqrt{\text{var}(\tau)}} = \frac{1}{\sqrt{N \text{var}(\tau)}} \sum_{i=1}^N (\tau_i - E(\tau)) \\ &= \frac{1}{\sqrt{N \text{var}(\tau)}} \sum_{i=1}^N \left( \frac{M_{12i}^\circ}{\sqrt{M_{22i}^\circ}} - E(\tau) \right) + O_p\left(\frac{\sqrt{N}}{T}\right), \end{aligned}$$

where the first term on the right-hand side converges to a standard normal distribution as  $N \rightarrow \infty$ . Thus, for  $\tau_{IPS}$  to be asymptotically normal, one needs to assume that  $\frac{\sqrt{N}}{T} \rightarrow 0$  as  $N, T \rightarrow \infty$ , as otherwise the  $O_p(\sqrt{N}/T)$  remainder will not disappear (see also IPS 2003). A similar result applies to  $\tau_{LLC}$ . The point here is that if we use the sequential limit method where  $N$  is treated as fixed in the first step then this remainder is  $O_p(1/T)$ , which vanishes as  $T \rightarrow \infty$ . The sequential limit method therefore breaks down unless  $\frac{\sqrt{N}}{T} \rightarrow 0$ . But if this result holds in the current very restrictive case with normal innovations, it is expected to hold also under more relaxed conditions. In practice this means that in general one needs  $T \gg N$  for these tests to work properly.

Figures 4 and 5 illustrate this point by plotting the size of the LLC and IPS tests at the 5% level when  $T$  is set as a function of  $N$ . If  $T = N$  the condition that  $\frac{\sqrt{N}}{T} \rightarrow 0$  is satisfied, while if  $T = \sqrt{N}$ , then the condition is violated. The model includes a constant but otherwise there are no nuisance parameters to correct for. In contrast to the case when  $T = N$ , in which both tests tend to perform well, we see that setting  $T = \sqrt{N}$  generally leads to substantial size distortions, especially for the IPS test.



**FIGURE 4** Size of  $\tau_{LLC}$  when  $T$  is set as a function of  $N$ . (Figure available in color online.)



**FIGURE 5** Size of  $\tau_{IPS}$  when  $T$  is set as a function of  $N$ . (Figure available in color online.)

## 6. CONCLUDING REMARKS

This paper points to a number of facts that are oftentimes overlooked when considering the problem of testing for a unit root in panel data. Suppose for example that one would like to test the null hypothesis that

the variable  $y_{it}$  has a unit root versus the alternative that it is stationary with a nonzero mean, a very common research scenario. The by far most common way of carrying out such a test is to use the traditional DF approach of applying LS to an intercept-augmented autoregression. Being so common it is easy to get the impression that demeaning in this way is the best way to accommodate the nonzero mean in  $y_{it}$ . But this is not true. Indeed, as we show in the paper the inclusion of the intercept introduces a bias in the estimated AR coefficient, which then has to be corrected somehow. However, in so doing we find that the resulting corrected IPS and LLC tests are likely to suffer from low power and may even become inconsistent in some circumstances. As a response to this a few alternative demeaning procedures are suggested.

In this example, although applying traditional time series techniques leads to a larger computational burden and poorer small-sample performance, usually there are no fundamental shortcomings or flawed inference. Unfortunately, this is not always the case. Quite on the contrary, we find that ignoring these facts will in many cases lead to serious side-effects, including a complete breakdown of the whole test procedure. One example of such a situation is when  $y_{it}$  is contaminated by cross-section dependence in the form of common factors, in which case a failure to account for these factors can cause the test statistic to become divergent.

The implication is that one should be careful not to approach the testing problem from a too narrow and stylized perspective. In particular, we believe that the usual practice of looking at the problem from mainly a time series perspective can be deceptive in its simplicity, typically ignoring many important issues such as cross-sectional dependence, incidental trends and joint limit restrictions.

## ACKNOWLEDGEMENTS

Previous versions of this paper were presented at the 5th Nordic Econometric Meeting in Lund and seminars in Amsterdam, Athens, Gothenburg, Maastricht, Paris, and Thessaloniki. The authors would like to thank conference and seminar participants, and in particular Uwe Hassler, Jean-Pierre Urbain, Franz Palm, and four anonymous referees for helpful comments and suggestions. Thank you also to the Jan Wallander and Tom Hedelius Foundation for financial support under research grant number W2006-0068:1.

## REFERENCES

- Anderson, T. W., Hsiao, C. (1982). Formulation and estimation of dynamic models using panel data. *Journal of Econometrics* 76:47–82.
- Andrews, D. W. K. (1991). Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica* 59:817–858.

- Arellano, M., Bond, S. (1991). Tests of specification for panel data: Monte Carlo evidence and an application to employment equations. *Review of Economic Studies* 58:277–297.
- Arellano, M., Bover, O. (1995). Another look at the instrumental variable estimation of error-component models. *Journal of Econometrics* 68:29–51.
- Arellano, M., Hahn, J. (2007). Understanding bias in nonlinear panel models: some recent developments. In: Blundell, R., Newey, W., Persson, T., (Eds.) *Advances in Economics and Econometrics*. Ninth World Congress. Vol. III. Cambridge, UK: Cambridge University Press, pp. 381–409.
- Bai, J., Ng, S. (2004). A panic attack on unit roots and cointegration. *Econometrica* 72:1127–1177.
- Bai, J., Kao, C. (2006). On the estimation and inference of a panel cointegration model with cross-sectional dependence. In: Baltagi, B. (Ed.) *Contributions to Economic Analysis*. Cambridge, UK: Elsevier, pp. 3–30.
- Baltagi, B., Kao, C. (2000). Nonstationary panels, cointegration in panels, and dynamic panels: A survey. *Advances in Econometrics* 15:7–52.
- Banerjee, A. (1999). Panel data unit roots and cointegration: An overview. *Oxford Bulletin of Economics and Statistics* 61:607–630.
- Banerjee, A., Marcellino, M., Osbat, C. (2004). Testing for PPP: Should we use panel methods? *Empirical Economics* 30:77–91.
- Blundell, R., Bond, S. (1998). Initial conditions and moment restrictions in dynamic panel data models. *Journal of Econometrics* 87:115–143.
- Bond, S., Nauges, C., Windmeijer, F. (2005). *Unit Roots: Identification and Testing in Micro Panels*. CeMMAP Working Paper, London.
- Bowman, D. (1999). Efficient Tests for Autoregressive Unit Roots in Panel Data. International Finance Discussion Paper 646, Board of Governors of the Federal Reserve System.
- Breitung, J. (1997). Testing for unit roots in panel data using a gmm approach. *Statistical Papers* 38:253–269.
- Breitung, J. (2000). The local power of some unit root tests for panel data. In: Baltagi, B., (Ed.) *Advances in Econometrics: Nonstationary Panels, Panel Cointegration, and Dynamic Panels*. 15: 161–178. Amsterdam: JAI.
- Breitung, J., Meyer, W. (1994). Testing for unit roots in panel data: are wages on different bargaining levels cointegrated? *Applied Economics* 26:353–361.
- Breitung, J., Das, S. (2005). Panel unit root tests under cross sectional dependence. *Statistica Neerlandica* 59:414–433.
- Breitung, J., Das, S. (2008). Testing for unit roots in panels with a factor structure. *Econometric Theory* 24:88–108.
- Breitung, J., Pesaran, M. H. (2008). Unit roots and cointegration in panels. In: Mátyás, L., Sevestre, P. (Eds.) *The Econometrics of Panel Data*. Springer, pp. 279–322.
- Choi, I. (2001). Unit root tests for panel data. *Journal of International Money and Finance* 20:249–272.
- Choi, I. (2006). Nonstationary panels. *Palgrave Handbooks of Econometrics*. 1:511–539. New York: Palgrave Macmillan.
- Dickey, D. A., Fuller, W. A. (1979). Distribution of the estimates for autoregressive time series with a unit root. *Journal of the American Statistical Association* 74:427–431.
- Elliott, G., Rothenberg, T. J., Stock, J. H. (1996). Efficient tests for an autoregressive unit root. *Econometrica* 64:813–836.
- Hahn, J., Kuersteiner, G. (2002). Asymptotically unbiased inference for a dynamic panel model with fixed effects when both  $N$  and  $T$  are large. *Econometrica* 70:1639–57.
- Han, C., Phillips, P. C. B. (2010). GMM estimation for dynamic panels with fixed effects and strong instruments at unity. *Econometric Theory* 26:119–151.
- Harvey, A., Bates, D. (2005). *Multivariate Unit Root Tests, Stability and Convergence*. DAE Working Paper No. 301, University of Cambridge.
- Harris, R. D. F., Tzavalis, E. (1999). Inference for unit roots in dynamic panels where the time dimension is fixed. *Journal of Econometrics* 91:201–226.
- Harris, D., Harvey, D. I., Leybourne, S. J., Sakkas, N. D. (2010). local asymptotic power of the im-pesaran-shin panel unit root test and the impact of initial observations. *Econometric Theory* 26:311–324.
- Im, K. S., Pesaran, M. H., Shin, Y. (2003). Testing for unit roots in heterogeneous panels. *Journal of Econometrics* 115:53–74.

- Kruiniger, H. (2008). Maximum likelihood estimation and inference methods for the covariance stationary panel AR(1)/unit root model. *Journal of Econometrics* 144:447–464.
- Kruiniger, H. (2009). GMM estimation and inference in dynamic panel data models with persistent data. *Econometric Theory* 25:1348–1391.
- Levin, A., Lin, C. (1992). *Unit Root Tests in Panel Data: Asymptotic and Finite-sample Properties*. University of California working paper, pp. 92–23.
- Levin, A., Lin, C., Chu, C.-J. (2002). Unit root tests in panel data: Asymptotic and finite-sample properties. *Journal of Econometrics* 108:1–24.
- Lopez, C. (2009). A panel unit root tests with good power in small samples. *Econometric Reviews* 28:295–313.
- Madsen, E. (2010). Unit Root inference in panel data models where the time-series dimension is fixed: A comparison of different tests. *Econometrics Journal* 13:63–94.
- Moon, H. R., Perron, B. (2004). Testing for unit root in panels with dynamic factors. *Journal of Econometrics* 122:81–126.
- Moon, H. R., Perron, B. (2008). Asymptotic local power of pooled *t*-ratio tests for unit roots in panels with fixed effects. *Econometrics Journal* 11:80–104.
- Moon, H. R., Phillips, P. C. B. (1999). Maximum likelihood estimation in panels with incidental trends. *Oxford Bulletin of Economics and Statistics* 61:771–748.
- Moon, H. R., Phillips, P. C. B. (2004). GMM estimation of autoregressive roots near unity with panel data. *Econometrica* 72:467–522.
- Moon, H. R., Perron, B., Phillips, P. C. B. (2006). On the breitung test for panel unit roots and local asymptotic power. *Econometric Theory* 22:1179–1190.
- Moon, H. R., Perron, B., Phillips, P. C. B. (2007). Incidental trends and the power of panel unit root tests. *Journal of Econometrics* 141:416–459.
- Newey, W. K., West, K. D. (1987). A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica* 55:703–708.
- Newey, W. K., West, K. D. (1994). Autocovariance lag selection in covariance matrix estimation. *Review of Economic Studies* 61:613–653.
- Nickell, S. (1981). Biases in dynamic models with fixed effects. *Econometrica* 49:1417–1426.
- Pesaran, M. H. (2007). A simple panel unit root test in the presence of cross section dependence. *Journal of Applied Econometrics* 22:265–312.
- Phillips, P. C. B. (1987). Time series regression with a unit root. *Econometrica* 55:277–301.
- Phillips, P. C. B., Moon, H. R. (1999). Linear regression limit theory of nonstationary panel data. *Econometrica* 67:1057–1111.
- Phillips, P. C. B., Sul, D. (2003). Dynamic panel estimation and homogeneity testing under cross section dependence. *Econometrics Journal* 6:217–259.
- O'Connell, P. G. J. (1998). The overvaluation of purchasing power parity. *Journal of International Economics* 44:1–19.
- Ploberger, W., Phillips, P. C. B. (2002). *Optimal Testing for Unit Roots in Panel Data*. Unpublished manuscript.
- Quah, D. (1994). Exploiting cross-section variations for unit root inference in dynamic panels. *Economics Letters* 44:9–19.
- Schmidt, P., Phillips, P. C. B. (1992). LM tests for a unit root in the presence of deterministic trends. *Oxford Bulletin of Economics and Statistics* 54:257–287.
- Westerlund, J. (2010). A note on the use of the LLC panel unit root test. *Empirical Economics* 37: 517–531.