

TCSS 343 - Assignment 3

Jake McKenzie

July 17, 2018

In this problem use the Master Theorem to find and prove tight bounds for these recurrences (6 points each).

To solve this problem I used the master theorem taken from CLRS. I used a bit of a stronger statement than what was stated in the theorem but they are equivalent mathematically due to the nature of limits. I will include the theorem for the reader's benefit. To check my work I decided to include what I obtained using the Akra-Bazzi method.

Master Theorem

The Master Theorem applies to recurrences of the following form:

$$T(n) = aT(n/b) + f(n)$$

where $a \geq 1$ and $b > 1$ are constants and $f(n)$ is an asymptotically positive function.

There are 3 cases:

1. If $f(n) = O(n^{\log_b a - \epsilon})$ for some constant $\epsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$.
2. If $f(n) = \Theta(n^{\log_b a} \log^k n)$ with $k \geq 0$, then $T(n) = \Theta(n^{\log_b a} \log^{k+1} n)$.
3. If $f(n) = \Omega(n^{\log_b a + \epsilon})$ with $\epsilon > 0$, and $f(n)$ satisfies the regularity condition, then $T(n) = \Theta(f(n))$.

Regularity condition: $af(n/b) \leq cf(n)$ for some constant $c < 1$ and all sufficiently large n .

1.

$$T(n) = \begin{cases} c & \text{if } n < 8 \\ 16T(n/8) + n \log n & \text{if } n \geq 8 \end{cases}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{n \log n}{n^{\log_8 16 + \epsilon}} &= \lim_{n \rightarrow \infty} \frac{n \log n}{n^{\frac{4}{3} + \frac{2}{3}}} \\ &= \lim_{n \rightarrow \infty} \left(\frac{n}{n}\right) \left(\frac{\log n}{n}\right) \rightarrow 0 \\ T(n) &\in \Theta(n^{\frac{4}{3}}) \end{aligned}$$

By Akra-Bazzi I obtain:

$$\begin{aligned} 16(1/8)^p &= 1 \\ p &= \frac{4}{3} \end{aligned}$$

$$\begin{aligned}
T(n) &\in \Theta(n^p(1 + \int_1^n \frac{u \log u}{u^{p+1}} du)) \\
&\in \Theta(-9n + 10n^{\frac{4}{3}} - 3n \log n) \\
&\in \Theta(n^{\frac{4}{3}})
\end{aligned}$$

2.

$$T(n) = \begin{cases} c & \text{if } n < 4 \\ 8T(n/4) + n \log n & \text{if } n \geq 4 \end{cases}$$

By Master Method I obtain:

$$\begin{aligned}
\lim_{n \rightarrow \infty} \frac{n^{\frac{1}{3}}}{n^{\log_8 2 + \varepsilon}} &= \lim_{n \rightarrow \infty} \frac{n^{\frac{1}{3}}}{n^{\frac{1}{3}}} \\
&= \lim_{n \rightarrow \infty} 1 \rightarrow 1 \\
&\in \Theta(n^{\frac{1}{3}} \log n)
\end{aligned}$$

By Akra-Bazzi I obtain:

$$\begin{aligned}
2(1/8)^p &= 1 \\
p &= \frac{1}{3}
\end{aligned}$$

$$\begin{aligned}
T(n) &\in \Theta(n^p(1 + \int_1^n \frac{u^{\frac{1}{3}}}{u^{p+1}} du)) \\
&\in \Theta(n^{\frac{1}{3}} + n^{\frac{1}{3}} \log n) \\
&\in \Theta(n^{\frac{1}{3}} \log n)
\end{aligned}$$