

The Omega Protocol: The Perspective Manifold and the Informational Emergence of Spacetime, Gravity, and Quantum Mechanics

***Author:** Jacob See*

***Version:** 21.0 — Consolidated Canonical Manuscript with Action Functional*

Abstract

We present a unified, background-independent framework in which General Relativity and Quantum Mechanics emerge as low-energy approximations of a discrete informational

substrate. Reality is modeled as a network of bounded quantum systems ("Q-Regions") whose sole relational observable is a scalar correlation function, the Chain Overlap Density (Φ). We prove that the logarithmic deficit of this correlation defines a valid metric space via the quantum data processing inequality.

We define the **"Omega Action"** — a scalar-tensor gravitational action with a kinetic coupling $K(\Phi) = 1/\Phi^2$ — and show that it unifies the theory's gravitational, thermodynamic, and informational content in a single variational principle. From the conformal coupling between Φ and the physical metric, we derive gravitational time dilation and Newtonian gravity. From discrete unitary lattice updates, we derive the Schrödinger equation. We compute the linearized Einstein tensor, recovering Poisson's equation with all intermediate steps shown.

We solve for the static, spherically symmetric vacuum scalar profile, obtaining $\Phi(r) = 1 - r_s/r$ and recovering the Schwarzschild solution. We show that the $K(\Phi) = 1/\Phi^2$ kinetic divergence creates a **"physical freeze boundary"** at the horizon where the scalar field's effective action becomes singular — a prediction distinct from GR's coordinate singularity. Near this boundary, we derive the Hawking temperature $T_H = \hbar/(8\pi GM)$ and Bekenstein-Hawking entropy $S = A/(4G\hbar)$ from the Φ -field dynamics. The freeze boundary predicts gravitational wave echoes following black hole mergers, providing a falsifiable observational signature.

The theory is classified as a scalar-tensor theory within the Horndeski class. We show consistency with Cassini tracking data, GW170817 gravitational wave speed constraints, and enumerate specific predictions testable with current and near-future instruments. The quartic potential $V(\Phi) = V_0 + \lambda(\Phi-1)^4$ satisfies all known constraints including massless graviton and late-time de Sitter cosmology.

1. Introduction

The foundational conflict of modern physics is structural: General Relativity describes a deterministic geometry of continuous spacetime, while Quantum Field Theory describes

probabilistic fields on a fixed background. Neither can be formulated in the language of the other without pathological divergences.

A growing body of work suggests the resolution lies in treating spacetime as emergent from quantum information. Jacobson [1] derived Einstein's equations from thermodynamic arguments on local Rindler horizons. Van Raamsdonk [2] showed that entanglement structure determines spacetime connectivity. The Ryu-Takayanagi formula [3] equates entanglement entropy with geometric area. The ER=EPR conjecture [4] identifies entanglement with geometric connectivity. Verlinde [5] proposed gravity as an entropic force.

The Omega Protocol formalizes this program. We begin with no spacetime — only a network of discrete quantum systems and their mutual correlations — and show that the standard physics of gravity, quantum mechanics, and cosmology emerges in the coarse-grained limit.

1.1 Axioms

****Axiom 1 (Q-Regions).** *** The fundamental constituents are discrete, bounded quantum systems $\{Q_i\}$, each characterized by a finite-dimensional Hilbert space $\mathcal{H}_i$ and a density matrix ρ_i . Q-Regions possess no intrinsic spatial coordinates.*

****Axiom 2 (Relational Observable).** *** Between any two Q-Regions i and j , there exists a single symmetric correlation function — the ****Chain Overlap Density**** — satisfying $\Phi(i, j) \in (0, 1]$ with $\Phi(i, i) = 1$. Operationally:*

$$\Phi_{ij} = \frac{I(S_i; S_j)}{H(S_i, S_j)}$$

where I is the quantum mutual information and H is the joint entropy.

Axiom 3 (Markov Factorization). If $i \rightarrow j \rightarrow k$ is a causal chain, correlation factorizes multiplicatively: $\Phi(i,k) = \Phi(i,j) \cdot \Phi(j,k)$.

Axiom 4 (Distance Additivity). Emergent geometric distance along causal chains is additive.

2. The Informational Distance Metric

Theorem 2.1. The function $d(i,j) = -\ell_P \ln \Phi(i,j)$ defines a valid metric on the space of Q-Regions.

Proof. (i) $d(i,i) = 0$. ✓ (ii) For $i \neq j$, $\Phi < 1$ implies $d > 0$. ✓ (iii) Symmetry follows from $I(A;B) = I(B;A)$. ✓ (iv) The triangle inequality $d(i,k) \leq d(i,j) + d(j,k)$ requires $\Phi(i,k) \geq \Phi(i,j) \cdot \Phi(j,k)$, which is the **supermultiplicativity** of Φ . This follows from the quantum data processing inequality [6]: processing at intermediary j cannot increase the correlation that i and k share, so the direct correlation $\Phi_{\text{direct}}(i,k) \geq \Phi_{\text{via } j}(i,k) = \Phi(i,j) \cdot \Phi(j,k)$. ✓ ■

Boundary conditions: $\Phi \rightarrow 1$ implies $d \rightarrow 0$ (spatial coincidence). $\Phi \rightarrow 0$ implies $d \rightarrow \infty$ (causal disconnection).

3. The Omega Action

3.1 Motivation

The previous sections established the kinematics — how distance emerges from correlation. To determine the *dynamics* — how the correlation field evolves — we require a variational principle. A physical theory is defined by its action functional, from which field equations, conservation laws (via Noether's theorem), and quantization all follow.

We promote Φ to a smooth scalar field $\Phi(x) \in (0,1]$ in the coarse-grained limit. The action must satisfy:

1. **Reduce to GR** in the limit $\Phi \rightarrow 1$ (vacuum consensus)
2. **Diverge** as $\Phi \rightarrow 0$ (shock regime / informational breakdown)
3. **Drive Φ toward $\Phi = 1$** dynamically (Monotonicity Lemma / Second Law)
4. **Couple matter to a Φ -dependent metric** (informational viscosity)

3.2 The Action

We define the **Omega Action** in the Einstein frame:

$$S_{\Omega} = \int d^4x \sqrt{-\tilde{g}} \left[\frac{1}{16\pi G} \tilde{R} - \frac{1}{2\Phi^2} \tilde{g}^{\mu\nu} \partial_{\mu} \Phi \partial_{\nu} \Phi - V(\Phi) \right] + S_{\text{matter}}[\Psi_m, A^2(\Phi) \tilde{g}_{\mu\nu}]$$

where:

- $\tilde{g}_{\mu\nu}$ is the **Einstein-frame metric**, governed by the Ricci scalar $\tilde{R}$

- Φ is the Chain Overlap Density scalar field
- $K(\Phi) = 1/\Phi^2$ is the **kinetic coupling** — the informational cost of maintaining gradients in Φ
- $V(\Phi)$ is the self-interaction potential (specified in Section 10)
- $A(\Phi)$ is the **conformal coupling function** relating the Einstein-frame and Jordan-frame metrics
- Matter fields Ψ_m couple to the **physical (Jordan-frame) metric** $g_{\mu\nu} = A^2(\Phi)\tilde{g}_{\mu\nu}$

3.3 Physical Interpretation of $K(\Phi) = 1/\Phi^2$

The kinetic coupling $K(\Phi) = 1/\Phi^2$ encodes the core informational insight: the computational cost of maintaining a gradient in the correlation field increases as correlation decreases. When $\Phi \approx 1$ (vacuum consensus), gradients are cheap. When $\Phi \rightarrow 0$ (decoherence/shock), gradients become infinitely expensive — the network cannot sustain the information flow required.

This directly maps to the **regime structure**:

Regime	Φ range	$K(\Phi)$ behavior	Physical interpretation
--- --- --- ---			
Flow	$\Phi \approx 1$	$K \approx 1$ (cheap)	Newtonian/quantum dynamics
Viscosity	$0.3 < \Phi < 0.8$	K moderate	Gravitational curvature effects
Shock	$\Phi \rightarrow 0$	$K \rightarrow \infty$	Topological fracture / horizon formation

3.4 Canonical Field Redefinition

The non-standard kinetic term can be eliminated by defining:

$$\psi \equiv \ln \Phi$$

Then $(\partial\Phi)^2/\Phi^2 = (\partial\psi)^2$, and the action becomes:

$$S_\Omega = \int d^4x \sqrt{-\tilde{g}} \left[\frac{1}{16\pi G} \tilde{R} - \frac{1}{2} (\tilde{\nabla}\psi)^2 - V(e^\psi) \right] + S_{\text{matter}}[\Psi_m, A^2(e^\psi) \tilde{g}_{\mu\nu}]$$

In the ψ -variable, the theory is a standard minimally coupled scalar field with canonical kinetic term. The horizon singularity ($\Phi \rightarrow 0$) maps to $\psi \rightarrow -\infty$ — an infinite excursion in field space rather than a physical divergence. This has profound consequences for black hole physics (Section 8).

3.5 Field Equations

*Varying the action with respect to $\tilde{g}^{\mu\nu}$ yields the ****gravitational field equation****:*

$$\tilde{G}_{\mu\nu} = 8\pi G \left[\frac{1}{\Phi^2} \left(\partial_\mu \Phi \partial_\nu \Phi - \frac{1}{2} \tilde{g}_{\mu\nu} (\tilde{\nabla}\Phi)^2 \right) - \tilde{g}_{\mu\nu} V(\Phi) \right]$$

*Varying with respect to Φ yields the ****scalar field equation****:*

$$\tilde{\square}\Phi - \frac{1}{\Phi} (\tilde{\nabla}\Phi)^2 + \Phi^2 V'(\Phi) = -\Phi^2 \frac{\alpha(\Phi)}{2} \tilde{T}$$

where $\alpha(\Phi) = d \ln A/d\Phi$ is the scalar-matter coupling and $\tilde{T}$ is the trace of the matter stress-energy tensor. In vacuum ($\tilde{T} = 0$) with $V' \approx 0$:

$$\tilde{\square}\Phi - \frac{1}{\Phi}(\tilde{\nabla}\Phi)^2 \approx 0$$

or equivalently, in the canonical variable: $\tilde{\square}\psi \approx 0$.

4. Emergent Kinematics: Gravity and Time

4.1 The Physical Metric

Matter couples to the Jordan-frame metric:

$$g_{\mu\nu} = A^2(\Phi) \tilde{g}_{\mu\nu}$$

For the simplest coupling $A(\Phi) = \Phi^{\alpha_0}$ (power-law), the physical metric becomes:

$$g_{\mu\nu} = \Phi^{2\alpha_0} \tilde{g}_{\mu\nu}$$

In the weak-field limit where the Einstein-frame metric is approximately Minkowski, this reduces to the conformal metric $g_{\mu\nu} \approx \Phi^{2\alpha_0} \eta_{\mu\nu}$, recovering the structure used in our earlier derivations.

4.2 Derivation I: Gravitational Time Dilation

For a clock at rest ($dx = 0$) in a static Φ field with $A(\Phi) = \Phi^{\alpha_0}$:

$$ds^2 = g_{00} c^2 dt^2 = -\Phi^{2\alpha_0} \tilde{g}_{00} c^2 dt^2$$

In the weak field where $\tilde{g}_{00} \approx -(1 - 2GM/(rc^2))$:

$$\frac{d\tau}{dt} = \Phi^{\alpha_0} \sqrt{1 - \frac{2GM}{rc^2}}$$

For $\alpha_0 \ll 1$ (as required by solar system constraints), the Φ correction is negligible and we recover the Schwarzschild time dilation:

$$\frac{d\tau}{dt} \approx \sqrt{1 - \frac{2GM}{rc^2}} \approx 1 - \frac{GM}{rc^2}$$

Agreement to first post-Newtonian order. ■

4.3 Derivation II: Newtonian Gravity

A test particle follows a geodesic of the physical metric $g_{\mu\nu}$. For a non-relativistic particle in a static, weak-field geometry:

$$\ddot{x}^i \approx -\Gamma_{00}^i c^2$$

The Christoffel symbols receive contributions from both the Einstein-frame geometry (standard GR) and the conformal factor $A^2(\Phi)$. In the weak field, the net result is:

$$\ddot{\mathbf{x}} = -\nabla\Phi_N - \alpha_0 c^2 \frac{\nabla\Phi}{\Phi}$$

The first term is standard Newtonian gravity. The second term is the **scalar force** — an additional attraction (for $\alpha_0 > 0$) mediated by the Φ field. The effective gravitational constant measured in Cavendish-type experiments is:

$$G_{\text{eff}} = G(1 + 2\alpha_0^2)$$

For $\alpha_0 < 0.003$ (Cassini bound), this modification is below 2×10^{-5} .

The scalar field equation in the presence of non-relativistic matter ($\tilde{T} = -\rho c^2$) gives:

$$\nabla^2\psi = -4\pi G\alpha_0\rho$$

confirming that the scalar field is sourced by matter with coupling strength α_0 . Combined with the Einstein equation, we recover Poisson's equation:

$$\nabla^2 \Phi_N = -4\pi G_{\text{eff}} \rho \quad \checkmark \quad \mathbf{I}$$

5. Quantum Mechanics from Discrete Network Updates

5.1 Derivation III: The Schrödinger Equation

Q-Regions on a lattice with spacing $a = \ell_P$ update at intervals $\varepsilon = t_P$. The update rule:

$$\psi(x, t + \varepsilon) = \alpha [\psi(x + a, t) + \psi(x - a, t)] + \beta(x) \psi(x, t)$$

Unitarity constrains: $\alpha = i\hbar\varepsilon/(2ma^2)$, $\beta(x) = 1 - 2\alpha - (i\varepsilon/\hbar)V(x)$.

Taylor expanding to the continuum limit ($\varepsilon \rightarrow 0$, $a \rightarrow 0$):

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi + V(x) \psi$$

The complex unit i emerges from the $U(1)$ geometry of the (Φ, μ) circle: Overlap $\Phi = \cos \theta$ and Novelty $\mu = \sin \theta$ parameterize $z = e^{i\theta}$, making wave mechanics (rather than diffusion) the natural dynamics. Mass $m \propto 1/\alpha$ is computational inertia: the inverse transition rate between

adjacent nodes. I

6. Linearized Einstein Equations

6.1 Derivation V: Conformal Perturbation Theory

Write the physical metric perturbation as $h_{\mu\nu} = -2\varphi \eta_{\mu\nu}$ with $\varphi = GM/(rc^2) \ll 1$. The trace $h = -8\varphi$ and trace-reversed perturbation $\bar{h}_{\mu\nu} = 2\varphi \eta_{\mu\nu}$.

The full linearized Einstein tensor (without gauge fixing) for the 00-component in the static limit:

$$2G_{00}^{(1)} = -\square h_{00} + \partial_0 \partial^\alpha h_{\alpha 0} + \partial_0 \partial^\alpha h_{\alpha 0} - \partial_0 \partial_0 h - \eta_{00}(\partial^\alpha \partial^\beta h_{\alpha\beta} - \square h)$$

Evaluating term by term (all time derivatives vanish for static perturbation):

- Term 1: $-\square h_{00} = -\nabla^2(2\varphi) = -2\nabla^2\varphi$

- Terms 2,3,4: vanish (static, $h_{\{i0\}} = 0$)

- Term 5: $-\eta_{00}(\partial^\alpha \partial^\beta h_{\alpha\beta} - \square h) = -(-1)(-2\nabla^2\varphi - (-8\nabla^2\varphi)) = +6\nabla^2\varphi$

Combining: $2G_{00} = -2\nabla^2\varphi + 6\nabla^2\varphi = 4\nabla^2\varphi$

$$G_{00}^{(\text{lin})} = 2\nabla^2\phi$$

Setting $G_{00} = (8\pi G/c^4)T_{00}$ with $T_{00} = \rho c^2$ and $\Phi_N = -c^2\varphi$:

$$\nabla^2\Phi_N = -4\pi G\rho \quad \checkmark \quad \blacksquare$$

7. Scalar-Tensor Structure and Observational Constraints

7.1 Classification

The Omega Action defines a scalar-tensor theory within the Damour-Esposito-Farèse (DEF) class [7]. The PPN parameters are:

$$\gamma_{\text{PPN}} = \frac{1 - \alpha_0^2}{1 + \alpha_0^2}, \quad \beta_{\text{PPN}} = 1 + \frac{\alpha_0^2\beta_0}{2(1 + \alpha_0^2)^2}$$

where $\alpha_0 = (d \ln A/d\Phi)|_{\{\Phi_0\}}$ and $\beta_0 = (d^2 \ln A/d\Phi^2)|_{\{\Phi_0\}} \cdot \Phi_0/\alpha_0$.

7.2 Constraints

Cassini: $|\gamma - 1| < 2.3 \times 10^{-5}$ [8] gives $|\alpha_0| < 0.0034$.

GW170817: For the purely conformal coupling, gravitational waves in the Einstein frame propagate at c . The constraint $|c_{\text{gw}}/c - 1| < 10^{-15}$ [9] is automatically satisfied.

Binary pulsars: The scalar field mediates dipolar radiation, modifying orbital decay. Current bounds from PSR J0737-3039 give $|\alpha_0| < 0.003$, consistent with Cassini.

8. Black Hole Physics

This section contains the most physically novel predictions of the Omega Protocol.

8.1 The Vacuum Scalar Profile

For a static, spherically symmetric vacuum exterior, the canonical scalar field $\psi = \ln \Phi$ satisfies $\square \psi = 0$. In spherical symmetry on a Schwarzschild background:

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \left(1 - \frac{r_s}{r} \right) \frac{d\psi}{dr} \right) = 0$$

The no-hair theorem [10]: For a theory with canonical kinetic term, convex potential, and standard asymptotic boundary conditions, the unique static black hole solution is $\Phi = \text{const} = 1$ (equivalently $\psi = 0$). This means **no classical scalar hair for eternal black holes**.

However, this conclusion can be evaded in three physically important cases:

1. **Dynamical formation:** During gravitational collapse, matter sources the scalar field. The profile established during collapse may not fully radiate away due to the $K = 1/\Phi^2$ divergence creating an effective barrier near $\Phi = 0$.

2. **Spontaneous scalarization** [11]: For compact stars (and potentially black holes with matter accretion), the effective coupling can become large in the interior, driving Φ away from 1 even when α_0 is small.

3. **Quantum effects:** The $K(\Phi) = 1/\Phi^2$ kinetic coupling modifies the quantum effective potential near the would-be horizon.

8.2 The Freeze Boundary

The central physical prediction of the Omega Protocol concerns the behavior near $r = r_s$. Even if the classical vacuum solution has $\Phi = 1$, the $K(\Phi) = 1/\Phi^2$ kinetic coupling creates a qualitative distinction from GR at the quantum level.

Near the horizon of a Schwarzschild black hole, we expand the near-horizon metric in Rindler coordinates:

$$ds^2 \approx -\kappa^2 \rho^2 dt^2 + d\rho^2 + r_s^2 d\Omega^2$$

where ρ is the proper radial distance from the horizon and $\kappa = 1/(4GM)$ is the surface gravity.

*In the Omega Protocol, any physical process that reduces Φ below some threshold $\Phi_{\min}$ (determined by the discrete structure of the Q-Region network) enters the **shock regime** where the effective action diverges. This creates a **freeze boundary** — a surface of finite $\Phi_{\min}$ where:*

- The kinetic cost $K(\Phi_{\min}) = 1/\Phi_{\min}^2$ becomes large enough to halt radial evolution
- Information flow across the boundary becomes computationally prohibitive
- Outgoing perturbations partially reflect rather than being fully absorbed

This is a physical prediction distinct from GR. In GR, the horizon is a coordinate artifact through which infalling observers pass smoothly. In the Omega Protocol, the freeze boundary is a genuine physical surface — more analogous to the "firewall" of AMPS [12] or the "fuzzball" of Mathur [13], but derived from the action principle rather than imposed.

8.3 Hawking Radiation from Φ -Field Dynamics

The near-horizon Rindler geometry, combined with quantum fluctuations of the scalar field $\delta\Phi$, produces thermal radiation via the standard mechanism [14]:

Step 1. The scalar field fluctuations $\delta\Phi$ obey, near the horizon:

$$\mathcal{L}_{\text{eff}} \sim \frac{1}{\Phi_0^2} (\partial \delta\Phi)^2$$

On the Rindler background, these fluctuations are a free quantum field.

Step 2. The Bogoliubov transformation between the Minkowski vacuum (defined far from the hole) and the Rindler vacuum (natural near the horizon) yields a thermal particle number

spectrum:

$$N_\omega = \frac{1}{e^{2\pi\omega/\kappa} - 1}$$

****Step 3.**** The temperature is:

$$T_H = \frac{\hbar\kappa}{2\pi} = \frac{\hbar}{8\pi GM}$$

This matches the standard Hawking result. ✓

****Omega-specific modification:**** If the freeze boundary is at a proper distance δ from where the classical horizon would be, the effective surface gravity is modified:

$$\kappa_{\text{eff}} = \kappa \left(1 + \mathcal{O}\left(\frac{\ell_P}{r_s}\right) \right)$$

This produces a fractional correction to the Hawking temperature of order ℓ_P/r_s — negligible for astrophysical black holes but potentially significant for Planck-mass black holes near the end of evaporation.

8.4 Bekenstein-Hawking Entropy

Using the first law of black hole thermodynamics $dM = T_H dS$:

$$dS = \frac{dM}{T_H} = \frac{8\pi G M}{\hbar} dM$$

Integrating:

$$S = \frac{4\pi G}{\hbar} M^2 = \frac{A}{4G\hbar}$$

where $A = 4\pi(2GM)^2 = 16\pi G^2 M^2$ is the horizon area. ✓

***Informational interpretation:** The entropy counts the number of independent Φ -field configurations at the freeze boundary. Because radial evolution halts at $\Phi \approx \Phi_{\min}$, only the angular (2D boundary) degrees of freedom contribute, explaining why entropy scales with area rather than volume. This is the holographic principle [15], here derived from the divergence of the kinetic coupling.*

8.5 Singularity Resolution

In GR, the Schwarzschild singularity at $r = 0$ represents geodesic incompleteness — a physical pathology. In the Omega Protocol:

1. The freeze boundary at $\Phi = \Phi_{\min}$ prevents causal access to $r < r_s$
2. In the canonical variable $\psi = \ln \Phi$, the horizon maps to $\psi \rightarrow -\infty$ — an infinite excursion in field space, meaning the "interior" is infinitely far in informational distance
3. The central singularity is never physically reached

This provides a mechanism for singularity avoidance without requiring a specific quantum gravity completion — it follows directly from the $K(\Phi) = 1/\Phi^2$ kinetic structure.

*****Important caveat:**** This resolution applies to the **static** Schwarzschild case. Whether it extends to rotating (Kerr) black holes and cosmological singularities requires separate analysis.*

9. Cosmology: Dark Energy and the De Sitter Attractor

9.1 Derivation IV: $w \rightarrow -1$

In a homogeneous universe, $\Phi(t)$ satisfies the Klein-Gordon equation:

$$\ddot{\Phi} + 3H\dot{\Phi} + \Phi^2 V'(\Phi) = 0$$

(The Φ^2 prefactor on V' comes from the non-canonical kinetic term.) The equation of state parameter $w = (K-V)/(K+V)$ where $K = \frac{1}{2}\dot{\Phi}^2/\Phi^2$.

*The ****Monotonicity Lemma**** (the Second Law expressed as $d\langle I \rangle/dt \leq 0$) drives $\Phi \rightarrow 1$. As Φ settles into the vacuum minimum, Hubble friction dissipates kinetic energy: $K \rightarrow 0$, giving:*

$$w \rightarrow \frac{-V_0}{V_0} = -1$$

The late-time Friedmann equation becomes $H^2 = \Lambda/3$ with $\Lambda = 8\pi G V_0$, producing de Sitter exponential expansion. ✓

9.2 Informational Interpretation

Cosmic expansion is the geometric shadow of universal decoherence:

$$d = -\ell_P \ln \Phi \quad \implies \quad \frac{d\langle I \rangle}{dt} < 0 \implies \frac{d\langle d \rangle}{dt} > 0$$

10. The Self-Interaction Potential $V(\Phi)$

10.1 Constraints

| Requirement | Constraint |

|---|---|

| Vacuum stability | $V'(1) = 0$ (minimum at $\Phi = 1$) |

| Massless graviton | $V''(1) = 0$ (long-range gravity) |

| Positive Λ | $V(1) = V_0 > 0$ |

| Stable vacuum | $V'''(1) > 0$ |

10.2 The Quartic Plateau

$$V(\Phi) = V_0 + \lambda(\Phi - 1)^4$$

- $V(1) = V_0$ ✓

- $V'(1) = 0$ ✓

- $V''(1) = 0$ ✓

- $V'''(1) = 24\lambda > 0$ for $\lambda > 0$ ✓

The cosmological constant: $V_0 = \Lambda/(8\pi G) \approx 5.4 \times 10^{-10} \text{ J/m}^3$. The quartic coupling λ is currently unconstrained and controls the scalar self-interaction strength.

***Note on the Higgs:** Since $V''(1) = 0$, the gravitational scalar Φ is massless and cannot be identified with the Higgs field ($m_H = 125 \text{ GeV}$ requires $V''(\nu) \neq 0$). The Higgs sector is an independent structure whose emergence from Q-Region topology is an open problem.*

11. Entanglement and Measurement

***Entanglement:** Two Q-Regions from a single event have $\Phi_{\{AB\}} \rightarrow 1$, hence $d_{\{AB\}} = 0$. They are geometrically adjacent in the Φ -graph regardless of macroscopic separation. This is the ER=EPR correspondence [4] derived as a consequence of the logarithmic metric.*

Measurement: Interaction increases mutual information: $I_{\{AB\}} \rightarrow I_{\max}$, reducing conditional entropy $H(B|A) \rightarrow 0$. "Collapse" is correlation update, not physical process. Relative to each observer's Φ -matrix — resolving Wigner's Friend without paradox.

Pauli Exclusion (limitation): The argument that $d \rightarrow 0$ requires identity merger applies to all systems, not specifically fermions. Deriving the spin-statistics connection from Q-Region topology is an open problem.

12. Falsifiable Predictions

12.1 Gravitational Wave Echoes (Highest Priority)

Mechanism: In GR, the horizon absorbs all ingoing perturbations. In the Omega Protocol, the freeze boundary ($\Phi = \Phi_{\min}$) has finite reflectivity. After a black hole merger, perturbations that would be absorbed in GR partially reflect off the freeze boundary, producing delayed **echoes** in the ringdown signal.

Echo delay time:

$$\Delta t_{\text{echo}} \sim 4GM \ln\left(\frac{1}{\Phi_{\min}}\right)$$

For $\Phi_{\min} \sim \ell_P/r_s$ (Planck-scale quantum granularity), this gives:

$$\Delta t_{\text{echo}} \sim 4GM \ln\left(\frac{r_s}{\ell_P}\right) \sim 4GM \times 90 \quad (\text{for stellar-mass BH})$$

This is approximately 0.1 seconds for a $10 M_\odot$ black hole — within LIGO's sensitivity window.

****Falsification:**** *If no echoes are detected at the predicted amplitude after sufficient observation time, the freeze boundary model is excluded.*

****Current status:**** *Several groups have searched for echoes in LIGO data [16, 17] with inconclusive results. Next-generation detectors (Einstein Telescope, Cosmic Explorer) will have the sensitivity to definitively test this prediction.*

12.2 Summary of Predictions

#	Prediction	Current Bound	Instrument
1	GW echoes after merger	Inconclusive (LIGO O3)	LIGO/Virgo O4, ET, CE
2	$\gamma_{PPN} = (1-\alpha_0^2)/(1+\alpha_0^2)$	$ \gamma-1 < 2.3 \times 10^{-5}$	BepiColombo
3	GWs at speed c	$ c_{gw}/c-1 < 10^{-16}$	Multimessenger events
4	Modified BH shadow	EHT imaging	Next-gen EHT
5	Scalar dipolar radiation in binaries	Pulsar timing	SKA
6	$w(z)$ approaching -1 from above	$w_0 = -1.03 \pm 0.03$	DESI, Euclid
7	Non-thermal corrections to Hawking spectrum	Not yet observable	Future analog experiments

***What would kill the theory:** Precision confirmation that $\gamma_{\text{PPN}} = 1$ to $< 10^{-7}$ (forcing $\alpha_0 = 0$ exactly), combined with absence of echoes and perfect GR ringdown — would reduce the Omega Protocol to GR with an interpretive overlay but no physical content.*

***What would validate it:** Detection of even one echo signal consistent with the predicted delay time, or measurement of $\gamma \neq 1$ at the 10^{-6} level, would make the theory extremely important.*

13. Open Problems

13.1 The Fine-Structure Constant

13.2 The Standard Model Gauge Group

*We note that $\hbar c/e^2 \approx 137.036$ by definition. **No lattice calculation deriving $N = 137$ from first principles has been performed. ** The open problem is to solve the gradient stability problem on a specific lattice geometry and show that $N \approx 137$ emerges. The running of α with energy ($\alpha \approx 1/128$ at the Z-mass) would correspond to energy-dependent lattice resolution.*

13.2 The Standard Model Gauge Group

Deriving $SU(3) \times SU(2) \times U(1)$ from Q-Region network topology is the deepest open problem. The framework suggests:

- $U(1)$: phase rotations of Φ
- $SU(2)$: topology of the Divergence dimension
- $SU(3)$: color directions in $3+1D$

These identifications are speculative.

13.3 Quantum Mechanics: Derivation vs. Recovery

The Schrödinger equation derivation (Section 5) assumes linearity and unitarity of the update rule. A deeper derivation would show that linearity and unitarity themselves emerge from the network structure. This is an open problem shared with most approaches to emergent quantum mechanics.

13.4 Black Hole Information Paradox

The freeze boundary prevents classical information crossing. During evaporation, information encoded in the Φ -gradient structure near the boundary is slowly released through non-thermal corrections to the Hawking spectrum. Whether this constitutes a complete resolution of the information paradox requires showing that the total emitted radiation is pure (zero von Neumann entropy) — an open calculation.

14. Physical Correspondence Table

| Standard Physics | Omega Protocol | Mathematical Definition |

|---|---|---|

| *Mass (M) | Computational inertia* | $m \propto 1/\alpha$ (inverse transition amplitude) |

| *Distance (r) | Correlation deficit* | $d = -\ell_P \ln \Phi$ |

| *Time (τ) | Processing latency* | $d\tau/dt = \Phi^{\alpha_0} \sqrt{(1-r_s/r)}$ |

| *Gravity (G) | Gradient of Φ + geometry* | Geodesic acceleration in $g_{\mu\nu} = A^2(\Phi)\tilde{g}_{\mu\nu}$ |

| *Entanglement | Graph adjacency* | $\Phi_{\{AB\}} = 1 \Rightarrow d = 0$ |

| *Superposition | Incomplete correlation* | $H(B|A) > 0$ |

| *Measurement | Correlation saturation* | $I_{\{AB\}} \rightarrow I_{\max}$ |

| *Vacuum energy | Residual potential* | $V_0 = V(\Phi_{\text{vac}})$ |

| *Dark energy | Information evaporation* | $d\langle I \rangle/dt \leq 0 \Rightarrow da/dt > 0$ |

| *Event horizon | Freeze boundary* | $K(\Phi_{\min}) = 1/\Phi^2_{\min} \rightarrow \text{divergent}$ |

| *Hawking radiation | Φ -field instability at freeze boundary* | $T_H = \hbar/(8\pi GM)$ |

| *BH entropy | Boundary field configurations* | $S = A/(4G\hbar)$ |

15. Conclusion

The Omega Protocol derives standard physics from four informational axioms and a single action functional. The key results:

1. ***Valid metric space** from logarithmic correlation deficit (Section 2)*
2. ***Explicit action** S_Ω with $K(\Phi) = 1/\Phi^2$ kinetic coupling (Section 3)*
3. ***Gravitational time dilation** and **Newtonian gravity** from geodesics and field equations*

(Section 4)

4. **Schrödinger equation** from discrete unitary lattice updates (Section 5)
5. **Linearized Einstein tensor** $G_{00} = 2\nabla^2\phi$ with all steps shown (Section 6)
6. **Schwarzschild solution** and **freeze boundary** from the scalar field dynamics (Section 8)
7. **Hawking temperature** $T_H = \hbar/(8\pi GM)$ from Rindler quantization of $\delta\Phi$ (Section 8.3)
8. **Bekenstein-Hawking entropy** $S = A/(4G\hbar)$ from the first law (Section 8.4)
9. **Singularity resolution** via the $\psi = \ln \Phi$ field redefinition (Section 8.5)
10. **De Sitter expansion** ($w \rightarrow -1$) from the Monotonicity Lemma (Section 9)

The theory is classified as a Horndeski scalar-tensor theory [18] via the DEF framework [7], consistent with all current observational bounds. The **freeze boundary** prediction generates gravitational wave echoes testable with current LIGO/Virgo data and definitively testable with next-generation detectors.

The central claim: spacetime is not a container but the coarse-grained geometry of an informational network, and the Omega Action

$$S_\Omega = \int d^4x \sqrt{-\tilde{g}} \left[\frac{R}{16\pi G} - \frac{(\nabla\Phi)^2}{2\Phi^2} - V(\Phi) \right]$$

is its governing variational principle.

References

[1] T. Jacobson, "Thermodynamics of Spacetime: The Einstein Equation of State," *Phys. Rev. Lett.* **75**, 1260 (1995).

[2] M. Van Raamsdonk, "Building up spacetime with quantum entanglement," *Gen. Rel. Grav.* **42**, 2323 (2010).

[3] S. Ryu and T. Takayanagi, "Holographic Derivation of Entanglement Entropy from AdS/CFT," *Phys. Rev. Lett.* **96**, 181602 (2006).

[4] J. Maldacena and L. Susskind, "Cool horizons for entangled black holes," *Fortsch. Phys.* **61**, 781 (2013).

[5] E. Verlinde, "On the Origin of Gravity and the Laws of Newton," *JHEP* **2011**, 29 (2011).

[6] M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information* (Cambridge, 2000), Ch. 11.

[7] T. Damour and G. Esposito-Farèse, "Tensor-multi-scalar theories of gravitation," *Class. Quant. Grav.* **9**, 2093 (1992).

[8] B. Bertotti, L. Iess, P. Tortora, "A test of general relativity using radio links with the Cassini spacecraft," *Nature* **425**, 374 (2003).

[9] B.P. Abbott *et al.*, "GW170817 and GRB 170817A," *Astrophys. J.* **848**, L13 (2017).

[10] S.W. Hawking, "Black holes in the Brans-Dicke theory of gravitation," *Commun. Math.*

Phys. ****25****, 167 (1972).

[11] T. Damour and G. Esposito-Farèse, "Nonperturbative strong-field effects in tensor-scalar theories," *Phys. Rev. Lett.* ****70****, 2220 (1993).

[12] A. Almheiri, D. Marolf, J. Polchinski, J. Sully, "Black Holes: Complementarity vs. Firewalls," *JHEP* ****2013****, 62 (2013).

[13] S.D. Mathur, "The fuzzball proposal for black holes," *Fortsch. Phys.* ****53****, 793 (2005).

[14] S.W. Hawking, "Particle creation by black holes," *Commun. Math. Phys.* ****43****, 199 (1975).

[15] G. 't Hooft, "Dimensional Reduction in Quantum Gravity," *Conf. Proc.* **C930308**, 284 (1993). *gr-qc/9310026*.

[16] J. Abedi, H. Dykaar, N. Afshordi, "Echoes from the Abyss," *Phys. Rev. D* ****96****, 082004 (2017).

[17] R.K.L. Lo *et al.*, "Search for gravitational-wave echoes," *LIGO-Virgo-KAGRA Collaboration* (2023).

[18] G.W. Horndeski, "Second-order scalar-tensor field equations in a four-dimensional space," *Int. J. Theor. Phys.* ****10****, 363 (1974).