

22.01

Electric Fields

“Action At a Distance”

- Particles that don’t physically “touch” can stil have electrostatic forces exchanged
 - How does that work if the particles aren’t “touching”?
 - * **Answer:** *electric fields*

About This Chapter

- *Three* goals of this chapter
 1. Define **electric field**
 2. Learn about analytic methods of describing electric fields
 3. Learn about how electric fields can affect charged particles

What is an “Electric Field”?

- **Field** = an object where each element in some specified **domain** is *uniquely* mapped to another **value**
 - Very similar to the concept behind a *function*
 - **Domain** = the space over which the field is described
 - The **value** can be **scalar** or **vector**
 - * **Scalar** = a mathematical object that specifies *magnitude*
 - **Fields** where the *associated values* are scalars are called a **scalar field**
 - * **Vector** = a mathematical object that specifies *magnitude* and *direction*
 - **Fields** where the *associated values* are vectors are called a **vector field**
 - More abstractly, a **vector** is just a mathematical object that contains many **scalar** values
 - * **Scalars** and **vectors** each have systems of operators that define how arithmetic works within their world and between
 - Examples
 - * Temperature field in an oven
 - * Pressure field in a pool

- **Electric Field** = a **vector field** that maps individual points in space to electrostatic force per unit charge
 - Mathematically, it looks like this

$$\vec{E} = \frac{\vec{F}}{q_0}$$

- where q_0 is an extremely small, positive charge, and \vec{F} is the electrostatic force exerted on the particle of charge q_0
- *Notice* that, since q_0 is a *positive* charge, \vec{E} and \vec{F} *must* point in the same direction
- The SI unit for electric field is $\frac{N}{C}$, which is a **vector** object

Procedure For Figuring Out \vec{E}

1. Take a particle of a very small, positive charge q_0
2. Place that particle at a point \vec{P} near some charged object O
3. Determine the electrostatic force between O and the particle of charge q_0 through empirical means
 - Perhaps measure acceleration and use newtonian mechanics to find \vec{F}
4. Calculate \vec{E} at \vec{P} by the following equation

$$\vec{E}_{\vec{P}} = \frac{\vec{F}_{\vec{P}}}{q_0}$$

Why Does q_0 Need to be Small?

- The purpose of q_0 is to detect the strength of \vec{E} at any given point
 - If q_0 were large, it would have a non-negligible affect on the electric field is is trying to measure!

Electric Field Lines

- Micheal Faraday came up with the idea
- **Electric Field Lines** = a way of visualizing the details of the electric field around an object
 - Basically just a series of vectors that float in space
 - The direction of the **electric field line** is the same as that of the **electrostatic force**
 - * Result of the mathematical definition of \vec{E}

- Two rules
 1. The electric field vector must be tangent to the electric field line through that point and in the same direction
 2. If the electric field vectors have tails that lie in a plane perpendicular to said electric field vectors, then the magnitude of \vec{E} is visually present by the relative density of electric field vectors, not by the magnitude of them
 - **Uniform Electric Field** = an electric field where all vectors point in the same direction
 - **Nonuniform Electric Field** = an electric field where vector direction varies from point to point
-

22.02

Electric Field Due to a Point Charge

- Because the strength of the electric field at any given point is

$$\vec{E} = \frac{\vec{F}}{q_0}$$

- we can substitute our particle of charge q_0 to get a formula to use:

$$|\vec{E}| = \frac{\frac{1}{4\pi\epsilon_0} \frac{qq_0}{r^2}}{q_0}$$

$$|\vec{E}| = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2}$$

- This formula allows to calculate the *magnitude* of the electric field at any given point
 - Note that this formula doesn't use q_0 ; we can calculate \vec{E} independent of any empirical data
- Because forces obey superposition (the permissibility of treating a vector sum as representative of the whole), we can demonstrate that the electric field obeys superposition

$$\vec{E} = \frac{\sum \vec{F}_i}{q_0}$$

$$\vec{E} = \frac{\vec{F}_1 + \vec{F}_2 + \vec{F}_3 + \dots + \vec{F}_n}{q_0}$$

$$\vec{E} = \frac{\vec{F}_1}{q_0} + \frac{\vec{F}_2}{q_0} + \dots + \frac{\vec{F}_n}{q_0}$$