




Learning Phenotypes and Dynamic Patient Representations via RNN Regularized Collective Non-negative Tensor Factorization


Kejing Yin¹, Dong Qian¹, William K. Cheung¹, Benjamin C. M. Fung², Jonathan Poon³

¹Hong Kong Baptist University ²McGill University ³Hong Kong Hospital Authority
{cskjin, dongqian, william}@comp.hkbu.edu.hk, ben.fung@mcgill.ca, jonathan@ha.org.hk


Introduction



❖ Phenotyping using Tensor CP Factorization




- **CP factorization** approximates the interaction tensor with the sum of rank-one tensors.



- Each rank-one tensor is defined as a candidate of the resulting phenotypes.
- The patient factor can be used as patient representations.

❖ Limitation: Temporal Dynamics Unexplored



- The temporal dynamics are simply ignored.
- Consequently, disease states appear at different times are **mixed together**.
- Especially in inpatients data and ICU cases.

❖ Adding Time as a Dimension?

- **Aligning Patients:** Patients have different Length-of-Stay (LoS), thus very difficult to represent all patients with one temporal tensor.

- **Phenotypes being dynamic:** With a temporal tensor for all patients, the global temporal patterns are embedded into the phenotypes, making it difficult to interpret.


❖ Our Goal

- **Avoid aligning patients.**
- **Keep the phenotypes static.**
- **Embed the individual disease progression into the dynamic patient representation.**
- **Explore the global temporal patterns as well.**

Related Works

- **Marble**^[2]: Poisson non-negative tensor factorization model for computational phenotyping task, with a bias tensor to capture the global information.
- **Rubik**^[3]: Incorporating guidance of existing medical knowledge, and adding pairwise constraint to make phenotypes distinct.
- **SPARTan**^[6]: Represent the EHR data using an “irregular” tensor and solve for the factors using PARAFAC2 decomposition.
- However, this model does not explicitly consider the interaction between the diagnoses and medications.

Proposed Method




Experiments and Results


- **Data Set:** MIMIC-III, an open-source, large-scale, de-identified ICU patients related EHR data set.
- Extract a subset containing 4,590 adult patients with LoS longer than 7 days.
- **Baselines:** **Rubik** (one of the state-of-the-art NTF computational phenotyping models), **CP-APR** (widely used Poisson non-negative tensor factorization model).

Phenotype Comparison: Phenotypes derived by our proposed model (top) and Rubik (bottom)

	Phenotype 1	Phenotype 4	Phenotype 9	
Clinically Highly Relevant	Chronic kidney disease (CKD) (0.536)	Other forms of chronic ischemic heart disease (0.507) Cardiac dysrhythmias (0.372) Essential hypertension (0.024)	Other diseases of lung (0.876)	Acute Respiratory Failure (classified “other disease of lung”)
Chronic Disease	RBC (Urine) (0.200) Osmolality, Measured (Blood) (0.117) Protein/Creatinine Ratio (Urine) (0.069) Hydromorphone (0.336) Phenylephrine (0.038) Aspirin (0.033)	Hematocrit (Blood) (0.072) Red Blood Cells (Blood) (0.071) Hemoglobin (Blood) (0.070) Acetaminophen (0.188) Metoclopramide (0.102) Insulin Human Regular (0.070)	pO2 (Blood Gas) (0.253) pCO2 (Blood Gas) (0.237) pH (Blood Gas) (0.215) Acetaminophen (0.113) Insulin (0.099) Bisacodyl (0.089)	
	Phenotype 1	Phenotype 2	Phenotype 3	
	Other diseases of lung (0.045) Septicemia (0.040) Certain adverse effects not elsewhere classified (0.039)	Other diseases of lung (0.040) Acute kidney failure (0.036) Certain adverse effects not elsewhere classified (0.032)	Acute kidney failure (0.039) Other diseases of lung (0.037) Cardiac dysrhythmias (0.033)	Dominated by acute diseases.
	Glucose(Blood) (0.019) Red Blood Cells(Blood) (0.019) Hematocrit(Blood) (0.019) Vancomycin (0.017) Insulin (0.015) Potassium Chloride (0.015)	Hematocrit(Blood) (0.017) Red Blood Cells(Blood) (0.017) Glucose(Blood) (0.017) Vancomycin (0.013) Potassium Chloride (0.013) Pantoprazole Sodium (0.012)	Glucose(Blood) (0.018) Hematocrit(Blood) (0.018) Red Blood Cells(Blood) (0.018) Vancomycin (0.015) Potassium Chloride (0.014) Heparin (0.014)	



Visualization and Interpretation of Dynamic Patient Representations



Contributions

- We proposed CNTF to jointly learn the dynamic patient representations and the static globally shared phenotypes.
- RNN-based regularization and HITF model are integrated to model the time dependency and incorporate the non-temporal modalities.
- The proposed model can derive clinically meaningful and interpretable phenotypes, and better separate the disease states appearing at different time.

References

- [1] Chi, Eric C., and Tamara G. Kolda. "On tensors, sparsity, and nonnegative factorizations." *SIAM Journal on Matrix Analysis and Applications* 33.4 (2012): 1272-1299.
- [2] Ho, Joyce C., Joydeep Ghosh, and Jimeng Sun. "Marble: high-throughput phenotyping from electronic health records via sparse nonnegative tensor factorization." *Proceedings of the 20th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*. ACM, 2014.
- [3] Wang, Yichen, et al. "Rubik: Knowledge guided tensor factorization and completion for health data analytics." *Proceedings of the 21th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*. ACM, 2015.
- [4] Kim, Yejin, et al. "Discriminative and distinct phenotyping by constrained tensor factorization." *Scientific Reports* 7.1 (2017): 1114.
- [5] Gunasekar, Suriya, et al. "Phenotyping using Structured Collective Matrix Factorization of Multi-source EHR Data." *arXiv preprint arXiv:1609.04466* (2016).
- [6] Yin, Kejing, et al. "Joint Learning of Phenotypes and Diagnosis-Medication Correspondence via Hidden Interaction Tensor Factorization." *IJCAI*. 2018.
- [7] Perros, Ioakeim, et al. "SPARTan: Scalable PARAFAC2 for large & sparse data." *Proceedings of the 23rd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*. ACM, 2017.