













Interpolation of Supernova Time Series Spectra with Optimal Transport

Jacob Krag Nørgaard, Gustav Gudmandsen, Anna Sophia Maxen
and Lucas Hallgren

A novel optimal transport-based approach for interpolating spectral time series

Paving the way for photometric classification of supernovae

Mauricio Ramirez^{1,2,★}, Giuliano Pignata^{3,2}, Francisco Förster^{2,4,5,6}, Santiago González-Gaitán⁷,
Claudia P. Gutiérrez^{10,11}, Bastian Ayala^{1,2}, Guillermo Cabrera-Vives^{2,12,13}, Márcio Catelan^{2,8,9},
Alejandra M. Muñoz Arancibia^{2,5}, and Jonathan Pineda-García^{1,2}



Importance

- Expect to observe many SNe in the future
- Type classification and photometric properties -> light curve fitting
- Importance of SNe in astronomy




Optimal Transport theory

- OT plan to get from initial to final distribution, $x_i \rightarrow y_i$
- Minimizing total cost


$$C_T = \sum_i^n c(x_i, T(x_i))$$

- What is cost?



Visual illustration


- Wasserstein distances
- Weighted by α
- Comparing with L2







The Wasserstein distance

"(Finding the Wasserstein barycenter) is like finding a middle point, not in terms of physical distance, but in terms of how much you would have to change each distribution to reach this middle point."



Results







Conclusion

- Transportation Theory and OT has a wide range of uses
- This paper highlighted one method, accomplished through interpolation, for astrophysical purposes



Reference

M. Ramirez et al., A novel optimal transport-based approach for interpolating spectral time series: Paving the way for photometric classification of supernovae
691:A33, November 2024. doi: 10.1051/0004-6361/202449170