

The drainage of the Baltic Ice Lake and a new Scandinavian reference ¹⁰Be production rate

Arjen P Stroeven¹, Jakob Heyman¹, Derek Fabel², Svante Björck³, Marc W Caffee⁴, Ola Fredin⁵, Jonathan M Harbor⁶

- 1. Department of Physical Geography and Quaternary Geology / Bolin Centre for Climate Research, Stockholm University, Sweden
- 2. School of Geographical and Earth Sciences, University of Glasgow, UK
- 3. Department of Geology, Lund University, Sweden
- 4. Department of Physics and Astronomy / PRIME Lab, Purdue University, USA
- 5. Geological Survey of Norway, Trondheim, Norway
- 6. Department of Earth, Atmospheric, and Planetary Sciences, Purdue University, USA

INTRODUCTION

The Baltic Ice Lake drained at the end of Younger Dryas when the southern margin of the Fennoscandian Ice Sheet retreated past the northern end of Mount Billingen in SC Sweden (Figs. 1, 2). The rapid drainage (reducing the water level of the Baltic Ice Lake by c. 25 m) scoured a c. 3 km wide and 20 m deep zone and left gravelly deposits west of northern Mount Billingen (Fig. 2).



Figure 1. The Baltic Ice Lake and the Fennoscandian Ice Sheet at the end of the Younger Dryas. The location of the Billingen study area and the four production rate calibration sites in Norway are marked.

BILLINGEN ¹⁰Be PROD RATE

We calibrate the reference ¹⁰Be production rate by matching ¹⁰Be concentrations from bedrock surfaces, boulders, and cobbles (Fig. 3) against an independently derived calibration age of 11,673 ± 100 years before sampling in 2003 based on the clay varve record and correlation to the GRIP ice core record.

We use the CRONUS calculator code (Balco et al. 2008) to derive reference ¹⁰Be production rates for the five CRONUS production rate scaling schemes and the two LSD (Lifton et al. 2014) production rate scaling schemes. All samples were submerged under water for the initial 129-940 years and we therefore model the production rate reduction during emergence through the water column (Fig. 4).




Figure 2. Topographic map of the Mount Billingen area based on a LiDAR DEM with sample locations and the width of the Baltic Ice Lake catastrophic drainage path marked with blue arrows.



Figure 3. Flood-scoured and streamlined bedrock surfaces (a, b). Sampling bedrock surfaces (c, e) and a boulder (d).




Figure 4. Sample emergence through water after the Baltic Ice Lake drainage event.

NORWAY RECALIBRATION

We recalibrate the reference ¹⁰Be production rate for four sites in Norway (Figs. 1, 5; Fenton et al. 2011; Goehring et al. 2012) using the CRONUS and the LSD production rate scaling schemes.




Figure 5. Individual sample and location reference ¹⁰Be production rates as probability density plots using the nuclide specific LSDn scaling scheme for five locations in Scandinavia. We use the well-clustered production rate from the bedrock surfaces for Billingen and we exclude outliers for the two western Norway locations to yield a $\chi^2_R < 1.5$.

SCANDINAVIAN PROD RATE

We combine the well-clustered Billingen bedrock surface reference ¹⁰Be production rate with the recalibrated reference production rates from three sites in Norway (one site excluded as an outlier) to derive a tightly clustered regional Scandinavian reference ¹⁰Be production rate (Fig. 6).




Figure 6. Local reference ¹⁰Be production rates for the LSD_n scaling scheme and arithmetic mean and standard deviation (horizontal black line and gray band). The Grøtlandsura reference production rate is excluded as an outlier. The production rates at Russenes (Fenton et al., 2011), Halsnøy and Oldedalen (Goehring et al., 2012b), and Mount Billingen (this study) for the LSD_n scaling scheme average to a new Scandinavian reference ¹⁰Be production rate of 3.95 ± 0.10 atoms g⁻¹ yr⁻¹.

OUTCOME

- Billingen reference ¹⁰Be production rate:
LSDn: 4.02 ± 0.18 atoms g⁻¹ yr⁻¹
CRONUS Lm: 4.19 ± 0.20 atoms g⁻¹ yr⁻¹
- Billingen reference ¹⁰Be production rate:
LSDn: 3.95 ± 0.10 atoms g⁻¹ yr⁻¹
CRONUS Lm: 4.13 ± 0.11 atoms g⁻¹ yr⁻¹

References

Balco G, Stone JO, Lifton NA, Dunai TJ, 2008. A complete and easily accessible means of calculating surface exposure ages or erosion rates from ¹⁰Be and ²⁶Al measurements. *Quaternary Geochronology* 3, 174-195.

Fenton CR, Hermanns RL, Blikra LH, Kubik PW, Bryant C, Niedermann S, Meixner A, Goethals MM, 2011. Regional ¹⁰Be production rate calibration for the past 12 ka deduced from the radiocarbon-dated Grøtlandsura and Russenes rock avalanches at 69° N, Norway. *Quaternary Geochronology* 6, 437-452.

Goehring BM, Lohne ØS, Mangerud J, Svendsen JI, Gyllencreutz R, Schaefler J, Finkel R, 2012. Late glacial and Holocene ¹⁰Be production rates for western Norway. *Journal of Quaternary Science* 27, 89-96.

Lifton N, Sato T, Dunai TJ, 2014. Scaling in situ cosmogenic nuclide production rates using analytical approximations to atmospheric cosmic-ray fluxes. *Earth and Planetary Sciences Letters* 386, 149-160.