

IDMA 202S: WEEK 6

POSETS AND GRAPHS

SRIKANTH SRINIVASAN

Algorithms & Complexity Section

DIKVU

srsy@di.ku.dk

Last time

- Relation $R \subseteq A \times B$
- Relation on A: $R \subseteq A \times A$

Last time

- Relation $R \subseteq A \times B$
- Relation on A: $R \subseteq A \times A$
- Representing relations as matrices & directed graphs (digraphs)

Last time

- Relation $R \subseteq A \times B$
- Relation on A: $R \subseteq A \times A$
- Representing relations as matrices & directed graphs (digraphs)
- Types of relations: Symmetric, Reflexive, Transitive
- Equivalence

Last time

- Relation $R \subseteq A \times B$
- Relation on A: $R \subseteq A \times A$
- Representing relations as matrices & directed graphs (digraphs)
- Types of relations: Symmetric, Reflexive, Transitive
- Equivalence
- Partially Ordered Sets (Posets)


Partial Order

→ (Total or Linear) Order: Relation R that allows us to compare all elements




Partial Order

- (Total or Linear) Order: Relation R that allows us to compare all elements
- Partial Order: can compare some elements




Partial Order

- (Total or Linear) Order: Relation R that allows us to compare all elements
- Partial Order: can compare some elements



Partial Order

- (Total or Linear) Order: Relation R that allows us to compare all elements
- Partial Order: can compare some elements
- Reflexive: aRa for all $a \in A$
- Antisymmetric:
 $aRb \wedge bRa \Rightarrow a=b$
- Transitive:
 $aRb \wedge bRc \Rightarrow aRc$



Examples

(\mathbb{N}, \leq) $(\mathbb{N}, =)$ $(P(\{1,2,3\}), \subseteq)$ $(\{1,2,3\}, =)$

Reflexive?

$$aRa$$

Antisymmetric?

$$aRb \wedge bRa$$

$$\Rightarrow a=b$$

Transitive?

$$aRb \wedge bRc$$

$$\Rightarrow aRc$$

Total?

$$aRb \vee bRa$$

Examples

(\mathbb{N}, \leq) $(\mathbb{N}, =)$ $(P(\{1,2,3\}), \subseteq)$ $(\{1,2,3\}, =)$

Reflexive?

$$aRa$$

Antisymmetric?

$$aRb \wedge bR a$$

$$\Rightarrow a=b$$

Transitive?

$$aRb \wedge bRc$$

$$\Rightarrow aRc$$

Total?

$$aRb \text{ OR } bRa$$

Examples

(\mathbb{N}, \leq) $(\mathbb{N}, =)$ $(P(\{1,2,3\}), \subseteq)$ $(\{1,2,3\}, =)$

Reflexive?

$$aRa$$

Antisymmetric?

$$aRb \wedge bR a$$

$$\Rightarrow a=b$$

Transitive?

$$aRb \wedge bRc$$

$$\Rightarrow aRc$$

Total?

$$aRb \text{ OR } bRa$$

Examples

(\mathbb{N}, \leq) $(\mathbb{N}, =)$ $(P(\{1,2,3\}), \subseteq)$ $(\{1,2,3\}, =)$

Reflexive?

✓ ✓ ✓

aRa

Antisymmetric?

✓ ✓ ✓

$aRb \& bRb$

$\Rightarrow a=b$

Transitive?

✓ ✓ ✓

$aRb \& bRc$

$\Rightarrow aRc$

Total?

✓ ✗ ✗

$aRb \text{ OR } bRa$

Examples

(\mathbb{N}, \leq) $(\mathbb{N}, =)$ $(P(\{1,2,3\}), \subseteq)$ $(\{1,2,3\}, =)$

Reflexive? $\checkmark \quad \checkmark \quad \checkmark \quad \checkmark$

aRa

Antisymmetric? $\checkmark \quad \checkmark \quad \checkmark \quad \checkmark$

$aRb \& bRb$

$\Rightarrow a=b$

Transitive? $\checkmark \quad \checkmark \quad \checkmark \quad \checkmark$

$aRb \& bRc$

$\Rightarrow aRc$

Total? $\checkmark \quad \times \quad \times \quad \times$

$aRb \text{ OR } bRa$

Product Posets


(A, R_1) , (B, R_2) posets

$(A \times B, R_3)$

$(a_1, b_1) R_3 (a_2, b_2)$ if

$a_1 R_1 a_2 \& b_1 R_1 b_2$

This is a poset.




Representing finite Posets

$(P(\{1, 2, 3\}), \subseteq)$

Representing finite Posets



$(P(\{1, 2, 3\}), \subseteq)$



Digraph



Representing finite Posets

$(P(\{1, 2, 3\}), \subseteq)$




Representing finite Posets

$(P(\{1, 2, 3\}), \subseteq)$



- No selfloops
- Remove arrows
(edges go up)
- Remove edges implied
by transitivity

Lattices (A, R) a poset




Lattices (A, R) a poset

Upper bound for $B \subseteq A$

$a \in A$ s.t. bRa for all $b \in B$.

Eg: $B = \{\emptyset, \{1\}, \{2\}\}$

Upper bounds: $\{\{1, 2\}, \{\{1, 2\}, \emptyset\}\}$



Lattices (A, R) a poset

Upper bound for $B \subseteq A$

$a \in A$ s.t. bRa for all $b \in B$.

Eg: $B = \{\emptyset, \{1\}, \{2\}\}$


Upper bounds: $\{\{1, 2\}\}, \{\{1\}, \{2\}\}$

Least Upper bound (LUB)

for B : $a \in A$ s.t.

- Upper bound for B

- $a' \in A$ an upper bound for
 $B \Rightarrow aRa'$



Lattices (A, R) a poset

Upper bound for $B \subseteq A$

$a \in A$ s.t. bRa for all $b \in B$.

Eg: $B = \{\emptyset, \{1\}, \{2\}\}$

Upper bounds: $\{\{1, 2\}\}, \{\{1\}, \{2\}\}$

Least Upper bound (LUB)


for B : $a \in A$ s.t.

- Upper bound for B

- $a' \in A$ an upper bound for

$$B \Rightarrow aRa'$$

Similar: Greatest lower bound (GLB)



Lattices (A, R) a poset

Examples

$(P(\{1, 2, 3\}), \subseteq)$ (\mathbb{N}, \leq) $(\mathbb{N}, |)$ $(\{1, 2, 3\}, =)$

LUB

(b_1, \dots, b_r)

GLB

(b_1, \dots, b_r)

Lattices (A, R) a poset

Examples

$$(P(\{1, 2, 3\}), \subseteq) \quad (\mathbb{N}, \leq) \quad (\mathbb{N}, |) \quad (\{1, 2, 3\}, =)$$

LUB $b_1 \vee \dots \vee b_r$

$$(\{b_1, \dots, b_r\})$$

GLB $b_1 \wedge b_2 \wedge \dots \wedge b_r$

$$(\{b_1, \dots, b_r\})$$

Lattices (A, R) a poset

Examples

$$(P(\{1, 2, 3\}), \subseteq) \quad (\mathbb{N}, \leq) \quad (\mathbb{N}, |) \quad (\{1, 2, 3\}, =)$$

LUB
 $(\{b_1, \dots, b_r\})$

$$b_1 \cup \dots \cup b_r$$

$$\max \{b_{1r}, b_r\}$$

GLB
 $(\{b_1, \dots, b_r\})$

$$b_1 \cap b_2 \cap \dots \cap b_r$$

$$\min \{b_1, \dots, b_r\}$$

Lattices (A, R) a poset

Examples

$$(P(\{1, 2, 3\}), \subseteq)$$

LUB

$$(\{b_1, \dots, b_r\})$$

$b_1 \cup \dots \cup b_r$

max
 $\{b_{1r}, b_r\}$

lcm
 $\{b_1, \dots, b_r\}$

GLB

$$(\{b_1, \dots, b_r\})$$

$b_1 \cap b_2 \cap \dots \cap b_r$

min
 $\{b_1, \dots, b_r\}$

gcd
 $\{b_1, \dots, b_r\}$

Lattices (A, R) a poset

Examples

$$(P(\{1, 2, 3\}), \subseteq)$$

LUB

$$(\{b_1, \dots, b_r\})$$

$$b_1 \cup \dots \cup b_r$$

$$\max \{b_{1r}, b_r\}$$

$$\text{lcm} \{b_1, \dots, b_r\}$$

may not exist

GLB

$$(\{b_1, \dots, b_r\})$$

$$b_1 \cap b_2 \cap \dots \cap b_r$$

$$\min \{b_1, \dots, b_r\}$$

$$\text{gcd} \{b_1, \dots, b_r\}$$

may not exist

Lattices (A, R) a poset

Examples

$$(P(\{1, 2, 3\}), \subseteq) \quad (\mathbb{N}, \leq) \quad (IN, |) \quad (\{1, 2, 3\}, =)$$

LUB

$$(\{b_1, \dots, b_r\})$$

$b_1 \vee \dots \vee b_r$

max
 $\{b_{1r}, \dots, b_r\}$

lcm
 $\{b_1, \dots, b_r\}$

may not
exist

GLB

$$(\{b_1, \dots, b_r\})$$

$b_1 \wedge b_2 \wedge \dots \wedge b_r$

min
 $\{b_1, \dots, b_r\}$

gcd
 $\{b_1, \dots, b_r\}$

may not
exist

Obs: LUBs & GLBs unique (if they exist)

Lattice: Poset with LUBs & GLBs.

Graphs

Directed graph

$$G = (V, E)$$

V - vertices

E - edges, each edge
an ordered pair (u, v)


Graphs

Directed graph

$$G = (V, E)$$

V - vertices

E - edges, each edge
an ordered pair (u, v)



$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, a), (a, d), (d, c)\}$$


Graphs

Directed graph

$$G = (V, E)$$

V - vertices

E - edges, each edge
an ordered pair (u, v)




$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, a), (a, d), (d, c)\}$$

Undirected graph

E - each edge an "unordered
pair", i.e. a set of two vertices



$$V = \{a, b, c, d\}$$

$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$


Graphs

Directed graph

$$G = (V, E)$$

V - vertices

E - edges, each edge
an ordered pair (u, v)




$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, a), (a, d), (d, c)\}$$

Undirected graph

E - each edge an "unordered
pair", i.e. a set of two vertices




$$V = \{a, b, c, d\}$$

Simple - no self loops or
graph parallel edges


$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Neighbours, degree



$$V = \{a, b, c, d\}$$


$$E = \{(a, b), (b, c), (c, a), (a, d), (d, c)\}$$



$$V = \{a, b, c, d\}$$

$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Neighbours, degree




$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, d), (a, d)\}$$

Neighbours of $u \in V$
 $N(u) = \{v \in V \mid \{u, v\} \in E\}$

Degree $\deg(u) = |N(u)|$



$$V = \{a, b, c, d\}$$

$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$


Neighbours, degree

Out-neighbours of $u \in V$

$$N_{out}(u) = \{v \mid (u, v) \in E\}$$

Out degree

$$\deg_{out}(u) = |N_{out}(u)|$$




$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, a), (b, c), (c, a), (c, d), (d, c)\}$$

Neighbours of $u \in V$

$$N(u) = \{v \in V \mid \{u, v\} \in E\}$$

Degree $\deg(u) = |N(u)|$



$$V = \{a, b, c, d\}$$

$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Neighbours, degree

Out-neighbours of $u \in V$

$$N_{out}(u) = \{v \mid (u, v) \in E\}$$

Out degree

$$\deg_{out}(u) = |N_{out}(u)|$$

In-neighbours

$$N_{in}(u) = \{v \mid (v, u) \in E\}$$


In-degree

$$\deg_{in}(u) = |N_{in}(u)|$$

Neighbours of $u \in V$


$$N(u) = \{v \in V \mid \{u, v\} \in E\}$$

Degree $\deg(u) = |N(u)|$



$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, a), (a, d), (d, c)\}$$




$$V = \{a, b, c, d\}$$

$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Neighbours, degree


$$\sum_{u \in V} \deg_{out}(u) = |E|$$

$$\sum_{u \in V} \deg_{in}(u) = |E|$$



$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, d), (d, a)\}$$



$$V = \{a, b, c, d\}$$


$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Neighbours, degree

$$\sum_{u \in V} \deg_{out}(u) = |E|$$


$$\sum_{u \in V} \deg_{in}(u) = |E|$$

$$\frac{1}{2} \sum_{u \in V} \deg(u) = |E|$$



$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, d), (d, a)\}$$



$$V = \{a, b, c, d\}$$


$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Paths & Simple Paths

Path - sequence of vertices


(v_0, v_1, \dots, v_n) s.t.

$\forall i \in \{0, \dots, n-1\} : (v_i, v_{i+1}) \in E$



$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, d), (d, a)\}$$



$$V = \{a, b, c, d\}$$

$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Paths & Simple Paths


Path - sequence of vertices

$$(v_0, v_1, \dots, v_n) \text{ s.t.}$$

$$\forall i \in \{0, \dots, n-1\}: (v_i, v_{i+1}) \in E$$


Simple Path - vertices are

not repeated



$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, a), (a, d), (d, c)\}$$



$$V = \{a, b, c, d\}$$

$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Paths & Simple Paths

Path - sequence of vertices


(v_0, v_1, \dots, v_n) s.t.

$\forall i \in \{0, \dots, n-1\} : (v_i, v_{i+1}) \in E$


Simple Path - vertices are not repeated

(a, b, c, d) - simple path

(a, b, c, a, d) - Path, not simple



$$E = \{(a, b), (b, c), (c, a), (a, d), (d, c)\}$$



$$E = \{\{a, b\}, \{b, c\}, \{a, c\}, \{a, d\}, \{d, c\}\}$$

Paths & Simple Paths

Path - sequence of vertices

(v_0, v_1, \dots, v_n) s.t.


$\forall i \in \{0, \dots, n-1\} : (v_i, v_{i+1}) \in E$

Simple Path - vertices are not repeated

(a, b, c, d) - Simple path


(a, b, c, a, d) - Path, not simple

Cycle - $v_0 = v_n$ & no repeated edges.



$$V = \{a, b, c, d\}$$

$$E = \{(a, b), (b, c), (c, d), (d, a)\}$$



$$V = \{a, b, c, d\}$$



$$E = \{\{a, b\}, \{b, c\}, \{c, d\}, \{d, a\}\}$$

Connectedness

$G = (V, E)$ Undirected graph

G is connected if

$\forall u, v \in V : \exists \text{ path } (v_0, \dots, v_n) \quad v_0 = u \& v_n = v$



Undirected Tree

$G = (V, E)$ undirected tree if it is connected
& contains no cycles (acyclic).

Undirected Tree

$G = (V, E)$ undirected tree if it is connected & contains no cycles (acyclic).

Theorem G finite graph is a tree if & only if:

- ① G connected & acyclic
- ② G connected & $|E| = |V| - 1$
- ③ G acyclic & $|E| = |V| - 1$
- ④ $\forall u \in V$: there is a unique simple path between u & v

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ $\forall u, v : \text{there is a unique simple path b/w } u, v$

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ ~~True~~: there is a unique
acyclic simple path b/w u, v

$\textcircled{4} \Rightarrow \textcircled{1}$ or $\neg \textcircled{1} \Rightarrow \neg \textcircled{4}$

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ $\forall u, v : \text{there is a unique simple path b/w } u, v$
acyclic

$\textcircled{4} \Rightarrow \textcircled{1}$ or $\neg \textcircled{1} \Rightarrow \neg \textcircled{4}$

Assume $\neg \textcircled{1}$. Then either

$\rightarrow \textcircled{6}$ disconnected

or

$\rightarrow \textcircled{6}$ has a cycle

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ $\forall u, v : \text{there is a unique simple path b/w } u, v$
acyclic

$\textcircled{4} \Rightarrow \textcircled{1}$ or $\neg \textcircled{1} \Rightarrow \neg \textcircled{4}$

Assume $\neg \textcircled{1}$. Then either

$\rightarrow \textcircled{6}$ disconnected $\Rightarrow \exists u, v : \text{no path b/w } u, v$
or

$\rightarrow \textcircled{6}$ has a cycle

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ $\exists u \neq v : \text{there is a unique simple path b/w } u, v$
acyclic

$\textcircled{4} \Rightarrow \textcircled{1}$ or $\neg \textcircled{1} \Rightarrow \neg \textcircled{4}$

Assume $\neg \textcircled{1}$. Then either

$\rightarrow G$ disconnected $\Rightarrow \exists u \neq v : \text{no path b/w } u, v$
or

$\rightarrow G$ has a cycle $\Rightarrow \exists u \neq v : \text{two distinct simple paths b/w } u \& v$

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ $\forall u \neq v$: there is a unique simple path b/w u, v
acyclic

$\textcircled{4} \Rightarrow \textcircled{1}$ or $\neg \textcircled{1} \Rightarrow \neg \textcircled{4}$


Assume $\neg \textcircled{1}$. Then either

$\rightarrow G$ disconnected $\Rightarrow \exists u \neq v$: no path b/w u, v

or

$\rightarrow G$ has a cycle $\Rightarrow \exists u \neq v$: two distinct simple paths b/w u, v

because



\rightarrow Shortest cycle in G does not repeat vertices

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ ~~Hypo~~: there is a unique
acyclic simple path blue u, v

$\textcircled{1} \Rightarrow \textcircled{4}$. or $\neg \textcircled{4} \Rightarrow \neg \textcircled{1}$

Assume $\neg \textcircled{4}$.

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{4}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ ~~Hypo~~: there is a unique acyclic

simple path b/w u, v

$\textcircled{1} \Rightarrow \textcircled{4}$. or $\neg \textcircled{4} \Rightarrow \neg \textcircled{1}$

Assume $\neg \textcircled{4}$. Then either

$\rightarrow \exists_{u \neq v}$: no simple path b/w u, v

$\rightarrow \exists_{u \neq v}$: two simple paths b/w u, v

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{4}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ ~~Hypo~~: there is a unique
acyclic simple path b/w u, v

$\textcircled{1} \Rightarrow \textcircled{4}$. or $\neg \textcircled{4} \Rightarrow \neg \textcircled{1}$ (Shortest paths are
↑ simple.)

Assume $\neg \textcircled{4}$. Then either

$\rightarrow \exists u \neq v$: no simple path b/w $u, v \Rightarrow$

$\rightarrow \exists u \neq v$: no path b/w $u, v \Rightarrow G$ disconnected

$\rightarrow \exists u \neq v$: two simple paths b/w u, v

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{4}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ ~~Hypo~~: there is a unique
acyclic simple path b/w u, v

$\textcircled{1} \Rightarrow \textcircled{4}$. or $\neg \textcircled{4} \Rightarrow \neg \textcircled{1}$ (Shortest paths are
simple.)

Assume $\neg \textcircled{4}$. Then either

$\rightarrow \exists u \neq v$: no simple path b/w $u, v \Rightarrow$

$\rightarrow \exists u \neq v$: no path b/w $u, v \Rightarrow G$ disconnected

$\exists u \neq v$: no path b/w $u, v \Rightarrow G$ disconnected

$\rightarrow \exists u \neq v$: two simple paths b/w $u, v \Rightarrow G$
has a cycle.

Proof of Theorem $(\textcircled{1} \Leftrightarrow \textcircled{4})$

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{4}$ $\exists u, v : \text{there is a unique simple path b/w } u, v$
acyclic

$\textcircled{1} \Rightarrow \textcircled{4}$. or $\neg \textcircled{4} \Rightarrow \neg \textcircled{1}$ (Shortest paths are
 \uparrow simple.)

Assume $\neg \textcircled{4}$. Then either

$\rightarrow \exists u, v : \text{no simple path b/w } u, v \Rightarrow$
 $\rightarrow \exists u, v : \text{no path b/w } u, v \Rightarrow G \text{ disconnected}$
 $\exists u, v : \text{no path b/w } u, v \Rightarrow$
 $\rightarrow \exists u, v : \text{two simple paths b/w } u, v \Rightarrow G$
 \uparrow
 $\rightarrow \exists u, v : \text{two simple paths b/w } u, v \Rightarrow G$
has a cycle.

Find closest such u, v & consider
shortest & second shortest path. Concatenate
them.

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1} G = (V, E)$ connected &
acyclic $\textcircled{2} G$ connected &
 $|E| = |V| - 1$

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{2}$ G connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{2}$ G connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Induction: Assume for n . Say $|V| = n$.

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{2}$ G connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Induction: Assume for n . Say $|V| = n$.

$\textcircled{1} \Rightarrow \exists v \in V : \deg(v) = 1$

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{2}$ G connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Induction: Assume for n . Say $|V| = n$.

$\textcircled{1} \Rightarrow \exists v \in V : \deg(v) = 1 \longrightarrow$ If every vertex
has degree ≥ 2 , G
has a cycle.

If any vertex has
degree 0, G is
disconnected.

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1} G = (V, E)$ connected & $\textcircled{2} G$ connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Induction: Assume for n . Say $|V| = n$.

$\textcircled{1} \Rightarrow \exists v \in V : \deg(v) = 1 \longrightarrow$ If every vertex
has degree ≥ 2 , G
has a cycle.

Consider $G' = (V \setminus \{v\}, E \setminus \{(v, u)\})$
where u is the only nbr. of v .

G' is connected & acyclic

$$\Rightarrow |E| - 1 = n - 1$$

$$\Rightarrow |E| = n$$

If any vertex has
degree 0, G is
disconnected.

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{2}$ G connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Induction: Assume for n . Say $|V| = n + 1$.

$\textcircled{2} \Rightarrow \exists v \in V : \deg(v) = 1$

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1}$ $G = (V, E)$ connected & $\textcircled{2}$ G connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Induction: Assume for n . Say $|V| = n + 1$.

$\textcircled{2} \Rightarrow \exists v \in V : \deg(v) = 1 \longrightarrow \sum_{u \in V} \deg(u) = 2|E| = 2n$.

If any vertex has degree 0, G is disconnected.

Proof of Theorem ($\textcircled{1} \Leftrightarrow \textcircled{2}$)

$\textcircled{1} G = (V, E)$ connected & $\textcircled{2} G$ connected &
acyclic $|E| = |V| - 1$

Induction on $n = |V|$.

Base case: $n = 1$. $|E| = 0 = n - 1$.

Induction: Assume for n . Say $|V| = n + 1$.

$\textcircled{2} \Rightarrow \exists v \in V : \deg(v) = 1 \longrightarrow \sum_{u \in V} \deg(u) = 2|E| = 2n$.

Consider $G' = (V \setminus \{v\}, E \setminus \{(v, u)\})$

where u is the only nbr. of v .

G' is connected & $|E \setminus \{(v, u)\}| = n - 1$

$\Rightarrow G'$ is connected & acyclic
 \Rightarrow So is G .

If any vertex has degree 0, G is disconnected.

Spanning Trees

$$G = (V, E)$$

Subgraph G' of G : obtained by removing some subset of edges and/or vertices.

Spanning Trees

$$G = (V, E)$$

Subgraph G' of G : obtained by removing some subset of edges and/or vertices.



Spanning tree: Subgraph $G' = (V, E')$ that is a tree.

Spanning Trees

$$G = (V, E)$$

Subgraph G' of G : obtained by removing some subset of edges and/or vertices.

Spanning tree: Subgraph $G' = (V, E')$ that is a tree.



Spanning trees of G

Directed Tree $T = (V, E)$ directed graph

→ Special vertex v_0 (root)

→ For $v \in V \setminus \{v_0\}$, unique walk from v_0 to v .

→ No walk from v_0 to v_0 .

Directed Tree $T = (V, E)$ directed graph

→ Special vertex v_0 (root)

→ For $v \in V \setminus \{v_0\}$, unique walk from v_0 to v .

→ No walk from v_0 to v_0 .


Thm: (T, v_0) a directed tree

① T contains no cycles

② v_0 is the unique root.

③ $\deg_{in}(v_0) = 0$

$\deg_{in}(v) = 1 \quad \forall v \in V \setminus \{v_0\}$



Directed Tree $T = (V, E)$ directed graph

→ Special vertex v_0 (root)

→ For $v \in V \setminus \{v_0\}$, unique walk from v_0 to v .

→ No walk from v_0 to v_0 .

Thm: (T, v_0) a directed tree


① T contains no cycles

② v_0 is the unique root.

③ $\deg_{in}(v_0) = 0$

$\deg_{in}(v) = 1 \quad \forall v \in V \setminus \{v_0\}$

Thm: $|E| = \sum_{v \in V} \deg_{in}(v) = |V| - 1$.




Directed Tree $T = (V, E)$ directed graph

→ Special vertex v_0 (root)


→ For $v \in V \setminus \{v_0\}$, unique walk from v_0 to v .

→ No walk from v_0 to v_0 .




Directed Tree $T = (V, E)$ directed graph

- Special vertex v_0 (root)
- For $v \in V \setminus \{v_0\}$, unique walk from v_0 to v .
- No walk from v_0 to v_0 .



Directed Tree $T = (V, E)$ directed graph

- Special vertex v_0 (root)
- For $v \in V \setminus \{v_0\}$, unique walk from v_0 to v .
- No walk from v_0 to v_0 .




Directed Tree $T = (V, E)$ directed graph

- Special vertex v_0 (root)
- For $v \in V \setminus \{v_0\}$, unique walk from v_0 to v .
- No walk from v_0 to v_0 .

Binary tree: ≤ 2 children

Ternary,
 n -ary





Outline

- Record of relations
- Diagram resp
- Reflexive → Antisymmetris
- Symmetric → Transitivity
- Par
- Examples & non-examples
- Product parts
- Points via Hausdorffian
- Categories
- Examples

Graphs

- Defn of graph
- Vertices, adjacency, degree
- paths & cycles
- Trees, directed
- Paths + alg
- Terminology
- Undirected trees
- Graphs
- Spanning trees