

Supercritical Space-Width Trade-offs for Resolution

Jakob Nordström

KTH Royal Institute of Technology

SAT and Interactions
Schloss Dagstuhl
September 19–23, 2016

Joint work with Christoph Berkholz

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

1. $x \vee y$
2. $x \vee \bar{y} \vee z$
3. $\bar{x} \vee z$
4. $\bar{y} \vee \bar{z}$
5. $\bar{x} \vee \bar{z}$

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

1.	$x \vee y$	Axiom
2.	$x \vee \bar{y} \vee z$	Axiom
3.	$\bar{x} \vee z$	Axiom
4.	$\bar{y} \vee \bar{z}$	Axiom
5.	$\bar{x} \vee \bar{z}$	Axiom
6.	$x \vee \bar{y}$	Res(2, 4)
7.	x	Res(1, 6)
8.	\bar{x}	Res(3, 5)
9.	\perp	Res(7, 8)

The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ **annotated list** or
- ▶ directed acyclic graph (DAG)

- | | | |
|-----------|---------------------------|------------------|
| 1. | $x \vee y$ | Axiom |
| 2. | $x \vee \bar{y} \vee z$ | Axiom |
| 3. | $\bar{x} \vee z$ | Axiom |
| 4. | $\bar{y} \vee \bar{z}$ | Axiom |
| 5. | $\bar{x} \vee \bar{z}$ | Axiom |
| 6. | $x \vee \bar{y}$ | Res(2, 4) |
| 7. | x | Res(1, 6) |
| 8. | \bar{x} | Res(3, 5) |
| 9. | \perp | Res(7, 8) |

The Resolution Proof System

Goal: refute **unsatisfiable** CNF


- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ annotated list or
- ▶ **directed acyclic graph (DAG)**



The Resolution Proof System

Goal: refute **unsatisfiable** CNF

- ▶ Start with **axiom** clauses in formula
- ▶ Derive new clauses by **resolution rule**


$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

- ▶ Done when empty clause \perp derived

Can represent refutation/proof as

- ▶ annotated list or
- ▶ **directed acyclic graph (DAG)**

Tree-like resolution if DAG is tree



Resolution Size/Length and Width

Length of proof = # clauses (9 in our example)

Length of refuting F = min length over all proofs for F

Resolution Size/Length and Width

Length of proof = # clauses (9 in our example)

Length of refuting F = min length over all proofs for F

Size should strictly speaking measure # symbols

But for resolution don't care too much about linear factors here

Set size = length

Resolution Size/Length and Width

Length of proof = # clauses (9 in our example)

Length of refuting F = min length over all proofs for F

Size should strictly speaking measure # symbols

But for resolution don't care too much about linear factors here

Set size = length

Width of proof = # literals in largest clause (3 in our example)

Width of refuting F = min width over all proofs for F

Width at most linear, so here obviously care about linear factors

Resolution Space

Space = amount of memory needed
when performing refutation

- | | | |
|----|-------------------------|-----------|
| 1. | $x \vee y$ | Axiom |
| 2. | $x \vee \bar{y} \vee z$ | Axiom |
| 3. | $\bar{x} \vee z$ | Axiom |
| 4. | $\bar{y} \vee \bar{z}$ | Axiom |
| 5. | $\bar{x} \vee \bar{z}$ | Axiom |
| 6. | $x \vee \bar{y}$ | Res(2, 4) |
| 7. | x | Res(1, 6) |
| 8. | \bar{x} | Res(3, 5) |
| 9. | \perp | Res(7, 8) |

Resolution Space

Space = amount of memory needed
when performing refutation

Can be measured in different ways:

- ▶ clause space (our focus)
- ▶ total space

- | | | |
|----|-------------------------|-----------|
| 1. | $x \vee y$ | Axiom |
| 2. | $x \vee \bar{y} \vee z$ | Axiom |
| 3. | $\bar{x} \vee z$ | Axiom |
| 4. | $\bar{y} \vee \bar{z}$ | Axiom |
| 5. | $\bar{x} \vee \bar{z}$ | Axiom |
| 6. | $x \vee \bar{y}$ | Res(2, 4) |
| 7. | x | Res(1, 6) |
| 8. | \bar{x} | Res(3, 5) |
| 9. | \perp | Res(7, 8) |

Resolution Space

Space = amount of memory needed
when performing refutation

Can be measured in different ways:

- ▶ clause space (our focus)
- ▶ total space

Clause space at step t : # clauses at
steps $\leq t$ used at steps $\geq t$

Total space at step t : Count also literals

- | | | |
|----|-------------------------|-----------|
| 1. | $x \vee y$ | Axiom |
| 2. | $x \vee \bar{y} \vee z$ | Axiom |
| 3. | $\bar{x} \vee z$ | Axiom |
| 4. | $\bar{y} \vee \bar{z}$ | Axiom |
| 5. | $\bar{x} \vee \bar{z}$ | Axiom |
| 6. | $x \vee \bar{y}$ | Res(2, 4) |
| 7. | x | Res(1, 6) |
| 8. | \bar{x} | Res(3, 5) |
| 9. | \perp | Res(7, 8) |

Resolution Space

Space = amount of memory needed
when performing refutation

Can be measured in different ways:

- ▶ clause space (our focus)
- ▶ total space

Clause space at step t : # clauses at
steps $\leq t$ used at steps $\geq t$

Total space at step t : Count also literals

Example: Clause space at step 7

- | | | |
|----|-------------------------|-----------|
| 1. | $x \vee y$ | Axiom |
| 2. | $x \vee \bar{y} \vee z$ | Axiom |
| 3. | $\bar{x} \vee z$ | Axiom |
| 4. | $\bar{y} \vee \bar{z}$ | Axiom |
| 5. | $\bar{x} \vee \bar{z}$ | Axiom |
| 6. | $x \vee \bar{y}$ | Res(2, 4) |
| 7. | x | Res(1, 6) |
| 8. | \bar{x} | Res(3, 5) |
| 9. | \perp | Res(7, 8) |

Resolution Space

Space = amount of memory needed
when performing refutation


Can be measured in different ways:

- ▶ clause space (our focus)
- ▶ total space

Clause space at step t : # clauses at
steps $\leq t$ used at steps $\geq t$

Total space at step t : Count also literals

Example: Clause space at step 7



Resolution Space

Space = amount of memory needed
when performing refutation


Can be measured in different ways:

- ▶ clause space (our focus)
- ▶ total space

Clause space at step t : # clauses at
steps $\leq t$ used at steps $\geq t$

Total space at step t : Count also literals

Example: Clause space at step 7 is 5



Resolution Space

Space = amount of memory needed
when performing refutation


Can be measured in different ways:

- ▶ clause space (our focus)
- ▶ total space

Clause space at step t : # clauses at
steps $\leq t$ used at steps $\geq t$

Total space at step t : Count also literals

Example: Clause space at step 7 is 5
Total space at step 7 is 9



Resolution Space

Space = amount of memory needed
when performing refutation

Can be measured in different ways:

- ▶ clause space (our focus)
- ▶ total space

Clause space at step t : # clauses at
steps $\leq t$ used at steps $\geq t$


Total space at step t : Count also literals

Example: Clause space at step 7 is 5

Total space at step 7 is 9

Space of proof = max over all steps

Space of refuting F = min over all proofs



Upper Bounds on Resolution Complexity Measures

Worst-case upper bounds for resolution refutations of formula
(from now on assume $n = \# \text{variables}$):

Upper Bounds on Resolution Complexity Measures

Worst-case upper bounds for resolution refutations of formula
(from now on assume $n = \# \text{variables}$):

Size / length # derivation steps $\mathcal{O}(2^n)$

Upper Bounds on Resolution Complexity Measures

Worst-case upper bounds for resolution refutations of formula
(from now on assume $n = \# \text{variables}$):

Size / length	# derivation steps	$\mathcal{O}(2^n)$
Width	max # literals in a clause	$\mathcal{O}(n)$

Upper Bounds on Resolution Complexity Measures

Worst-case upper bounds for resolution refutations of formula
(from now on assume $n = \# \text{variables}$):

Size / length	# derivation steps	$\mathcal{O}(2^n)$
Width	max # literals in a clause	$\mathcal{O}(n)$
Clause space	max # clauses in memory	$\mathcal{O}(n)$

Upper Bounds on Resolution Complexity Measures

Worst-case upper bounds for resolution refutations of formula
(from now on assume $n = \# \text{variables}$):

Size / length	# derivation steps	$\mathcal{O}(2^n)$
Width	max # literals in a clause	$\mathcal{O}(n)$
Clause space	max # clauses in memory	$\mathcal{O}(n)$
Total space	total size of memory	$\mathcal{O}(n^2)$

Upper Bounds on Resolution Complexity Measures

Worst-case upper bounds for resolution refutations of formula
(from now on assume $n = \# \text{variables}$):

Size / length	# derivation steps	$\mathcal{O}(2^n)$
Width	max # literals in a clause	$\mathcal{O}(n)$
Clause space	max # clauses in memory	$\mathcal{O}(n)$
Total space	total size of memory	$\mathcal{O}(n^2)$

This talk: focus on width and clause space

Upper Bounds on Resolution Complexity Measures

Worst-case upper bounds for resolution refutations of formula
(from now on assume $n = \# \text{variables}$):

Size / length	# derivation steps	$\mathcal{O}(2^n)$
Width	max # literals in a clause	$\mathcal{O}(n)$
Clause space	max # clauses in memory	$\mathcal{O}(n)$
Total space	total size of memory	$\mathcal{O}(n^2)$

This talk: focus on width and clause space
But results translate to total space by:

$$\text{clause space} \leq \text{total space} \leq \text{clause space} \cdot \text{width}$$

Lower Bounds via Resolution Width

For n -variable k -CNFs (k constant) it holds that:

$$\text{width} \leq \Omega(\text{clause space}) \quad [\text{Atserias \& Dalmau '03}]$$

Lower Bounds via Resolution Width

For n -variable k -CNFs (k constant) it holds that:

$$\text{width} \leq \Omega(\text{clause space}) \quad [\text{Atserias \& Dalmau '03}]$$

$$\text{width}^2 \leq \Omega(\text{total space}) \quad [\text{Bonacina '16}]$$

Lower Bounds via Resolution Width

For n -variable k -CNFs (k constant) it holds that:

$$\text{width} \leq \Omega(\text{clause space}) \quad [\text{Atserias \& Dalmau '03}]$$

$$\text{width}^2 \leq \Omega(\text{total space}) \quad [\text{Bonacina '16}]$$

$$\text{width}^2 \leq \Omega(n \log(\text{size})) \quad [\text{Ben-Sasson \& Wigderson '99}]$$

Lower Bounds via Resolution Width

For n -variable k -CNFs (k constant) it holds that:

$$\text{width} \leq \Omega(\text{clause space}) \quad [\text{Atserias \& Dalmau '03}]$$

$$\text{width}^2 \leq \Omega(\text{total space}) \quad [\text{Bonacina '16}]$$

$$\text{width}^2 \leq \Omega(n \log(\text{size})) \quad [\text{Ben-Sasson \& Wigderson '99}]$$

In particular, $\text{width} = \Omega(n) \implies \text{size} = \Omega(2^n)$

Lower Bounds via Resolution Width

For n -variable k -CNFs (k constant) it holds that:

$$\text{width} \leq \Omega(\text{clause space}) \quad [\text{Atserias \& Dalmau '03}]$$

$$\text{width}^2 \leq \Omega(\text{total space}) \quad [\text{Bonacina '16}]$$

$$\text{width}^2 \leq \Omega(n \log(\text{size})) \quad [\text{Ben-Sasson \& Wigderson '99}]$$

In particular, $\text{width} = \Omega(n) \implies \text{size} = \Omega(2^n)$

So clearly **width key measure**—but not the answer to every question

Lower Bounds via Resolution Width

For n -variable k -CNFs (k constant) it holds that:

$$\text{width} \leq \Omega(\text{clause space}) \quad [\text{Atserias \& Dalmau '03}]$$

$$\text{width}^2 \leq \Omega(\text{total space}) \quad [\text{Bonacina '16}]$$

$$\text{width}^2 \leq \Omega(n \log(\text{size})) \quad [\text{Ben-Sasson \& Wigderson '99}]$$

In particular, $\text{width} = \Omega(n) \implies \text{size} = \Omega(2^n)$

So clearly **width key measure**—but not the answer to every question

- Can have width $\Theta(\sqrt{n})$ and still size $\text{poly}(n)$
[Bonet & Galesi '99]

Lower Bounds via Resolution Width

For n -variable k -CNFs (k constant) it holds that:

$$\text{width} \leq \Omega(\text{clause space}) \quad [\text{Atserias \& Dalmau '03}]$$

$$\text{width}^2 \leq \Omega(\text{total space}) \quad [\text{Bonacina '16}]$$

$$\text{width}^2 \leq \Omega(n \log(\text{size})) \quad [\text{Ben-Sasson \& Wigderson '99}]$$

In particular, $\text{width} = \Omega(n) \implies \text{size} = \Omega(2^n)$

So clearly **width key measure**—but not the answer to every question

- ▶ Can have width $\Theta(\sqrt{n})$ and still size $\text{poly}(n)$
[Bonet & Galesi '99]
- ▶ Can have width $\mathcal{O}(1)$ and still clause space $\Omega(n/\log n)$
[Ben-Sasson & Nordström '08]

Upper Bounds via Resolution Width

$$\text{size} \leq n^{\mathcal{O}(\text{width})}$$

Upper Bounds via Resolution Width

$$\text{size} \leq n^{\mathcal{O}(\text{width})}$$

$$\text{time to find a refutation} \leq n^{\mathcal{O}(\text{width})}$$

for $w \leftarrow 3 \dots n$ **do**

 Resolve all clauses & keep resolvents with at most w literals

 If \perp has been derived, then output **UNSAT**

end for

Output **SAT**

Upper Bounds via Resolution Width

$$\text{size} \leq n^{\mathcal{O}(\text{width})}$$

$$\text{time to find a refutation} \leq n^{\mathcal{O}(\text{width})}$$

for $w \leftarrow 3 \dots n$ **do**

 Resolve all clauses & keep resolvents with at most w literals

 If \perp has been derived, then output **UNSAT**

end for

Output **SAT**

Algorithm (and resolution proof) requires $n^{\mathcal{O}(\text{width})}$ space!

Upper Bounds via Resolution Width

$$\text{size} \leq n^{\mathcal{O}(\text{width})}$$

$$\text{time to find a refutation} \leq n^{\mathcal{O}(\text{width})}$$

for $w \leftarrow 3 \dots n$ **do**

 Resolve all clauses & keep resolvents with at most w literals

 If \perp has been derived, then output **UNSAT**

end for

Output **SAT**

Algorithm (and resolution proof) requires $n^{\mathcal{O}(\text{width})}$ space!

[Ben-Sasson '02] exhibited formulas

- ▶ refutable in width $\mathcal{O}(1)$ and clause space $\mathcal{O}(1)$
- ▶ width $\mathcal{O}(1) \implies$ clause space $\Omega(n / \log n)$

Upper Bounds via Resolution Width

$$\text{size} \leq n^{\mathcal{O}(\text{width})}$$

$$\text{time to find a refutation} \leq n^{\mathcal{O}(\text{width})}$$

for $w \leftarrow 3 \dots n$ **do**

 Resolve all clauses & keep resolvents with at most w literals

 If \perp has been derived, then output **UNSAT**

end for

Output **SAT**

Algorithm (and resolution proof) requires $n^{\mathcal{O}(\text{width})}$ space!

[Ben-Sasson '02] exhibited formulas

- ▶ refutable in width $\mathcal{O}(1)$ and clause space $\mathcal{O}(1)$
- ▶ width $\mathcal{O}(1) \implies$ clause space $\Omega(n / \log n)$

Which bound is closer to the truth?

Upper Bounds via Resolution Width

$$\text{size} \leq n^{\mathcal{O}(\text{width})}$$

$$\text{time to find a refutation} \leq n^{\mathcal{O}(\text{width})}$$

for $w \leftarrow 3 \dots n$ **do**

 Resolve all clauses & keep resolvents with at most w literals

 If \perp has been derived, then output **UNSAT**

end for

Output **SAT**

Algorithm (and resolution proof) requires $n^{\mathcal{O}(\text{width})}$ space!

[Ben-Sasson '02] exhibited formulas

- ▶ refutable in width $\mathcal{O}(1)$ and clause space $\mathcal{O}(1)$
- ▶ width $\mathcal{O}(1) \implies$ clause space $\Omega(n / \log n)$

Which bound is closer to the truth?

Recall: can always do clause space $\mathcal{O}(n)$

A Supercritical Space-Width Tradeoff

Theorem

For any $\varepsilon > 0$ and $6 \leq w \leq n^{\frac{1}{2}-\varepsilon}$ exist n -variable w -CNFs F_n s.t.

1. Resolution can refute F_n in width w
2. Any width- w refutation of F_n requires clause space $n^{\Omega(w)}$

A Supercritical Space-Width Tradeoff

Theorem

For any $\varepsilon > 0$ and $6 \leq w \leq n^{\frac{1}{2}-\varepsilon}$ exist n -variable w -CNFs F_n s.t.

1. Resolution can refute F_n in width w
2. Any width- w refutation of F_n requires clause space $n^{\Omega(w)}$

Space lower bound $n^{\Omega(w)}$ holds for all proofs up to width $o(w \log n)$

A Supercritical Space-Width Tradeoff

Theorem

For any $\varepsilon > 0$ and $6 \leq w \leq n^{\frac{1}{2}-\varepsilon}$ exist n -variable w -CNFs F_n s.t.

1. Resolution can refute F_n in width w
2. Any width- w refutation of F_n requires clause space $n^{\Omega(w)}$

Space lower bound $n^{\Omega(w)}$ holds for all proofs up to width $o(w \log n)$

Proof outline

Use hardness condensation approach in [Razborov '16]:

1. Start with formula that requires nearly linear clause space
2. Reduce the number of variables from n to $n^{1/w}$
3. But maintain space lower bound for small-width proofs

A Supercritical Space-Width Tradeoff

Theorem

For any $\varepsilon > 0$ and $6 \leq w \leq n^{\frac{1}{2}-\varepsilon}$ exist n -variable w -CNFs F_n s.t.

1. Resolution can refute F_n in width w
2. Any width- w refutation of F_n requires clause space $n^{\Omega(w)}$

Space lower bound $n^{\Omega(w)}$ holds for all proofs up to width $o(w \log n)$

Proof outline

Use hardness condensation approach in [Razborov '16]:

1. Start with formula that requires nearly linear clause space
2. Reduce the number of variables from n to $n^{1/w}$
3. But maintain space lower bound for small-width proofs

Key components:

- ▶ Expander graphs
- ▶ XORification (substitution with exclusive or)

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}
- ▶ There are instances I_n such that:
 - ▶ \exists solutions S_1, S_2 with $\varphi(S_1) = \text{small}'$ and $\psi(S_2) = \text{small}''$

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}
- ▶ There are instances I_n such that:
 - ▶ \exists solutions S_1, S_2 with $\varphi(S_1) = \text{small}'$ and $\psi(S_2) = \text{small}''$
 - ▶ Any solution S with $\varphi(S)$ even medium-small must have $\psi(S)$ approach critical value ψ_{crit}

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}
- ▶ There are instances I_n such that:
 - ▶ \exists solutions S_1, S_2 with $\varphi(S_1) = \text{small}'$ and $\psi(S_2) = \text{small}''$
 - ▶ Any solution S with $\varphi(S)$ even medium-small must have $\psi(S)$ approach critical value ψ_{crit}
 - ▶ Conversely, $\psi(S)$ medium-small $\implies \varphi(S) \approx \varphi_{\text{crit}}$

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}
- ▶ There are instances I_n such that:
 - ▶ \exists solutions S_1, S_2 with $\varphi(S_1) = \text{small}'$ and $\psi(S_2) = \text{small}''$
 - ▶ Any solution S with $\varphi(S)$ even medium-small must have $\psi(S)$ approach critical value ψ_{crit}
 - ▶ Conversely, $\psi(S)$ medium-small $\implies \varphi(S) \approx \varphi_{\text{crit}}$

Supercritical setting for trade-offs:

- ▶ Any S with $\varphi(S)$ medium-small must have $\psi(S) \gg \psi_{\text{crit}}$

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}
- ▶ There are instances I_n such that:
 - ▶ \exists solutions S_1, S_2 with $\varphi(S_1) = \text{small}'$ and $\psi(S_2) = \text{small}''$
 - ▶ Any solution S with $\varphi(S)$ even medium-small must have $\psi(S)$ approach critical value ψ_{crit}
 - ▶ Conversely, $\psi(S)$ medium-small $\implies \varphi(S) \approx \varphi_{\text{crit}}$

Supercritical setting for trade-offs:

- ▶ Any S with $\varphi(S)$ medium-small must have $\psi(S) \gg \psi_{\text{crit}}$
- ▶ Optimizing φ pushes ψ up into supercritical regime above worst case!

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}
- ▶ There are instances I_n such that:
 - ▶ \exists solutions S_1, S_2 with $\varphi(S_1) = \text{small}'$ and $\psi(S_2) = \text{small}''$
 - ▶ Any solution S with $\varphi(S)$ even medium-small must have $\psi(S)$ approach critical value ψ_{crit}
 - ▶ Conversely, $\psi(S)$ medium-small $\implies \varphi(S) \approx \varphi_{\text{crit}}$

Supercritical setting for trade-offs:

- ▶ Any S with $\varphi(S)$ medium-small must have $\psi(S) \gg \psi_{\text{crit}}$
- ▶ Optimizing φ pushes ψ up into supercritical regime above worst case!
- ▶ **Very** strong trade-offs—Razborov refers to them as “ultimate”

What Do You Mean “Supercritical”?!

Typical setting for trade-off results:

- ▶ Have two complexity measures φ and ψ
- ▶ Worst-case (usually trivial) upper bounds φ_{crit} and ψ_{crit}
- ▶ There are instances I_n such that:
 - ▶ \exists solutions S_1, S_2 with $\varphi(S_1) = \text{small}'$ and $\psi(S_2) = \text{small}''$
 - ▶ Any solution S with $\varphi(S)$ even medium-small must have $\psi(S)$ approach critical value ψ_{crit}
 - ▶ Conversely, $\psi(S)$ medium-small $\implies \varphi(S) \approx \varphi_{\text{crit}}$

Supercritical setting for trade-offs:


- ▶ Any S with $\varphi(S)$ medium-small must have $\psi(S) \gg \psi_{\text{crit}}$
- ▶ Optimizing φ pushes ψ up into supercritical regime above worst case!
- ▶ **Very** strong trade-offs—Razborov refers to them as “ultimate”
- ▶ We feel “supercritical” is more descriptive

Expanders

Very well-connected so-called **expander graphs** play leading role in many proof complexity lower bounds

Expanders

Very well-connected so-called **expander graphs** play leading role in many proof complexity lower bounds




Clause-variable incidence graph (CVIG)

- ▶ Clauses on the left
- ▶ Variables on the right
- ▶ Edge if variable \in clause (ignore signs)

Expanders

Very well-connected so-called **expander graphs** play leading role in many proof complexity lower bounds



Clause-variable incidence graph (CVIG)


- ▶ Clauses on the left
- ▶ Variables on the right
- ▶ Edge if variable \in clause (ignore signs)

If **CVIG well-connected**, then **lower bounds** for

- ▶ width, size, and space in resolution
[Ben-Sasson & Wigderson '99, Ben-Sasson & Galesi '03]
- ▶ degree and size in polynomial calculus
[Impagliazzo et al. '99, Alekhovich & Razborov '01]

Expanders

Very well-connected so-called **expander graphs** play leading role in many proof complexity lower bounds



Clause-variable incidence graph (CVIG)

- ▶ Clauses on the left
- ▶ Variables on the right
- ▶ Edge if variable \in clause (ignore signs)

If **CVIG well-connected**, then **lower bounds** for

- ▶ width, size, and space in resolution
[Ben-Sasson & Wigderson '99, Ben-Sasson & Galesi '03]
- ▶ degree and size in polynomial calculus
[Impagliazzo et al. '99, Alekhovich & Razborov '01]

Can also define more general graphs that capture “underlying combinatorial structure” and extend results [Mikša & Nordström '15]

XORification

Modify F to $F[\oplus_2]$ by substituting $x_1 \oplus x_2$ for every variable x

XORification

Modify F to $F[\oplus_2]$ by substituting $x_1 \oplus x_2$ for every variable x

$$\begin{aligned} & \bar{x} \vee y \\ & \Downarrow \\ & \neg (x_1 \oplus x_2) \vee (y_1 \oplus y_2) \\ & \Downarrow \\ & (x_1 \vee \bar{x}_2 \vee y_1 \vee y_2) \\ & \wedge (x_1 \vee \bar{x}_2 \vee \bar{y}_1 \vee \bar{y}_2) \\ & \wedge (\bar{x}_1 \vee x_2 \vee y_1 \vee y_2) \\ & \wedge (\bar{x}_1 \vee x_2 \vee \bar{y}_1 \vee \bar{y}_2) \end{aligned}$$

XORification

Modify F to $F[\oplus_2]$ by substituting $x_1 \oplus x_2$ for every variable x

$$\begin{aligned} & \bar{x} \vee y \\ & \Downarrow \\ & \neg (x_1 \oplus x_2) \vee (y_1 \oplus y_2) \\ & \Downarrow \\ & (x_1 \vee \bar{x}_2 \vee y_1 \vee y_2) \\ & \wedge (x_1 \vee \bar{x}_2 \vee \bar{y}_1 \vee \bar{y}_2) \\ & \wedge (\bar{x}_1 \vee x_2 \vee y_1 \vee y_2) \\ & \wedge (\bar{x}_1 \vee x_2 \vee \bar{y}_1 \vee \bar{y}_2) \end{aligned}$$

Used to prove, e.g.:

- $\text{width} \geq w$ for $F \implies \text{size} \geq \exp(\Omega(w))$ for $F[\oplus_2]$
[Ben-Sasson '02] (credited to [Alekhnovich & Razborov])

XORification

Modify F to $F[\oplus_2]$ by substituting $x_1 \oplus x_2$ for every variable x

$$\begin{aligned} & \bar{x} \vee y \\ & \Downarrow \\ & \neg (x_1 \oplus x_2) \vee (y_1 \oplus y_2) \\ & \Downarrow \\ & (x_1 \vee \bar{x}_2 \vee y_1 \vee y_2) \\ & \wedge (x_1 \vee \bar{x}_2 \vee \bar{y}_1 \vee \bar{y}_2) \\ & \wedge (\bar{x}_1 \vee x_2 \vee y_1 \vee y_2) \\ & \wedge (\bar{x}_1 \vee x_2 \vee \bar{y}_1 \vee \bar{y}_2) \end{aligned}$$

Used to prove, e.g.:

- ▶ $\text{width} \geq w$ for $F \implies \text{size} \geq \exp(\Omega(w))$ for $F[\oplus_2]$
[Ben-Sasson '02] (credited to [Alekhnovich & Razborov])
- ▶ $\# \text{ vars in memory} \geq s$ for $F \implies \text{clause space} \geq \Omega(s)$ for $F[\oplus_2]$
[Ben-Sasson & Nordström '08]

Intuition for XORification Lower Bounds

How to construct resolution refutation π of $F[\oplus_2]$?

Intuition for XORification Lower Bounds

How to construct resolution refutation π of $F[\oplus_2]$?

Naive idea: Simulate resolution refutation π' of F
(using substitution on previous slide)

Intuition for XORification Lower Bounds

How to construct resolution refutation π of $F[\oplus_2]$?

Naive idea: Simulate resolution refutation π' of F
(using substitution on previous slide)

Seems like a bad idea—XORification causes bad blow-up

- ▶ linear in $\#$ variables in memory
- ▶ exponential in width

Intuition for XORification Lower Bounds

How to construct resolution refutation π of $F[\oplus_2]$?

Naive idea: Simulate resolution refutation π' of F
(using substitution on previous slide)

Seems like a bad idea—XORification causes bad blow-up

- ▶ linear in $\#$ variables in memory
- ▶ exponential in width

Nevertheless, can prove (sort of) this is the best resolution can do

Intuition for XORification Lower Bounds

How to construct resolution refutation π of $F[\oplus_2]$?

Naive idea: Simulate resolution refutation π' of F
(using substitution on previous slide)

Seems like a bad idea—XORification causes bad blow-up

- ▶ linear in $\#$ variables in memory
- ▶ exponential in width

Nevertheless, can prove (sort of) this is the best resolution can do

Intuition behind proof


- ▶ Given resolution refutation π of $F[\oplus_2]$
- ▶ Extract the refutation π' of F that π is simulating
- ▶ Prove that extraction preserves complexity measures of interest

Pebbling Formulas

Encode **pebble games** on DAGs

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$




- sources are true
- truth propagates upwards
- but sink is false

Pebbling Formulas

Encode **pebble games** on DAGs

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$




- ▶ **sources are true**
- ▶ truth propagates upwards
- ▶ but sink is false

Pebbling Formulas

Encode **pebble games** on DAGs

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$




- sources are true
- **truth propagates upwards**
- but sink is false

Pebbling Formulas

Encode **pebble games** on DAGs

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$




- sources are true
- truth propagates upwards
- but sink is false

Pebbling Formulas

Encode **pebble games** on DAGs

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$




- sources are true
- **truth propagates upwards**
- but sink is false

Pebbling Formulas

Encode **pebble games** on DAGs

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$




- sources are true
- truth propagates upwards
- but sink is false

Pebbling Formulas

Encode **pebble games on DAGs**

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$



- sources are true
- truth propagates upwards
- but sink is false

Written in CNF as explained before, e.g.


$$\begin{aligned}u_1 \oplus u_2 &= (u_1 \vee u_2) \wedge (\bar{u}_1 \vee \bar{u}_2) \\ \neg(z_1 \oplus z_2) &= (z_1 \vee \bar{z}_2) \wedge (\bar{z}_1 \vee z_2)\end{aligned}$$

Pebbling Formulas

Encode **pebble games on DAGs**

[Ben-Sasson & Wigderson '99]

1. $u_1 \oplus u_2$
2. $v_1 \oplus v_2$
3. $w_1 \oplus w_2$
4. $(u_1 \oplus u_2) \wedge (v_1 \oplus v_2) \rightarrow (x_1 \oplus x_2)$
5. $(v_1 \oplus v_2) \wedge (w_1 \oplus w_2) \rightarrow (y_1 \oplus y_2)$
6. $(x_1 \oplus x_2) \wedge (y_1 \oplus y_2) \rightarrow (z_1 \oplus z_2)$
7. $\neg(z_1 \oplus z_2)$



- sources are true
- truth propagates upwards
- but sink is false

Written in CNF as explained before, e.g.

$$\begin{aligned}u_1 \oplus u_2 &= (u_1 \vee u_2) \wedge (\bar{u}_1 \vee \bar{u}_2) \\ \neg(z_1 \oplus z_2) &= (z_1 \vee \bar{z}_2) \wedge (\bar{z}_1 \vee z_2)\end{aligned}$$

Easy to refute pebbling formulas in size $\mathcal{O}(n)$ and width $\mathcal{O}(1)$

Pebbling space lower bounds \Rightarrow clause space lower bounds

[Ben-Sasson & Nordström '08, '11]

XOR Substitution with Recycling (1/2)

Suppose

- ▶ F CNF formula over variables U
- ▶ $\mathcal{G} = (U \dot{\cup} V, E)$ bipartite graph

Substituted formula $F[\mathcal{G}]$ over variables V :

- ▶ replace every $u \in U$ by $\bigoplus_{v \in N(u)} v$


XOR Substitution with Recycling (1/2)

Suppose

- ▶ F CNF formula over variables U
- ▶ $\mathcal{G} = (U \dot{\cup} V, E)$ bipartite graph

Substituted formula $F[\mathcal{G}]$ over variables V :

- ▶ replace every $u \in U$ by $\bigoplus_{v \in N(u)} v$




XOR Substitution with Recycling (1/2)

Suppose

- ▶ F CNF formula over variables U
- ▶ $\mathcal{G} = (U \dot{\cup} V, E)$ bipartite graph

Substituted formula $F[\mathcal{G}]$ over variables V :

- ▶ replace every $u \in U$ by $\bigoplus_{v \in N(u)} v$



$$\bar{u}_1 \vee u_3 \longrightarrow \neg (v_1 \oplus v_2) \vee (v_5 \oplus v_6)$$


XOR Substitution with Recycling (1/2)

Suppose

- ▶ F CNF formula over variables U
- ▶ $\mathcal{G} = (U \dot{\cup} V, E)$ bipartite graph

Substituted formula $F[\mathcal{G}]$ over variables V :

- ▶ replace every $u \in U$ by $\bigoplus_{v \in N(u)} v$




XOR Substitution with Recycling (1/2)

Suppose

- ▶ F CNF formula over variables U
- ▶ $\mathcal{G} = (U \dot{\cup} V, E)$ bipartite graph


Substituted formula $F[\mathcal{G}]$ over variables V :

- ▶ replace every $u \in U$ by $\bigoplus_{v \in N(u)} v$




$$\overline{u_2} \vee u_5 \quad \longrightarrow \quad \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

XOR Substitution with Recycling (2/2)



$$\overline{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$


XOR Substitution with Recycling (2/2)



$$\overline{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$


XOR Substitution with Recycling (2/2)



$$\bar{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$


XOR Substitution with Recycling (2/2)



$$\overline{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$
 - ▶ $F[\mathcal{G}]$ refutable in width $\leq w$


XOR Substitution with Recycling (2/2)



$$\bar{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$
 - ▶ $F[\mathcal{G}]$ refutable in width $\leq w$ ✓


XOR Substitution with Recycling (2/2)



$$\bar{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$
 - ▶ $F[\mathcal{G}]$ refutable in width $\leq w$ ✓
 - ▶ space of width- w refutation of $F[\mathcal{G}] \gtrsim$
space of refutation of $F = \Omega(n/\log n) = |V|^{\Omega(w)}$


XOR Substitution with Recycling (2/2)



$$\bar{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$
 - ▶ $F[\mathcal{G}]$ refutable in width $\leq w$ ✓
 - ▶ space of width- w refutation of $F[\mathcal{G}] \gtrsim$
 space of refutation of $F = \Omega(n/\log n) = |V|^{\Omega(w)}$?

XOR Substitution with Recycling (2/2)




$$\bar{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$

$$u_6 \longrightarrow (v_4 \oplus v_5)$$

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$
 - ▶ $F[\mathcal{G}]$ refutable in width $\leq w$ ✓
 - ▶ space of width- w refutation of $F[\mathcal{G}] \gtrsim$
space of refutation of $F = \Omega(n/\log n) = |V|^{\Omega(w)}$?

XOR Substitution with Recycling (2/2)



$$\bar{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$


$$u_6 \longrightarrow (v_4 \oplus v_5)$$

$$\bar{u}_7 \longrightarrow \neg (v_4 \oplus v_5)$$



- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$
 - ▶ $F[\mathcal{G}]$ refutable in width $\leq w$ ✓
 - ▶ space of width- w refutation of $F[\mathcal{G}] \gtrsim$
 space of refutation of $F = \Omega(n/\log n) = |V|^{\Omega(w)}$?

XOR Substitution with Recycling (2/2)



$$\bar{u}_2 \vee u_5 \longrightarrow \neg (v_2 \oplus v_3 \oplus v_4) \vee (v_4 \oplus v_6)$$


$$u_6 \longrightarrow (v_4 \oplus v_5)$$

$$\bar{u}_7 \longrightarrow \neg (v_4 \oplus v_5)$$

Solution: Use expander graphs!

- ▶ Apply to pebbling formulas F in [Ben-Sasson & Nordström '08]
 - ▶ refutable in width 6
 - ▶ require space $\Omega(n/\log n)$
- ▶ \mathcal{G} **expander** with left-degree $\leq w/6$, $|U|=n$, and $|V|=n^{\mathcal{O}(1/w)}$
 - ▶ $F[\mathcal{G}]$ refutable in width $\leq w$ ✓
 - ▶ space of width- w refutation of $F[\mathcal{G}] \gtrsim$
space of refutation of $F = \Omega(n/\log n) = |V|^{\Omega(w)}$ ✓

Bipartite Boundary Expander




$\mathcal{G} = (U \dot{\cup} V, E)$ is (d, r, c) -boundary expander if

- ▶ left-degree $\leq d$
- ▶ for every $U' \subseteq U$, $|U'| \leq r$ it holds that $|\partial(U')| \geq c|U'|$

$$\partial(U') := \{v \in N(U') : |N(v) \cap U'| = 1\}$$

Bipartite Boundary Expander



$\mathcal{G} = (U \dot{\cup} V, E)$ is (d, r, c) -boundary expander if


- ▶ left-degree $\leq d$
- ▶ for every $U' \subseteq U$, $|U'| \leq r$ it holds that $|\partial(U')| \geq c|U'|$

$$\partial(U') := \{v \in N(U') : |N(v) \cap U'| = 1\}$$

Example

- ▶ left-degree $d = 3$
- ▶ expanding set size $r = 3$
- ▶ expansion factor $c = 1$

Bipartite Boundary Expander



$\mathcal{G} = (U \dot{\cup} V, E)$ is (d, r, c) -boundary expander if


- ▶ left-degree $\leq d$
- ▶ for every $U' \subseteq U$, $|U'| \leq r$ it holds that $|\partial(U')| \geq c|U'|$

$$\partial(U') := \{v \in N(U') : |N(v) \cap U'| = 1\}$$

Example

- ▶ left-degree $d = 3$
- ▶ expanding set size $r = 3$
- ▶ expansion factor $c = 1$

Bipartite Boundary Expander



$\mathcal{G} = (U \dot{\cup} V, E)$ is (d, r, c) -boundary expander if


- ▶ left-degree $\leq d$
- ▶ for every $U' \subseteq U$, $|U'| \leq r$ it holds that $|\partial(U')| \geq c|U'|$

$$\partial(U') := \{v \in N(U') : |N(v) \cap U'| = 1\}$$

Example

- ▶ left-degree $d = 3$
- ▶ expanding set size $r = 3$
- ▶ expansion factor $c = 1$

Bipartite Boundary Expander



$\mathcal{G} = (U \dot{\cup} V, E)$ is (d, r, c) -boundary expander if

- ▶ left-degree $\leq d$
- ▶ for every $U' \subseteq U$, $|U'| \leq r$ it holds that $|\partial(U')| \geq c|U'|$

$$\partial(U') := \{v \in N(U') : |N(v) \cap U'| = 1\}$$

Example

- ▶ left-degree $d = 3$
- ▶ expanding set size $r = 3$
- ▶ expansion factor $c = 1$

Lemma ([Razborov '16])

For $\varepsilon > 0$ and n, d with $d \leq |V|^{\frac{1}{2}-\varepsilon}$, $|U| = n$, $|V| = n^{O(1/d)}$ there are $(d, r, 2)$ -boundary expanders \mathcal{G} with $r = d \log n$

Sketch of Proof Sketch


Look at clauses \mathcal{D} in memory in width- w refutation of $F[\mathcal{G}]$

Recover clauses \mathcal{C} in memory in “simulated refutation” of F

Sketch of Proof Sketch

Look at clauses \mathcal{D} in memory in width- w refutation of $F[\mathcal{G}]$

Recover clauses \mathcal{C} in memory in “simulated refutation” of F



$U \sim F$

$V \sim F[\mathcal{G}]$

Must have $N(\text{Vars}(\mathcal{C})) \subseteq \text{Vars}(\mathcal{D})$

$$\text{Ker}(V') := \{u \in U : N(u) \subseteq V'\}$$


$$|V'| \leq r \implies |\text{Ker}(V')| \leq |V'|$$

(since left vertex sets expand a lot)

Sketch of Proof Sketch

Look at clauses \mathcal{D} in memory in width- w refutation of $F[\mathcal{G}]$

Recover clauses \mathcal{C} in memory in “simulated refutation” of F



Must have $N(\text{Vars}(\mathcal{C})) \subseteq \text{Vars}(\mathcal{D})$

$$\text{Ker}(V') := \{u \in U : N(u) \subseteq V'\}$$

$$|V'| \leq r \implies |\text{Ker}(V')| \leq |V'|$$

(since left vertex sets expand a lot)


Example

$$V' = \{v_3, \dots, v_8\}, \text{Ker}(V') = \{u_6, u_7, u_{12}\}$$

Sketch of Proof Sketch

Look at clauses \mathcal{D} in memory in width- w refutation of $F[\mathcal{G}]$

Recover clauses \mathcal{C} in memory in “simulated refutation” of F



Must have $N(\text{Vars}(\mathcal{C})) \subseteq \text{Vars}(\mathcal{D})$

$$\text{Ker}(V') := \{u \in U : N(u) \subseteq V'\}$$

$$|V'| \leq r \implies |\text{Ker}(V')| \leq |V'|$$

(since left vertex sets expand a lot)


Example

$$V' = \{v_3, \dots, v_8\}, \text{Ker}(V') = \{u_6, u_7, u_{12}\}$$

Sketch of Proof Sketch

Look at clauses \mathcal{D} in memory in width- w refutation of $F[\mathcal{G}]$

Recover clauses \mathcal{C} in memory in “simulated refutation” of F



Must have $N(\text{Vars}(\mathcal{C})) \subseteq \text{Vars}(\mathcal{D})$

$$\text{Ker}(V') := \{u \in U : N(u) \subseteq V'\}$$

$$|V'| \leq r \implies |\text{Ker}(V')| \leq |V'|$$

(since left vertex sets expand a lot)

Example

$$V' = \{v_3, \dots, v_8\}, \text{Ker}(V') = \{u_6, u_7, u_{12}\}$$


Locally looks almost like XORification without recycling, so previous proof might work...

And give bound in terms of $|U| \gg |V|$

Sketch of Proof Sketch

Look at clauses \mathcal{D} in memory in width- w refutation of $F[\mathcal{G}]$

Recover clauses \mathcal{C} in memory in “simulated refutation” of F



Must have $N(\text{Vars}(\mathcal{C})) \subseteq \text{Vars}(\mathcal{D})$

$$\text{Ker}(V') := \{u \in U : N(u) \subseteq V'\}$$

$$|V'| \leq r \implies |\text{Ker}(V')| \leq |V'|$$

(since left vertex sets expand a lot)

Example

$$V' = \{v_3, \dots, v_8\}, \text{Ker}(V') = \{u_6, u_7, u_{12}\}$$

Locally looks almost like XORification without recycling, so previous proof might work...

And give bound in terms of $|U| \gg |V|$

Actual details very different

On the Method of Hardness Condensation

Introduced in [Razborov JACM '16] to show that **treelike** resolution refutations of **width w** can require **doubly exponential size $2^{n^{\Omega(w)}}$**

On the Method of Hardness Condensation

Introduced in [Razborov JACM '16] to show that **treelike** resolution refutations of **width w** can require **doubly exponential size $2^{n^{\Omega(w)}}$**

Has also been used to establish

- ▶ Tradeoffs between width and rank for Lovász-Schrijver linear programming hierarchy [Razborov ECCC '16]

On the Method of Hardness Condensation

Introduced in [Razborov JACM '16] to show that **treelike** resolution refutations of **width w** can require **doubly exponential size $2^{n^{\Omega(w)}}$**

Has also been used to establish

- ▶ Tradeoffs between width and rank for Lovász-Schrijver linear programming hierarchy [Razborov ECCC '16]
- ▶ Quantifier depth lower bounds for finite variable fragments of first-order logic [Berkholz & Nordström LICS '16]

On the Method of Hardness Condensation

Introduced in [Razborov JACM '16] to show that **treelike** resolution refutations of **width w** can require **doubly exponential size $2^{n^{\Omega(w)}}$**

Has also been used to establish

- ▶ Tradeoffs between width and rank for Lovász-Schrijver linear programming hierarchy [Razborov ECCC '16]
- ▶ Quantifier depth lower bounds for finite variable fragments of first-order logic [Berkholz & Nordström LICS '16]

Where else can this technique be useful?

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Open question 1

Similar tradeoffs for degree vs. space in polynomial calculus?

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Open question 1

Similar tradeoffs for degree vs. space in polynomial calculus?

- ▶ Weaknesses: non-constant width and **huge** size blow-up

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Open question 1

Similar tradeoffs for degree vs. space in polynomial calculus?

- ▶ Weaknesses: non-constant width and **huge** size blow-up
- ▶ Inherent for XORification with large arity

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Open question 1

Similar tradeoffs for degree vs. space in polynomial calculus?

- ▶ Weaknesses: non-constant width and **huge** size blow-up
- ▶ Inherent for XORification with large arity

Open question 2

Are there supercritical tradeoffs for 3-CNFs?

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Open question 1

Similar tradeoffs for degree vs. space in polynomial calculus?

- ▶ Weaknesses: non-constant width and **huge** size blow-up
- ▶ Inherent for XORification with large arity

Open question 2

Are there supercritical tradeoffs for 3-CNFs?

- ▶ Probably yes, unless $\text{PSPACE} = \text{EXPTIME}$

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Open question 1

Similar tradeoffs for degree vs. space in polynomial calculus?

- ▶ Weaknesses: non-constant width and **huge** size blow-up
- ▶ Inherent for XORification with large arity

Open question 2

Are there supercritical tradeoffs for 3-CNFs?

- ▶ Probably yes, unless $\text{PSPACE} = \text{EXPTIME}$
- ▶ Can search for small-space refutations in PSPACE, but finding refutations in given width EXPTIME-complete [[Berkholz '12](#)]

Concluding Remarks

- ▶ We exhibit **supercritical space-width trade-offs** for resolution
- ▶ Minimizing width can make space go way above linear (worst-case “critical” bound)

Open question 1

Similar tradeoffs for degree vs. space in polynomial calculus?

- ▶ Weaknesses: non-constant width and **huge** size blow-up
- ▶ Inherent for XORification with large arity

Open question 2

Are there supercritical tradeoffs for 3-CNFs?

- ▶ Probably yes, unless $\text{PSPACE} = \text{EXPTIME}$
- ▶ Can search for small-space refutations in PSPACE, but finding refutations in given width EXPTIME-complete [[Berkholz '12](#)]

Thank you for your attention!