

# Proof Complexity as a Computational Lens: Lecture 22

## Size-Space Trade-offs for Cutting Planes

Jakob Nordström

University of Copenhagen and Lund University

January 30, 2026



# Recap: Configuration-Style Proofs

- Proof system operates with formulas of some syntactic form
- Proof/refutation is “presented on blackboard”
- Derivation steps:
  - Write down axiom clauses of CNF formula being refuted (as encoded by proof system)
  - Infer new lines by deductive rules of proof system
  - Erase lines not currently needed (to save space on blackboard)
- Refutation ends when (explicit) contradiction is derived

# Cutting Planes (CP)

Clauses interpreted as linear inequalities

E.g.,  $x \vee y \vee \bar{z} \rightsquigarrow x + y + (1 - z) \geq 1 \rightsquigarrow x + y - z \geq 0$

Proof system also works for any system of 0–1 linear inequalities with integer coefficients

# Cutting Planes (CP)

Clauses interpreted as linear inequalities

E.g.,  $x \vee y \vee \bar{z} \rightsquigarrow x + y + (1 - z) \geq 1 \rightsquigarrow x + y - z \geq 0$

Proof system also works for any system of 0–1 linear inequalities with integer coefficients

*Variable axioms*  $\frac{}{0 \leq x \leq 1}$

*Addition*  $\frac{\sum a_i x_i \geq A \quad \sum b_i x_i \geq B}{\sum (a_i + b_i) x_i \geq A + B}$

*Multiplication*  $\frac{\sum a_i x_i \geq A}{\sum c a_i x_i \geq c A} \quad c \in \mathbb{N}^+$

*Division*  $\frac{\sum c a_i x_i \geq A}{\sum a_i x_i \geq \lceil A/c \rceil} \quad c \in \mathbb{N}^+$

**Goal:** Derive  $0 \geq 1 \Leftrightarrow$  formula/system of inequalities unsatisfiable

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} \vee x_{1,2}$
2.  $x_{2,1} \vee x_{2,2}$
3.  $x_{3,1} \vee x_{3,2}$
4.  $\bar{x}_{1,1} \vee \bar{x}_{2,1}$
5.  $\bar{x}_{1,1} \vee \bar{x}_{3,1}$
6.  $\bar{x}_{2,1} \vee \bar{x}_{3,1}$
7.  $\bar{x}_{1,2} \vee \bar{x}_{2,2}$
8.  $\bar{x}_{1,2} \vee \bar{x}_{3,2}$
9.  $\bar{x}_{2,2} \vee \bar{x}_{3,2}$

## Pigeonhole principle (PHP)

" $n + 1$  pigeons don't fit into  $n$  holes"

Variables  $x_{i,j}$  = "pigeon  $i$  goes into hole  $j$ "

$x_{i,1} \vee x_{i,2} \vee \dots \vee x_{i,n}$  every pigeon  $i$  gets a hole

$\bar{x}_{i,j} \vee \bar{x}_{i',j}$  no hole  $j$  gets two pigeons  $i \neq i'$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
  2.  $x_{2,1} + x_{2,2} \geq 1$
  3.  $x_{3,1} + x_{3,2} \geq 1$
  4.  $-x_{1,1} - x_{2,1} \geq -1$
  5.  $-x_{1,1} - x_{3,1} \geq -1$
  6.  $-x_{2,1} - x_{3,1} \geq -1$
  7.  $-x_{1,2} - x_{2,2} \geq -1$
  8.  $-x_{1,2} - x_{3,2} \geq -1$
  9.  $-x_{2,2} - x_{3,2} \geq -1$
- 



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$

2.  $x_{2,1} + x_{2,2} \geq 1$

3.  $x_{3,1} + x_{3,2} \geq 1$

4.  $-x_{1,1} - x_{2,1} \geq -1$

5.  $-x_{1,1} - x_{3,1} \geq -1$

6.  $-x_{2,1} - x_{3,1} \geq -1$

7.  $-x_{1,2} - x_{2,2} \geq -1$

8.  $-x_{1,2} - x_{3,2} \geq -1$

9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

$$-x_{1,1} - x_{2,1} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

$$-x_{1,1} - x_{2,1} \geq -1$$

$$-x_{1,1} - x_{3,1} \geq -1$$



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

$$-x_{1,1} - x_{2,1} \geq -1$$

$$-x_{1,1} - x_{3,1} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

$$\begin{aligned} -x_{1,1} - x_{2,1} &\geq -1 \\ -x_{1,1} - x_{3,1} &\geq -1 \\ -2x_{1,1} - x_{2,1} - x_{3,1} &\geq -2 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

**Erase** the line  $-x_{1,1} - x_{3,1} \geq -1$

$$\begin{aligned} &-x_{1,1} - x_{2,1} \geq -1 \\ &\textcolor{blue}{-x_{1,1} - x_{3,1} \geq -1} \\ &-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

$$\begin{aligned} -x_{1,1} - x_{2,1} &\geq -1 \\ -2x_{1,1} - x_{2,1} - x_{3,1} &\geq -2 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

$$\begin{aligned} -x_{1,1} - x_{2,1} &\geq -1 \\ -2x_{1,1} - x_{2,1} - x_{3,1} &\geq -2 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

$$-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

$$\begin{aligned} &-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2 \\ &-x_{2,1} - x_{3,1} \geq -1 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

$$-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$$

$$-x_{2,1} - x_{3,1} \geq -1$$



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

$$\begin{aligned} &-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2 \\ &-x_{2,1} - x_{3,1} \geq -1 \\ &-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

**Erase** the line  $-x_{2,1} - x_{3,1} \geq -1$

$$-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$$

$$-x_{2,1} - x_{3,1} \geq -1$$

$$-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

**Erase** the line  $-x_{2,1} - x_{3,1} \geq -1$

$$\begin{aligned} -2x_{1,1} - x_{2,1} - x_{3,1} &\geq -2 \\ -2x_{1,1} - 2x_{2,1} - 2x_{3,1} &\geq -3 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

**Erase** the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

$$\begin{aligned} -2x_{1,1} - x_{2,1} - x_{3,1} &\geq -2 \\ -2x_{1,1} - 2x_{2,1} - 2x_{3,1} &\geq -3 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

**Erase** the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

$$-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

$$-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 4:  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

$$-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

**Erase** the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

$$-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 5:  $-x_{1,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

**Erase** the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

$$1. \quad x_{1,1} + x_{1,2} \geq 1$$

$$2. \quad x_{2,1} + x_{2,2} \geq 1$$

$$3. \quad x_{3,1} + x_{3,2} \geq 1$$

$$4. \quad -x_{1,1} - x_{2,1} \geq -1$$

$$5. \quad -x_{1,1} - x_{3,1} \geq -1$$

$$6. \quad -x_{2,1} - x_{3,1} \geq -1$$

$$7. \quad -x_{1,2} - x_{2,2} \geq -1$$

$$8. \quad -x_{1,2} - x_{3,2} \geq -1$$

$$9. \quad -x_{2,2} - x_{3,2} \geq -1$$

## History of derivation steps

Add to get  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,1} - x_{3,1} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} \geq -1$$

$$-x_{1,2} - x_{3,2} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} \geq -1$$

$$-x_{1,2} - x_{3,2} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,1} - x_{2,1} \geq -1$

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} \geq -1$$

$$-x_{1,2} - x_{3,2} \geq -1$$

$$-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} \geq -1$$

$$-x_{1,2} - x_{3,2} \geq -1$$

$$-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 6:  $-x_{2,1} - x_{3,1} \geq -1$

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} \geq -1$$

$$-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} \geq -1$$

$$-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$



# Example: Cutting planes Refutation of Pigeonhole Principle

$$1. \quad x_{1,1} + x_{1,2} \geq 1$$

$$2. \quad x_{2,1} + x_{2,2} \geq 1$$

$$3. \quad x_{3,1} + x_{3,2} \geq 1$$

$$4. \quad -x_{1,1} - x_{2,1} \geq -1$$

$$5. \quad -x_{1,1} - x_{3,1} \geq -1$$

$$6. \quad -x_{2,1} - x_{3,1} \geq -1$$

$$7. \quad -x_{1,2} - x_{2,2} \geq -1$$

$$8. \quad -x_{1,2} - x_{3,2} \geq -1$$

$$9. \quad -x_{2,2} - x_{3,2} \geq -1$$

## History of derivation steps

Add to get  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

**Erase** the line  $-x_{1,2} - x_{2,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2 \\ &-x_{2,2} - x_{3,2} \geq -1 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2 \\ &-x_{2,2} - x_{3,2} \geq -1 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-2x_{1,1} - x_{2,1} - x_{3,1} \geq -2$

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2 \\ &-x_{2,2} - x_{3,2} \geq -1 \\ &-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2 \\ &-x_{2,2} - x_{3,2} \geq -1 \\ &-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Divide to get  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2 \\ &-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

**Erase** the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &\textcolor{blue}{-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2} \\ &-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-2x_{1,1} - 2x_{2,1} - 2x_{3,1} \geq -3$

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

**Erase** the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

$$\begin{aligned} -x_{1,1} - x_{2,1} - x_{3,1} &\geq -1 \\ -2x_{1,2} - 2x_{2,2} - 2x_{3,2} &\geq -3 \end{aligned}$$



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 7:  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3 \\ &-x_{1,2} - x_{2,2} - x_{3,2} \geq -1 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

**Erase** the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

$$\begin{aligned} &-x_{1,1} - x_{2,1} - x_{3,1} \geq -1 \\ &\textcolor{blue}{-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3} \\ &-x_{1,2} - x_{2,2} - x_{3,2} \geq -1 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 8:  $-x_{1,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

**Erase** the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

$$1. \quad x_{1,1} + x_{1,2} \geq 1$$

$$2. \quad x_{2,1} + x_{2,2} \geq 1$$

$$3. \quad x_{3,1} + x_{3,2} \geq 1$$

$$4. \quad -x_{1,1} - x_{2,1} \geq -1$$

$$5. \quad -x_{1,1} - x_{3,1} \geq -1$$

$$6. \quad -x_{2,1} - x_{3,1} \geq -1$$

$$7. \quad -x_{1,2} - x_{2,2} \geq -1$$

$$8. \quad -x_{1,2} - x_{3,2} \geq -1$$

$$9. \quad -x_{2,2} - x_{3,2} \geq -1$$

## History of derivation steps

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

$$1. \quad x_{1,1} + x_{1,2} \geq 1$$

$$2. \quad x_{2,1} + x_{2,2} \geq 1$$

$$3. \quad x_{3,1} + x_{3,2} \geq 1$$

$$4. \quad -x_{1,1} - x_{2,1} \geq -1$$

$$5. \quad -x_{1,1} - x_{3,1} \geq -1$$

$$6. \quad -x_{2,1} - x_{3,1} \geq -1$$

$$7. \quad -x_{1,2} - x_{2,2} \geq -1$$

$$8. \quad -x_{1,2} - x_{3,2} \geq -1$$

$$9. \quad -x_{2,2} - x_{3,2} \geq -1$$

## History of derivation steps

Add to get  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

**Erase** the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,2} - x_{2,2} \geq -1$

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

**Erase** the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

$$1. \quad x_{1,1} + x_{1,2} \geq 1$$

$$2. \quad x_{2,1} + x_{2,2} \geq 1$$

$$3. \quad x_{3,1} + x_{3,2} \geq 1$$

$$4. \quad -x_{1,1} - x_{2,1} \geq -1$$

$$5. \quad -x_{1,1} - x_{3,1} \geq -1$$

$$6. \quad -x_{2,1} - x_{3,1} \geq -1$$

$$7. \quad -x_{1,2} - x_{2,2} \geq -1$$

$$8. \quad -x_{1,2} - x_{3,2} \geq -1$$

$$9. \quad -x_{2,2} - x_{3,2} \geq -1$$

## History of derivation steps

Write down axiom 9:  $-x_{2,2} - x_{3,2} \geq -1$

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} \geq 1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

$$1. \quad x_{1,1} + x_{1,2} \geq 1$$

$$2. \quad x_{2,1} + x_{2,2} \geq 1$$

$$3. \quad x_{3,1} + x_{3,2} \geq 1$$

$$4. \quad -x_{1,1} - x_{2,1} \geq -1$$

$$5. \quad -x_{1,1} - x_{3,1} \geq -1$$

$$6. \quad -x_{2,1} - x_{3,1} \geq -1$$

$$7. \quad -x_{1,2} - x_{2,2} \geq -1$$

$$8. \quad -x_{1,2} - x_{3,2} \geq -1$$

$$9. \quad -x_{2,2} - x_{3,2} \geq -1$$

## History of derivation steps

Add to get  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} \geq 1$$

$$x_{2,1} + x_{2,2} \geq 1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} \geq 1$$

$$x_{2,1} + x_{2,2} \geq 1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} \geq 1$$

$$x_{2,1} + x_{2,2} \geq 1$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} \geq 1$$

$$x_{2,1} + x_{2,2} \geq 1$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-2x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} \geq 1$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} \geq 1$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Divide to get  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

**Erase** the line  $x_{1,1} + x_{1,2} \geq 1$

$$\begin{aligned} -x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} &\geq -2 \\ x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} &\geq 2 \end{aligned}$$

# Example: Cutting planes Refutation of Pigeonhole Principle

$$1. \quad x_{1,1} + x_{1,2} \geq 1$$

$$2. \quad x_{2,1} + x_{2,2} \geq 1$$

$$3. \quad x_{3,1} + x_{3,2} \geq 1$$

$$4. \quad -x_{1,1} - x_{2,1} \geq -1$$

$$5. \quad -x_{1,1} - x_{3,1} \geq -1$$

$$6. \quad -x_{2,1} - x_{3,1} \geq -1$$

$$7. \quad -x_{1,2} - x_{2,2} \geq -1$$

$$8. \quad -x_{1,2} - x_{3,2} \geq -1$$

$$9. \quad -x_{2,2} - x_{3,2} \geq -1$$

## History of derivation steps

Erase the line  $-2x_{1,2} - 2x_{2,2} - 2x_{3,2} \geq -3$

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

$$x_{3,1} + x_{3,2} \geq 1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

$$x_{3,1} + x_{3,2} \geq 1$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Add to get  $-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

$$x_{3,1} + x_{3,2} \geq 1$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

Erase the line  $x_{3,1} + x_{3,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

$$x_{3,1} + x_{3,2} \geq 1$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,2} - x_{2,2} - x_{3,2} \geq -1$

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

Erase the line  $x_{3,1} + x_{3,2} \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

Erase the line  $x_{3,1} + x_{3,2} \geq 1$

**Erase** the line  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$$



# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Erase the line  $-x_{1,1} - x_{2,1} - x_{3,1} \geq -1$

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

Erase the line  $x_{3,1} + x_{3,2} \geq 1$

**Erase** the line  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

Erase the line  $x_{3,1} + x_{3,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Add to get  $0 \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$$

# Example: Cutting planes Refutation of Pigeonhole Principle

1.  $x_{1,1} + x_{1,2} \geq 1$
2.  $x_{2,1} + x_{2,2} \geq 1$
3.  $x_{3,1} + x_{3,2} \geq 1$
4.  $-x_{1,1} - x_{2,1} \geq -1$
5.  $-x_{1,1} - x_{3,1} \geq -1$
6.  $-x_{2,1} - x_{3,1} \geq -1$
7.  $-x_{1,2} - x_{2,2} \geq -1$
8.  $-x_{1,2} - x_{3,2} \geq -1$
9.  $-x_{2,2} - x_{3,2} \geq -1$

## History of derivation steps

Write down axiom 1:  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 2:  $x_{2,1} + x_{2,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Erase the line  $x_{2,1} + x_{2,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} \geq 1$

Write down axiom 3:  $x_{3,1} + x_{3,2} \geq 1$

Add to get  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$

Erase the line  $x_{3,1} + x_{3,2} \geq 1$

Erase the line  $x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} \geq 2$

Add to get  $0 \geq 1$

$$-x_{1,1} - x_{2,1} - x_{3,1} - x_{1,2} - x_{2,2} - x_{3,2} \geq -2$$

$$x_{1,1} + x_{1,2} + x_{2,1} + x_{2,2} + x_{3,1} + x_{3,2} \geq 3$$

$$0 \geq 1$$

# Complexity Measures for Cutting Planes

**Length** = total # lines/inequalities in refutation

**Size** = sum also size of coefficients

**Line space** = max # lines in memory during refutation

**Total space** = max # bits in memory (sum also size of coefficients)

# Size Lower Bounds for Cutting Planes

## Clique-colouring formulas

“A graph with an  $m$ -clique is not  $(m-1)$ -colourable”

Exponential lower bound via interpolation and circuit complexity [Pud97]

Technique very specifically tied to structure of formula

# Size Lower Bounds for Cutting Planes

## Clique-colouring formulas

“A graph with an  $m$ -clique is not  $(m-1)$ -colourable”

Exponential lower bound via interpolation and circuit complexity [Pud97]

Technique very specifically tied to structure of formula

## Random $\mathcal{O}(\log n)$ -CNF formulas

“Large number of randomly sampled clauses can be satisfied”

Exponential lower bound via bottleneck counting argument [Sok24]

Very intriguing new technique! (Or circuit lower bound in disguise?)

# What About Line Space in Cutting Planes?

## Pebbling formulas

“Possible to get from sources to sink in connected directed acyclic graph”

Short cutting planes refutations of (lifted) pebbling formulas on certain DAGs must have large line space [HN12, GP18]  
(and such short refutations do exist)

# What About Line Space in Cutting Planes?

## Pebbling formulas

“Possible to get from sources to sink in connected directed acyclic graph”

Short cutting planes refutations of (lifted) pebbling formulas on certain DAGs must have large line space [HN12, GP18]  
(and such short refutations do exist)

## Tseitin formulas

“Sum of degrees of vertices in graph is even”

Short refutations of (lifted) Tseitin formulas on expanders must have large line space [GP18]  
Not clear whether such short refutations exist. . .



# Size-Space Trade-offs for Cutting Planes?

- **Surprise:** Cutting planes can refute any CNF in **line space 5** (!) [GPT15]  
(But coefficients will be exponentially large)

# Size-Space Trade-offs for Cutting Planes?

- **Surprise:** Cutting planes can refute any CNF in **line space 5** (!) [GPT15]  
(But coefficients will be exponentially large)
- Plug into [HN12, GP18]  $\Rightarrow$  get trade-off of sorts

# Size-Space Trade-offs for Cutting Planes?

- **Surprise:** Cutting planes can refute any CNF in **line space 5** (!) [GPT15]  
(But coefficients will be exponentially large)
- Plug into [HN12, GP18]  $\Rightarrow$  get trade-off of sorts
- But “constant-space” proofs with exponential-size coefficients somehow doesn't feel quite right. . .

# Size-Space Trade-offs for Cutting Planes?

- **Surprise:** Cutting planes can refute any CNF in **line space 5** (!) [GPT15]  
(But coefficients will be exponentially large)
- Plug into [HN12, GP18]  $\Rightarrow$  get trade-off of sorts
- But “constant-space” proofs with exponential-size coefficients somehow doesn't feel quite right. . .

What about “true” trade-offs?

Are there **trade-offs** where the space-efficient cutting planes refutations have **small coefficients**? (Say, of polynomial or even constant size)

# Focus of This Lecture (and Next Lecture)

## Theorem (Informal sample)

*There are families of 6-CNF formulas  $\{F_N\}_{N=1}^{\infty}$  of size  $\Theta(N)$  such that:*

# Focus of This Lecture (and Next Lecture)

## Theorem (Informal sample)

*There are families of 6-CNF formulas  $\{F_N\}_{N=1}^{\infty}$  of size  $\Theta(N)$  such that:*

- 1  *$F_N$  can be refuted by cutting planes with constant-size coefficients in size  $\mathcal{O}(N)$  and total space  $\mathcal{O}(N^{2/5})$*

# Focus of This Lecture (and Next Lecture)

## Theorem (Informal sample)

*There are families of 6-CNF formulas  $\{F_N\}_{N=1}^{\infty}$  of size  $\Theta(N)$  such that:*

- 1  *$F_N$  can be refuted by cutting planes with constant-size coefficients in size  $\mathcal{O}(N)$  and total space  $\mathcal{O}(N^{2/5})$*
- 2  *$F_N$  can be refuted by cutting planes with constant-size coefficients in total space  $\mathcal{O}(N^{1/40})$  and size  $\exp(\mathcal{O}(N^{1/40}))$*

# Focus of This Lecture (and Next Lecture)

## Theorem (Informal sample)

*There are families of 6-CNF formulas  $\{F_N\}_{N=1}^{\infty}$  of size  $\Theta(N)$  such that:*

- ①  *$F_N$  can be refuted by cutting planes with constant-size coefficients in size  $\mathcal{O}(N)$  and total space  $\mathcal{O}(N^{2/5})$*
- ②  *$F_N$  can be refuted by cutting planes with constant-size coefficients in total space  $\mathcal{O}(N^{1/40})$  and size  $\exp(\mathcal{O}(N^{1/40}))$*
- ③ *Any cutting planes refutation even with coefficients of unbounded size in line space less than  $N^{1/20-\epsilon}$  requires length  $\exp(\Omega(N^{1/40}))$*



# Focus of This Lecture (and Next Lecture)

## Theorem (Informal sample)

*There are families of 6-CNF formulas  $\{F_N\}_{N=1}^{\infty}$  of size  $\Theta(N)$  such that:*

- ①  *$F_N$  can be refuted by cutting planes with constant-size coefficients in size  $\mathcal{O}(N)$  and total space  $\mathcal{O}(N^{2/5})$*
- ②  *$F_N$  can be refuted by cutting planes with constant-size coefficients in total space  $\mathcal{O}(N^{1/40})$  and size  $\exp(\mathcal{O}(N^{1/40}))$*
- ③ *Any cutting planes refutation even with coefficients of unbounded size in line space less than  $N^{1/20-\epsilon}$  requires length  $\exp(\Omega(N^{1/40}))$*

- Upper bounds for # bits; lower bounds for # lines/inequalities

# Focus of This Lecture (and Next Lecture)

## Theorem (Informal sample)

*There are families of 6-CNF formulas  $\{F_N\}_{N=1}^{\infty}$  of size  $\Theta(N)$  such that:*

- ❶  *$F_N$  can be refuted by cutting planes with constant-size coefficients in size  $\mathcal{O}(N)$  and total space  $\mathcal{O}(N^{2/5})$*
- ❷  *$F_N$  can be refuted by cutting planes with constant-size coefficients in total space  $\mathcal{O}(N^{1/40})$  and size  $\exp(\mathcal{O}(N^{1/40}))$*
- ❸ *Any cutting planes refutation even with coefficients of unbounded size in line space less than  $N^{1/20-\epsilon}$  requires length  $\exp(\Omega(N^{1/40}))$*

- Upper bounds for # bits; lower bounds for # lines/inequalities
- Hold uniformly for resolution, polynomial calculus, and cutting planes

# Focus of This Lecture (and Next Lecture)

## Theorem (Informal sample)

*There are families of 6-CNF formulas  $\{F_N\}_{N=1}^{\infty}$  of size  $\Theta(N)$  such that:*

- ①  *$F_N$  can be refuted by cutting planes with constant-size coefficients in size  $\mathcal{O}(N)$  and total space  $\mathcal{O}(N^{2/5})$*
- ②  *$F_N$  can be refuted by cutting planes with constant-size coefficients in total space  $\mathcal{O}(N^{1/40})$  and size  $\exp(\mathcal{O}(N^{1/40}))$*
- ③ *Any cutting planes refutation even with coefficients of unbounded size in line space less than  $N^{1/20-\epsilon}$  requires length  $\exp(\Omega(N^{1/40}))$*

- Upper bounds for # bits; lower bounds for # lines/inequalities
- Hold uniformly for resolution, polynomial calculus, and cutting planes
- Even for **semantic** proofs where anything implied by blackboard inferred in single step

# Outline of Proof

Proof is by carefully constructed chain of delicate reductions

# Outline of Proof

Proof is by carefully constructed chain of delicate reductions  
(a.k.a. the kitchen sink)

# Outline of Proof

Proof is by carefully constructed chain of delicate reductions  
(a.k.a. the kitchen sink)

- 1 Short, space-efficient proof  $\Rightarrow$  efficient communication protocol for falsified clause search problem

# Outline of Proof

Proof is by carefully constructed chain of delicate reductions  
(a.k.a. the kitchen sink)

- ① Short, space-efficient proof  $\Rightarrow$  efficient communication protocol for falsified clause search problem
- ② Crucial twists:
  - Study real communication model
  - Consider round efficiency of protocols

# Outline of Proof

Proof is by carefully constructed chain of delicate reductions  
(a.k.a. the kitchen sink)

- ① Short, space-efficient proof  $\Rightarrow$  efficient communication protocol for falsified clause search problem
- ② Crucial twists:
  - Study real communication model
  - Consider round efficiency of protocols
- ③ Protocol for composed search problem  $\Rightarrow$  parallel decision tree via simulation theorem



# Outline of Proof

Proof is by carefully constructed chain of delicate reductions  
(a.k.a. the kitchen sink)

- ① Short, space-efficient proof  $\Rightarrow$  efficient communication protocol for falsified clause search problem
- ② Crucial twists:
  - Study real communication model
  - Consider round efficiency of protocols
- ③ Protocol for composed search problem  $\Rightarrow$  parallel decision tree via simulation theorem
- ④ Parallel decision tree for pebbling formulas  $Peb_G$   
 $\Rightarrow$  pebbling strategy for Dymond–Tompkins game on graph  $G$

# Outline of Proof

Proof is by carefully constructed chain of delicate reductions  
(a.k.a. the kitchen sink)

- ① Short, space-efficient proof  $\Rightarrow$  efficient communication protocol for falsified clause search problem
- ② Crucial twists:
  - Study real communication model
  - Consider round efficiency of protocols
- ③ Protocol for composed search problem  $\Rightarrow$  parallel decision tree via simulation theorem
- ④ Parallel decision tree for pebbling formulas  $Peb_G$   
 $\Rightarrow$  pebbling strategy for Dymond–Tompkins game on graph  $G$
- ⑤ Construct graphs  $G$  with strong round-cost trade-offs for Dymond–Tompkins pebbling

# Outline of Proof

Proof is by carefully constructed chain of delicate reductions  
(a.k.a. the kitchen sink)

- ① Short, space-efficient proof  $\Rightarrow$  efficient communication protocol for falsified clause search problem [HN12]
- ② Crucial twists:
  - Study real communication model [Kra98, BEGJ00]
  - Consider round efficiency of protocols
- ③ Protocol for composed search problem  $\Rightarrow$  parallel decision tree [Val75] via simulation theorem à la [RM99, GPW15]
- ④ Parallel decision tree for pebbling formulas  $Peb_G$  [BW01]  $\Rightarrow$  pebbling strategy for Dymond–Tompkins game on graph  $G$  [DT85]
- ⑤ Construct graphs  $G$  with strong round-cost trade-offs for Dymond–Tompkins pebbling inspired by [CS82, LT82, BN11, Nor12]

# Deterministic Communication

- Two players:
  - Alice with private input  $x$
  - Bob with private input  $y$
  - Both deterministic but with unbounded computational powers

# Deterministic Communication

- Two players:
  - Alice with private input  $x$
  - Bob with private input  $y$
  - Both deterministic but with unbounded computational powers
- Task: Compute  $f(x, y)$  by communicating according to protocol (= tree)

# Deterministic Communication

- Two players:
  - Alice with private input  $x$
  - Bob with private input  $y$
  - Both deterministic but with unbounded computational powers
- Task: Compute  $f(x, y)$  by communicating according to protocol (= tree)
- Method: Each node  $v$  of protocol tree specifies:
  - Either Alice sends a bit  $a_v \in \{0, 1\}$
  - Or Bob sends a bit  $b_v \in \{0, 1\}$
  - A round starts when the sender changes

# Deterministic Communication

- Two players:
  - Alice with private input  $x$
  - Bob with private input  $y$
  - Both deterministic but with unbounded computational powers
- Task: Compute  $f(x, y)$  by communicating according to protocol (= tree)
- Method: Each node  $v$  of protocol tree specifies:
  - Either Alice sends a bit  $a_v \in \{0, 1\}$
  - Or Bob sends a bit  $b_v \in \{0, 1\}$
  - A round starts when the sender changes
- When they reach any leaf, both Alice and Bob should have learned  $f(x, y)$

# Deterministic Communication

- Two players:
  - Alice with private input  $x$
  - Bob with private input  $y$
  - Both deterministic but with unbounded computational powers
- Task: Compute  $f(x, y)$  by communicating according to protocol (= tree)
- Method: Each node  $v$  of protocol tree specifies:
  - Either Alice sends a bit  $a_v \in \{0, 1\}$
  - Or Bob sends a bit  $b_v \in \{0, 1\}$
  - A round starts when the sender changes
- When they reach any leaf, both Alice and Bob should have learned  $f(x, y)$
- Function  $f$  solved by  $r$ -round deterministic communication in cost  $c$  if  $\exists$  protocol tree such that along any path from root
  - # rounds  $\leq r$
  - total # bits sent  $\leq c$



# Real Communication [Kra98]

- Same players **Alice** with **private input  $x$**  and **Bob** with **private input  $y$**

# Real Communication [Kra98]

- Same players Alice with private input  $x$  and Bob with private input  $y$
- Task: Compute  $f(x, y)$  by sending real numbers to referee

# Real Communication [Kra98]

- Same players Alice with private input  $x$  and Bob with private input  $y$
- Task: Compute  $f(x, y)$  by sending real numbers to referee
- Method: In each round  $v$ 
  - Alice sends  $a_{v,1}(x), \dots, a_{v,c_v}(x) \in \mathbb{R}^{c_v}$
  - Bob sends  $b_{v,1}(y), \dots, b_{v,c_v}(y) \in \mathbb{R}^{c_v}$
  - Referee announces results of comparisons  $a_{v,i}(x) \leq b_{v,i}(y)$  for  $i \in [c_v]$

# Real Communication [Kra98]

- Same players Alice with private input  $x$  and Bob with private input  $y$
- Task: Compute  $f(x, y)$  by sending real numbers to referee
- Method: In each round  $v$ 
  - Alice sends  $a_{v,1}(x), \dots, a_{v,c_v}(x) \in \mathbb{R}^{c_v}$
  - Bob sends  $b_{v,1}(y), \dots, b_{v,c_v}(y) \in \mathbb{R}^{c_v}$
  - Referee announces results of comparisons  $a_{v,i}(x) \leq b_{v,i}(y)$  for  $i \in [c_v]$
- Function  $f$  solved by  $r$ -round real communication in cost  $c$  if  $\exists$  protocol such that
  - # rounds  $\leq r$
  - total # comparisons made by referee  $\leq c$

# Real Communication [Kra98]

- Same players Alice with private input  $x$  and Bob with private input  $y$
- Task: Compute  $f(x, y)$  by sending real numbers to referee
- Method: In each round  $v$ 
  - Alice sends  $a_{v,1}(x), \dots, a_{v,c_v}(x) \in \mathbb{R}^{c_v}$
  - Bob sends  $b_{v,1}(y), \dots, b_{v,c_v}(y) \in \mathbb{R}^{c_v}$
  - Referee announces results of comparisons  $a_{v,i}(x) \leq b_{v,i}(y)$  for  $i \in [c_v]$
- Function  $f$  solved by  $r$ -round real communication in cost  $c$  if  $\exists$  protocol such that
  - # rounds  $\leq r$
  - total # comparisons made by referee  $\leq c$
- Strictly stronger than standard deterministic communication  
(EQUALITY solved with real communication in 1 round with cost 2)

# Falsified Clause Search Problem

Falsified clause search problem  $\text{Search}(F)$

**Set-up:** Fixed (and unsatisfiable) CNF formula  $F$

**Input:** Assignment  $\alpha$  to  $\text{Vars}(F)$

**Output:** Clause  $C \in F$  such that  $\alpha$  falsifies  $C$

# Falsified Clause Search Problem

Falsified clause search problem  $Search(F)$

**Set-up:** Fixed (and unsatisfiable) CNF formula  $F$

**Input:** Assignment  $\alpha$  to  $Vars(F)$

**Output:** Clause  $C \in F$  such that  $\alpha$  falsifies  $C$

For any standard proof system, refutation  $\pi : F \vdash \perp$  (viewed as DAG) can be used to solve  $Search(F)$ :

- Start at sink (labelled by  $\perp$ )
- Walk backwards along nodes falsified by  $\alpha$
- Axiom clause  $C \in F$  labelling source node is valid answer

# Falsified Clause Search Problem (Communication Version)

Falsified clause search problem  $\text{Search}(F)$  for Alice and Bob

**Set-up:** Fixed (and unsatisfiable) CNF formula  $F$   
And (devious) partition of  $\text{Vars}(F)$  between Alice and Bob

**Input:** Assignment  $\alpha$  to  $\text{Vars}(F)$  split between Alice and Bob

**Task:** Alice and Bob should communicate to find clause  $C \in F$  such that  $\alpha$  falsifies  $C$



# Falsified Clause Search Problem (Communication Version)

Falsified clause search problem  $\text{Search}(F)$  for Alice and Bob

**Set-up:** Fixed (and unsatisfiable) CNF formula  $F$   
And (devious) partition of  $\text{Vars}(F)$  between Alice and Bob

**Input:** Assignment  $\alpha$  to  $\text{Vars}(F)$  split between Alice and Bob

**Task:** Alice and Bob should communicate to find clause  $C \in F$  such that  $\alpha$  falsifies  $C$

Actually, communication protocol should compute not function but **relation** — will mostly ignore this distinction


# Succinct Refutations Yield Efficient Protocols

Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



# Succinct Refutations Yield Efficient Protocols


Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard

# Succinct Refutations Yield Efficient Protocols


Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard

# Succinct Refutations Yield Efficient Protocols


Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard

# Succinct Refutations Yield Efficient Protocols


Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard

# Succinct Refutations Yield Efficient Protocols


Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard

# Succinct Refutations Yield Efficient Protocols

Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$




Use binary search to find transition from true to false blackboard

Must happen when  $C \in F$  written down — answer to  $Search(F)$



# Succinct Refutations Yield Efficient Protocols

Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$




Use binary search to find transition from true to false blackboard

Must happen when  $C \in F$  written down — answer to  $\text{Search}(F)$

Refutation length  $L \Rightarrow$  **evaluate  $\log L$  blackboards**

# Succinct Refutations Yield Efficient Protocols

Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard


Must happen when  $C \in F$  written down — answer to  $Search(F)$

Refutation length  $L \Rightarrow$  **evaluate  $\log L$  blackboards**

Refutation line space  $s \Rightarrow$  **max  $s$  bits of communication per blackboard**

# Succinct Refutations Yield Efficient Protocols

Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard

Must happen when  $C \in F$  written down — answer to  $\text{Search}(F)$


Refutation length  $L \Rightarrow$  **evaluate  $\log L$  blackboards**

Refutation line space  $s \Rightarrow$  **max  $s$  bits of communication per blackboard**

Only one round per blackboard evaluation

# Succinct Refutations Yield Efficient Protocols

Evaluate blackboard configurations of a refutation of  $F$  under  $\alpha$



Use binary search to find transition from true to false blackboard

Must happen when  $C \in F$  written down — answer to  $\text{Search}(F)$

Refutation length  $L \Rightarrow$  **evaluate  $\log L$  blackboards**

Refutation line space  $s \Rightarrow$  **max  $s$  bits of communication per blackboard**

Only one round per blackboard evaluation

(Alice and Bob simply evaluate their parts of each inequality and ask referee to compare)

# Lifting of Functions

Construct hard communication problems by “hardness amplification”  
using **lifting** or **composition**

# Lifting of Functions

Construct hard communication problems by “hardness amplification”  
using **lifting** or **composition**


Start with function  $f : \{0, 1\}^m \rightarrow \{0, 1\}$

$$f\left(\begin{array}{|c|c|c|} \hline & & \\ \hline \end{array}\right)$$

# Lifting of Functions



Construct hard communication problems by “hardness amplification”  
using **lifting** or **composition**

Start with function  $f : \{0, 1\}^m \rightarrow \{0, 1\}$



Construct new function on inputs

$x \in [\ell]^m$  and  $y \in \{0, 1\}^{\ell m}$



# Lifting of Functions


Construct hard communication problems by “hardness amplification”  
using **lifting** or **composition**

Start with function  $f : \{0, 1\}^m \rightarrow \{0, 1\}$

Construct new function on inputs

$x \in [\ell]^m$  and  $y \in \{0, 1\}^{\ell m}$

Alice's  $x$ -variables determine...



$$f\left(\begin{array}{|c|c|c|}\hline & & \\ \hline\end{array}\right)$$



# Lifting of Functions

Construct hard communication problems by “hardness amplification”  
using **lifting** or **composition**


Start with function  $f : \{0, 1\}^m \rightarrow \{0, 1\}$

Construct new function on inputs

$x \in [\ell]^m$  and  $y \in \{0, 1\}^{\ell m}$

Alice's  $x$ -variables determine...

... which of Bob's  $y$ -bits to feed to  $f$



# Lifting of Functions

Construct hard communication problems by “hardness amplification”  
using **lifting** or **composition**

Start with function  $f : \{0, 1\}^m \rightarrow \{0, 1\}$

Construct new function on inputs


$x \in [\ell]^m$  and  $y \in \{0, 1\}^{\ell m}$

Alice's  $x$ -variables determine...

... which of Bob's  $y$ -bits to feed to  $f$

Length- $\ell$  lifting of  $f$  defined as

$Lift_\ell(f)(x, y) := f(y_{1,x_1}, \dots, y_{m,x_m})$



# Lifting of Functions

Construct hard communication problems by “hardness amplification”  
using **lifting** or **composition**

Start with function  $f : \{0, 1\}^m \rightarrow \{0, 1\}$

Construct new function on inputs

$x \in [\ell]^m$  and  $y \in \{0, 1\}^{\ell m}$


Alice's  $x$ -variables determine...

... which of Bob's  $y$ -bits to feed to  $f$

Length- $\ell$  lifting of  $f$  defined as


$Lift_\ell(f)(x, y) := f(y_{1,x_1}, \dots, y_{m,x_m})$

Building on ideas from e.g. [She08, BHP10]



# Simulation of Protocols by Parallel Decision Trees [Val75]

Each node  $t$  in tree labelled by variables  $V_t$ ;  
has  $2^{|V_t|}$  outgoing edges




# Simulation of Protocols by Parallel Decision Trees [Val75]

Each node  $t$  in tree labelled by variables  $V_t$ ;

has  $2^{|V_t|}$  outgoing edges

Parallel decision tree:

- uses  $\# \text{ queries} = \max \sum |V_t|$  along any path




# Simulation of Protocols by Parallel Decision Trees [Val75]

Each node  $t$  in tree labelled by variables  $V_t$ ;

has  $2^{|V_t|}$  outgoing edges

Parallel decision tree:

- uses **# queries** =  $\max \sum |V_t|$  along any path
- has **depth** = length of longest path




# Simulation of Protocols by Parallel Decision Trees [Val75]

Each node  $t$  in tree labelled by variables  $V_t$ ;

has  $2^{|V_t|}$  outgoing edges

Parallel decision tree:

- uses **# queries** =  $\max \sum |V_t|$  along any path
- has **depth** = length of longest path
- **solves search problem**  $S \subseteq \{0, 1\}^m \times Q$  if  $\forall \alpha \in \{0, 1\}^m$  path defined by  $\alpha$  ends in leaf with  $q$  s.t.  $(\alpha, q) \in S$




# Simulation of Protocols by Parallel Decision Trees [Val75]

Each node  $t$  in tree labelled by variables  $V_t$ ;

has  $2^{|V_t|}$  outgoing edges

Parallel decision tree:

- uses **# queries** =  $\max \sum |V_t|$  along any path
- has **depth** = length of longest path
- **solves search problem**  $S \subseteq \{0, 1\}^m \times Q$  if  $\forall \alpha \in \{0, 1\}^m$  path defined by  $\alpha$  ends in leaf with  $q$  s.t.  $(\alpha, q) \in S$
- Can be simulated by Alice & Bob to solve lifted problem (easy)






# Simulation of Protocols by Parallel Decision Trees [Val75]

Each node  $t$  in tree labelled by variables  $V_t$ ;

has  $2^{|V_t|}$  outgoing edges

Parallel decision tree:

- uses **# queries** =  $\max \sum |V_t|$  along any path
- has **depth** = length of longest path
- **solves search problem**  $S \subseteq \{0, 1\}^m \times Q$  if  $\forall \alpha \in \{0, 1\}^m$  path defined by  $\alpha$  ends in leaf with  $q$  s.t.  $(\alpha, q) \in S$
- Can be simulated by Alice & Bob to solve lifted problem (easy)



## Simulation theorem of protocol by decision tree (hard direction)

Let  $S$  search problem with domain  $\{0, 1\}^m$  and let  $\ell = m^{3+\epsilon}$ ,  $\epsilon > 0$ . Then:

$\exists$   $r$ -round real communication protocol in cost  $c$  solving  $Lift_\ell(S)$

$\Rightarrow \exists$  depth- $r$  parallel decision tree solving  $S$  with  $\mathcal{O}(c/\log \ell)$  queries.

# Where to Get Formulas with Trade-off Properties?

Questions about time-space trade-offs fundamental in theoretical computer science


Well-studied (and well-understood) for **pebble games** modelling calculations described by DAGs

In particular, for **black-white pebble game** investigated by [CS76] and many others

# Pebbling Contradictions

CNF formulas encoding black-white pebble game played on DAG  $G$

1.  $u_1$
2.  $u_2$
3.  $u_3$
4.  $\bar{u}_1 \vee \bar{u}_2 \vee u_4$
5.  $\bar{u}_2 \vee \bar{u}_3 \vee u_5$
6.  $\bar{u}_4 \vee \bar{u}_5 \vee u_6$
7.  $\bar{u}_6$




- sources are true
- truth propagates upwards
- but sink is false

# Pebbling Contradictions

CNF formulas encoding black-white pebble game played on DAG  $G$

1.  $u_1$
2.  $u_2$
3.  $u_3$
4.  $\bar{u}_1 \vee \bar{u}_2 \vee u_4$
5.  $\bar{u}_2 \vee \bar{u}_3 \vee u_5$
6.  $\bar{u}_4 \vee \bar{u}_5 \vee u_6$
7.  $\bar{u}_6$




- sources are true
- truth propagates upwards
- but sink is false

# Pebbling Contradictions

CNF formulas encoding black-white pebble game played on DAG  $G$

1.  $u_1$
2.  $u_2$
3.  $u_3$
4.  $\bar{u}_1 \vee \bar{u}_2 \vee u_4$
5.  $\bar{u}_2 \vee \bar{u}_3 \vee u_5$
6.  $\bar{u}_4 \vee \bar{u}_5 \vee u_6$
7.  $\bar{u}_6$




- sources are true
- truth propagates upwards
- but sink is false

# Pebbling Contradictions

CNF formulas encoding black-white pebble game played on DAG  $G$

1.  $u_1$
2.  $u_2$
3.  $u_3$
4.  $\bar{u}_1 \vee \bar{u}_2 \vee u_4$
5.  $\bar{u}_2 \vee \bar{u}_3 \vee u_5$
6.  $\bar{u}_4 \vee \bar{u}_5 \vee u_6$
7.  $\bar{u}_6$




- sources are true
- truth propagates upwards
- but sink is false

# Pebbling Contradictions

CNF formulas encoding black-white pebble game played on DAG  $G$

1.  $u_1$
2.  $u_2$
3.  $u_3$
4.  $\bar{u}_1 \vee \bar{u}_2 \vee u_4$
5.  $\bar{u}_2 \vee \bar{u}_3 \vee u_5$
6.  $\bar{u}_4 \vee \bar{u}_5 \vee u_6$
7.  $\bar{u}_6$




- sources are true
- truth propagates upwards
- but sink is false

# Pebbling Contradictions

CNF formulas encoding black-white pebble game played on DAG  $G$

1.  $u_1$
2.  $u_2$
3.  $u_3$
4.  $\bar{u}_1 \vee \bar{u}_2 \vee u_4$
5.  $\bar{u}_2 \vee \bar{u}_3 \vee u_5$
6.  $\bar{u}_4 \vee \bar{u}_5 \vee u_6$
7.  $\bar{u}_6$




- sources are true
- truth propagates upwards
- but sink is false



# Pebbling Contradictions

CNF formulas encoding black-white pebble game played on DAG  $G$

1.  $u_1$
2.  $u_2$
3.  $u_3$
4.  $\bar{u}_1 \vee \bar{u}_2 \vee u_4$
5.  $\bar{u}_2 \vee \bar{u}_3 \vee u_5$
6.  $\bar{u}_4 \vee \bar{u}_5 \vee u_6$
7.  $\bar{u}_6$



- sources are true
- truth propagates upwards
- but sink is false

Appeared in various contexts in e.g. [RM99, BEGJ00, BW01]

Used in [Nor06, NH08, BN08, BN11, BNT13] to study space and size-space trade-offs in resolution and polynomial calculus

Formulas inherit some DAG properties, but not enough — make them harder by lifting!

# Lifted CNF Formulas

Given

- CNF formula  $F$  over variables  $u_1, \dots, u_n$
- lift length  $\ell \in \mathbb{N}^+$

the **lifted formula**  $Lift_\ell(F)$  has

- **selector variables**  $\{x_{i,j}\}_{i \in [n], j \in [\ell]}$
- **main variables**  $\{y_{i,j}\}_{i \in [n], j \in [\ell]}$
- for every  $i \in [n]$  an **auxiliary clause**

$$x_{i,1} \vee x_{i,2} \vee \dots \vee x_{i,\ell}$$

- for every  $C = u_{i_1} \vee \dots \vee u_{i_s} \vee \bar{u}_{i_{s+1}} \vee \dots \vee \bar{u}_{i_t}$  in  $F$  and  $(j_1, \dots, j_t) \in [\ell]^t$  a **main clause**

$$\bar{x}_{i_1,j_1} \vee y_{i_1,j_1} \vee \dots \vee \bar{x}_{i_s,j_s} \vee y_{i_s,j_s} \vee \bar{x}_{i_{s+1},j_{s+1}} \vee \bar{y}_{i_{s+1},j_{s+1}} \vee \dots \vee \bar{x}_{i_t,j_t} \vee \bar{y}_{i_t,j_t}$$

# Toy Example Lifted Pebbling Contradiction (Lift Length 2)

$$\begin{aligned}
 & (x_{1,1} \vee x_{1,2}) \\
 & \wedge (x_{2,1} \vee x_{2,2}) \\
 & \wedge (x_{3,1} \vee x_{3,2}) \\
 & \wedge (x_{4,1} \vee x_{4,2}) \\
 & \wedge (x_{5,1} \vee x_{5,2}) \\
 & \wedge (x_{6,1} \vee x_{6,2}) \\
 & \wedge (\overline{x}_{1,1} \vee y_{1,1}) \\
 & \wedge (\overline{x}_{1,2} \vee y_{1,2}) \\
 & \wedge (\overline{x}_{2,1} \vee y_{2,1}) \\
 & \wedge (\overline{x}_{2,2} \vee y_{2,2}) \\
 & \wedge (\overline{x}_{3,1} \vee y_{3,1}) \\
 & \wedge (\overline{x}_{3,2} \vee y_{3,2}) \\
 & \wedge (\overline{x}_{1,1} \vee \overline{y}_{1,1} \vee \overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{4,1} \vee y_{4,1}) \\
 & \wedge (\overline{x}_{1,1} \vee \overline{y}_{1,1} \vee \overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{4,2} \vee y_{4,2}) \\
 & \wedge (\overline{x}_{1,1} \vee \overline{y}_{1,1} \vee \overline{x}_{2,2} \vee \overline{y}_{2,2} \vee \overline{x}_{4,1} \vee y_{4,1}) \\
 & \wedge (\overline{x}_{1,1} \vee \overline{y}_{1,1} \vee \overline{x}_{2,2} \vee \overline{y}_{2,2} \vee \overline{x}_{4,2} \vee y_{4,2}) \\
 & \wedge (\overline{x}_{1,2} \vee \overline{y}_{1,2} \vee \overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{4,1} \vee y_{4,1}) \\
 & \wedge (\overline{x}_{1,2} \vee \overline{y}_{1,2} \vee \overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{4,2} \vee y_{4,2}) \\
 & \wedge (\overline{x}_{1,2} \vee \overline{y}_{1,2} \vee \overline{x}_{2,2} \vee \overline{y}_{2,2} \vee \overline{x}_{4,1} \vee y_{4,1}) \\
 & \wedge (\overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{3,1} \vee \overline{y}_{3,1} \vee \overline{x}_{5,1} \vee y_{5,1}) \\
 & \wedge (\overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{3,1} \vee \overline{y}_{3,1} \vee \overline{x}_{5,2} \vee y_{5,2}) \\
 & \wedge (\overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{3,2} \vee \overline{y}_{3,2} \vee \overline{x}_{5,1} \vee y_{5,1}) \\
 & \wedge (\overline{x}_{2,1} \vee \overline{y}_{2,1} \vee \overline{x}_{3,2} \vee \overline{y}_{3,2} \vee \overline{x}_{5,2} \vee y_{5,2}) \\
 & \wedge (\overline{x}_{2,2} \vee \overline{y}_{2,2} \vee \overline{x}_{3,1} \vee \overline{y}_{3,1} \vee \overline{x}_{5,1} \vee y_{5,1}) \\
 & \wedge (\overline{x}_{2,2} \vee \overline{y}_{2,2} \vee \overline{x}_{3,1} \vee \overline{y}_{3,1} \vee \overline{x}_{5,2} \vee y_{5,2}) \\
 & \wedge (\overline{x}_{2,2} \vee \overline{y}_{2,2} \vee \overline{x}_{3,2} \vee \overline{y}_{3,2} \vee \overline{x}_{5,1} \vee y_{5,1}) \\
 & \wedge (\overline{x}_{2,2} \vee \overline{y}_{2,2} \vee \overline{x}_{3,2} \vee \overline{y}_{3,2} \vee \overline{x}_{5,2} \vee y_{5,2}) \\
 & \wedge (\overline{x}_{4,1} \vee \overline{y}_{4,1} \vee \overline{x}_{5,1} \vee \overline{y}_{5,1} \vee \overline{x}_{6,1} \vee y_{6,1}) \\
 & \wedge (\overline{x}_{4,1} \vee \overline{y}_{4,1} \vee \overline{x}_{5,1} \vee \overline{y}_{5,1} \vee \overline{x}_{6,2} \vee y_{6,2}) \\
 & \wedge (\overline{x}_{4,1} \vee \overline{y}_{4,1} \vee \overline{x}_{5,2} \vee \overline{y}_{5,2} \vee \overline{x}_{6,1} \vee y_{6,1}) \\
 & \wedge (\overline{x}_{4,1} \vee \overline{y}_{4,1} \vee \overline{x}_{5,2} \vee \overline{y}_{5,2} \vee \overline{x}_{6,2} \vee y_{6,2}) \\
 & \wedge (\overline{x}_{4,2} \vee \overline{y}_{4,2} \vee \overline{x}_{5,1} \vee \overline{y}_{5,1} \vee \overline{x}_{6,1} \vee y_{6,1}) \\
 & \wedge (\overline{x}_{4,2} \vee \overline{y}_{4,2} \vee \overline{x}_{5,1} \vee \overline{y}_{5,1} \vee \overline{x}_{6,2} \vee y_{6,2}) \\
 & \wedge (\overline{x}_{4,2} \vee \overline{y}_{4,2} \vee \overline{x}_{5,2} \vee \overline{y}_{5,2} \vee \overline{x}_{6,1} \vee y_{6,1}) \\
 & \wedge (\overline{x}_{4,2} \vee \overline{y}_{4,2} \vee \overline{x}_{5,2} \vee \overline{y}_{5,2} \vee \overline{x}_{6,2} \vee y_{6,2}) \\
 & \wedge (\overline{x}_{6,1} \vee \overline{y}_{6,1}) \\
 & \wedge (\overline{x}_{6,2} \vee \overline{y}_{6,2})
 \end{aligned}$$

# Lifted Pebbling Contradictions and the Simulation Theorem

Plug in the simulation theorem:

- From  $r$ -round real communication protocol in cost  $c$  solving  $\text{Search}(\text{Lift}_\ell(\text{Peb}_G))$
- Get depth- $r$  parallel decision tree solving  $\text{Search}(\text{Peb}_G)$  with  $\mathcal{O}(c/\log \ell)$  queries

# Lifted Pebbling Contradictions and the Simulation Theorem

Plug in the simulation theorem:

- From  $r$ -round real communication protocol in cost  $c$  solving  $\text{Search}(\text{Lift}_\ell(\text{Peb}_G))$
- Get depth- $r$  parallel decision tree solving  $\text{Search}(\text{Peb}_G)$  with  $\mathcal{O}(c/\log \ell)$  queries

So now we need to understand decision trees for pebbling formulas  $\text{Peb}_G$ !

# Lifted Pebbling Contradictions and the Simulation Theorem

Plug in the simulation theorem:


- From  $r$ -round real communication protocol in cost  $c$  solving  $\text{Search}(\text{Lift}_\ell(\text{Peb}_G))$
- Get depth- $r$  parallel decision tree solving  $\text{Search}(\text{Peb}_G)$  with  $\mathcal{O}(c/\log \ell)$  queries

So now we need to understand decision trees for pebbling formulas  $\text{Peb}_G$ !

Study pebble game on graph  $G$ , but other game than black-white pebbling


# From Parallel Decision Trees to Dymond–Tompa Games

- From [DT85]; more recently studied in [Cha13, CLNV15]



# From Parallel Decision Trees to Dymond–Tompa Games


- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**






# From Parallel Decision Trees to Dymond–Tompa Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler** places **pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger** either **jumps** to **newly pebbled** vertex (always in 1st round) **or stays**




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler** places **pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger** either **jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tomba Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler** places **pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger** either **jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)



# From Parallel Decision Trees to Dymond–Tompkins Games


- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)






# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)




# From Parallel Decision Trees to Dymond–Tompkins Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler places pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger either jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)



# From Parallel Decision Trees to Dymond–Tomba Games

- From [DT85]; more recently studied in [Cha13, CLNV15]
- Two players **Pebbler** and **Challenger**
- In each round
  - **Pebbler** places **pebbles** on subset of vertices (including sink in 1st round)
  - **Challenger** either **jumps** to **newly pebbled** vertex (always in 1st round) **or stays**
- Pebbler wins at end of round when **Challenger on vertex with all predecessors pebbled** (or on source vertex)



## Lemma

$\exists$  depth- $r$  parallel decision tree for  $\text{Search}(Peb_G)$  with  $\leq c$  queries  
 $\Rightarrow$  Pebbler wins  $r$ -round Dymond–Tomba game on  $G$  in cost  $\leq c + 1$

# Putting the Pieces Together

Prove round-cost trade-offs for Dymond–Tompkins games on graphs  $G$   
(hacking graph constructions from [CS82, LT82, Nor12])

# Putting the Pieces Together

Prove round-cost trade-offs for Dymond–Tompkins games on graphs  $G$   
(hacking graph constructions from [CS82, LT82, Nor12])



Depth-query trade-offs for parallel decision trees for  $\text{Search}(Peb_G)$

# Putting the Pieces Together

Prove round-cost trade-offs for Dymond–Tompkins games on graphs  $G$   
(hacking graph constructions from [CS82, LT82, Nor12])



Depth-query trade-offs for parallel decision trees for  $\text{Search}(Peb_G)$



Real communication round-cost trade-offs for  $\text{Lift}(\text{Search}(Peb_G))$



# Putting the Pieces Together

Prove round-cost trade-offs for Dymond–Tompkins games on graphs  $G$   
(hacking graph constructions from [CS82, LT82, Nor12])



Depth-query trade-offs for parallel decision trees for  $\text{Search}(Peb_G)$



Real communication round-cost trade-offs for  $\text{Lift}(\text{Search}(Peb_G))$



Same communication lower bounds hold for  $\text{Search}(\text{Lift}(Peb_G))$ ,  
i.e., search problem for lifted formulas in [BHP10]

# Putting the Pieces Together

Prove round-cost trade-offs for Dymond–Tompkins games on graphs  $G$   
(hacking graph constructions from [CS82, LT82, Nor12])



Depth-query trade-offs for parallel decision trees for  $\text{Search}(Peb_G)$



Real communication round-cost trade-offs for  $\text{Lift}(\text{Search}(Peb_G))$



Same communication lower bounds hold for  $\text{Search}(\text{Lift}(Peb_G))$ ,  
i.e., search problem for lifted formulas in [BHP10]



Cutting planes length-space trade-off for  $\text{Lift}(Peb_G)$

## Some Interesting Questions

### **Communication complexity**

- Smaller length of lift?
- Simulation theorems for stronger communication models (in particular, for so-called DAG-like communication)

# Some Interesting Questions

## Communication complexity

- Smaller length of lift?
- Simulation theorems for stronger communication models (in particular, for so-called DAG-like communication)

## Proof complexity

- Better Dymond–Tompia trade-offs?
- Reduction to black-white pebbling instead of Dymond–Tompia?
- Supercritical size-space trade-offs for Tseitin formulas à la [BNT13, BBI16]?
- Line space lower bounds for CP with bounded coefficients (strengthening [GPT15])

# Some Interesting Questions

## Communication complexity

- Smaller length of lift?
- Simulation theorems for stronger communication models (in particular, for so-called DAG-like communication)

## Proof complexity

- Better Dymond–Tompia trade-offs?
- Reduction to black-white pebbling instead of Dymond–Tompia?
- Supercritical size-space trade-offs for Tseitin formulas à la [BNT13, BBI16]?
- Line space lower bounds for CP with bounded coefficients (strengthening [GPT15])

## And more to come in future lectures...

- But now it is time to switch to the board and do some proper proofs!

# References I

- [BBI16] Paul Beame, Chris Beck, and Russell Impagliazzo. Time-space tradeoffs in resolution: Superpolynomial lower bounds for superlinear space. *SIAM Journal on Computing*, 45(4):1612–1645, August 2016. Preliminary version in *STOC '12*.
- [BEGJ00] María Luisa Bonet, Juan Luis Esteban, Nicola Galesi, and Jan Johannsen. On the relative complexity of resolution refinements and cutting planes proof systems. *SIAM Journal on Computing*, 30(5):1462–1484, 2000. Preliminary version in *FOCS '98*.
- [BHP10] Paul Beame, Trinh Huynh, and Toniann Pitassi. Hardness amplification in proof complexity. In *Proceedings of the 42nd Annual ACM Symposium on Theory of Computing (STOC '10)*, pages 87–96, June 2010.
- [BN08] Eli Ben-Sasson and Jakob Nordström. Short proofs may be spacious: An optimal separation of space and length in resolution. In *Proceedings of the 49th Annual IEEE Symposium on Foundations of Computer Science (FOCS '08)*, pages 709–718, October 2008.
- [BN11] Eli Ben-Sasson and Jakob Nordström. Understanding space in proof complexity: Separations and trade-offs via substitutions. In *Proceedings of the 2nd Symposium on Innovations in Computer Science (ICS '11)*, pages 401–416, January 2011.

# References II

- [BNT13] Chris Beck, Jakob Nordström, and Bangsheng Tang. Some trade-off results for polynomial calculus. In *Proceedings of the 45th Annual ACM Symposium on Theory of Computing (STOC '13)*, pages 813–822, May 2013.
- [BW01] Eli Ben-Sasson and Avi Wigderson. Short proofs are narrow—resolution made simple. *Journal of the ACM*, 48(2):149–169, March 2001. Preliminary version in *STOC '99*.
- [Cha13] Siu Man Chan. Just a pebble game. In *Proceedings of the 28th Annual IEEE Conference on Computational Complexity (CCC '13)*, pages 133–143, June 2013.
- [CLNV15] Siu Man Chan, Massimo Lauria, Jakob Nordström, and Marc Vinyals. Hardness of approximation in PSPACE and separation results for pebble games (Extended abstract). In *Proceedings of the 56th Annual IEEE Symposium on Foundations of Computer Science (FOCS '15)*, pages 466–485, October 2015.
- [CS76] Stephen A. Cook and Ravi Sethi. Storage requirements for deterministic polynomial time recognizable languages. *Journal of Computer and System Sciences*, 13(1):25–37, 1976. Preliminary version in *STOC '74*.
- [CS82] David A. Carlson and John E. Savage. Extreme time-space tradeoffs for graphs with small space requirements. *Information Processing Letters*, 14(5):223–227, 1982.

# References III

- [DT85] Patrick W. Dymond and Martin Tompa. Speedups of deterministic machines by synchronous parallel machines. *Journal of Computer and System Sciences*, 30(2):149–161, April 1985. Preliminary version in *STOC '83*.
- [GP18] Mika Göös and Toniann Pitassi. Communication lower bounds via critical block sensitivity. *SIAM Journal on Computing*, 47(5):1778–1806, October 2018. Preliminary version in *STOC '14*.
- [GPT15] Nicola Galesi, Pavel Pudlák, and Neil Thapen. The space complexity of cutting planes refutations. In *Proceedings of the 30th Annual Computational Complexity Conference (CCC '15)*, volume 33 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 433–447, June 2015.
- [GPW15] Mika Göös, Toniann Pitassi, and Thomas Watson. Deterministic communication vs. partition number. In *Proceedings of the 56th Annual IEEE Symposium on Foundations of Computer Science (FOCS '15)*, pages 1077–1088, October 2015.
- [HN12] Trinh Huynh and Jakob Nordström. On the virtue of succinct proofs: Amplifying communication complexity hardness to time-space trade-offs in proof complexity (Extended abstract). In *Proceedings of the 44th Annual ACM Symposium on Theory of Computing (STOC '12)*, pages 233–248, May 2012.
- [Kra98] Jan Krajíček. Interpolation by a game. *Mathematical Logic Quarterly*, 44(4):450–458, 1998.



## References IV

- [LT82] Thomas Lengauer and Robert Endre Tarjan. Asymptotically tight bounds on time-space trade-offs in a pebble game. *Journal of the ACM*, 29(4):1087–1130, October 1982. Preliminary version in *STOC '79*.
- [NH08] Jakob Nordström and Johan Håstad. Towards an optimal separation of space and length in resolution (Extended abstract). In *Proceedings of the 40th Annual ACM Symposium on Theory of Computing (STOC '08)*, pages 701–710, May 2008.
- [Nor06] Jakob Nordström. Narrow proofs may be spacious: Separating space and width in resolution (Extended abstract). In *Proceedings of the 38th Annual ACM Symposium on Theory of Computing (STOC '06)*, pages 507–516, May 2006.
- [Nor12] Jakob Nordström. On the relative strength of pebbling and resolution. *ACM Transactions on Computational Logic*, 13(2):16:1–16:43, April 2012. Preliminary version in *CCC '10*.
- [Pud97] Pavel Pudlák. Lower bounds for resolution and cutting plane proofs and monotone computations. *Journal of Symbolic Logic*, 62(3):981–998, September 1997.
- [RM99] Ran Raz and Pierre McKenzie. Separation of the monotone NC hierarchy. *Combinatorica*, 19(3):403–435, March 1999. Preliminary version in *FOCS '97*.

# References V

- [She08] Alexander A. Sherstov. The pattern matrix method for lower bounds on quantum communication. In *Proceedings of the 40th Annual ACM Symposium on Theory of Computing (STOC '08)*, pages 85–94, May 2008.
- [Sok24] Dmitry Sokolov. Random  $(\log n)$ -CNF are hard for cutting planes (again). In *Proceedings of the 56th Annual ACM Symposium on Theory of Computing (STOC '24)*, pages 2008–2015, June 2024.
- [Val75] Leslie G. Valiant. Parallelism in comparison problems. *SIAM Journal on Computing*, 4(3):348–355, March 1975.