Checklist for Follow-Up on Sterilization of Towels and Replacement with Disposable Sets for Cardiac Surgery

N	Step	Achieved	Not
0			Achieved
1.	Pre-cleaning and inspection		
	✓ Thoroughly cleaned to remove dirt, debris, or organic matter		
	 ✓ Towels should be visually inspected to ensure they are free of any tears, holes, or fraying or foreign materials ✓ Identify stains or discoloration ✓ Isolate any towels with defects, mark defective items with a clear label ("Unfit for Use") ✓ Record the quantity, type of defect, and batch details 		X X X X
2.	Packaging ✓ Properly folding and wrapping towels ensures uniform sterilization ✓ Items should be packed to allow adequate steam penetration and prevent air pockets	Needs more specific practice	
3.	Sterilization methods ✓ Steam sterilization requires autoclaving at 121°C (250°F) for minimum 30 minutes	Not sure	

ı

4.	 ✓ Or Prevcum Autoclave: 132°C for at least 4 mins ✓ Ensuring that sterilization cycles include a drying phase to prevent residual moisture Quality control ✓ Each sterilized pack should include indicators (Check sterilization tape indicators for color changes), to confirm that the sterilization parameters were met. 	√	X
5.	Storage ✓ Once sterilized, towels should be stored in a clean, dry, designated sterile area ✓ Limiting handling to prevent contamination and ensuring that packages remain intact until they are needed for surgical use		X
6.	Replacement with disposable sets ✓ Use high-quality, single-use disposable sets that meet surgical standards (ISO 11607) ✓ Verify expiration dates and packaging integrity before use ✓ Store Properly: - Maintain disposable sets in a clean, dry, and sterile environment - Rotate stock to avoid expired items		