

## \* 3. Matrices \*

### \* Types of matrix:

- (1) Row
  - (11) complex
  - (2) column
  - (12) symmetric
  - (3) square
  - (13) skew symmetric
  - (4) Diagonal
  - (14) orthogonal [ $A^T = A^{-1}$ ]  
OR  $AA^{-1} = A^T A = I$
  - (5) scalar
  - (15) conjugate matrix
  - (6) Identity
  - (7) Null
  - (8) Upper diagonal
  - (9) Lower diagonal
  - (10) Real
  - (16) Hermitian matrix
  - (17) Skew Hermitian
  - (18) unitary matrix
  - (19) idempotent matrix
  - (20) involutory matrix
  - (21) nilpotent matrix
- $A = \begin{bmatrix} 1+i & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$   $\bar{A} = A^C = \begin{bmatrix} 1-i & 2 & 3 \\ 0 & 1 & 2 \\ -2i & 0 & 0 \end{bmatrix}$

### \* Definitions :-

- (1) complex number: The matrix  $A = [a_{ij}]_{m \times n}$  is called complex matrix if atleast one entry of complex number.
- (2) conjugate matrix: The matrix obtained from any given matrix  $A = [a_{ij}]_{m \times n}$  on replacing it's elements by the corresponding conjugate complex number is called the conjugate matrix and it is denoted by  $\bar{A}$  or  $A^C$ .

$$\text{eg: } A = \begin{bmatrix} -i & 3i & 7+4i \\ 2 & 4i & 10 \end{bmatrix} \quad A^C = \begin{bmatrix} i & -3i & 7-4i \\ 2 & -4i & 10 \end{bmatrix}$$

(3) Hermitian matrix : A matrix  $A = [a_{ij}]_{m \times n}$  is called Hermitian matrix if

$$(A^c)^T = (A^T)^c = A = A^0 + A^H$$

(4) skew Hermitian matrix : A matrix  $A = [a_{ij}]_{m \times n}$  is called skew Hermitian matrix if

$$(A^c)^T = (A^T)^c = -A = A^0 + A^H$$

(5) unitary matrix : A matrix  $A = [a_{ij}]_{m \times n}$  is called unitary matrix if

$$(A^c)^T = (A^T)^c = A^0 = A^H = A^{-1}$$

$$A^H A = A A^H = I$$

(6) Idempotent matrix : if a matrix  $A = [a_{ij}]_{m \times n}$  is a square matrix such that  $A^2 = A$  then the given matrix is called idempotent matrix.

(7) Involutory matrix : if a matrix  $A = [a_{ij}]_{m \times n}$  is a square matrix such that  $A^2 = I$  then the given matrix is called involutory matrix.

(8) nilpotent matrix : if a matrix  $A = [a_{ij}]_{m \times n}$  is a square matrix such that  $A^m = 0$ , where  $m$  is positive integer, then the given matrix is called nilpotent matrix.

## \* classwork Examples

1. Find the determinant of  $\begin{pmatrix} 1 & -2 \\ 3 & 0 \end{pmatrix} = 0 + 6 = 6$

2. Find the determinant of  $\begin{pmatrix} 1 & -2 & 3 \\ 1 & -1 & 2 \\ 2 & 1 & 3 \end{pmatrix}$   
 $= 1(-3-2) + 2(3-4) + 3(1+2)$   
 $= -5 - 2 + 9$   
 $= 2$

3. Find the determinant of  $\begin{pmatrix} 2 & 0 & 0 & 3 \\ 4 & -3 & 1 & 2 \\ 3 & 1 & 2 & 1 \\ 0 & -4 & 0 & 7 \end{pmatrix}$   
 $= 2 \begin{pmatrix} -3 & 1 & 2 \\ 1 & -2 & 1 \\ 0 & 0 & 7 \end{pmatrix} - 3 \begin{pmatrix} 4 & -3 & 1 \\ 3 & 1 & 2 \\ 0 & -4 & 0 \end{pmatrix}$   
 $= 2 [(-3)(14) - 1(7+4) + 2(8)] - 3 [(4)(8) + 3(0) + 1(-12)]$   
 $= 2 (-42 - 11 + 16) - 3 (32 - 12)$   
 $= -74 - 60$   
 $= -134$

4. Prove that the  $\det \begin{pmatrix} a^2 & a & bc \\ b^2 & b & ca \\ c^2 & c & ab \end{pmatrix} = -\det \begin{pmatrix} 1 & 1 & 1 \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{pmatrix}$

LHS =  $\det \begin{pmatrix} a^2 & a & bc \\ b^2 & b & ca \\ c^2 & c & ab \end{pmatrix}$

$$= \frac{\det 1}{abc} \begin{pmatrix} a^3 & a^2 & abc \\ b^3 & b^2 & abc \\ c^3 & c^2 & abc \end{pmatrix} \quad (\because R_1 \rightarrow aR_1, R_2 \rightarrow bR_2, R_3 \rightarrow cR_3)$$

$$= \frac{\det 1}{abc} \begin{pmatrix} a^3 & a^2 & 1 \\ b^3 & b^2 & 1 \\ c^3 & c^2 & 1 \end{pmatrix} \quad (\because P_3 \rightarrow 1/c_3)$$

$$= \det \begin{pmatrix} a^3 & b^3 & c^3 \\ a^2 & b^2 & c^2 \\ 1 & 1 & 1 \end{pmatrix} \quad (\because (A \Rightarrow A^{-1}))$$

$$= -\det \begin{pmatrix} 1 & 1 & 1 \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{pmatrix} \quad (R_1 \Rightarrow R_3, R_3 \Rightarrow R_1)$$

- RHS

5. Find the determinant of

$$\begin{pmatrix} 2 & 0 & -1 & 3 \\ 0 & -3 & 0 & 2 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$= 2 \begin{pmatrix} -3 & 0 & 2 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{pmatrix} + 1 \begin{pmatrix} 0 & -3 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} - 3 \begin{pmatrix} 0 & -3 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

$$= 2(0+0+0) + 1(0) - 3(0)$$

$$= 0 + 0 + 0$$

$$= 0$$

6. Find the determinant of

$$\begin{pmatrix} 9 & 9 & 12 \\ 1 & 3 & -4 \\ 1 & -9 & 12 \end{pmatrix}$$

$$= 9(36 + 36) - 9(12 + 4) + 12(9 - 3)$$

$$= 648 - 144 + 72$$

$$= 576$$

\* Practice examples

(1) Find the determinant of  $\begin{pmatrix} 0 & 2 & 7 \\ -1 & 5 & 0 \\ 8 & -3 & 2 \end{pmatrix}$

$$= -2(-2 - 0) + 7(3 - 40)$$

$$= -4 - 259$$

$$= -255$$

(2) Find the determinant of  $\begin{pmatrix} \sin\theta & -\cos\theta \\ \cos\theta & \sin\theta \end{pmatrix}$

$$= -\sin^2\theta + \cos^2\theta$$

$$= \cos^2\theta - \sin^2\theta$$

$$= \cos 2\theta$$

(3) Find the determinant of  $\begin{pmatrix} 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 7 \\ -2 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \end{pmatrix}$

$$= 5 \begin{pmatrix} 0 & 0 & 7 \\ -2 & 0 & 0 \\ 0 & 3 & 0 \end{pmatrix}$$

$$= 5[7(-6)]$$

$$= 5(-42)$$

$$= -210$$

(4) Show that the  $\det \begin{pmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{pmatrix} = (a-b)(b-c)(c-a)$ .

$$= \begin{vmatrix} 0 & 1 & 0 \\ a-b & b & c-b \\ a^2-b^2 & b^2 & c^2-b^2 \end{vmatrix} \quad (\because C_1 \rightarrow C_1 - C_2) \\ \quad (\because C_3 \rightarrow C_3 - C_2)$$

$$= \begin{vmatrix} 0 & 1 & 0 \\ a-b & b & c-b \\ (a-b)(c+b) & b^2 & (c-b)(c+b) \end{vmatrix}$$

$$= (a-b)(c-b) \begin{vmatrix} 0 & 1 & 0 \\ 1 & b & 1 \\ a+b & b^2 & c+b \end{vmatrix} \quad (\because C_1 \rightarrow \frac{1}{a-b} C_1)$$

$$\qquad \qquad \qquad C_3 \rightarrow \frac{1}{c-b} C_3$$

$$= (a-b)(c-b) [-1(c+b-a-b)]$$

$$= (a-b)(c-b)(a-c)$$

$$= (a-b)(b-c)(c-a)$$

= RHS

$$(5) \text{ prove that the det } \begin{pmatrix} -a^2 & ab & ac \\ ba & -b^2 & bc \\ ac & bc & -c^2 \end{pmatrix} = 4a^2b^2c^2.$$

$$= abc \begin{vmatrix} -a & b & c \\ a & -b & c \\ a & b & -c \end{vmatrix} \quad (\because R_1 \rightarrow 1/a R_1)$$

$$\qquad \qquad \qquad R_2 \rightarrow 1/b R_2$$

$$\qquad \qquad \qquad R_3 \rightarrow 1/c R_3)$$

$$= a^2b^2c^2 \begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} \quad (\because C_1 \rightarrow 1/a C_1)$$

$$\qquad \qquad \qquad C_2 \rightarrow 1/b C_2$$

$$\qquad \qquad \qquad C_3 \rightarrow 1/c C_3)$$

$$= a^2b^2c^2 [-1(1-1) - 1(-1-1) + 1(1+1)]$$

$$= a^2b^2c^2 (2+2)$$

$$= 4a^2b^2c^2$$

= RHS

(6) Show that the  $\det \begin{pmatrix} 3a & b-a & c-a \\ a-b & 3b & c-b \\ a-c & b-c & 3c \end{pmatrix} = 3(a+b+c)(ab+bc+ca)$ .

$$\text{LHS} = \begin{vmatrix} 3a & b-a & c-a \\ a-b & 3b & c-b \\ a-c & b-c & 3c \end{vmatrix}$$

$$= \begin{vmatrix} a+b+c & b-a & c-a \\ a+b+c & 3b & c-b \\ a+b+c & b-c & 3c \end{vmatrix} \quad (\because C_1 \rightarrow C_1 + C_2 + C_3)$$

$$= (a+b+c) \begin{vmatrix} 1 & b-a & c-a \\ 1 & 3b & c-b \\ 1 & b-c & 3c \end{vmatrix} \quad (\because C_1 \rightarrow 1 \cdot C_1)$$

$$= (a+b+c) \begin{vmatrix} 0 & c-a & -2c-a \\ 0 & 2b+c & -2c-b \\ 1 & b-c & -3c \end{vmatrix}$$

$$= (a+b+c) [ 1((c-a)(-2c-b)) + (2c+a)(2b+c) ]$$

$$= (a+b+c) (-2c^2 - bc + 2ac + ab + 4bc + 2c^2 + 2ab + ac)$$

$$= (a+b+c) (3bc + 3ac + 3ab)$$

$$= 3(a+b+c)(ab+bc+ca)$$

$$= \text{RHS}$$

### \* Rank of a matrix:

→ A number 'r' is said to be the rank of a matrix 'A' if

- (i) there is at least one non-zero minor of order r.
- (ii) all the minors of order greater than r are zero.

Rank of a matrix is denoted by

$$\text{r}(A) \text{ or } \text{R}(A)$$

### \* Property :

(1) A matrix A is a null matrix or zero matrix if and only if rank of A = 0.

(2) Let A be an  $m \times n$  matrix then rank of A  $\leq \min\{m, n\}$

$$(3) \text{r}(A) = \text{r}(A^T)$$

### \* Classwork Examples :

(1) find the rank of  $\begin{pmatrix} 2 & 3 & 4 \\ 3 & 5 & 7 \end{pmatrix}$  using minors.

Given matrix is of order  $2 \times 3$

From the property,

$$s(A) \leq \min \{2, 3\}$$

$$s(A) \leq 2$$

all possible submatrix of order 2

$$\begin{bmatrix} 2 & 3 \\ 3 & 5 \end{bmatrix}, \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}, \begin{bmatrix} 2 & 4 \\ 3 & 7 \end{bmatrix}$$

det of  $\begin{bmatrix} 2 & 3 \\ 3 & 5 \end{bmatrix}$  is non zero

$\therefore$  rank of the given matrix is 2.

$$s(A) = 2$$

2. Find the rank of  $\begin{pmatrix} 1 & 3 & -4 \\ -1 & -3 & 4 \\ 2 & 6 & -8 \end{pmatrix}$  using minors.

given matrix is order of  $3 \times 3$

from the property

$$s(A) \leq \{\min \{3 \times 3\}\}$$

$$s(A) \leq 3$$

all possible submatrix of order 3

$$\text{here } \det A = 0$$

$\therefore s(A)$  is not 3.

now, all possible submatrix of order 2.

$$\begin{bmatrix} -3 & 4 \\ 6 & -8 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ 2 & -8 \end{bmatrix}, \begin{bmatrix} -1 & -3 \\ 2 & 6 \end{bmatrix}, \begin{bmatrix} 3 & -4 \\ 6 & -8 \end{bmatrix}, \begin{bmatrix} 1 & -4 \\ 2 & -8 \end{bmatrix},$$

$$\begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}, \begin{bmatrix} 3 & -4 \\ -3 & 4 \end{bmatrix}, \begin{bmatrix} 1 & -4 \\ -1 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 3 \\ -1 & -3 \end{bmatrix}$$

Here all submatrix of order 2 has  $\det 0$

$$\therefore s(A) \neq 2.$$

Now, all possible submatrix of order 1.

$$[1], [3], [-4], [-1], [-3], [4], [2], [6], [-8]$$

Det of above submatrix are non-zero sign of the given matrix is 1.

$$S(A) = 1$$

4. Find the rank of  $\begin{pmatrix} 2 & 1 & 5 & -1 \\ -1 & 2 & 5 & 3 \\ 3 & 2 & 9 & -1 \end{pmatrix}$  using minors

Given matrix is of order  $3 \times 4$ .

from the property.

$$S(A) \leq \min\{3, 4\}$$

$$S(A) \leq 3$$

now all the possible submatrix of order 3,

$$\begin{bmatrix} 2 & 1 & 5 \\ -1 & 2 & 5 \\ 3 & 2 & 9 \end{bmatrix}, \begin{bmatrix} 2 & 1 & -1 \\ -1 & 2 & 3 \\ 3 & 2 & -1 \end{bmatrix}, \begin{bmatrix} 2 & 5 & -1 \\ -1 & 5 & 3 \\ 3 & 9 & -1 \end{bmatrix}, \begin{bmatrix} 1 & 5 & -1 \\ 2 & 5 & 3 \\ 2 & 9 & -1 \end{bmatrix}$$

Here all submatrix of order 3 has det 0.

$$\therefore S(A) \neq 3.$$

now all possible submatrix of order 2.

$$\begin{bmatrix} 2 & 5 \\ 2 & 9 \end{bmatrix}, \begin{bmatrix} 5 & 3 \\ 9 & -7 \end{bmatrix}, \begin{bmatrix} 2 & 3 \\ 2 & -1 \end{bmatrix}, \text{ and so on.}$$

Det of above submatrix are non-zero. Rank

of the given matrix is 2.

$$S(A) = 2$$

Find the rank of  $\begin{pmatrix} 2 & 0 & 0 & 3 \\ 0 & -3 & 0 & 2 \\ 0 & 2 & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  using minors.

$$\begin{pmatrix} 2 & 0 & 0 & 3 \\ 0 & -3 & 0 & 2 \\ 0 & 2 & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Given matrix is of order  $4 \times 4$ .

From the property,

$$S(A) \leq \min \{4, 4\}$$

$$S(A) \leq 4$$

Now all the possible submatrix of order 4.

$$\text{Here } \det A = 0$$

$$\therefore S(A) \neq 4$$

Now all the possible submatrix of order 3.

$$\begin{bmatrix} -3 & 0 & 2 \\ 2 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 2 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -3 & 2 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -3 & 0 \\ 0 & 2 & \frac{1}{2} \\ 0 & 0 & 0 \end{bmatrix},$$

$$\begin{bmatrix} 0 & 0 & 3 \\ 2 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 3 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 3 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & \frac{1}{2} \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 3 \\ -3 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 3 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 3 \\ 0 & -3 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 3 \\ -3 & 0 & 2 \\ 2 & \frac{1}{2} & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 3 \\ 0 & 0 & 2 \\ 0 & \frac{1}{2} & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 3 \\ 0 & -3 & 2 \\ 0 & 2 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 2 & \frac{1}{2} \end{bmatrix}$$

$$\det \text{ of } \begin{bmatrix} 2 & 0 & 3 \\ 0 & 0 & 2 \\ 0 & \frac{1}{2} & 0 \end{bmatrix} = 2(-14) - 3(0)$$

$$\begin{bmatrix} 0 & 0 & 2 \\ 0 & \frac{1}{2} & 0 \end{bmatrix} \neq 0$$

$\therefore$  rank of the given matrix is 3.

$$S(A) = 3.$$

6. Find the rank of  $\begin{pmatrix} 1 & -1 & 2 \\ 2 & 2 & -4 \\ -5 & 5 & -10 \\ 3 & -3 & 6 \end{pmatrix}$  using minors.

Given matrix is of order  $4 \times 3$ .

From the property

$$S(A) \leq \min \{4, 3\}$$

$$S(A) \leq 3$$

Now all possible submatrix of order 3.

$$\begin{bmatrix} 1 & -1 & 2 \\ 2 & 2 & -4 \\ -5 & 5 & -10 \end{bmatrix}, \begin{bmatrix} 2 & 2 & -4 \\ -5 & 5 & -10 \\ 3 & -3 & 6 \end{bmatrix}, \begin{bmatrix} 1 & -1 & 2 \\ -5 & 5 & -10 \\ 3 & -3 & 6 \end{bmatrix}, \begin{bmatrix} 1 & -1 & 2 \\ 2 & 2 & -4 \\ 3 & -3 & 6 \end{bmatrix}$$

here det of matrix is 0

$$\therefore S(A) \neq 3.$$

Now all possible submatrix of order 2.

$$\begin{bmatrix} 1 & -1 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} -1 & 2 \\ 2 & -4 \end{bmatrix}, \begin{bmatrix} 2 & 2 \\ -5 & 5 \end{bmatrix}, \begin{bmatrix} 2 & -4 \\ 5 & -10 \end{bmatrix}, \begin{bmatrix} -5 & 5 \\ 3 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 5 & -10 \\ -3 & 6 \end{bmatrix} \text{ and so on.}$$

here det of matrix is 0.

$$\therefore S(A) \neq 2.$$

Now all possible submatrix of order 1.

$$[1], [-1], [2], [2], [2], [-4], [-5], [5], [-10], [3], [-3], [6]$$

∴ here det of matrix is non-zero.

$$\therefore S(A) = 1.$$

3. Find the rank of  $\begin{pmatrix} 1 & 1 & a \\ 1 & a & 1 \\ a & 1 & 1 \end{pmatrix}$  using minors.

$$\det A = 1(a-1) - 1(1-a) + a(1-a^2)$$

$$= a - 1 - 1 + a + a - a^3$$

$$= -a^3 + 3a - 2$$

$$= a^3 - 3a + 2$$

$$a^2 + a - 2 = ?$$

$$\begin{array}{c|cc} a & a-1 & a^2-3a+2 \\ \hline & & a^3-a^2 \\ & & \hline & & \end{array}$$

$$\begin{array}{r} a^2-3a+2 \\ \hline a^2-a \\ \hline -2a+2 \\ \hline -2a+2 \\ \hline 0 \end{array}$$

$$\begin{aligned} a^2 + a - 2 &= a^2 + 2a - a - 2 \\ &= a(a+2) - (a+2) \\ &= (a+1)(a+2) \end{aligned}$$

$$\text{if } a=1 \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} = 1(1-1) - 1(1-1) + 1(1-1) \\ S(A) \neq 3$$

$$\text{Now all submatrix of order } 2, \quad \det A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \quad S(A) \neq 0$$

$\therefore$  now all submatrix of order  $1, \quad S(A) = 1$ .

$$\text{if } a = -2 \quad A = \begin{pmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{pmatrix} = 1(-2-1) - 1(1+2) + 2(1-4) \\ = -3 - 3 + 6 \\ = 0 \quad S(A) \neq 3$$

$$\text{Submatrix of order } 2 \quad \det \begin{pmatrix} 1 & 1 \\ 1 & -2 \end{pmatrix} \neq 0 \quad S(A) = 2$$

\* Practice examples :-

(I) Determine the rank of  $\begin{pmatrix} 3 & 5 & 1 \\ 2 & -2 & 4 \\ 7 & 1 & 9 \end{pmatrix}$  using minors.

Given matrix is of order  $3 \times 3$ .

From the property

$$S(A) \leq \min \{ 3, 3 \}$$

$$S(A) \leq 3$$

Now all possible submatrix of order 3.

$$\begin{aligned} \det A &= 3(18 - 4) - 5(18 - 28) + 7(2 + 14) \\ &= -66 + 50 + 16 \\ &= 0 \end{aligned}$$

$$S(A) \neq 3$$

Now all possible submatrix of order 2.

$$* \begin{bmatrix} 3 & 5 \\ 2 & -2 \end{bmatrix}, \begin{bmatrix} 5 & 1 \\ -2 & 4 \end{bmatrix}, \begin{bmatrix} 2 & -2 \\ 7 & 1 \end{bmatrix}, \begin{bmatrix} 5 & 1 \\ 1 & 9 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 7 & 9 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 5 \\ 7 & 1 \end{bmatrix}, \begin{bmatrix} 5 & 1 \\ -2 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 1 \\ 2 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 5 \\ 2 & -2 \end{bmatrix}$$

$$\therefore \det A \neq 0$$

$$\therefore S(A) = 2$$

2. Determine the rank of  $\begin{pmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{pmatrix}$  using minors.

$$\begin{aligned}\det A &= 1(bc^3 - cb^3) - 1(ac^3 - ca^3) + 1(cb^3 - ba^3) \\ &= bc^3 - cb^3 - ac^3 + ca^3 + cb^3 - ba^3 \\ &= ca^3 - ba^3 - cb^3 + ab^3 + bc^3 - ac^3 \\ &= a^3(c-b) + b^3(a-c) + c^3(b-a)\end{aligned}$$

$$\begin{pmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{pmatrix}$$

$$(c-a)(b-a)^2 = 0$$

$$(a-b)(b-a)^2 = 0$$

$$(a-b)(c-a)^2 = 0$$

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

$$\begin{pmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{pmatrix}$$

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

Rank of matrix is 2. Hence rank of matrix is 2.

3. Determine the rank of  $\begin{pmatrix} 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -3 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$  using minors.

given matrix is of order  $4 \times 4$   
from the property

$$S(A) \leq \min \{4 \times 4\}$$

$$S(A) \leq 4$$

$$\text{det of matrix} = 2 \begin{pmatrix} 0 & 0 & -1 \\ 0 & -3 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

$$= 2 (0 - 1 (-3))$$

$$= 2 (3)$$

$$= 6$$

$$\neq 0$$

rank of the given matrix is 4.

$$S(A) = 4.$$

4. Determine the rank of  $\begin{pmatrix} -1 & -2 & -1 \\ 6 & 12 & 6 \\ 5 & 10 & 5 \end{pmatrix}$  using minors

given matrix is of order  $3 \times 3$ .

from the property

$$S(A) \leq \min \{3 \times 3\}$$

$$S(A) \leq 3$$

Now,  $\det A$  is 0

$$\therefore S(A) \neq 3.$$

Now all possible submatrix of order 2.

$$\begin{bmatrix} -1 & -2 \\ 6 & 12 \end{bmatrix} \quad \begin{bmatrix} -2 & -1 \\ 12 & 6 \end{bmatrix} \quad \begin{bmatrix} 6 & 12 \\ 5 & 10 \end{bmatrix} \quad \begin{bmatrix} 12 & 6 \\ 10 & 5 \end{bmatrix} \quad \begin{bmatrix} -1 & -2 \\ 5 & 10 \end{bmatrix}$$

$$\begin{bmatrix} -2 & -1 \\ 10 & 5 \end{bmatrix}$$

$\therefore$  hence  $\det$  of the matrix is 0.

$$\therefore S(A) \neq 3$$

now all possible submatrices of order 1.

$[-1], [-2], [-1], [6], [12], [6], [5], [10], [5]$

here det of matrix is non zero.

$\therefore$  rank of the matrix is 1.

$$\therefore S(A) = 1.$$

(5) Determine the rank of  $\begin{pmatrix} 1 & 2 & -1 \end{pmatrix}$  using minors.

$\begin{pmatrix} 1 & 2 & -1 \\ 5 & 10 & -5 \end{pmatrix}$

given matrix is of order  $3 \times 3$ .

From the property

$$S(A) \leq \min \{2 \times 3\}$$

$$S(A) \leq 2$$

now, all possible submatrix of order 2.

$\begin{bmatrix} 1 & 2 \\ 5 & 10 \end{bmatrix}, \begin{bmatrix} 2 & -1 \\ 10 & -5 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 5 & -5 \end{bmatrix}$

$\therefore$  here det of submatrix is 0.

$$\therefore S(A) \neq 2.$$

now, all possible submatrix of order 1.

$[1], [2], [-1], [5], [10], [-5]$

here det of submatrix is non zero.

$$\therefore S(A) = 1.$$

(6) Determine the rank of  $\begin{pmatrix} 0 & 1 & -3 & -1 \\ 0 & 0 & 1 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & 2 & 0 \end{pmatrix}$  using minors

$\begin{pmatrix} 0 & 1 & -3 & -1 \\ 0 & 0 & 1 & 1 \\ 3 & 1 & 0 & 2 \\ 1 & 1 & 2 & 0 \end{pmatrix}$

given matrix is of order  $4 \times 4$ .

From the property

$$S(A) \leq \min \{4 \times 4\}$$

$$S(A) \leq 4.$$

det of matrix  $A = -1 \begin{pmatrix} 0 & 1 & 1 \\ 3 & 0 & 2 \\ 1 & 1 & 0 \end{pmatrix} + 3 \begin{pmatrix} 0 & 0 & 1 \\ 3 & 1 & 2 \\ 1 & 1 & 0 \end{pmatrix}$

$= -1 \begin{pmatrix} 0 & 0 & 1 \\ 3 & 1 & 0 \\ 1 & 1 & 2 \end{pmatrix} +$

$$\begin{aligned} &= -1 [-1(-2) + 1(3)] + 3 [1(3-1)] - 1 [1(3-1)] \\ &= -1 [5] + 3 [2] - 1 [2] \\ &= -5 + 5 - 2 \\ &= -1 \\ &\neq 0 \end{aligned}$$

here det of matrix is non-zero

$\therefore$  rank of the matrix is 4.

$$\therefore S(A) = 4.$$

### \* Nullity :

$\rightarrow$  Nullity of an  $m \times n$  matrix A is,  $n - r(A)$ .

where  $n$  is number of column.

It is denoted by  $n(r(A))$ .

\* Leading entry: The first nonzero entry of  $i^{th}$  row is known as the leading entry of  $i^{th}$  row.  
And the column containing the leading entry of  $i^{th}$  row is denoted by  $C(i)$ .

### \* Examples :-

①  $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 2 & 3 \end{bmatrix}$

②  $\begin{bmatrix} 0 & 3 & 4 \\ 2 & 1 & 3 \\ 0 & 0 & 3 \end{bmatrix}$

\* A matrix  $A$  is said to be in row-echelon form if

- (i) all zero rows are at the bottom of the matrix.
- (ii) all leading entries must be 1.
- (iii) leading entries moves from left to right if we go down. i.e.  $\ell(i) < \ell(i+1)$ .

e.g.  $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$

zero row at bottom.  
all leading entries are 1.

\* Upper triangular matrix with leading entries 0 and 1 will be in row echelon form.

\* Classwork Examples :

1. Reduce the matrix  $\begin{pmatrix} 1 & -1 & 2 & 3 & -2 \\ 0 & 1 & -1 & -6 & 6 \\ 0 & -2 & 2 & 4 & -4 \end{pmatrix}$  to row-

echelon / reduced row-echelon form and hence determine the rank.

Row echelon form :

$R_3 \rightarrow R_3 + 2R_2$

$$\begin{pmatrix} 1 & -1 & 2 & 3 & -2 \\ 0 & 1 & -1 & -6 & 6 \\ 0 & 0 & 0 & -8 & 8 \end{pmatrix}$$

$$R_3 \rightarrow \left(\frac{-1}{8}\right) R_3$$

$$\begin{pmatrix} 1 & -1 & 2 & -3 & -2 \\ 0 & 1 & -7 & -6 & 6 \\ 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

Reduced row echelon form:

$$\Rightarrow \begin{pmatrix} 1 & -1 & 2 & 3 & -2 \\ 0 & 1 & -1 & -6 & 6 \\ 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

$$R_1 \rightarrow R_1 + R_2$$

$$\begin{pmatrix} 1 & 0 & 1 & -3 & 4 \\ 0 & 1 & -1 & -6 & 6 \\ 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

$$R_1 \rightarrow R_1 + 3R_3$$

$$R_2 \rightarrow R_2 + 6R_3$$

$$\begin{pmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

$\text{r}(A) = \text{no. of non-zero rows in reduced row-echelon form.}$

$$\text{r}(A) = 3.$$

\* Reduced row-echelon form: A matrix  $A$  is said to be in reduced row-echelon form if

- (i) the matrix is in row-echelon form.
- (ii) all the entries in a column containing the leading entry must be zero other than the leading entry.

e.g.  $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$

3. Reduce the matrix  $\begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 5 \\ 1 & 5 & 5 & 7 \\ 8 & 1 & 14 & 17 \end{pmatrix}$  to row-echelon/

reduced row-echelon form and hence determine the rank.

$$R_2 \rightarrow R_2 - 2R_1$$

$$R_4 \rightarrow R_4 - 8R_1$$

$$\left( \begin{array}{cccc} 1 & 2 & 3 & 4 \\ 0 & -3 & -2 & -3 \\ 0 & 3 & 2 & 3 \\ 0 & -15 & -10 & -15 \end{array} \right)$$

$$R_2 \rightarrow -\frac{1}{3}R_2$$

$$\left( \begin{array}{cccc} 1 & 2 & 3 & 4 \\ 0 & 1 & \frac{2}{3} & 1 \\ 0 & 3 & 2 & 3 \\ 0 & -15 & -10 & -15 \end{array} \right)$$

$$R_3 \rightarrow R_3 - 3R_2$$

$$R_4 \rightarrow R_4 + 15R_2$$

$$\left( \begin{array}{cccc} 1 & 2 & 3 & 4 \\ 0 & 1 & 2/3 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

Reduced row echelon form:

$$R_1 \rightarrow R_1 - 2R_2$$

$$\left( \begin{array}{cccc} 1 & 0 & 5/3 & 2 \\ 0 & 1 & 2/3 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$$S(A) = 2.$$

4. Reduce the matrix  $\begin{pmatrix} 3 & 1 & 7 & 9 \\ 1 & 2 & 4 & 6 \\ 4 & -1 & 7 & 5 \end{pmatrix}$  to row-echelon form.

$$\left( \begin{array}{cccc} 3 & 1 & 7 & 9 \\ 1 & 2 & 4 & 6 \\ 4 & -1 & 7 & 5 \end{array} \right)$$

Reduced row echelon form and hence determine the rank.  
 $R_1 \leftrightarrow R_2, R_3 \leftrightarrow R_4$

$$\left( \begin{array}{ccc} 1 & 2 & 4 \\ 3 & 1 & 7 \\ 4 & -1 & 5 \\ 4 & -1 & 7 \end{array} \right)$$

$$R_2 \rightarrow R_2 - 3R_1$$

$$R_3 \rightarrow R_3 - 4R_1$$

$$R_4 \rightarrow R_4 - 4R_1$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 4 & 8 \\ 0 & -5 & -5 & 8 \\ 0 & -9 & -11 & 8 \\ 0 & -9 & -9 & 8 \end{array} \right)$$

$$R_2 \rightarrow R_2 - 5R_1$$

$$\left( \begin{array}{ccc|c} 1 & 2 & 4 & 8 \\ 0 & 1 & 1 & 8 \\ 0 & -9 & -11 & 8 \\ 0 & -9 & -9 & 8 \end{array} \right) \left( \begin{array}{ccc|c} 1 & 2 & 4 & 8 \\ 0 & 1 & 1 & 8 \\ 0 & 0 & -10 & 8 \\ 0 & 0 & 0 & 8 \end{array} \right)$$

$$R_1 \rightarrow R_1 - 2R_2$$

$$R_3 \rightarrow R_3 - R_4$$

$$R_4 \rightarrow R_4 + 9R_2$$

$$\left( \begin{array}{ccc|c} 1 & 0 & 2 & 8 \\ 0 & 1 & 1 & 8 \\ 0 & 0 & -2 & 8 \\ 0 & 0 & 0 & 8 \end{array} \right) \left( \begin{array}{ccc|c} 1 & 0 & 2 & 8 \\ 0 & 1 & 1 & 8 \\ 0 & 0 & 0 & 8 \\ 0 & 0 & 0 & 8 \end{array} \right)$$

$$R_3 \rightarrow R_3 - 2R_4$$

$$\left( \begin{array}{ccc|c} 1 & 0 & 2 & 8 \\ 0 & 1 & 1 & 8 \\ 0 & 0 & 1 & 8 \\ 0 & 0 & 0 & 8 \end{array} \right) \left( \begin{array}{ccc|c} 1 & 0 & 0 & 8 \\ 0 & 1 & 0 & 8 \\ 0 & 0 & 1 & 8 \\ 0 & 0 & 0 & 8 \end{array} \right)$$

$$R_1 \rightarrow R_1 - 2R_3$$

$$\therefore SCA = 3.$$

$$R_2 \rightarrow R_2 - R_3$$

2. Reduce the matrix  $\begin{pmatrix} 1 & 3 & 5 \\ 2 & -1 & 4 \\ -2 & 8 & 2 \end{pmatrix}$  to row-echelon form and hence determine the rank.

$$R_2 \rightarrow R_2 - 2R_1$$

$$R_3 \rightarrow R_3 + R_2$$

$$\begin{pmatrix} 1 & 3 & 5 \\ 0 & -7 & -6 \\ 0 & 7 & 6 \end{pmatrix}$$

$$R_2 \rightarrow \frac{1}{-7} R_2$$

$$-7 \quad R_2 \rightarrow R_2 + 7R_1$$

$$\begin{pmatrix} 1 & 3 & 5 \\ 0 & 1 & 6/7 \\ 0 & 7 & 6 \end{pmatrix}$$

$$R_1 \rightarrow R_1 - 3R_2$$

$$R_3 \rightarrow R_3 - 7R_2$$

$$\begin{pmatrix} 1 & 0 & 17/7 \\ 0 & 1 & 6/7 \\ 0 & 0 & 0 \end{pmatrix}$$

$$S(A) = 2$$

6. Reduce the matrix  $\begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & 2 \\ 2 & 2 & 3 \end{pmatrix}$  to row-echelon / reduced row-echelon form and hence determine the rank.

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - 2R_1$$

$$\begin{pmatrix} 1 & 1 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

$$R_1 \rightarrow R_1 - R_2$$

$$\begin{pmatrix} 0 & 0 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

$$R_3 \rightarrow \frac{1}{-1} R_3$$

$$R_1 \rightarrow R_1 - 2R_3$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\therefore \text{rank}(A) = 3.$$

↳ Solution of a system of linear eq<sup>n</sup> by crissus elimination and crissus Jordan Methods.

System of L.E. :

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \quad \text{①}$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \quad \text{②}$$

⋮

⋮

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

Matrix form

$$\left[ \begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & x_1 \\ a_{21} & a_{22} & \dots & a_{2n} & x_2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & x_n \end{array} \right] = \left[ \begin{array}{c} b_1 \\ b_2 \\ \vdots \\ b_m \end{array} \right]$$

↳ HOMOGENIOUS

if  $b_i = 0$  then its known as homogenous system.

↳ co-efficient matrix :

The matrix  $A = \left[ \begin{array}{cccc} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{array} \right]$  is called

co-efficient matrix of eq<sup>n</sup> ①

↳ Augmented Matrix :

The matrix  $(A/b)$

$\left[ \begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & & & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right]$  is called augmented matrix.

↳ solution of homogeneous eq<sup>n</sup>.

- A homogeneous system always has at least 1 solution namely which is known as a trivial solution.
- If rank of  $A = n$  then trivial solution is the only solution of the system.
- If rank of  $A < n$  then the system is infinitely many solution.

\* Homogeneous system :  $b = 0$

\* non-Homogeneous system :  $b \neq 0$

\* Solution of non homogeneous eq<sup>n</sup>.

- If  $\text{r}(A/b) \neq \text{r}(A)$ , then the system is said to be inconsistent. i.e the system does not have any solution.
- If  $\text{r}(A/b) = \text{r}(A) = n$ , then the system has unique solution.
- If  $\text{r}(A/b) = \text{r}(A) < n$ , then the system has infinitely many solutions.

1. solve the system:

$$2x + y - z = 4$$

$$x - y + 2z = -2$$

$$-x + 2y - z = 2$$

by criss cross elimination / Gauss-Jordan method, if is consistent.

$$(A|b) = \left( \begin{array}{ccc|c} 2 & 1 & -1 & 4 \\ 1 & -1 & 2 & -2 \\ -1 & 2 & -1 & 2 \end{array} \right)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 2 & -2 \\ 2 & 1 & -1 & 4 \\ -1 & 2 & -1 & 2 \end{array} \right)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 2 & -2 \\ 0 & 3 & -5 & 8 \\ 0 & 1 & 1 & 0 \end{array} \right) \quad (R_2 \rightarrow R_2 - 2R_1) \quad (R_3 \rightarrow R_1 + R_3)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 2 & -2 \\ 0 & 1 & 1 & 0 \\ 0 & 3 & -5 & 8 \end{array} \right) \quad (R_2 \leftrightarrow R_3)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 2 & -2 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & -8 & 8 \end{array} \right) \quad (R_3 \rightarrow R_3 - 3R_2)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 2 & -2 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & -1 & -1 \end{array} \right) \quad (R_3 \rightarrow -R_3)$$

Note that

$$\det(A) = \det(A/B) = 3$$

Gauss elimination method;

$$x - y + 2z = -2$$

$$y + z = 0$$

$$\boxed{z = -1}$$

$$\boxed{y = 1}$$

$$x - 1 - 2 = -2$$

$$\boxed{x = 1}$$

$$(x, y, z) = (1, 1, -1)$$

Gauss-Jordan method

$$\sim \left( \begin{array}{ccc|c} 1 & 0 & 3 & -2 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & -1 \end{array} \right) \quad (R_1 \rightarrow R_1 + R_2)$$

$$\sim \left( \begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \end{array} \right) \quad (R_2 \rightarrow R_2 - R_3) \quad (R_1 \rightarrow R_1 - 3R_3)$$

$$(x, y, z) = (1, 1, -1)$$

2. solve the system:

$$2x + z = 3,$$

$$x - y + z = 1,$$

$$4x - 2y + 3z = 3$$

by Gauss elimination / Gauss-Jordan method, if it is consistent.

$$(A|b) = \left( \begin{array}{ccc|c} 2 & 0 & 1 & 3 \\ 1 & -1 & 1 & 1 \\ 4 & -2 & 3 & 3 \end{array} \right)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 2 & 0 & 1 & 3 \\ 4 & -2 & 3 & 3 \end{array} \right) \quad (R_1 \leftrightarrow R_2)$$

$$\times \sim \left( \begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 1 & 0 & 1/2 & 3/2 \\ 0 & & & \end{array} \right) \quad (R_2 \rightarrow 1/2 R_2) \\ \quad (R_3 \rightarrow R_3 - 4R_1)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 0 & 2 & -1 & 1 \\ 0 & 2 & -1 & -1 \end{array} \right) \quad (R_2 \rightarrow R_2 - 2R_1) \\ \quad (R_3 \rightarrow R_3 - 4R_1)$$

$$\sim \left( \begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 0 & 1 & -1/2 & 1/2 \\ 0 & 2 & -1 & -1 \end{array} \right) \quad (R_2 \rightarrow 1/2 R_2)$$

$$\sim \left( \begin{array}{ccc|c} 1 & 0 & 1/2 & 3/2 \\ 0 & 1 & -1/2 & 1/2 \\ 0 & 0 & 0 & -2 \end{array} \right) \quad (R_1 \rightarrow R_1 + R_2) \\ \quad (R_3 \rightarrow R_3 - 2R_2)$$

$$\sim \left( \begin{array}{ccc|c} 1 & 0 & 1/2 & 3/2 \\ 0 & 1 & -1/2 & 1/2 \\ 0 & 0 & 0 & 1 \end{array} \right) \quad (\therefore R_3 \rightarrow -1/2 R_3)$$

Note that  $\det(A) = 2 \cdot \det(A|b) = 3$

Gauss Jordan method :-

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & 1/2 & 3/2 \\ 0 & 1 & -1/2 & 1/2 \\ 0 & 0 & 0 & 1 \end{array} \right] \quad R_1 \rightarrow R_1 + R_2$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & -\frac{1}{2} & 0 \\ 0 & 1 & -\frac{1}{2} & 0 \\ 0 & 0 & 0 & 1 \end{array} \right] \quad R_1 \rightarrow R_1 - \frac{1}{2}R_2, \quad R_2 \rightarrow R_2 - \frac{1}{2}R_3$$

$$\therefore \text{rref}(A) = I$$

$$\therefore \text{rref}(A/b) = I$$

$$\therefore \text{rref}(A/b) \neq \text{rref}(A)$$

$\therefore$  The system is inconsistent.

3. Solve the system:

$$4x - 3y + 9z + 6w = 0$$

$$2x + 3y + 3z + 6w = 6$$

$$4x - 21y - 39z - 6w = -24$$

by crasss elimination / crasss - Jordan method, if is consistent.

$$(A/b) = \left[ \begin{array}{cccc|c} 4 & -3 & -9 & 6 & 0 \\ 2 & 3 & 3 & 6 & 6 \\ 4 & -21 & -39 & -6 & -24 \end{array} \right]$$

$$\sim \left[ \begin{array}{cccc|c} 1 & -\frac{3}{4} & -\frac{9}{4} & \frac{6}{4} & 0 \\ 2 & 3 & 3 & 6 & 6 \\ 4 & -21 & -39 & -6 & -24 \end{array} \right] \quad (R_1 \rightarrow \frac{1}{4}R_1)$$

$$\sim \left[ \begin{array}{cccc|c} 1 & -\frac{3}{4} & -\frac{9}{4} & \frac{6}{4} & 0 \\ 0 & \frac{9}{2} & \frac{15}{2} & 3 & 6 \\ 0 & 18 & 30 & -12 & -24 \end{array} \right] \quad (R_2 \rightarrow R_2 + (-2)R_1) \quad (R_3 \rightarrow R_3 + (-4)R_1)$$

$$\sim \left[ \begin{array}{cccc|c} 1 & -\frac{3}{4} & -\frac{9}{4} & \frac{6}{4} & 0 \\ 0 & 1 & \frac{5}{3} & \frac{2}{3} & \frac{4}{3} \\ 0 & 18 & 30 & -12 & -24 \end{array} \right] \quad (R_2 \rightarrow \frac{(-2)}{9}R_2)$$

$$\sim \left[ \begin{array}{cccc|c} 1 & 0 & -1 & 2 & 1 \\ 0 & 1 & 15/9 & 2/3 & 9/13 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad CR_1 \rightarrow R_1 + (-\frac{3}{4})R_2$$

$$(R_3 \rightarrow R_3 + (-15)R_2)$$

Note that  $\det(A) = \det(A/b) = 2$

By Gauss Jordan Method,

$$x - z + 2w = 1$$

$$y + \frac{15}{9}z + \frac{2}{3}w = \frac{4}{3}$$

Take  $z = k_1$  and  $w = k_2$

$$x = 1 + z - 2w$$

$$= 1 + k_1 - 2k_2$$

and

$$y = \frac{4}{3} - \frac{15}{9}z - \frac{2}{3}w$$

$$= \frac{4}{3} - \frac{15}{9}k_1 - \frac{2}{3}k_2$$

4.  $3x + y + 2z = 0$

$$x - y + 3z = 0$$

$$x + 5y - 4z = 0$$

$$(A/b) = \left[ \begin{array}{ccc|c} 3 & 1 & 2 & 0 \\ 1 & -2 & 3 & 0 \\ 1 & 5 & -4 & 0 \end{array} \right]$$

$$\sim \left[ \begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ 3 & 1 & 2 & 0 \\ 1 & 5 & -4 & 0 \end{array} \right] \quad (R_1 \leftrightarrow R_2)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ 0 & 7 & -7 & 0 \\ 0 & 7 & -7 & 0 \end{array} \right] \quad (R_2 \rightarrow R_2 + (-3)R_1) \\ (R_3 \rightarrow R_3 + (-1)R_1)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 7 & -7 & 0 \end{array} \right] \quad (R_2 \rightarrow (\frac{1}{7})R_2)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \quad (R_1 \rightarrow R_1 + (2)R_2) \\ (R_3 \rightarrow R_3 + (-7)R_2)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \quad (R_3 \rightarrow (\frac{1}{8}R_3))$$

Note that  $\det(A) = \det(A/b) = 2$

By Gauss Jordan Method,

$$x + z = 0$$

$$y - z = 0$$

$$\text{take } z = +1$$

$$x = -1$$

$$y = 1$$

$$\{(-1, 1, 1) / k \in \mathbb{R}\}$$

$$\{k(-1, 1, 1) / k \in \mathbb{R}\}$$

5. Consider the system of linear equation

$$x + y + z = 6$$

$$x + 2y + 3z = 10$$

$$x + 2y + \lambda z = 9$$

find the value of  $\lambda$  and  $\mu$  so that the system (a) has a unique soln (b) has infinite number of solution (c) is inconsistent.

$$(A|b) = \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 1 & 2 & 3 & 10 \\ 1 & 2 & \lambda & u \end{array} \right]$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 1 & 2 & 3 & 10 \\ 0 & 0 & \lambda-3 & u-10 \end{array} \right] \quad (R_3 \rightarrow R_3 - R_2)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & \lambda-3 & u-10 \end{array} \right] \quad (R_2 \rightarrow R_2 - R_1)$$

→ If  $\lambda = 3$  &  $u \neq 10$  then  $\text{r}_1(A) = 2$  &  $\text{r}_1(A|b) = 3$   
 $\therefore$  The system is inconsistent.

→ If  $\lambda \neq 3$  &  $u \in \mathbb{R}$  then  $\text{r}_1(A) = \text{r}_1(A|b) = 3$   
 $\therefore$  The system is unique soln.

→ If  $\lambda = 3$  &  $u = 10$  then  $\text{r}_1(A) = \text{r}_1(A|b) = 2 < 3$ .  
 $\therefore$  The system infinite solution.

H.W. Investigate for what value of  $\lambda$  and  $u$

$$x + 2y + z = 8$$

$$2x + 2y + \lambda z = 13$$

$$3x + 4y + \lambda z = u$$

have (i) unique soln, (ii) infinite soln (iii)  
 $\therefore$  inconsistent soln.

6. find the value of  $k$ , so that the equation

$$x + y + 3z = 0$$

$$4x + 3y + kz = 0$$

$$2x + y + 2z = 0$$

have a non-trivial soln.

$$A = \begin{pmatrix} 1 & 1 & 3 \\ 4 & 3 & k \\ 2 & 1 & 2 \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 1 & 3 \\ 0 & -1 & k-12 \\ 0 & -1 & -4 \end{pmatrix} \quad (R_2 \rightarrow R_2 - 4R_1) \quad (R_3 \rightarrow R_3 - 2R_1)$$

$$\sim \begin{pmatrix} 1 & 1 & 3 \\ 0 & 1 & 12-k \\ 0 & 1 & 4 \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 1 & 3 \\ 0 & 1 & 4 \\ 0 & 1 & 12-k \end{pmatrix} \quad (R_2 \leftrightarrow R_3)$$

$$\sim \begin{pmatrix} 1 & 1 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 8-k \end{pmatrix} \quad (R_3 \rightarrow R_3 - R_2)$$

if  $k = 8$  then the rank of  $A = 2 < 3$

so the system has non-trivial soln.

7. By applying Kirchhoff's law to a circuit we obtain the following equations:

$$7i_1 + 9i_2 = 3$$

$$5i_1 + 7i_2 = 1$$

Where  $i_1$  and  $i_2$  represents currents. Find the values of  $i_1$  and  $i_2$  by Gauss elimination - Jordan method. If it is consistent.

$$(A|b) = \left[ \begin{array}{cc|c} 7 & 9 & 3 \\ 5 & 7 & 1 \end{array} \right]$$

$$(R_1 \rightarrow R_1)$$

$$\sim \left[ \begin{array}{cc|c} 1 & 9/7 & 3/7 \\ 5 & 7 & 1 \end{array} \right]$$

$$(R_2 \rightarrow R_2 - 5R_1)$$

$$\sim \left[ \begin{array}{cc|c} 1 & 9/7 & 3/7 \\ 0 & 4/7 & -8/7 \end{array} \right]$$

$$(R_2 \rightarrow \frac{R_2}{4/7})$$

$$\sim \left[ \begin{array}{cc|c} 1 & 9/7 & 3/7 \\ 0 & 1 & -2 \end{array} \right]$$

$$\sim \left[ \begin{array}{cc|c} 1 & 9/7 & 3/7 \\ 0 & 1 & -2 \end{array} \right]$$

$$(R_1 \rightarrow R_1 - \frac{9}{7}R_2)$$

$$\sim \left[ \begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & -2 \end{array} \right]$$

$$\therefore I_1 = 3$$

$$I_2 = -2$$

$$I_2 = -2$$

$$I_1 + 9I_2 = 3$$

$$+ +$$

$$I_1 + \frac{9(-2)}{7} = \underline{\underline{3}}$$

$$+ I_1 - 18 = 3$$

$$+ I_1 = 21$$

$$\boxed{I_1 = 3}$$

10. A pulley system gives the following equations

$$\ddot{x}_1 + \ddot{x}_2 = 0$$

$$2\ddot{x}_2 = 20 - T$$

$$5\ddot{x}_2 = 50 - T$$

where  $\ddot{x}_1, \ddot{x}_2$  represent acceleration and  $T$  represents tension in the rope. Determine  $\ddot{x}_1, \ddot{x}_2$  and  $T$  by criss-cross elimination / criss-cross - Tordan method. If it is consistent.

$$\rightarrow (A|b) = \left( \begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 2 & 0 & 1 & 20 \\ 0 & 5 & 1 & 50 \end{array} \right)$$

$$= \left( \begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & -2 & 1 & 20 \\ 0 & 5 & 1 & 50 \end{array} \right) \quad (R_2 \rightarrow R_2 - 2R_1)$$

$$= \left( \begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & -1/2 & -10 \\ 0 & 5 & 1 & 50 \end{array} \right) \quad (R_2 \rightarrow \underline{R_2}, (-2))$$

$$= \left( \begin{array}{ccc|c} 1 & 0 & +1/2 & +10 \\ 0 & 1 & -1/2 & -10 \\ 0 & 0 & 7/2 & 100 \end{array} \right) \quad (R_1 \rightarrow R_1 - R_2) \quad (R_3 \rightarrow R_3 - 5R_2)$$

$$= \left( \begin{array}{ccc|c} 1 & 0 & +1/2 & +10 \\ 0 & 1 & -1/2 & -10 \\ 0 & 0 & 1 & 200/7 \end{array} \right) \quad (R_3 \rightarrow \frac{2}{7}R_3)$$

$$= \left( \begin{array}{ccc|c} 1 & 0 & 0 & -30/7 \\ 0 & 1 & 0 & 30/7 \\ 0 & 0 & 1 & 200/7 \end{array} \right) \quad (R_1 \rightarrow R_1 - (\frac{1}{2})R_3) \quad (R_2 \rightarrow R_2 + (\frac{1}{2})R_3)$$

$$\therefore (x, y, z) = \left( \frac{-30}{7}, \frac{30}{7}, \frac{200}{7} \right)$$

Ex. Investigate for what value of  $\lambda$  cmd it  
 $x + 2y + z = 8$

$$2x + 2y + 2z = 13$$

$$3x + 4y + \lambda z = 0$$

have (i) unique soln (ii) infinite soln  
 (iii) inconsistent soln.

$$(A/b) = \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 8 \\ 2 & 2 & 2 & 13 \\ 3 & 4 & \lambda & 0 \end{array} \right]$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 8 \\ 0 & -2 & 0 & -3 \\ 0 & -2 & \lambda-3 & 0-24 \end{array} \right] \quad R_2 \rightarrow R_2 - 2R_1$$

$$R_3 \rightarrow R_3 - 3R_1$$

$$\cancel{X} \sim \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 8 \\ 0 & 1 & 0 & 3/2 \\ 0 & -2 & \lambda-3 & 0-24 \end{array} \right] \quad R_2 \rightarrow \frac{1}{-2} R_2$$

$$\cancel{X} \sim \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 8 \\ 0 & 1 & 0 & 3/2 \\ 0 & -2 & \lambda-3 & 0-24 \end{array} \right] \quad (R_1 \rightarrow R_1 + R_3) \\ (R_3 \rightarrow R_3 + R_1)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 8 \\ 0 & -2 & 0 & -3 \\ 0 & 0 & \lambda-3 & 0-24 \end{array} \right] \quad (R_3 \rightarrow R_3 - R_2)$$

$$R_2 \rightarrow \frac{R_2}{-2}$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 8 \\ 0 & 1 & 0 & 3/2 \\ 0 & 0 & \lambda-3 & \lambda-21 \end{array} \right]$$

→  $\det(A) = 3$  if  $\lambda \neq 3$  &  $\det(A/b) = 3$  if  $\lambda \neq 3$  so for unique soln we have  $\lambda \neq 3$ .

- if  $\lambda = 3$ ,  $b_1 = 21$  then soln has infinitely many soln.

→ If  $\lambda = 3$ ,  $b_1 \neq 21$  then soln has no soln.

$$II. \quad -4y + 2x + 5z = 15000$$

$$-2z + 3x + y = 1000$$

$$-x + 3y + z = 4000$$

$$2x - 4y + 5z = 15000$$

$$3x + y - 2z = 1000$$

$$-x + 3y + z = 4000$$

$$(A/b) = \left[ \begin{array}{ccc|c} 2 & -4 & 5 & 15000 \\ 3 & 1 & -2 & 1000 \\ -1 & 3 & 1 & 4000 \end{array} \right]$$

$$\cancel{\times} = \left[ \begin{array}{ccc|c} 1 & -2 & 5/2 & 7500 \\ 3 & 1 & -2 & 1000 \\ -1 & 3 & 1 & 4000 \end{array} \right] \quad (R_1 \rightarrow \frac{1}{2} R_1)$$

$$\cancel{\times} = \left[ \begin{array}{ccc|c} 1 & -2 & 5/2 & 7500 \\ 0 & 7 & -19/2 & -21500 \\ 0 & 1 & 7/2 & 11500 \end{array} \right] \quad (R_2 \rightarrow R_2 - 3R_1) \quad (R_3 \rightarrow R_3 + R_1)$$

~~X~~ = 
$$\left[ \begin{array}{ccc|c} 1 & -2 & \frac{5}{12} & 7500 \\ 0 & 1 & -\frac{19}{14} & -21500 \\ 0 & 1 & \frac{7}{12} & 11500 \end{array} \right] \quad (R_2 \rightarrow R_2 - F_2)$$

~~X~~ = 
$$\left[ \begin{array}{ccc|c} 1 & 0 & -\frac{3}{114} & \frac{95500}{17} \\ 0 & 1 & -\frac{19}{14} & -\frac{19500}{17} \\ 0 & 0 & \frac{34}{17} & \frac{102000}{17} \end{array} \right] \quad (R_1 \rightarrow R_1 + 2F_2) \quad (R_3 \rightarrow R_3 - F_2)$$

~~X~~ = 
$$\left[ \begin{array}{cc|c} 1 & 0 & -\frac{3}{114} \\ 0 & 0 & \frac{95500}{17} \end{array} \right] \quad (R_2 \rightarrow -\frac{19}{19} R_2)$$

$\sim \left[ \begin{array}{ccc|c} 1 & -3 & -1 & -4000 \\ 3 & 1 & -2 & 1000 \\ 2 & -4 & 5 & 15000 \end{array} \right] \quad (R_1 \rightarrow -R_1)$

$\sim \left[ \begin{array}{ccc|c} 1 & -3 & -1 & -4000 \\ 0 & 10 & 1 & 13000 \\ 0 & 2 & 7 & 23000 \end{array} \right] \quad (R_2 \rightarrow R_2 - 3R_1) \quad (R_3 \rightarrow R_3 - 2R_1)$

$\sim \left[ \begin{array}{ccc|c} 1 & -3 & -1 & -4000 \\ 0 & 1 & \frac{1}{10} & 1300 \\ 0 & 2 & 7 & 23000 \end{array} \right] \quad (R_2 \rightarrow R_2) \quad 10$

$\sim \left[ \begin{array}{ccc|c} 1 & 0 & -\frac{7}{110} & -100 \\ 0 & 1 & \frac{1}{10} & 1300 \\ 0 & 0 & \frac{34}{15} & 20400 \end{array} \right] \quad R_1 \rightarrow R_1 + 3R_2 \quad R_3 \rightarrow R_3 - 2R_2$

$\sim \left[ \begin{array}{ccc|c} 1 & 0 & -\frac{7}{110} & -100 \\ 0 & 1 & \frac{1}{10} & 1300 \\ 0 & 0 & 1 & 3000 \end{array} \right] \quad R_3 \rightarrow 5 R_3 \quad 34$

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 2000 \\ 0 & 1 & 0 & 1000 \\ 0 & 0 & 1 & 3000 \end{array} \right] \quad R_1 \rightarrow R_1 + \frac{7}{10} R_3$$

$$R_2 \rightarrow R_2 - \frac{1}{10} R_3$$

$$\therefore (x, y, z) = (2000, 1000, 3000)$$

12.  $9a + 3b + c = 64 \quad v(t) = at^2 + bt + c, \quad 0 \leq t \leq 100$   
 $8Gc + 6b + c = 133 \quad v(3) = 64 \quad v(9) = 208$   
 $81c + 9b + c = 208 \quad v(6) = 133$

$$(A | b) = \left[ \begin{array}{ccc|c} 9 & 3 & 1 & 64 \\ 81 & 9 & 1 & 208 \\ 36 & 6 & 1 & 133 \end{array} \right]$$

$$\sim \left[ \begin{array}{ccc|c} 9 & 3 & 1 & 64 \\ 0 & -6 & -3 & -123 \\ 0 & -18 & -8 & -368 \end{array} \right] \quad (R_3 \rightarrow R_3 - 9R_1)$$

$$(R_2 \rightarrow R_2 - 4R_1)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 1/3 & 1/9 & 64/9 \\ 0 & -6 & -3 & -123 \\ 0 & -18 & -8 & -368 \end{array} \right] \quad (R_1 \rightarrow \frac{R_1}{9})$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 1/3 & 1/9 & 64/9 \\ 0 & 1 & 1/2 & 123/16 \\ 0 & -18 & -8 & -368 \end{array} \right] \quad (R_2 \rightarrow \frac{R_2}{-6})$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & -1/18 & 5/18 \\ 0 & 1 & 1/2 & 123/16 \\ 0 & 0 & 1 & 1 \end{array} \right] \quad (R_1 \rightarrow R_1 - \frac{1}{3} R_2)$$

$$(R_3 \rightarrow R_3 + 18R_2)$$

$$\sim \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 1/3 \\ 0 & 1 & 0 & 20 \\ 0 & 0 & 1 & 1 \end{array} \right] \quad (R_1 \rightarrow R_1 + \frac{1}{18} R_3)$$

$$(R_2 \rightarrow R_2 - 1/2 R_3)$$

$$(a, b, c) = (1/3, 20, 1)$$

$$\begin{aligned}v(15) &= at^2 + bt + c \\&= \left(\frac{1}{3}\right)(15)^2 + 20(15) + 1 \\&= 75 + 300 + 1\end{aligned}$$

$$v(15) = 376$$