

# Almond Model

Anna Abelman, Julia Dagum, Jaleise Hall

4/8/2021

Summary:

```
#read in the clim data
#clim <- read_csv(here("clim.csv"))
clim <- read.table("clim.txt", sep=" ", header=T)

source("almond_model.R")

temperature <- clim %>%
  filter(month == "2") %>%
  group_by(year) %>%
  summarize(
    avg = mean(tmin_c)
  )

rain <- clim %>%
  filter(month == "1") %>%
  group_by(year) %>%
  summarize(
    sum = sum(precip)
  )

df <- data.frame(temperature, rain) %>%
  select(-year.1)

almond_model(clim)

## [1] -0.3552237  9.2906757  68.9130633  15.4280698  20.2083803
## [6]  2.4820009 1919.9811511  3.5818399 329.6938750  27.8636956
## [11] -0.1436364  9.5999883 159.5119587  0.2450914 -0.2585997
## [16] -0.2367722 656.3724121  18.6324135  20.2007396 576.2821943
## [21]  0.7367438 153.7655092

almond_yield_anomaly <- data.frame(year = rain$year, anomaly = almond_model(clim)) %>%
  mutate(year = lubridate::ymd(year, truncated = 2L))

ggplot(data = almond_yield_anomaly, aes(y = anomaly, x = year)) +
  geom_line() +
  scale_x_date(date_breaks = "1 year",
    date_labels = "%Y",
    limits = as.Date(c("1989-01-01", "2010-01-01"))) +
  labs(x = "Year",
    y = expression("Anomaly (ton" ~acre^-1~ ")"),
```

```

title = "Annual Almond Yield Anomaly (1989 - 2010)" +
theme_minimal() +
theme(panel.grid.minor.x = element_blank(), # to remove minor x axis
plot.title = element_text(hjust = 0.5),
axis.text.x = element_text(angle = 60, vjust = 1, hjust = 1))

```




Figure 1: Almond yield anomaly in California for all years 1989 to 2010. The anomaly is calculated using the regression model:  $Y = -0.015T_{n,2} - 0.0046T_{n,2}^2 - 0.07P_1 + 0.0043P_1^2 + 0.28$