

Computer Algebra

Lecture 9

James Davenport

University of Bath

13 September 2018

Definition (Lecture 8 Slide 10)

The *derivative* of $f := \sum_{i=0}^n a_i x^i$ is $f' := \sum_{i=1}^n i a_i x^{i-1}$

Corollary

$$(f + g)' = f' + g'; (fg)' = f'g + fg'.$$

Definition (Alternative)

The derivative (with respect to x) satisfies $x' = 1$ and $(f + g)' = f' + g'; (fg)' = f'g + fg'$.

Corollary

The derivative of $f := \sum_{i=0}^n a_i x^i$ is $f' := \sum_{i=1}^n i a_i x^{i-1}$

Corollary

$$\left(\frac{p}{q}\right)' = \frac{p'q - pq'}{q^2}$$

Proof. [No calculus needed]

$$q \left(\frac{p}{q}\right) = p$$

$$q' \left(\frac{p}{q}\right) + q \left(\frac{p}{q}\right)' = p'$$

$$q \left(\frac{p}{q}\right)' = p' - q' \left(\frac{p}{q}\right) = \frac{p'q - pq'}{q}$$

$$\left(\frac{p}{q}\right)' = \frac{p'q - pq'}{q^2}$$

Notation (Fundamental Theorem of Calculus)

(Indefinite) integration is the inverse of differentiation, i.e.

$$F = \int f \Leftrightarrow F' = f. \quad (1)$$

“Where did dx go?”, but x is, from this point of view, merely that object such that $x' = 1$, i.e. $x = \int 1$.

Proposition

If $\theta = \text{RootOf}(p(z))$ with $p = \sum_{i=0}^n a_i z^i$, then

$$\theta' = -\frac{\sum_{i=0}^n a'_i \theta^i}{\sum_{i=0}^n i a_i \theta^{i-1}}.$$

Integration of Rational Functions

The integration of polynomials is trivial:

$$\int \sum_{i=0}^n a_i x^i = \sum_{i=0}^n \frac{1}{i+1} a_i x^{i+1}. \quad (2)$$

Since any rational expression $f(x) \in K(x)$ can be written as

$$f = p + \frac{q}{r} \text{ with } \begin{cases} p, q, r \in K[x] \\ \deg(q) < \deg(r) \end{cases}, \quad (3)$$

and p is always integrable by (2), we have proved

Proposition (Decomposition Lemma (rational expressions))

In the notation of (3), f is integrable if, and only if, q/r is.

$\frac{q}{r}$ is a proper rational.

Integration of Proper Rationals

In fact, the integration of proper rational expressions is conceptually trivial (we may as well assume r is monic, absorbing any constant factor in q):

- 1 perform a square-free decomposition of $r = \prod_{i=1}^n r_i^i$;
- 2 factorize each r_i completely, as $r_i(x) = \prod_{j=1}^{n_i} (x - \alpha_{i,j})$;
- 3 perform a partial fraction decomposition of q/r as

$$\frac{q}{r} = \frac{q}{\prod_{i=1}^n r_i^i} = \sum_{i=1}^n \frac{q_i}{r_i^i} = \sum_{i=1}^n \sum_{j=1}^{n_i} \sum_{k=1}^i \frac{\beta_{i,j,k}}{(x - \alpha_{i,j})^k}; \quad (4)$$

- 4 integrate this term-by-term, obtaining

$$\int \frac{q}{r} = \sum_{i=1}^n \sum_{j=1}^{n_i} \sum_{k=2}^i \frac{-\beta_{i,j,k}}{(k-1)(x - \alpha_{i,j})^{k-1}} + \sum_{i=1}^n \sum_{j=1}^{n_i} \beta_{i,j,1} \log(x - \alpha_{i,j}). \quad (5)$$

From a practical point of view, this approach has several snags:

- ① we have to factor r , and even the best algorithms from the previous chapter can be expensive;
- ② we have to factor each r_i into *linear factors*, which might necessitate the introduction of algebraic numbers to represent the roots of polynomials;
- ③ These steps might result in a complicated expression of what is otherwise a simple answer.

These are practical problems

$$\begin{aligned} & \int \frac{5x^4 + 60x^3 + 255x^2 + 450x + 274}{x^5 + 15x^4 + 85x^3 + 225x^2 + 274x + 120} dx \\ &= \log(x^5 + 15x^4 + 85x^3 + 225x^2 + 274x + 120) \\ &= \log(x + 1) + \log(x + 2) + \log(x + 3) + \log(x + 4) + \log(x + 5) \end{aligned} \tag{6}$$

is pretty straightforward, but adding 1 to the numerator gives

$$\begin{aligned} & \int \frac{5x^4 + 60x^3 + 255x^2 + 450x + 275}{x^5 + 15x^4 + 85x^3 + 225x^2 + 274x + 120} dx \\ &= \frac{5}{24} \log(x^{24} + 72x^{23} + \dots + 102643200000x + 9331200000) \\ &= \frac{25}{24} \log(x + 1) + \frac{5}{6} \log(x + 2) + \frac{5}{4} \log(x + 3) + \frac{5}{6} \log(x + 4) + \frac{25}{24} \log(x + 5) \end{aligned} \tag{7}$$

These are practical problems

Adding 1 to the denominator is pretty straightforward,

$$\int \frac{5x^4 + 60x^3 + 255x^2 + 450x + 274}{x^5 + 15x^4 + 85x^3 + 225x^2 + 274x + 121} dx = \log(x^5 + 15x^4 + 85x^3 + 225x^2 + 274x + 121), \quad (8)$$

but adding 1 to both gives

$$= 5 \sum_{\alpha} \alpha \ln \left(x + \frac{2632025698}{289} \alpha^4 - \frac{2086891452}{289} \alpha^3 + \frac{608708804}{289} \alpha^2 - \frac{4556915}{17} \alpha + \frac{3632420}{289} \right), \quad (9)$$

where

$$\alpha = \text{RootOf} (38569 z^5 - 38569 z^4 + 15251 z^3 - 2981 z^2 + 288 z - 11). \quad (10)$$

Hence the challenge is to produce an algorithm that achieves (6) and (8) simply, preferably gives us the second form of the answer in (7), *but* is still capable of solving (9).

We might also wonder where (9) came from: see later.

Hermite's Algorithm

We can write our integral as

$$\int \frac{q}{r} = \frac{s_1}{t_1} + \int \frac{s_2}{t_2}, \quad (11)$$

where the integral on the right-hand resolves itself as *purely* a sum of logarithms, i.e. is the $\sum_{i=1}^n \sum_{j=1}^{n_i} \beta_{i,j,1} \log(x - \alpha_{i,j})$ term.

Galois Theory shows that (11) can be written without any α_i : Can we find it without involving them?

We can perform a square-free decomposition of r as $\prod r_i^j$, and then a partial fraction decomposition to write

$$\frac{q}{r} = \sum \frac{q_i}{r_i^j} \quad (12)$$

and, since each term is a rational expression and therefore integrable, it suffices to integrate (12) term-by-term.

Hermite's Algorithm continued

Now r_i and r'_i are relatively prime, so, by Extended Euclid, there are polynomials a and b satisfying $ar_i + br'_i = 1$. Therefore

$$\int \frac{q_i}{r_i^i} = \int \frac{q_i(ar_i + br'_i)}{r_i^i} \quad (13)$$

$$= \int \frac{q_i a}{r_i^{i-1}} + \int \frac{q_i b r'_i}{r_i^i} \quad (14)$$

$$= \int \frac{q_i a}{r_i^{i-1}} + \int \frac{(q_i b / (i-1))'}{r_i^{i-1}} - \left(\frac{q_i b / (i-1)}{r_i^{i-1}} \right)' \quad (15)$$

$$= - \left(\frac{q_i b / (i-1)}{r_i^{i-1}} \right) + \int \frac{q_i a + (q_i b / (i-1))'}{r_i^{i-1}}, \quad (16)$$

and we have reduced the exponent of r_i by one.

When programming this method one may need to take care of the fact that, while $\frac{q_i}{r_i'}$ is a proper rational expression, $\frac{q_i b}{r_i' - 1}$ may not be, but the excess is precisely compensated for by the other term in (16).

Hence, at the cost of a square-free decomposition and a partial fraction decomposition, but *not* a factorization, we have found the rational part of the integral, i.e. performed the decomposition of (11).

In fact, we have done somewhat better, since the $\int \frac{s_2}{t_2}$ term will have been split into summands corresponding to the different r_i .

The Ostrogradski–Horowitz Algorithm

It follows from (16) that, in the notation of (12) $t_1 = \prod r_i^{j-1}$. Furthermore every factor of t_2 arises from the r_i , and is not repeated. Hence we can choose

$$t_1 = \gcd(r, r') \text{ and } t_2 = r/t_1. \quad (17)$$

Having done this, we can solve for the coefficients in s_1 and s_2 , and the resulting equations are linear in the unknown coefficients. More precisely, the equations become

$$q = s'_1 \frac{r}{t_1} - s_1 \frac{t'_1 t_2}{t_1} + s_2 t_1, \quad (18)$$

where the polynomial divisions are exact, and the linearity is now obvious. The programmer should note that s_1/t_1 is guaranteed to be in lowest terms, but s_2/t_2 is not (and indeed will be 0 if there is no logarithmic term)

The Trager–Rothstein Algorithm [Rot76, Tra76]

We still have to integrate the logarithmic part. (6)–(9) shows that this may, but need not, require algebraic numbers. How do we tell? The answer is provided by the following observation; if we write the integral of the logarithmic part as $\sum c_i \log v_i$, we can determine the equation satisfied by the c_i , i.e. the analogue of (10), by purely rational computations.

So write

$$\int \frac{s_2}{t_2} = \sum c_i \log v_i, \quad (19)$$

where we can assume

- ① $\frac{s_2}{t_2}$ is in lowest terms;
- ② the v_i are polynomials (using $\log \frac{f}{g} = \log f - \log g$);
- ③ the v_i are square-free (using $\log \prod f_i^i = \sum i \log f_i$);
- ④ the v_i are relatively prime (using $c \log pq + d \log pr = (c + d) \log p + c \log q + d \log r$);
- ⑤ the c_i are all different (using $c \log p + c \log q = c \log pq$);
- ⑥ the c_i generate the smallest possible extension of the original

The Trager–Rothstein Algorithm (continued)

(19) can be rewritten as

$$\frac{s_2}{t_2} = \sum c_i \frac{v'_i}{v_i}. \quad (20)$$

Hence $t_2 = \prod v_i$ and, writing $u_j = \prod_{i \neq j} v_i$, we can write (20) as

$$s_2 = \sum c_i v'_i u_i. \quad (21)$$

Furthermore, since $t_2 = \prod v_i$, $t'_2 = \sum v'_i u_i$. Hence

$$\begin{aligned} v_k &= \gcd(0, v_k) \\ &= \gcd\left(s_2 - \sum c_i v'_i u_i, v_k\right) \\ &= \gcd\left(s_2 - c_k v'_k u_k, v_k\right) \\ &\quad \text{since all the other } u_i \text{ are divisible by } v_k \\ &= \gcd\left(s_2 - c_k \sum v'_i u_i, v_k\right) \\ &\quad \text{for the same reason} \\ &= \gcd\left(s_2 - c_k t'_2, v_k\right). \end{aligned}$$

The Trager–Rothstein Algorithm (continued)

But if $l \neq k$,

$$\begin{aligned}\gcd(s_2 - c_k t'_2, v_l) &= \gcd\left(\sum c_i v'_i u_i - c_k \sum v'_i u_i, v_l\right) \\ &= \gcd(c_l v'_l u_l - c_k v'_l u_l, v_l) \\ &\quad \text{since all the other } u_i \text{ are divisible by } v_l \\ &= 1.\end{aligned}$$

Since $t_2 = \prod v_l$, we can put these together to deduce that

$$v_k = \gcd(s_2 - c_k t'_2, t_2). \quad (22)$$

Given c_k , this will tell us v_k . But we can deduce more from this: the c_k are precisely those numbers λ such that $\gcd(s_2 - \lambda t'_2, t_2)$ is non-trivial. Hence λ must be such that

$$P(\lambda) := \text{Res}_x(s_2 - \lambda t'_2, t_2) = 0. \quad (23)$$

If t_2 has degree n , $P(\lambda)$ is the determinant of a $2n - 1$ square matrix, n of whose rows depend linearly on λ , and thus is a polynomial of degree n in λ .

Complex Numbers?

The same process also leads to

$$\int \frac{1}{x^2 + 1} dx = \frac{i}{2} (\ln(1 - ix) - \ln(1 + ix)), \quad (24)$$

at which point the reader might complain “I asked to integrate a *real* function, but the answer is coming back in terms of *complex* numbers”. The answer is, of course, *formally correct*: differentiating the right-hand side of (24) yields

$$\frac{i}{2} \left(\frac{-i}{1 - ix} - \frac{i}{1 + ix} \right) = \frac{i}{2} \left(\frac{-i(1 + ix)}{1 + x^2} - \frac{i(1 - ix)}{1 + x^2} \right) = \frac{1}{1 + x^2} :$$

the issue is that the reader, *interpreting* the symbols \log etc. as the usual functions of calculus, is surprised.



M. Rothstein.

Aspects of Symbolic Integration and Simplification of Exponential and Primitive Functions.

PhD thesis, Univ. of Wisconsin, 1976.



B.M. Trager.

Algebraic Factoring and Rational Function Integration.

In R.D. Jenks, editor, *Proceedings SYMSAC 76*, pages 219–226, 1976.