

Billiards

Jonathan Allen, John Wang

Massachusetts Institute of Technology

November 22nd, 2013


Introduction

- Billiard ball bouncing in a square
- Assume no gravity or friction
- Examine sequence of side collisions

Example

Example


Examine the periodic sequence: 'abab'



Another Example

Example

Examine the periodic sequence: 'aaabaaab'




Presentation Outline

- 1 Introduction
 - Examples
 - Outline
 - Notation and Problem Statement
- 2 Lemmas
 - Tiling
 - 1-dimensional Problem
- 3 Algorithm
- 4 Future Research
 - Tileable Polygons
 - Non-Tileable Polygons
 - Circles

Notation

Definition


A table T is the unit square in \mathbb{R}^2 . A particle $p \in T$ begins at position $\bar{x}_0 \in T$ with velocity \bar{v} . When the particle reaches an edge of the table, velocity is reflected about the line perpendicular to the table's edge.



Notation

Definition

Opposite sides of the table are named a and b . **Primary side** (most collisions) is a , **secondary side** is b .



Notation

Definition

Collision string consists of the sides of the table that have been collided with for a given starting position and velocity.

Notation

Definition

Collision string consists of the sides of the table that have been collided with for a given starting position and velocity.

Definition

Primary substring is a subsequence from the collision string which contains the primary side collisions that occur between consecutive secondary side collisions.

Notation

Definition

Collision string consists of the sides of the table that have been collided with for a given starting position and velocity.

Definition

Primary substring is a subsequence from the collision string which contains the primary side collisions that occur between consecutive secondary side collisions.

Example

Collision string: 'aabaaabaabaaab', **Primary substrings:** 'aa', 'aaa'

Problem Statement

Problem: Characterize the properties of collision sequences.

- Given a sequence of a 's and b 's, determine if it is a valid collision sequence.
- Given a valid collision sequence, determine a possible starting position and velocity.

- 1 Introduction
 - Examples
 - Outline
 - Notation and Problem Statement

- 2 Lemmas
 - Tiling
 - 1-dimensional Problem

- 3 Algorithm

- 4 Future Research
 - Tileable Polygons
 - Non-Tileable Polygons
 - Circles

- 1 Introduction
 - Examples
 - Outline
 - Notation and Problem Statement

- 2 Lemmas
 - Tiling
 - 1-dimensional Problem



Representing Collision Strings

- Reflect squares about each side to create a tiling
- Solutions become lines in the plane
- Intersections become places where collisions occur

Representing Collision Strings

Example



Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Representing Collision Strings

Example



Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Representing Collision Strings

Example



Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Representing Collision Strings

Example



Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Representing Collision Strings

Example



Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Representing Collision Strings

Example



Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Representing Collision Strings

Example



Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Representing Collision Strings

Example

Tiling of $\vec{x}_0 = (0, 0.5)$ and $\vec{v} = (0.25, 0.25)$.



Periodicity of Rationals

We can represent a particle as a line in the plane $y = mx + y_0$ in the tiling. Periodicity occurs if $y_0 2k = mx + y_0$ for some $k \in \mathbb{N}$.

- If $m, y_0 \in \mathbb{Q}$, then all sequences are periodic.
- If m or y_0 are irrational, then the sequences are not periodic.

Maximum Differences between Primary Substring Lengths

Given a collision string, how different can primary substrings be?

Example

Is *abaaab* possible?

Maximum Differences between Primary Substring Lengths

Length of primary substring i is given by:

$$L(i, m, y_0) = \lfloor \frac{i - y_0}{m} \rfloor - \lfloor \frac{i - 1 - y_0}{m} \rfloor \quad (1)$$

Maximum Differences between Primary Substring Lengths

For $i, j \in \mathbb{N}$, $m \in [0, 1]$, $y_0 \in [0, 1]$, we will show:

$$\max_{i > j} L(i, m, y_0) - L(j, m, y_0) \leq 2 \quad (2)$$

Maximum Differences between Primary Substring Lengths

Lemma

$$\lfloor a - x \rfloor + \lfloor b - x \rfloor = \lfloor a \rfloor - \lfloor b \rfloor + (\lfloor \{a\} - \{x\} \rfloor - \lfloor \{b\} - \{x\} \rfloor) \quad (3)$$

$$L(i, m, y_0) - L(j, m, y_0) = \left(\left\lfloor \frac{i}{m} \right\rfloor - \left\lfloor \frac{i-1}{m} \right\rfloor \right) + \left(\left\lfloor \frac{j}{m} \right\rfloor - \left\lfloor \frac{j-1}{m} \right\rfloor \right) \quad (4)$$

$$+ \left(\left\lfloor \left\{ \frac{i}{m} \right\} - \left\{ \frac{y_0}{m} \right\} \right\rfloor - \left\lfloor \left\{ \frac{i-1}{m} \right\} - \left\{ \frac{y_0}{m} \right\} \right\rfloor \right) \quad (5)$$

$$- \left(\left\lfloor \left\{ \frac{j}{m} \right\} - \left\{ \frac{y_0}{m} \right\} \right\rfloor - \left\lfloor \left\{ \frac{j-1}{m} \right\} - \left\{ \frac{y_0}{m} \right\} \right\rfloor \right) \quad (6)$$

Maximum Differences between Primary Substring Lengths

Let $g(x, m, y_0) = \lfloor \{\frac{x}{m}\} - \{\frac{y_0}{m}\} \rfloor - \lfloor \{\frac{x-1}{m}\} - \{\frac{y_0}{m}\} \rfloor$.

$$L(i, m, y_0) - L(j, m, y_0) = \left\{ \frac{i-1}{m} \right\} - \left\{ \frac{i}{m} \right\} - \left(\left\{ \frac{j-1}{m} \right\} - \left\{ \frac{j}{m} \right\} \right) \quad (7)$$

$$+ g(i, m, y_0) - g(j, m, y_0) \quad (8)$$

Maximum Differences between Primary Substring Lengths

$$L(i, m, y_0) - L(j, m, y_0) = \left\{ \frac{i-1}{m} \right\} - \left\{ \frac{i}{m} \right\} - \left(\left\{ \frac{j-1}{m} \right\} - \left\{ \frac{j}{m} \right\} \right) \quad (9)$$

$$+ g(i, m, y_0) - g(j, m, y_0) \quad (10)$$

- If $g(x, m, y_0) = 1$, then:

$$\left\{ \frac{x-1}{m} \right\} < \left\{ \frac{y_0}{m} \right\} < \left\{ \frac{x}{m} \right\} \quad (11)$$

Which implies $0 < L(x, m, y_0) < 1$.

- If $g(x, m, y_0) = -1$, then:

$$\left\{ \frac{x}{m} \right\} < \left\{ \frac{y_0}{m} \right\} < \left\{ \frac{x-1}{m} \right\} \quad (12)$$

Which implies $-1 < L(x, m, y_0) < 0$.

- If $g(x, m, y_0) = 0$, then: $-1 < L(x, m, y_0) < 1$.

Maximum Differences between Primary Substring Lengths

$$\max_{i>j} L(i, m, y_0) - L(j, m, y_0) \leq 2 \quad (13)$$

Sequence Characterization

$$\begin{array}{cccccccccccc}
 \underbrace{aaa}_3 & b & \underbrace{aa}_2 & b & \underbrace{aa}_2 & b & \underbrace{aaa}_3 & b & \underbrace{aa}_2 & b & & \\
 & & & & & & \underbrace{aa}_2 & b & \underbrace{aaa}_3 & b & \underbrace{aa}_2 & b & \underbrace{aaa}_3 & \dots \\
 & & & & & & & & & & & & & \dots
 \end{array}$$

Sequence Characterization

$$\begin{array}{ccccccc}
 3 & \underbrace{22}_{2} & 3 & \underbrace{22}_{2} & 3 & \underbrace{2}_{1} & 3 \dots \\
 & & & & & & \dots
 \end{array}$$

- 1 Introduction
 - Examples
 - Outline
 - Notation and Problem Statement

- 2 Lemmas
 - Tiling
 - 1-dimensional Problem

- 3 Algorithm

- 4 Future Research
 - Tileable Polygons
 - Non-Tileable Polygons
 - Circles

- 1 Introduction
 - Examples
 - Outline
 - Notation and Problem Statement

- 2 Lemmas
 - Tiling
 - 1-dimensional Problem

- 1 Introduction
 - Examples
 - Outline
 - Notation and Problem Statement

- 2 Lemmas
 - Tiling
 - 1-dimensional Problem

- 3 Algorithm

- 4 Future Research
 - Tileable Polygons
 - Non-Tileable Polygons
 - Circles

- 1 Introduction
 - Examples
 - Outline
 - Notation and Problem Statement

- 2 Lemmas
 - Tiling
 - 1-dimensional Problem

Extensions to Tileable Polygons

Other Tileable Polygons:




Figure: Regular Hexagons




Figure: Equilateral Triangles

Extensions to Non-Tileable Polygons

- Irregular triangles
- Pentagons
- Octagons

Extensions to Circles

- Characterize how particle bounces around circle
- Analog to a , b might be sequence of collision points as you move around circle.

