

AL BARAKH ORGANICS

Farmer's Application & Crop Dosage Guide

Practical Field Application Guidelines for Maximum Yield & Cost Reduction

Recommended Field Dosage by Crop Type (40 KG Bags)

1. Field Crops (Wheat, Rice, Cotton, Maize, Sugarcane)

- Dosage Rate: 4 to 6 Bags (40 KG) per Acre
- Application Timing: Apply during land preparation / basal furrow before sowing.

2. Fruit Orchards (Mango, Citrus, Guava, Dates)

- Dosage Rate: 5 to 10 KG per Mature Tree
- Application Timing: Applied twice a year around the drip line (Pre-flowering & Post-harvest).

3. Vegetables & High-Value Tunnels (Tomato, Chili, Potato)

- Dosage Rate: 8 to 10 Bags (40 KG) per Acre
- Application Timing: Bed preparation and root zone furrow incorporation.

4. Plant Nurseries & Seedling Beds

- Dosage Rate: 25% to 30% Blend with Soil
- Application Timing: Mix thoroughly with soil/sand for potting and nursery germination trays.

5. Kitchen Gardens & Potted Plants (5 KG Bag)

- Dosage Rate: 2 to 3 Tablespoons per Pot
- Application Timing: Top-dress around plant roots once every 3 to 4 weeks, followed by light watering.

Key Rules for Successful Application:

1. Soil Moisture: Apply vermicompost when soil is slightly moist or irrigate within 24-48 hours.
2. Root Zone Placement: Incorporate 2-4 inches into the soil so living microbes are shielded from direct sun.
3. Chemical Reduction: Gradually reduce synthetic Urea and DAP by 40% to 60% in Year 1.
4. Storage: Always store vermicompost bags in cool shade; avoid direct sun exposure to keep microbes alive.

For Custom Soil Assessments & Bulk Delivery: 0316 8803363

Al Barakh Organics — Tando Ghulam Ali & Hyderabad Branches | Serving All Over Pakistan

Website: www.albarakhorganics.com | Mon - Sat: 09:00 - 17:00