

First Order Linear ODEs

James Arthur

September 19, 2020

Contents

1	Recap of previous modules	2
1.1	Notation:	2
1.2	Prerequisites:	2
2	Basic Defintions and Concepts	2
2.1	Classifications	2
2.2	Review of integration methods	2
2.3	Concepts	2
2.4	Famous models	3
3	Analytical Solutions	3
3.1	Seperation of variables	3
3.2	Reduction to seperable form	3
3.3	Exact ODEs and integrating factors	3

1 Recap of previous modules

A differentiable equation is an mathematical relation involving a derivative of a dependant variable w.r.t. single/many independant variables

1.1 Notation:

$$\frac{dx}{dt} = \dot{x}, \quad \frac{dy}{dx} = y', \quad f\left(x, y, \frac{dy}{dx}, \dots, \frac{d^ny}{dx^n}\right) = 0$$

$$f_{xy} = \frac{\partial^2 f}{\partial x \partial y}$$

1.2 Prerequisites:

Integration, ODEs and PDEs from Advanced Calculus.

2 Basic Defintions and Concepts

2.1 Classifications

1. Use of full or partial derivative
2. Coefficients are functions of independant variables only / constant, otherwise non Linear
3. The highest derivative is the order of the description DE
4. Degree is the highest derivative in rationalised form
5. An explicit solution is; $f = F(x, y, z, t)$ and implicit solution; $F(f, x, y, z, t) = 0$
6. Initial Value Problem (time) or boundary value problem (space).

2.2 Review of integration methods

1. List of commonly used integrals (link)
2. Polynomials, logarithms, trigonometric, inverse, hyperbolic and inverse hyperbolic trig.

2.3 Concepts

1. Given a DE, we want a solution.
2. A solution is a derived relation between the dependant and independant variables without any derivative term and defined in the interval / domain / region.
3. replacing the solution within the domain satisfies the description DE
4. Needs integration on one or two variables
5. Not always analytical and closed forms possible. (could use numerical integration, iterative solution schemes.)
1. Linear higher order ODEs (Laplace transforms)
 - (a) transforms linear ODEs in algebraic forms
 - (b) needs table of laplace, inverse laplace formula
2. the Geometric meaning is the slope of $y(x)$: $y'(x) = f(x_0, y_0)$ implies at a point (x_0, y_0) is the slope at $\frac{dy}{dx}$
3. There is something called a direction field that we can use to visualise the DE without solving it.
4. Curves of equal inclination $f(x, y) = c$ along which derivative is constant. Lots of parallel lines.
5. Limitation of direction field give an overall idea about the solution but have limited accuracy
6. Orthogonal trajectories are a family of curves that intersect another family of curves at right angles. For a curve $G(x, y, c)$ firstly find out $\frac{dy}{dx} = f(x, y)$. General solution of the orthogonal trajectory $\frac{dy}{dx} = \frac{-1}{f(x, y)}$.
7. existence is under what condition there is at least one Solution
8. uniqueness is what condition it has at most one Solution item the general solution contains the constants of integration
9. particular solution are when you use the initial / boundary conditions

2.4 Famous models

- Van der Pol oscillator

$$\begin{aligned}\dot{x} &= y \\ \dot{y} &= \mu(1 - x^2)y - x\end{aligned}$$

- Lorentz Attractor

$$\begin{aligned}\dot{x} &= \sigma(y - x) \\ \dot{y} &= x(\rho - z) - y \\ \dot{z} &= xy - \beta z\end{aligned}$$

Have to be careful between phase portrait and direction fields. Phase portraits are almost a guess and direction field as you have a solution at every point with a direction field.

3 Analytical Solutions

3.1 Seperation of variables

$$\begin{aligned}g(y)y' &= f(x) \\ \int g(y)dy &= \int f(x)dx + c\end{aligned}$$

3.2 Reduction to seperable form

$$\begin{aligned}y' &= f\left(\frac{y}{x}\right) \\ v + x\frac{dv}{dx} &= f(v) \quad \text{by letting } y = vx \\ \int \frac{dv}{f(v) - v} &= \int \frac{dx}{x} + c \\ &= \ln|x| + c\end{aligned}$$

3.3 Exact ODEs and integrating factors

Let us have an ODE: $M(x, y) + N(x, y)\frac{dy}{dx} = 0$ which can be written as $M(x, y)dx + N(x, y)dy$. We can then write a total differential as partial derivatives:

$$du = \frac{\partial u}{\partial x}dx + \frac{\partial u}{\partial y}dy = 0 \implies u(x, y) = c \quad (1)$$

Which then we can compare the two and we get:

$$M = \frac{\partial u}{\partial x}, \quad N = \frac{\partial u}{\partial y}$$

$$\implies \frac{\partial M}{\partial y} = \frac{\partial^2 u}{\partial y \partial x}, \quad \frac{\partial N}{\partial x} = \frac{\partial^2 u}{\partial x \partial y}$$

So then we can say for an ODE to be exact:

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

To solve the DE, take (1) and integrate with respect to u:

$$du = \frac{\partial u}{\partial x}dx + \frac{\partial u}{\partial y}dy = 0 \quad (1)$$

$$u = \int Mdx + K(y) = c$$