

Algebraic Topology

Part III preparatory notes based on Algebraic Topology by Allen Hatcher

Notes taken by James Arthur

Summer 2021

Part III requires a load of complex and interesting Mathematics that I have not covered thus far in my course and will not. These collections of notes are my thoughts on topics and needed intuition to bits of Mathematics. They aren't here to teach or to be used a stand alone text.

Contents

0	Underlying Notions	2
0.1	Cell Complexes	2

0 Underlying Notions

We are going to adopt the convention that every map is continuous unless otherwise stated.

We want to be able to talk about ways to transform some shapes into other shapes and think about them then being equal. We do this by considering an $I \in [0, 1]$ that we can parameterise a function by. So if we have two curves $s : X \rightarrow X$ and $t : X \rightarrow X$, we can define some $f : X \times I \rightarrow X$ such that,

$$\begin{cases} f_0(x) &= s(x) \\ f_1(x) &= t(x) \end{cases}$$

Definition 0.1 (Deformation Retraction). A Deformation retraction of a space X onto an $A \subset X$ is a family of maps, $f_t : X \rightarrow X$ $t \in I$, such that, $f_0 = \mathbb{1}$, $f_t(X) = A$ and $f_t|_A = \mathbb{1} \forall t$. The family f_t should be continuous in the sense that the associated map $X \times I \rightarrow X$, $(x, t) \mapsto f_t(x)$ is continuous.

Definition 0.2 (Mapping Cylinder). A mapping cylinder, M_f , for a $f : X \rightarrow Y$ is the quotient space of the disjoint union $(X \times I) \amalg Y$, obtained by identifying each $(x, 1) \in X \times I$ with $f(x) \in Y$.

A mapping cylinder deformation, of f , retracts to Y .

A deformation retraction $f_t : X \rightarrow X$ is just a special case of a homotopy, which is just simply any family of maps, $f_t : X \times I \rightarrow Y$, given by $F(x, t) = f_t(x)$ is continuous.

Definition 0.3 (Homotopic). One says two maps are homotopic if there exists a homotopy f_t connecting them and one writes $f_0 \simeq f_1$

Retractions are the topological analogue of projection operators.

Definition 0.4 (Homotopy Relative). A homotopy $f_t : X \rightarrow Y$ whose restriction to a subspace $A \subset X$ is independent of t is homotopy relative, written as homotopy rel.

Definition 0.5 (Homotopy Equivalence). A map $f : X \rightarrow Y$ is called a homotopy equivalence if there exists a $g : Y \rightarrow X$ such that $fg \simeq \mathbb{1}$ and $gf \simeq \mathbb{1}$.

Moreover, we can now say that X and Y are homotopy equivalent or have the same homotopy type.

Lemma 0.6. Any two spaces X and Y are homotopy equivalent if there exists some other space Z containing both X and Y are deformation retracts.

Definition 0.7 (Contractible). A space that has the homotopy type of a point is contractible.

This amounts to the identity map of this space to be nullhomotopic, that is homotopic to a constant map.

0.1 Cell Complexes

To construct a cell complex, let us follow these rules,

- (i) Start with a discrete set X^0 , whose points are regarded as 0-cells.
- (ii) inductively, form the n -skeleton X^n from X^{n-1} by attaching n -cells e_α^n via maps $\phi_\alpha : S^{n-1} \rightarrow X^{n-1}$ with a collection of n -disks D_α^n under the identifications $x \sim \phi_\alpha(x)$ for $x \in \partial D_\alpha^n$. Thus $X^n = X^{n-1} \amalg_\alpha e_\alpha^n$.
- (iii) We can either stop at a finite amount or continue indefinitely. in the latter, X is given a weak topology, A set $A \subset X$ is open if and only if $A \cap X^n$ is open in X^n for each n . This similarly works for closed.

X is a cell complex or a CW complex. If $X = X^n$ then X is said to be finite dimensional and the smallest such n is the dimension of X , the maximum dimension of cells of X .

Notation. e^n denotes an n -cell.

Definition 0.8 (Characteristic Map). Each cell e_α^n in a cell complex X has a characteristic map $\Phi_\alpha : D_\alpha^n \rightarrow X$ which extends ϕ_α and is a homeomorphism from the interior of D_α^n onto e_α^n . Namely, we can take ϕ_α to be the composition, $D_\alpha^n \hookrightarrow X^{n-1} \coprod_\alpha D_\alpha^n X^n \hookrightarrow X^n$, where the middle map is the quotient map defining X^n .

Definition 0.9 (Subcomplex). A subcomplex of a cell complex X is a closed subspace $A \subset X$ this is a union of cells of X .

Definition 0.10 (CW Pair). A complex X with a subcomplex A , (X, A) , is called a CW pair.