

# The GGCMI Phase II experiment: simulating and emulating global crop yield responses to changes in carbon dioxide, temperature, water, and nitrogen levels

James Franke<sup>a,b,\*</sup>, Joshua Elliott<sup>b,c</sup>, Christoph Müller<sup>d</sup>, Alexander Ruane<sup>e</sup>, Abigail Snyder<sup>f</sup>, Jonas Jägermeyr<sup>c,b,d,e</sup>, Juraj Balkovic<sup>g,h</sup>, Philippe Ciais<sup>i,j</sup>, Marie Dury<sup>k</sup>, Pete Falloon<sup>l</sup>, Christian Folberth<sup>g</sup>, Louis François<sup>k</sup>, Tobias Hank<sup>m</sup>, Munir Hoffmann<sup>n</sup>, Cesar Izaurralde<sup>o,p</sup>, Ingrid Jacquemin<sup>k</sup>, Curtis Jones<sup>o</sup>, Nikolay Khabarov<sup>g</sup>, Marian Koch<sup>n</sup>, Michelle Li<sup>b,l</sup>, Wenfeng Liu<sup>r,i</sup>, Stefan Olin<sup>s</sup>, Meridel Phillips<sup>e,t</sup>, Thomas Pugh<sup>u,v</sup>, Ashwan Reddy<sup>o</sup>, Xuhui Wang<sup>i,j</sup>, Karina Williams<sup>l</sup>, Florian Zabel<sup>m</sup>, Elisabeth Moyer<sup>a,b</sup>

<sup>a</sup>Department of the Geophysical Sciences, University of Chicago, Chicago, IL, USA

<sup>b</sup>Center for Robust Decision-making on Climate and Energy Policy (RDCEP), University of Chicago, Chicago, IL, USA

<sup>c</sup>Department of Computer Science, University of Chicago, Chicago, IL, USA

<sup>d</sup>Potsdam Institute for Climate Impact Research, Leibniz Association (Member), Potsdam, Germany

<sup>e</sup>NASA Goddard Institute for Space Studies, New York, NY, United States

<sup>f</sup>Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD, USA

<sup>g</sup>Ecosystem Services and Management Program, International Institute for Applied Systems Analysis, Laxenburg, Austria

<sup>h</sup>Department of Soil Science, Faculty of Natural Sciences, Comenius University in Bratislava, Bratislava, Slovak Republic

<sup>i</sup>Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, 91191 Gif-sur-Yvette, France

<sup>j</sup>Sino-French Institute of Earth System Sciences, College of Urban and Environmental Sciences, Peking University, Beijing, China

<sup>k</sup>Unité de Modélisation du Climat et des Cycles Biogéochimiques, UR SPHERES, Institut d'Astrophysique et de Géophysique, University of Liège, Belgium

<sup>l</sup>Met Office Hadley Centre, Exeter, United Kingdom

<sup>m</sup>Department of Geography, Ludwig-Maximilians-Universität, Munich, Germany

<sup>n</sup>Georg-August-University Göttingen, Tropical Plant Production and Agricultural Systems Modelling, Göttingen, Germany

<sup>o</sup>Department of Geographical Sciences, University of Maryland, College Park, MD, USA

<sup>p</sup>Texas AgriLife Research and Extension, Texas A&M University, Temple, TX, USA

<sup>q</sup>Department of Statistics, University of Chicago, Chicago, IL, USA

<sup>r</sup>EAWAG, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

<sup>s</sup>Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

<sup>t</sup>Earth Institute Center for Climate Systems Research, Columbia University, New York, NY, USA

<sup>u</sup>Karlsruhe Institute of Technology, IMK-IFU, 82467 Garmisch-Partenkirchen, Germany.

<sup>v</sup>School of Geography, Earth and Environmental Science, University of Birmingham, Birmingham, UK.

## Abstract

Concerns about food security under climate change have motivated efforts to better understand the future changes in yields by using detailed process-based models in agronomic sciences. Process-based crop models differ on many details affecting yields and considerable uncertainty remains in future yield projections. Phase II of the Global Gridded Crop Model Intercomparison (GGCMI), an activity of the Agricultural Model Intercomparison and Improvement Project (AgMIP), consists of a large simulation set with perturbations in atmospheric CO<sub>2</sub> concentrations, temperature, precipitation, and applied nitrogen inputs and constitutes a data-rich basis of projected yield changes across twelve models and five crops (maize, soy, rice, spring wheat, and winter wheat) using global gridded simulations. In this paper we present the simulation output database from Phase II of the GGCMI effort, a targeted experiment aimed at understanding the sensitivity to and interaction between multiple climate variables (as well as management) on yields, and illustrate some initial summary results from the model intercomparison project. We also present the construction of a simple “emulator or statistical representation of the simulated 30-year mean climatological output in each location for each crop and model. The emulator captures the mean-climatological response of the process-based models in a lightweight, computationally tractable form that facilitates model comparison as well as potential applications in subsequent modeling efforts such as integrated assessment.

**Keywords:** climate change, food security, model emulation, AgMIP, crop model

## 1. Introduction

2      Understanding crop yield response to a changing climate  
3      is critically important, especially as the global food produc-  
4      tion system will face pressure from increased demand over the  
5      next century. Climate-related reductions in supply could there-  
6      fore have severe socioeconomic consequences. Multiple stud-  
7      ies using different crop or climate models concur in predicting  
8      sharp yield reductions on currently cultivated cropland under  
9      business-as-usual climate scenarios, although their yield pro-  
10     jections show considerable spread (e.g. Porter et al. (IPCC),  
11     2014, Rosenzweig et al., 2014, Schauberger et al., 2017, and  
12     references therein). Modeling crop responses continues to be  
13     challenging, as crop growth is a function of complex interac-  
14     tions between climate inputs and management practices. Inter-  
15     comparison projects targeting model responses to important  
16     drivers are critical to improve future projections.

17     Computational models have been used to project crop yields  
18     since the 1950's, beginning with statistical models that attempt  
19     to capture the relationship between input factors and resultant  
20     yields (e.g. Heady, 1957, Heady & Dillon, 1961). These statisti-  
21     cal models were typically developed on a small scale for loca-  
22     tions with extensive histories of yield data. The emergence of  
23     electronic computers allowed development of numerical mod-  
24     els that simulate the process of photosynthesis and the biology  
25     and phenology of individual crops (first proposed by de Wit  
26     (1957) and Duncan et al. (1967) and attempted by Duncan  
27     (1972); for a history of crop model development see Rosen-  
28     zweig et al. (2014)). A half-century of improvement in both  
29     models and computing resources means that researchers can  
30     now run crop simulations for many years at high spatial res-  
31     olution on the global scale.

32     Both types of models continue to be used, and compara-

33     tive studies have concluded that when done carefully, both ap-  
34     proaches can provide similar yield estimates (e.g. Lobell &  
35     Burke, 2010, Moore et al., 2017, Roberts et al., 2017, Zhao  
36     et al., 2017). Models tend to agree broadly in major response  
37     patterns, including a reasonable representation of the spatial  
38     pattern in historical yields of major crops (e.g. Elliott et al.,  
39     2015, Müller et al., 2017) and projections of decreases in yield  
40     under future climate scenarios.

41     Process-based models do continue to struggle with some impor-  
42     tant details, including reproducing historical year-to-year  
43     variability (e.g. Müller et al., 2017), reproducing historical  
44     yields when driven by reanalysis weather (e.g. Glotter et al.,  
45     2014), and low sensitivity to extreme events (e.g. Glotter et al.,  
46     2015). These issues are driven in part by the diversity of new  
47     cultivars and genetic variants, which outstrips the ability of aca-  
48     demic modeling groups to capture them (e.g. Jones et al., 2017).  
49     Models also do not simulate many additional factors affecting  
50     production, including pests, diseases, and weeds. For these rea-  
51     sons, individual studies must generally re-calibrate models to  
52     ensure that short-term predictions reflect current cultivar mixes,  
53     and long-term projections retain considerable uncertainty (Wolf  
54     & Oijen, 2002, Jagtap & Jones, 2002, Iizumi et al., 2010, An-  
55     gulo et al., 2013, Asseng et al., 2013, 2015). Inter-model dis-  
56     crepancies can also be high in areas not yet cultivated (e.g.  
57     Challinor et al., 2014, White et al., 2011). Finally, process-  
58     based models present additional difficulties for high-resolution  
59     global studies because of their complexity and computational  
60     requirements. For economic impacts assessments, it is often  
61     impossible to integrate a set of process-based crop models di-  
62     rectly into an integrated assessment model to estimate the po-  
63     tential cost of climate change to the agricultural sector.

64     Nevertheless, process-based models are necessary for under-  
65     standing the global future yield impacts of climate change for  
66     many reasons. First, cultivation may shift to new areas, where

\*Corresponding author at: 4734 S Ellis, Chicago, IL 60637, United States.  
email: jfranke@uchicago.edu

no yield data are currently available and therefore statistical models cannot apply. Yield data are also often limited in the developing world, where future climate impacts may be the most critical. Finally, only process-based models can capture the growth response to novel conditions and practices that are not represented in historical data (e.g. Pugh et al., 2016, Roberts et al., 2017). These novel changes can include the direct fertilization effect of elevated CO<sub>2</sub>, and changes in management practices that may ameliorate climate-induced damages.

Interest has been rising in statistical emulation, which allows combining advantageous features of both statistical and process-based models. The approach involves constructing a statistical representation or “surrogate model” of complicated numerical simulations by using simulation output as the training data for a statistical model (e.g. O’Hagan, 2006, Conti et al., 2009). Emulation is particularly useful in cases where simulations are complex and output data volumes are large, and has been used in a variety of fields, including hydrology (e.g. Razavi et al., 2012), engineering (e.g. Storlie et al., 2009), environmental sciences (e.g. Ratto et al., 2012), and climate (e.g. Castruccio et al., 2014, Holden et al., 2014). For agricultural impacts studies, emulation of process-based models allows capturing key relationships between input variables in a lightweight, flexible form that is compatible with economic studies.

In the past decade, multiple studies have developed emulators of process-based crop simulations. Early studies proposing or describing potential crop yield emulators include Howden & Crimp (2005), Räisänen & Ruokolainen (2006), Lobell & Burke (2010), and Ferrise et al. (2011), who used a machine learning approach to predict Mediterranean wheat yields. Studies developing single-model emulators include Holzkämper et al. (2012) for the CropSyst model, Ruane et al. (2013) for the CERES wheat model, and Oyebamiji et al. (2015) for the




Figure 1: Presently cultivated area for rain-fed crops. Blue indicates grid cells with more than 20,000 hectares (~10% of the equatorial grid cells) of crop cultivated. Gray contour shows area with more than 10 hectares cultivated. Cultivated areas for maize, rice, and soy are taken from the MIRCA2000 (“monthly irrigated and rain-fed crop areas around the year 2000”) dataset (Portmann et al., 2010). Areas for winter and spring wheat areas are adapted from MIRCA2000 data and sorted by growing season. For analogous figure of irrigated crops, see Figure S1.

LPJmL model (for multiple crops, using multiple scenarios as a training set). More recently, emulators have begun to be used in the context of multi-model intercomparisons, with Blanc & Sultan (2015), Blanc (2017), Ostberg et al. (2018) and Misra et al. (2017) using them to analyze the five crop models of the Inter-Sectoral Impacts Model Intercomparison Project

(ISIMIP) (Warszawski et al., 2014), which simulated yields for maize, soy, wheat, and rice. Choices differ: Blanc & Sultan (2015) and Blanc (2017) base their emulation on historical simulations and a single future climate/emissions scenario (RCP8.5), and use local weather variables and yields in their regression but then aggregate across broad regions; Ostberg et al. (2018) consider multiple future climate scenarios, using global mean temperature change (and CO<sub>2</sub>) as regressors but then pattern-scale to emulate local yields; while Mistry et al. (2017) compare emulated and observed historical yields, using local weather data and a historical crop simulation. These efforts do share important common features: all emulate annual crop yields across the entire scenario or scenarios, and when future scenarios are considered, they are non-stationary, i.e. their input climate parameters evolve over time.

An alternative approach is to construct a training set of multiple stationary scenarios in which parameters are systematically varied. Such a “parameter sweep” offers several advantages for emulation over scenarios in which climate evolves over time. First, it allows separating the effects of different variables that impact yields but that are highly correlated in realistic future scenarios (e.g. CO<sub>2</sub> and temperature). Second, it allows making a distinction between year-over-year yield variations and climatological changes, which may involve different responses to the particular climate regressors used (e.g. Ruane et al., 2016). For example, if year-over-year yield variations are driven predominantly by variations in the distribution of temperatures throughout the growing season, and long-term climate changes are driven predominantly by shifts in means, then regressing on the mean growing season temperature will produce different yield responses at annual vs. climatological timescales.

Systematic parameter sweeps have begun to be used in crop model evaluation and emulation, with early efforts in 2015 (Makowski et al., 2015, Pirttioja et al., 2015), and several re-

cent studies in 2018 (Fronzek et al., 2018, Snyder et al., 2018, Ruiz-Ramos et al., 2018). All three studies sample multiple perturbations to temperature and precipitation (with Snyder et al. (2018) and Ruiz-Ramos et al. (2018) adding CO<sub>2</sub> as well), in 132, 99 and 220 different combinations, respectively, and take advantage of the structured training set to construct emulators (“response surfaces”) of climatological mean yields, omitting year-over-year variations. All are limited in some respects and focus on a limited number of sites. Fronzek et al. (2018) and Ruiz-Ramos et al. (2018) simulate only wheat (over many models) and Snyder et al. (2018) analyzes four crops (maize, wheat, rice, soy) for agricultural impacts experiments with the GCAM (Calvin et al., 2019) model.

In this paper we describe a new comprehensive dataset designed to expand the parameter sweep approach still further. The Global Gridded Crop Model Intercomparison (GGCMI) Phase II experiment involves running a suite of process-based crop models across historical conditions perturbed by a set of discrete steps in different input parameters, including an applied nitrogen dimension. The experimental protocol involves over 700 different parameter combinations for each model and crop, with simulations providing near-global coverage at a half degree spatial resolution. The experiment was conducted as part of the Agricultural Model Intercomparison and Improvement Project (AgMIP) (Rosenzweig et al., 2013, 2014), an international effort conducted under a framework similar to the Climate Model Intercomparison Project (CMIP) (Taylor et al., 2012, Eyring et al., 2016). The GGCMI protocol builds on the AgMIP Coordinated Climate-Crop Modeling Project (C3MP) (Ruane et al., 2014, McDermid et al., 2015) and will contribute to the AgMIP Coordinated Global and Regional Assessments (CGRA) (Ruane et al., 2018, Rosenzweig et al., 2018). GGCMI Phase II is designed to allow addressing goals such as understanding where highest-yield regions may shift

under climate change; exploring future adaptive management  
 176 strategies; understanding how interacting input drivers affect  
 crop yield; quantifying uncertainties across models and major  
 178 drivers; and testing strategies for producing lightweight em-  
 ulators of process-based models. In this paper, we describe  
 180 the GGCMI Phase II experiments, present initial results, and  
 demonstrate that it is tractable to emulation.

## 182 2. Simulation – Methods

GGCMI Phase II is the continuation of a multi-model com-  
 184 parison exercise begun in 2014. The initial Phase I compared  
 harmonized yields of 21 models for 19 crops over a 31-year  
 186 historical (1980-2010) scenario with a primary goal of model  
 evaluation (Elliott et al., 2015, Müller et al., 2017). Phase II  
 188 compares simulations of 12 models for 5 crops (maize, rice,  
 soybean, spring wheat, and winter wheat) over the same histor-  
 190 ical time series (1980-2010) used in Phase I, but with individ-  
 ual climate or management inputs adjusted from their historical  
 192 values. The reduced set of crops includes the three major global  
 cereals and the major legume and accounts for over 50% of hu-  
 194 man calories (in 2016, nearly 3.5 billion tons or 32% of total  
 global crop production by weight (Food and Agriculture Orga-  
 196 nization of the United Nations, 2018).

The guiding scientific rationale of GGCMI Phase II is to pro-  
 198 vide a comprehensive, systematic evaluation of the response  
 of process-based crop models to different values for carbon  
 200 dioxide, temperature, water, and applied nitrogen (collectively

known as “CTWN”). The dataset is designed to allow re-  
 202 searchers to:

- Enhance understanding of how models work by character-  
 204 izing their sensitivity to input climate and nitrogen drivers.
- Study the interactions between climate variables and nitro-  
 206 gen inputs in driving modeled yield impacts.
- Explore differences in crop response to warming across the  
 208 Earth’s climate regions.
- Provide a dataset that allows statistical emulation of crop  
 210 model responses for downstream modelers.
- Illustrate differences in potential adaptation via growing  
 212 season changes.

The experimental protocol consists of 9 levels for precipita-  
 214 tion perturbations, 7 for temperature, 4 for CO<sub>2</sub>, and 3 for ap-  
 plied nitrogen, for a total of 672 simulations for rain-fed agri-  
 216 culture and an additional 84 for irrigated (Table 1). For irri-  
 gated simulations, soil water is held at either field capacity or,  
 for those models that include water-log damage, at maximum  
 218 beneficial level. Temperature perturbations are applied as ab-  
 solute offsets from the daily mean, minimum, and maximum  
 220 temperature time series for each grid cell used as inputs. Pre-  
 cipitation perturbations are applied as fractional changes at the  
 222 grid cell level, and carbon dioxide and nitrogen levels are spec-  
 ified as discrete values applied uniformly over all grid cells.  
 Note that CO<sub>2</sub> changes are applied independently of changes  
 224 in climate variables, so that higher CO<sub>2</sub> is not associated with  
 higher temperatures. An additional, identical set of scenarios

Input variable	Abbr.	Tested range	Unit
CO <sub>2</sub>	C	360, 510, 660, 810	ppm
Temperature	T	-1, 0, 1, 2, 3, 4, 5*, 6	°C
Precipitation	W	-50, -30, -20, -10, 0, 10, 20, 30, (and W <sub>inf</sub> )	%
Applied nitrogen	N	10, 60, 200	kg ha <sup>-1</sup>

Table 1: GGCMI Phase II input variable test levels. Temperature and precipitation values indicate the perturbations from the historical, climatology. \* Only simulated by one model. W-percentage does not apply to the irrigated (W<sub>inf</sub>) simulations, which are all simulated at the maximum beneficial levels of water.

Model (Key Citations)	Maize	Soy	Rice	Winter Wheat	Spring Wheat	N Dim.	Simulations per Crop
<b>APSIM-UGOE</b> , Keating et al. (2003), Holzworth et al. (2014)	X	X	X	-	X	Yes	37
<b>CARAIB</b> , Dury et al. (2011), Pirttioja et al. (2015)	X	X	X	X	X	No	224
<b>EPIC-IIASA</b> , Balkovi et al. (2014)	X	X	X	X	X	Yes	35
<b>EPIC-TAMU</b> , Izaurrealde et al. (2006)	X	X	X	X	X	Yes	672
<b>JULES*</b> , Osborne et al. (2015), Williams & Falloon (2015), Williams et al. (2017)	X	X	X	-	X	No	224
<b>GEPIC</b> , Liu et al. (2007), Folberth et al. (2012)	X	X	X	X	X	Yes	384
<b>LPJ-GUESS</b> , Lindeskog et al. (2013), Olin et al. (2015)	X	-	-	X	X	Yes	672
<b>LPJmL</b> , von Bloh et al. (2018)	X	X	X	X	X	Yes	672
<b>ORCHIDEE-crop</b> , Valade et al. (2014)	X	-	X	-	X	Yes	33
<b>pDSSAT</b> , Elliott et al. (2014), Jones et al. (2003)	X	X	X	X	X	Yes	672
<b>PEPIC</b> , Liu et al. (2016a,b)	X	X	X	X	X	Yes	130
<b>PROMET*†</b> , Mauser & Bach (2015), Hank et al. (2015), Mauser et al. (2009)	X	X	X	X	X	Yes†	239
Totals	12	10	11	9	12	-	3993 (maize)

Table 2: Models included in GGCMI Phase II and the number of C, T, W, and N simulations that each performs for rain-fed crops (“Sims per Crop”), with 672 as the maximum. “N-Dim.” indicates whether the simulations include varying nitrogen levels. Two models provide only one nitrogen level, and †PROMET provides only two of the three nitrogen levels (and so is not emulated across the nitrogen dimension). All models provide the same set of simulations across all modeled crops, but some omit individual crops. (For example, APSIM does not simulate winter wheat.) Irrigated simulations are provided at the level of the other covariates for each model, i.e. an additional 84 simulations for fully-sampled models. Simulations are nearly global but their geographic extent can vary, even for different crops in an individual model, since some simulations omit regions far outside the currently cultivated area. In most cases, historical daily climate inputs are taken from the 0.5 degree NASA AgMERRA daily gridded re-analysis product specifically designed for agricultural modeling, with satellite-corrected precipitation (Ruane et al., 2015), but two models (marked with \*) require sub-daily input data and use alternative sources. See Elliott et al. (2015) for additional details.

228 (at the same C, T, W, and N levels) not shown or analyzed here  
229 simulate adaptive agronomy under climate change by varying  
230 the growing season for crop production. The resulting GGCMI  
231 Phase II dataset captures a distribution of crop responses over  
232 the potential space of future climate conditions.

The 12 models included in GGCMI Phase II are all mechanistic process-based crop models that are widely used in impacts assessments (Table 2). Although some models share a common base (e.g. the LPJ family or the EPIC family of models), they have subsequently developed independently. (For more details on model genealogy, see Figure S1 in Rosenzweig et al. (2014).) Differences in model structure mean that several key factors are not standardized across the experiment, including secondary soil nutrients, carry-over effects across growing years including residue management and soil moisture, and the extent of simulated area for different crops. Growing seasons are standardized across models (with assumptions based on Sacks et al. (2010) and Portmann et al. (2008, 2010)), but

vary by crop and by location on the globe. For example, maize is sown in March in Spain, in July in Indonesia, and in December in Namibia. All stresses are disabled other than factors related to nitrogen, temperature, and water (e.g. alkalinity and salinity). No additional nitrogen inputs, such as atmospheric deposition, are considered, but some model treatments of soil organic matter may allow additional nitrogen release through mineralization. See Rosenzweig et al. (2014), Elliott et al. (2015) and Müller et al. (2017) for further details on models and underlying assumptions.

The participating modeling groups provide simulations at any of four initially specified levels of participation, so the number of simulations varies by model, with some sampling only a part of the experiment variable space. Most modeling groups simulate all five crops in the protocol, but some omitted one or more. Table 2 provides details of coverage for each model. Note that the three models that provide less than 50 simulations are excluded from the emulator analysis.

264 Each model is run at 0.5 degree spatial resolution and cov-  
 265 ers all currently cultivated areas and much of the uncultivated  
 266 land area. (See Figure 1 for the present-day cultivated area of  
 267 rain-fed crops, and Figure S1 in the Supplemental Material for  
 268 irrigated crops.) Coverage extends considerably outside cur-  
 269 rently cultivated areas because cultivation will likely shift under  
 270 climate change. However, areas are not simulated if they are  
 271 assumed to remain non-arable even under an extreme climate  
 272 change; these regions include Greenland, far-northern Canada,  
 273 Siberia, Antarctica, the Gobi and Sahara Deserts, and central  
 274 Australia.


275 All models produce as output crop yields (tons  $\text{ha}^{-1}$  year $^{-1}$ )  
 276 for each 0.5 degree grid cell. Because both yields and yield  
 277 changes vary substantially across models and across grid cells,  
 278 we primarily analyze relative change from a baseline. We take  
 as the baseline the scenario with historical climatology (i.e. T

280 and P changes of 0), C of 360 ppm, and applied N at 200 kg  
 281  $\text{ha}^{-1}$ . We show absolute yields in some cases to illustrate geo-  
 282 graphic differences in yields.

283 The GGCMI Phase II simulations are designed for evalua-  
 284 ting changes in yield but not absolute yields, since they omit  
 285 detailed calibrations. To provide some validation of the skill of  
 286 the process-based models used, we repeat the validation exer-  
 287 cises of Müller et al. (2017) for GGCMI Phase I. See Appendix  
 288 A for details on simulation model validation.

### 3. Simulation – Results

289 Crop models in the GGCMI Phase II ensemble show broadly  
 290 consistent responses to climate and management perturbations  
 291 in most regions, with a strong negative impact of increased tem-  
 292 perature in all but the coldest regions. We illustrate this re-  
 293 sult for rain-fed maize in Figure 2, which shows yields across



294 Figure 2: Illustration of the distribution of regional yield changes across the multi-model ensemble, split by Köppen-Geiger climate regions (Rubel & Kottek, 2010). Note that ‘Equatorial’ and ‘Snow’ regions are sometimes referred to as ‘tropical’ and ‘cold-continental’ respectively. We show responses of a single crop (rain-fed maize) to applied uniform temperature perturbations, for three discrete precipitation perturbation levels (rain-fed (rf) -20%, rain-fed (0), and rain-fed +20%), with CO<sub>2</sub> and nitrogen held constant at baseline values (360 ppm and 200 kg  $\text{ha}^{-1}$   $\text{yr}^{-1}$ ). Y-axis is fractional change in the regional average climatological potential yield relative to the baseline. The figure shows all modeled land area; see Figure S6 in the supplemental material for only currently-cultivated land. Panel text gives the percentage of rain-fed maize presently cultivated in each climate zone (data from Portmann et al., 2010). Box-and-whiskers plots show distribution across models, with median marked. Box edges are first and third quartiles, i.e. box height is the interquartile range (IQR). Whiskers extend to maximum and minimum of simulations but are limited at 1.5·IQR; otherwise the outlier is shown. Models generally agree in most climate regions (other than cold continental), with projected changes larger than inter-model variance. Outliers in the tropics (strong negative impact of temperature increases) are the pDSSAT model; outliers in the high-rainfall case (strong positive impact of precipitation increases) are the JULES model. Inter-model variance increases in the case where precipitation is reduced, suggesting uncertainty in model response to water limitation. The right panel (Global) shows yield responses to an globally uniform temperature shift; note that these results are not directly comparable to simulations of more realistic climate scenarios with the same global mean change.




Figure 3: Illustration of the spatial pattern of potential yields and potential yield changes in the GGCMI Phase II ensemble, for three major crops. Left column (a) shows multi-model mean climatological yields for the baseline scenario for (top–bottom) rain-fed maize, soy, and rice. (For wheat see Figure S11 in the supplemental material.) White stippling indicates areas where these crops are not currently cultivated. Absence of cultivation aligns well with the lowest yield contour ( $0\text{--}2 \text{ ton ha}^{-1}$ ). Right column (b) shows the multi-model mean fractional yield change in the extreme  $T + 4 \text{ }^{\circ}\text{C}$  scenario (with other inputs at baseline values). Areas without hatching or stippling are those where confidence in projections is high: the multi-model mean fractional change exceeds two standard deviations of the ensemble. ( $\Delta > 2\sigma$ ). Hatching indicates areas of low confidence ( $\Delta < 1\sigma$ ), and stippling areas of medium confidence ( $1\sigma < \Delta < 2\sigma$ ). Crop model results in cold areas, where yield impacts are on average positive, also have the highest uncertainty.

all grid cells for the primary Köppen-Geiger climate regions  
296 (Rubel & Kottek, 2010). In warming scenarios with precipi-  
tation held constant, all models show decreases in maize yield  
298 in the ‘warm temperate’, ‘equatorial’ (tropical), and arid regions  
that account for nearly three-quarters of global maize produc-  
300 tion. These impacts are robust for even moderate climate per-  
turbations. In the ‘warm temperate’ zone, even a 1 degree tem-  
perature rise with other variables held fixed leads to a median  
302 yield reduction that outweighs the variance across models. A  
6 degree temperature rise results in median loss of ~25% of  
yields with a signal to noise ratio of nearly three to one. A no-  
304 table exception is the ‘snow’ (‘cold-continental’) region, where  
models disagree strongly, extending even to the sign of impacts.  
306

Other crops show similar responses to warming, with robust  
308 yield losses in warmer locations and high inter-model variance  
in the cold continental regions (Figure S7).  
310

The effects of rainfall changes on maize yields shown in Figure 2 are also as expected and are consistent across models.  
312 Increased rainfall mitigates the negative effect of higher tem-  
peratures by counteracting the increased evapo-transpiration to  
some degree, most strongly in arid regions. Decreased rain-  
fall amplifies yield losses and also increases inter-model vari-  
ance more strongly, suggesting that models have difficulty repre-  
314 senting crop response to water stress or increased evapo-  
transpiration due to warmer temperatures. We show only rain-  
fed maize here; see Figure S5 for the irrigated case. As ex-  
316

pected, irrigated crops are more resilient to temperature increases in all regions, especially so where water is limiting.

Mapping the distribution of baseline yields and yield changes shows the geographic dependencies that underlie these results. Figure 3 shows baseline and changes in the T+4 scenario for rain-fed maize, soy, and rice in the multi-model ensemble mean, with locations of model agreement marked. Absolute yield potentials show strong spatial variation, with much of the Earth’s surface area unsuitable for any of these crops. In general, models agree most on yield response in regions where yield potentials are currently high and therefore where crops are currently grown. Models show robust decreases in yields at low latitudes, and highly uncertain median increases at most high latitudes. For wheat crops see Figure S11; wheat projections are more uncertain, possible because calibration is especially important for wheat (e.g. Asseng et al., 2013).

#### 4. Emulation – Methods

As part of our demonstration of the properties of the GGCMI Phase II dataset, we construct an emulator of 30-year climatological mean yields. This approach is made possible by the structured set of simulations involving systematic perturbations. In the GGCMI Phase II dataset, the year-over-year responses are generally quantitatively distinct from (and larger than) climatological mean responses. In the example of Figure 4, responses to year-over-year temperature variations are 100% larger than those to long-term perturbations in the baseline case, and larger still under warmer conditions, rising to nearly 200% more in the T+6 case. The stronger year-over-year response under warmer conditions also manifests as a wider distribution of yields (Figure 5). As discussed previously, year-over-year and climatological responses can differ for many reasons including memory in the crop model, lurking covariants, and differing associated distributions of daily growing-season daily




Figure 4: Example showing distinction between crop yield responses to year-to-year and climatological mean temperature shifts. Figure shows irrigated maize for a representative high-yield region (nine adjacent grid cells in northern Iowa) from the pDSSAT model, for the baseline 1981-2010 historical climate (blue) and for the scenario of maximum temperature change (+6 K, red). Other variables are held at baseline values, and the choice of irrigated yields means that precipitation is not a factor. Open black circles mark climatological mean yield values for all six temperature scenarios ( $T-1, +0, +1, +2, +3, +4, +6$ ). Colored lines show total least squares linear regressions of year-over-year variations in each scenario. Black line shows the fit through the climatological mean values. Responses to year-over-year temperature variations (colored lines) are 100–200% larger than those to long-term climate perturbations, rising under warmer conditions.

weather (e.g. Ruane et al., 2016). Note that the GGCMI Phase II datasets do not capture one climatological factor, potential future distributional shifts, because all simulations are run with fixed offsets from the historical climatology. Prior work has suggested that mean changes are the dominant drivers of climatological crop yield shifts in non-arid regions (e.g. Glotter et al., 2014).

Emulation involves fitting individual regression models for each crop, simulation model, and 0.5 degree geographic pixel from the GGCMI Phase II dataset; the regressors are the applied constant perturbations in  $\text{CO}_2$ , temperature, water, and nitrogen (C, T, W, N). We regress 30-year climatological mean yields against a third-order polynomial in C, T, W, and N with interaction terms. (We aggregate the entire 30-year run in each case to




Figure 5: Example showing climatological mean yields and distribution of yearly yields for three 30-year scenarios. Figure shows irrigated maize for nine adjacent high-yield grid cells of Figure 4 from the pDSSAT model, for the baseline 1981-2010 historical climate (blue) and for scenarios with temperature shifted by T+3 (orange) and T+6 K (red), with other variables held at baseline values. The stronger year-over-year temperature response with higher temperatures seen in Figure 4 is manifested here as larger variance in annual yields even though the variance in climate drivers is identical. In this work we emulate not the year-over-year distributions but the climatological mean response (dashed vertical lines).

improve signal to noise ration in our model.) The higher-order terms are necessary to capture any nonlinear responses, which are well-documented in observations for temperature and water perturbations (e.g. Schlenker & Roberts (2009) for T and He et al. (2016) for W). We include interaction terms (both linear and higher-order) because past studies have shown them to be significant effects. For example, Lobell & Field (2007) and Tebaldi & Lobell (2008) showed that in real-world yields, the joint distribution in T and W is needed to explain observed yield variance. (C and N are fixed in these data.) Other observation-based studies have shown the importance of the interaction between water and nitrogen (e.g. Aulakh & Malhi, 2005), and between nitrogen and carbon dioxide (Osaki et al., 1992, Nakamura et al., 1997). To avoid overfitting or unstable parameter estimation, we apply a feature selection procedure (described below) that reduces the potential 34-term polynomial (for the

rain-fed case) to 23 terms.

We do not focus on comparing different functional forms in this study, and instead choose a relatively simple parametrization that allows for some interpretation of coefficients. Some prior studies have used more complex functional forms and larger numbers of parameters, e.g. 39 in Blanc & Sultan (2015) and Blanc (2017), who borrow information across space by fitting grid points simultaneously across a large region in a panel regression. The simple functional form used here allows emulation at the grid cell level. The emulation therefore indirectly includes any yield response to geographically distributed factors such as soil type, insolation, and the baseline climate itself. We hold the statistical specification constant across all crops and models to facilitate parameter by parameter simulation model comparison.

#### 4.1. Feature selection procedure

Although the GGCMI Phase II sampled variable space is large, it is still sufficiently limited that use of the full polynomial expression described above can be problematic. We therefore reduce the number of terms through a feature selection cross-validation process in which terms in the polynomial are tested for importance. In this procedure higher-order and interaction terms are added successively to the regression model; we then follow the reduction of the aggregate mean squared error with increasing terms and eliminate those terms that do not contribute significant reductions. See supplemental documents for more details. We select terms by applying the feature selection process to the three models that provided the complete set of 672 rain-fed simulations (pDSSAT, EPIC-TAMU, and LPJmL); the resulting choice of terms is then applied for all emulators.

Feature importance is remarkably consistent across all three models and across all crops (see Figure S4 in the supplemental material). The feature selection process results in a final polynomial in 23 terms, with 11 terms eliminated. We omit the  $N^3$

term, which cannot be fitted because we sample only three nitrogen levels. We eliminate many of the C terms: the cubic, the CT, CTN, and CWN interaction terms, and all higher order interaction terms in C. Finally, we eliminate two 2nd-order interaction terms in T and one in W. Implication of this choice include that nitrogen interactions are complex and important, and that water interaction effects are more nonlinear than those in temperature. The resulting statistical model (Equation 1) is used for all grid cells, models, and rain-fed crops. (The regressions for irrigated crops do not contain the W terms and the models that do not sample the nitrogen levels omit the N terms).

$$Y = K_1 \quad (1)$$

$$\begin{aligned} &+ K_2 C + K_3 T + K_4 W + K_5 N \\ &+ K_6 C^2 + K_7 T^2 + K_8 W^2 + K_9 N^2 \\ &+ K_{10} CW + K_{11} CN + K_{12} TW + K_{13} TN + K_{14} WN \\ &+ K_{15} T^3 + K_{16} W^3 + K_{17} TWN \\ &+ K_{18} T^2 W + K_{19} W^2 T + K_{20} W^2 N \\ &+ K_{21} N^2 C + K_{22} N^2 T + K_{23} N^2 W \end{aligned}$$

To fit the parameters  $K$ , we use a Bayesian Ridge probabilistic estimator (MacKay, 1991), which reduces volatility in parameter estimates when the sampling is sparse, by weighting parameter estimates towards zero. The Bayesian Ridge method is necessary to maintain a consistent functional form across all models and locations. We use the implementation of the Bayesian Ridge estimator from the scikit-learn package in Python (Pedregosa et al., 2011). In the GGCMI Phase II experiment, the most problematic fits are those for models that provided a limited number of cases or for low-yield geographic regions where some modeling groups did not run all scenarios. We do not attempt to emulate models that provided less than 50 simulations. The lowest number of simulations emulated across the full parameter space is then 130 (for the PEPIC

model). The resulting parameter matrices for all crop model emulators are available on request [give location?](#), as are the raw simulation data and a Python application to emulate yields. The yield output for a single GGCMI Phase II model that simulates all scenarios and all five crops is  $\sim 12.5$  GB; the emulator is  $\sim 100$  MB, a reduction by over two orders of magnitude.

## 5. Emulation – Results

Emulation provides not only a computational tool but a means of understanding and interpreting crop yield response across the parameter space. Emulation is only possible when crop yield responses are sufficiently smooth and continuous to allow fitting with a relatively simple functional form, but this condition largely holds in the GGCMI Phase II simulations. Responses are quite diverse across locations, crops, and models, but in most cases local responses are regular enough to permit emulation. We show illustrations of emulation fidelity in this section; for more detailed discussion see Appendix B.

Crop yield responses are geographically diverse, even in high-yield and high-cultivation areas. Figure 6 illustrates geographic diversity for a single crop and model (rain-fed maize in pDSSAT); this heterogeneity supports the choice of emulating at the grid cell level. Each panel in Figure 6 shows simulated yield output from scenarios varying only along a single dimension ( $\text{CO}_2$ , temperature, precipitation, or nitrogen addition), with other inputs held fixed at baseline levels, compared to the full 4D emulation across the parameter space. Yields evolve smoothly across the space sampled, and the polynomial fit captures the climatological response to perturbations. Crop yield responses generally follow similar functional forms across models, though with a large spread in magnitude likely due to the lack of calibration. Figure 7 illustrates inter-model diversity for a single crop and location (rain-fed maize in northern Iowa, also shown in Figure 6). Differences in response shape can lead




Figure 6: Illustration of spatial variations in yield response and emulation ability. We show rain-fed maize in the pDSSAT model in six example locations selected to represent high-cultivation areas around the globe. Legend includes hectares cultivated in each selected grid cell. Each panel shows variation along a single variable, with others held at baseline values. Dots show climatological mean yields and lines the results of the full 4D emulator of Equation 1. In general the climatological response surface is sufficiently smooth that it can be represented with the sampled variable space by the simple polynomial used in this work. Extrapolation can however produce misleading results. Nitrogen fits may not be realistic at intermediate values given limited sampling. For more detailed emulator assessment, see Appendix B.

to differences in the fidelity of emulation, though comparison  
here is complicated by the different sampling regimes across  
models. Note that models are most similar in their responses to  
temperature perturbations.

While the nitrogen dimension is important, it is also the most

problematic to emulate in this work because of its limited sampling. The GGCMI Phase II protocol specified only three nitrogen levels (10, 60 and 200 kg N y<sup>-1</sup> ha<sup>-1</sup>), so a third-order fit would be over-determined but a second-order fit can result in potentially unphysical results. Steep and nonlinear declines




Figure 7: Illustration of across-model variations in yield response. Figures shows simulations and emulations from six models for rain-fed maize in the same Iowa grid cell shown in Figure 6, with the same plot conventions. Models that do not simulate the nitrogen dimension are omitted for clarity. Note that models are uncalibrated, increasing spread in absolute yields. While most model responses can readily emulated with a simple polynomial, some response surfaces are more complicated (e.g. LPJ-GUESS here) and lead to emulation error, though error generally remains small relative to inter-model uncertainty. For more detailed emulator assessment, see Appendix A. As in Figure 6, extrapolation out of the sample space is problematic.




Figure 8: Global emulated damages for maize on currently cultivated lands for the GGCMI Phase II models emulated, for uniform temperature shifts with other inputs held at baseline. (The damage function is created from aggregating up emulated values at the grid cell level, not from a regression of global mean yields.) Lines are emulations for rain-fed (solid) and irrigated (dashed) crops; for comparison, dots are the simulated values for the rain-fed case. For most models, irrigated crops show a sharper reduction than do rain-fed because of the locations of cultivated areas: irrigated crops tend to be grown in warmer areas where impacts are more severe for a given temperature shift. (The exceptions are PROMET, JULES, and LPJmL.) For other crops and scenarios see Figures S16- S19 in the supplemental material.

in yield with lower nitrogen levels mean that some regressions imply a peak in yield between the 100 and 200 kg N  $y^{-1} ha^{-1}$  levels. While it is possible that over-application of nitrogen at the wrong time in the growing season could lead to reduced yields, these features are potentially an artifact of under sampling. In addition, the polynomial fit cannot capture the well-documented saturation effect of nitrogen application (e.g. Ingestad, 1977) as accurately as would be possible with a non-parametric model.

The emulation fidelity demonstrated here is sufficient to allow using emulated response surfaces to compare model responses and derive insight about impacts projections. Because the emulator or “surrogate model” transforms the discrete sim-

ulation sample space into a continuous response surface at any geographic scale, it can be used for a variety of applications, including construction of continuous damage functions. As an example, we show a damage function constructed from the 4D emulation, aggregated to global yield, with simulated values shown for comparison (Figure 8, which shows maize on currently cultivated land; see Figures S16- S19 for other crops and dimensions). The emulated values closely match simulations even at this aggregation level. Note that these functions are presented only as examples and do not represent true global projections, because they are developed from simulation data with a uniform temperature shift while increases in global mean temperature should manifest non-uniformly. The global coverage of the GGCMI Phase II simulations allows impacts modelers to apply arbitrary geographically-varying climate projections, as well as arbitrary aggregation masks, to develop damage functions for any climate scenario and any geopolitical or geographic level.

## 6. Conclusions and Discussion

The GGCMI Phase II experiment provides a database targeted to allow detailed study of crop yields from process-based models under climate change. the systematic input parameter variations are designed to facilitate not only comparing the sensitivities of process-based crop yield models to changing climate and management inputs but also evaluating the complex interactions between driving factors (CO<sub>2</sub>, temperature, precipitation, and applied nitrogen). Its global nature also allows identifying geographic shifts in high yield potential locations. We expect that the simulations will yield multiple insights in future studies, and show here a selection of preliminary results to illustrate their potential uses.

First, the GGCMI Phase II simulations allow identifying major areas of uncertainty. Across factors impacting yields, inter-

model uncertainty is greatest for the CO<sub>2</sub> fertilization and nitrogen response effects. (Note that CO<sub>2</sub> effects are small for maize, a C4 crop, in Figures 6–7; rice, wheat, and soy are C3 and show larger responses.) Across geographic regions, projections are most uncertain in the high latitudes where yields may increase, and most robust in low latitudes where yield impacts are largest.

Second, the GGCMI Phase II simulations allow understanding the way that climate-driven changes and locations of cultivated land combine to produce yield impacts. One counterintuitive result immediate apparent is that irrigated maize shows steeper yield reductions under warming than does rain-fed maize when considered only over currently cultivated land (Figure 8). The effect results from geographic differences in cultivation. In any given location, irrigation (or additional rainfall increases) crop resiliency to temperature increase (partly by reducing negative effects from increased evapo-transpiration), but irrigated maize is grown in warmer locations where the impacts of warming are more severe (Figures S5–S6). The same behavior holds for rice and winter wheat, but not for soy or spring wheat (Figures S8–S10). Irrigated wheat and maize are also more sensitive to nitrogen fertilization levels than are analogous non-irrigated crops, presumably because those rain-fed crops are limited by water as well as nitrogen availability (Figure S19). (Soy as an efficient atmospheric nitrogen-fixer is relatively insensitive to nitrogen, and rice is not generally grown in water-limited conditions).

Third, we show that even the relatively limited GGCMI Phase II sampling space allows emulation of the climatological response of crop models with a relatively simple reduced-form statistical model. The systematic parameter sampling in the GGCMI Phase II procedure provides information on the influence of multiple interacting factors in a way that single projections cannot, and emulating the resulting response surface then

produces a tool that can aid in both physical interpretation of the process-based models and in assessment of agricultural impacts under arbitrary climate scenarios. Emulating the climatological response isolates long-term impacts from any confounding factors that complicate year-over-year changes, and the use of simple functional forms offer the possibility of physical interpretation of parameter values. We anticipate that systematic parameter sampling will become the norm in future crop model intercomparison exercises.

While the GGCMI Phase II database should offer the foundation for multiple future studies, several cautions need to be noted. Because the simulation protocol was designed to focus on change in yield under climate perturbations and not on replicating real-world yields, the models are not formally calibrated so cannot be used for impacts projections unless in used in conjunction with historical data (or data products). Because the GGCMI Phase II simulations apply uniform perturbations to historical climate inputs, they do not sample changes in higher order moments, and cannot address the additional crop yield impacts of potential changes in climate variability. Although distributional changes in model projections are fairly uncertain at present, follow-on experiments may wish to consider them. Several recent studies have described procedures for generating simulations that combine historical data with model projections of changes not only in temperature and precipitation means but in their marginal distributions or temporal dependence (e.g. Leeds et al. (2015), Poppick et al. (2016), Chang et al. (2016) and Haugen et al. (2018)).

The GGCMI Phase II output dataset invites a broad range of potential future avenues of analysis. A major target area of research is studying the models themselves including: a detailed examination of interaction terms between the major input drivers, a robust quantification of the sensitivity of different models to the input drivers, and comparisons with field-

level experimental data. The parameter space tested in GGCMI Phase II will allow detailed investigations into yield variability and response to extremes under changing management and CO<sub>2</sub> levels and allow the study of geographic shifts in optimal growing regions for different crops. The output dataset also contains other runs and variables not analyzed or shown here. Runs include several which allowed adaptation to climate changes by altering growing seasons, and additional variables include above ground biomass, LAI, and root biomass (as many as 25 output variables for some models). Emulation studies that are possible include a more systematic evaluation of different statistical model specifications and formal calculation of uncertainties in derived parameters. The development of multi-model ensembles such as GGCMI Phase II provides a way to begin to better understand crop responses to a range of potential climate inputs, improve process based models, and explore the potential benefits of adaptive responses included shifting growing season, cultivar types and cultivar geographic extent.

## 7. Acknowledgments

We thank Michael Stein and Kevin Schwarzwald, who provided helpful suggestions that contributed to this work. This research was performed as part of the Center for Robust Decision-Making on Climate and Energy Policy (RDCEP) at the University of Chicago, and was supported through a variety of sources. RDCEP is funded by NSF grant #SES-1463644 through the Decision Making Under Uncertainty program. J.F. was supported by the NSF NRT program, grant #DGE-1735359. C.M. was supported by the MACMIT project (01LN1317A) funded through the German Federal Ministry of Education and Research (BMBF). C.F. was supported by the European Research Council Synergy grant #ERC-2013-SynG-610028 Imbalance-. P. P.F. and K.W. were supported by the Newton Fund through the Met Office Climate Science for Service Partnership Brazil

(CSSP Brazil). A.S. was supported by the Office of Science of the U.S. Department of Energy as part of the Multi-sector Dynamics Research Program Area. S.O. acknowledges support from the Swedish strong research areas BECC and MERGE together with support from LUCCI (Lund University Centre for studies of Carbon Cycle and Climate Interactions). Computing resources were provided by the University of Chicago Research Computing Center (RCC).

## 8. Appendix A: Simulations – Assessment

The Müller et al. (2017) procedure evaluates response to year-to-year temperature and precipitation variations in a control run driven by historical climate and compares it to detrended historical yields from the FAO (Food and Agriculture Organization of the United Nations, 2018) by calculating the Pearson product moment correlation coefficient. The procedure offers no means of assessing CO<sub>2</sub> fertilization, since CO<sub>2</sub> has been relatively constant over the historical data collection period. Nitrogen introduces some uncertainty into the analysis, since the GGCMI Phase II runs impose fixed, uniform nitrogen application levels that are not realistic for individual countries. We evaluate up to three control runs for each model, since some modeling groups provide historical runs for three different nitrogen levels.

Results are similar to those of GGCMI Phase I, with reasonable fidelity at capturing year-over-year variation, with differences by region and crop stronger than difference between models. (That is, Figure 9 shows more similarity in horizontal than vertical bars.) No single model is dominant, with each model providing near best-in-class performance in at least one location-crop combination. For example, maize in the United States is consistently well-simulated while maize in Mexico is problematic (mean Pearson correlation coefficients of 0.X and 0.X, respectively). In some cases, especially in the developing

world, low correlation coefficients may indicate not model failure but problems in FAO yield data. For example, models have greater apparent skill for rice in India than in the neighboring Pakistan and Bangladesh; this difference may be implausible as solely a model effect. In general, Pearson correlation coeffi-

cients in GGCMI Phase II are slightly below those of Phase I, likely because of unrealistic nitrogen levels and lack of country level calibration in some models. (Compare Figure 9 to Müller et al. (2017) Figures 1–4 and 6.) Note that in this methodology, simulations of crops with low year-to-year variability such




Figure 9: Time series of correlation coefficients between simulated crop yield and FAO data (Food and Agriculture Organization of the United Nations, 2018) at the country level. The top panels indicate two example cases: US maize (a good case), and rice in India (mixed case), both for the high nitrogen application case. The heatmaps illustrate the Pearson  $r$  correlation coefficient between the detrended simulation mean yield at the country level compared to the detrended FAO yield data for the top producing countries for each crop with continuous FAO data over the 1980–2010 period. Models that provided different nitrogen application levels are shown with low, med, and high label (models that did not simulate different nitrogen levels are analogous to a high nitrogen application level). The ensemble mean yield is also correlated with the FAO data (not the mean of the correlations). Wheat contains both spring wheat and winter wheat simulations. The Pearson  $r$  correlation coefficients are similar to those of GGCMI Phase I, with reasonable fidelity at capturing year-over-year variation, with differences by region and crop stronger than difference between models as indicated by more horizontal bars than vertical bars of the same color.

as irrigated rice and wheat will tend to score more poorly than  
662 those with higher variability.

Some models do show particular strength for particular  
664 crops. For example, the EPIC family of models, and especially the EPIC-TAMU model, perform particularly well for  
666 soy across all regions. In other cases a model has particular strength in only certain crop and region combinations. For example, The strongest correlation coefficient in Figure 9 is that  
668 for the pDSSAT model for maize in the U.S. (the example crop-model-location used in many example figures in this paper), but pDSSAT slightly under performs for maize in other regions.  
670 These model assessment results are similar to those for GGCMI Phase I in Müller et al. (2017).

## 674 9. Appendix B: Emulation – Assessment

No general criteria exist for defining an acceptable crop  
676 model emulator. For a multi-model comparison exercise like GGCMI Phase II, one reasonable criterion is what we term the  
678 “normalized error”, which compares the fidelity of an emulator for a given model and scenario to the inter-model uncertainty.  
680 We define the normalized error  $e$  for each scenario as the difference between the fractional yield change from the emulator and that in the original simulation, divided by the standard deviation  
682 of the multi-model spread (Equations 2 and 3):

$$F_{scn.} = \frac{Y_{scn.} - Y_{baseline}}{Y_{baseline}} \quad (2)$$

$$e_{scn.} = \frac{F_{em, scn.} - F_{sim, scn.}}{\sigma_{sim, scn.}} \quad (3)$$

684 Here  $F_{scn.}$  is the fractional change in a model’s mean emulated or simulated yield from a defined baseline, in some scenario (scn.) in C, T, W, and N space;  $Y_{scn.}$  and  $Y_{baseline}$  are the absolute emulated or simulated mean yields. To assess the ability of the polynomial emulation to capture the behavior of com-




Figure 10: Assessment of emulator performance over currently cultivated areas based on normalized error (Equations 3, 2). We show performance of all 9 models emulated, over all crops and all sampled T and P inputs, but with CO<sub>2</sub> and nitrogen held fixed at baseline values. Large columns are crops and large rows models; squares within are T,P scenario pairs. Colors denote the fraction of currently cultivated hectares ('area frac') for each crop with normalized area  $e$  less than 1 indicating the the error between the emulation and simulation less than one standard deviation of the ensemble simulation spread. Of the 756 scenarios with these CO<sub>2</sub> and N values, we consider only those for which all 9 models submitted data (Figure S3). JULES did not simulate spring wheat and LPJ-GUESS did not simulate rice and soy. Emulator performance is generally satisfactory, with some exceptions. Emulator failures (significant areas of poor performance) occur for individual crop-model combinations, with performance generally degrading for hotter and wetter scenarios.




Figure 11: Illustration of our test of emulator performance, applied to the CARAIB model for the T+4 scenario for rain-fed crops. Contour colors indicate the normalized emulator error  $e$ , where  $e > 1$  means that emulator error exceeds the multi-model standard deviation. White areas are those where crops are not simulated by this model. Models differ in their areas omitted, meaning the number of samples used to calculate the multi-model standard deviation is not spatially consistent in all locations. Emulator performance is generally good relative to model spread in areas where crops are currently cultivated (compare to Figure 1) and in temperate zones in general; emulation issues occur primarily in marginal areas with low yield potentials. For CARAIB, emulation of soy is more problematic, as was also shown in Figure 10.

plex process-based models, we evaluate the normalized emulator error. That is, for each grid cell, model, and scenario we evaluate the difference between the model yield and its emulation, normalized by the inter-model standard deviation in yield projections. The normalized error  $e$  is the difference between the emulated fractional change in yield and that actually simulated, normalized by  $\sigma_{sim}$ , the standard deviation in simulated fractional yields change  $F_{sim, scn}$  across all models. The emulator is fitted across all available simulation outputs, and then the error is calculated across the each of the simulation scenarios provided by all nine models (Figure S3).

This metric implies that emulation is generally satisfactory, with several distinct exceptions. Almost all model-crop combination emulators have normalized errors less than one over nearly all currently cultivated hectares (Figure 10), but some individual model-crop combinations are problematic (e.g. PROMET for rice and soy, JULES for soy and winter wheat, Figures S14–S15). Normalized errors for soy are somewhat higher across all models not because emulator fidelity is worse but because models agree more closely on yield changes for soy than for other crops (see Figure S16), lowering the denominator. Emulator performance often degrades in geographic locations where crops are not currently cultivated. Figure 11 shows a CARAIB case as an example, where emulator performance is satisfactory over cultivated areas for all crops other than soy, but uncultivated regions show some problematic areas (see also Figure S12).

This assessment procedure is relatively forgiving for several reasons. First, each emulation is evaluated against the simulation actually used to train the emulator. Had we used a spline interpolation the error would necessarily be zero. Second, the performance metric scales emulator fidelity not by the magnitude of yield changes but by the inter-model spread in those changes. The normalized error  $e$  for a model depends not only on the fi-

dentity of its emulator in reproducing a given simulation but on  
724 the particular suite of models considered in the intercompari-  
son exercise. Where models differ more widely, the standard  
726 for emulators becomes less stringent. This effect is readily seen  
when comparing assessments of emulator performance in sim-  
728 ulations at baseline CO<sub>2</sub> (Figure 10) with those at higher CO<sub>2</sub>  
levels (Figure S13) because models disagree on the magnitude  
730 of CO<sub>2</sub> fertilization. The rationale for this choice of assessment  
metric is to relate the fidelity of the emulation to an estimate of  
732 true uncertainty, which we take as the multi-model spread. We  
therefore do not provide a formal parameter uncertainty analy-  
734 sis, but note that the GGCMI Phase II dataset is well-suited to  
statistical exploration of emulation approaches and quantifica-  
736 tion of emulator fidelity. More rigorous emulator assessments  
that may be preformed in future work include: testing other sta-  
738 tistical specifications including non-parametric models, cross-  
validation procedures where the emulator is trained on some  
740 portion of data and tested on a held-out portion, and calculating  
standard error on emulator parameters.

## 742 10. References

- Angulo, C., Ritter, R., Lock, R., Enders, A., Fronzek, S., & Ewert, F. (2013).  
744 Implication of crop model calibration strategies for assessing regional im-  
pacts of climate change in europe. *Agric. For. Meteorol.*, *170*, 32 – 46.
- Asseng, S., Ewert, F., Martre, P., Ritter, R. P., B. Lobell, D., Cammarano, D.,  
746 A. Kimball, B., Ottman, M., W. Wall, G., White, J., Reynolds, M., D. Al-  
derman, P., Prasad, P. V. V., Aggarwal, P., Anothai, J., Basso, B., Bier-  
748 nath, C., Challinor, A., De Sanctis, G., & Zhu, Y. (2015). Rising tempera-  
tures reduce global wheat production. *Nature Climate Change*, *5*, 143–147.  
doi:10.1038/nclimate2470.
- Asseng, S., Ewert, F., Rosenzweig, C., Jones, J., Hatfield, J., Ruane, A.,  
752 J. Boote, K., Thorburn, P., Ritter, R. P., Cammarano, D., Brisson, N.,  
Basso, B., Martre, P., Aggarwal, P., Angulo, C., Bertuzzi, P., Biernath,  
754 C., Challinor, A., Doltra, J., & Wolf, J. (2013). Uncertainty in simu-  
lating wheat yields under climate change. *Nature Climate Change*, *3*, 827832.  
doi:10.1038/nclimate1916.
- Aulakh, M. S., & Malhi, S. S. (2005). Interactions of Nitrogen with Other  
758 Nutrients and Water: Effect on Crop Yield and Quality, Nutrient Use Ef-  
ficiency, Carbon Sequestration, and Environmental Pollution. *Advances in  
Agronomy*, *86*, 341 – 409.
- Balkovi, J., van der Velde, M., Skalsk, R., Xiong, W., Folberth, C., Khabarov,  
762 N., Smirnov, A., Mueller, N. D., & Obersteiner, M. (2014). Global wheat  
production potentials and management flexibility under the representative  
concentration pathways. *Global and Planetary Change*, *122*, 107 – 121.
- Blanc, E. (2017). Statistical emulators of maize, rice, soybean and wheat yields  
766 from global gridded crop models. *Agricultural and Forest Meteorology*, *236*,  
145 – 161.
- Blanc, E., & Sultan, B. (2015). Emulating maize yields from global gridded  
770 crop models using statistical estimates. *Agricultural and Forest Meteorol-  
ogy*, *214-215*, 134 – 147.
- von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., & Zaehle, S.  
772 (2018). Implementing the Nitrogen cycle into the dynamic global vegeta-  
tion, hydrology and crop growth model LPJmL (version 5.0). *Geoscientific  
Model Development*, *11*, 2789–2812.
- Calvin, K., Patel, P., Clarke, L., Asrar, G., Bond-Lamberty, B., Cui, R. Y.,  
774 Di Vittorio, A., Dorheim, K., Edmonds, J., Hartin, C., Hejazi, M., Horowitz,  
R., Iyer, G., Kyle, P., Kim, S., Link, R., McJeon, H., Smith, S. J., Snyder,  
776 A., Waldhoff, S., & Wise, M. (2019). Gcam v5.1: representing the linkages  
between energy, water, land, climate, and economic systems. *Geoscientific  
Model Development*, *12*, 677–698. doi:10.5194/gmd-12-677-2019.
- Castruccio, S., McInerney, D. J., Stein, M. L., Liu Crouch, F., Jacob, R. L.,  
782 & Moyer, E. J. (2014). Statistical Emulation of Climate Model Projections  
Based on Precomputed GCM Runs. *Journal of Climate*, *27*, 1829–1844.
- Challinor, A., Watson, J., Lobell, D., Howden, S., Smith, D., & Chhetri, N.  
784 (2014). A meta-analysis of crop yield under climate change and adaptation.  
*Nature Climate Change*, *4*, 287 – 291.
- Chang, W., Stein, M., Wang, J., Kotamarthi, V., & Moyer, E. (2016). Changes in  
786 spatio-temporal precipitation patterns in changing climate conditions. *Jour-  
nal of Climate*, *29*. doi:10.1175/JCLI-D-15-0844.1.
- Conti, S., Gosling, J. P., Oakley, J. E., & O'Hagan, A. (2009). Gaussian process  
788 emulation of dynamic computer codes. *Biometrika*, *96*, 663–676.
- Duncan, W. (1972). SIMCOT: a simulation of cotton growth and yield. In  
790 C. Murphy (Ed.), *Proceedings of a Workshop for Modeling Tree Growth,*  
Duke University, Durham, North Carolina (pp. 115–118). Durham, North  
Carolina.
- Duncan, W., Loomis, R., Williams, W., & Hanau, R. (1967). A model for  
792 simulating photosynthesis in plant communities. *Hilgardia*, (pp. 181–205).
- Dury, M., Hambuckers, A., Warnant, P., Henrot, A., Favre, E., Ouberoudous,  
794 M., & François, L. (2011). Responses of European forest ecosystems to  
21st century climate: assessing changes in interannual variability and fire  
intensity. *iForest - Biogeosciences and Forestry*, (pp. 82–99).
- Elliott, J., Kelly, D., Chryssanthacopoulos, J., Glotter, M., Jhunjhnuwala, K.,  
796 Best, N., Wilde, M., & Foster, I. (2014). The parallel system for integrating  
impact models and sectors (pSIMS). *Environmental Modelling and Soft-  
ware*, *62*, 509–516.
- Elliott, J., Müller, C., Deryng, D., Chryssanthacopoulos, J., Boote, K. J.,  
800 Büchner, M., Foster, I., Glotter, M., Heinke, J., Iizumi, T., Izaurralde, R. C.,  
Mueller, N. D., Ray, D. K., Rosenzweig, C., Ruane, A. C., & Sheffield, J.  
802 (2015). The Global Gridded Crop Model Intercomparison: data and mod-  
eling protocols for Phase 1 (v1.0). *Geoscientific Model Development*, *8*,  
261–277.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J.,  
804 & Taylor, K. E. (2016). Overview of the coupled model intercomparison  
project phase 6 (cmip6) experimental design and organization. *Geoscientific  
Model Development*, *9*, 1937–1958.
- Ferrise, R., Moriondo, M., & Bindi, M. (2011). Probabilistic assessments of cli-  
806 mate change impacts on durum wheat in the mediterranean region. *Natural  
Hazards and Earth System Sciences*, *11*, 1293–1302.
- Folberth, C., Gaiser, T., Abbaspour, K. C., Schulin, R., & Yang, H. (2012). Re-  
808 gionalization of a large-scale crop growth model for sub-Saharan Africa:  
Model setup, evaluation, and estimation of maize yields. *Agriculture,  
Ecosystems & Environment*, *151*, 21 – 33.
- Food and Agriculture Organization of the United Nations (2018). FAOSTAT  
810 database. URL: <http://www.fao.org/faostat/en/home>.
- Fronzek, S., Pirttioja, N., Carter, T. R., Bindi, M., Hoffmann, H., Palosuo, T.,  
812 Ruiz-Ramos, M., Tao, F., Trnka, M., Acutis, M., Asseng, S., Baranowski, P.,  
Basso, B., Bodin, P., Buis, S., Cammarano, D., Deligios, P., Destain, M.-F.,  
814 Dumont, B., Ewert, F., Ferrise, R., François, L., Gaiser, T., Hlavinka, P.,  
Jacquemin, I., Kersebaum, K. C., Kollas, C., Krzyszczak, J., Lorite, I. J.,  
Minet, J., Minguez, M. I., Montesino, M., Moriondo, M., Müller, C., Nen-  
del, C., Öztürk, I., Perego, A., Rodríguez, A., Ruane, A. C., Ruget, F., Sanna,  
816 M., Semenov, M. A., Slawinski, C., Strattonovich, P., Supit, I., Waha, K.,  
Wang, E., Wu, L., Zhao, Z., & Rötter, R. P. (2018). Classifying multi-model  
wheat yield impact response surfaces showing sensitivity to temperature and  
818 precipitation change. *Agricultural Systems*, *159*, 209–224.
- Glotter, M., Elliott, J., McInerney, D., Best, N., Foster, I., & Moyer, E. J. (2014).  
820 Evaluating the utility of dynamical downscaling in agricultural impacts pro-  
jections. *Proceedings of the National Academy of Sciences*, *111*, 8776–8781.
- Glotter, M., Moyer, E., Ruane, A., & Elliott, J. (2015). Evaluating the Sensitivity  
822 of Agricultural Model Performance to Different Climate Inputs. *Journal  
of Applied Meteorology and Climatology*, *55*, 151113145618001.

- Hank, T., Bach, H., & Mauser, W. (2015). Using a Remote Sensing-Supported Hydro-Agroecological Model for Field-Scale Simulation of Heterogeneous Crop Growth and Yield: Application for Wheat in Central Europe. *Remote Sensing*, 7, 3934–3965.
- Haugen, M., Stein, M., Moyer, E., & Sriver, R. (2018). Estimating changes in temperature distributions in a large ensemble of climate simulations using quantile regression. *Journal of Climate*, 31, 8573–8588. doi:10.1175/JCLI-D-17-0782.1.
- He, W., Yang, J., Zhou, W., Drury, C., Yang, X., D. Reynolds, W., Wang, H., He, P., & Li, Z.-T. (2016). Sensitivity analysis of crop yields, soil water contents and nitrogen leaching to precipitation, management practices and soil hydraulic properties in semi-arid and humid regions of Canada using the DSSAT model. *Nutrient Cycling in Agroecosystems*, 106, 201–215.
- Heady, E. O. (1957). An Econometric Investigation of the Technology of Agricultural Production Functions. *Econometrica*, 25, 249–268.
- Heady, E. O., & Dillon, J. L. (1961). *Agricultural production functions*. Iowa State University Press.
- Holden, P., Edwards, N., PH, G., Fraedrich, K., Lunkeit, F., E, K., Labriet, M., Kanudia, A., & F, B. (2014). Plasim-entsem v1.0: A spatiotemporal emulator of future climate change for impacts assessment. *Geoscientific Model Development*, 7, 433–451. doi:10.5194/gmd-7-433-2014.
- Holzkämper, A., Calanca, P., & Fuhrer, J. (2012). Statistical crop models: Predicting the effects of temperature and precipitation changes. *Climate Research*, 51, 11–21.
- Holzworth, D. P., Huth, N. I., deVoil, P. G., Zurcher, E. J., Herrmann, N. I., McLean, G., Chenu, K., van Oosterom, E. J., Snow, V., Murphy, C., Moore, A. D., Brown, H., Whish, J. P., Verrall, S., Fainges, J., Bell, L. W., Peake, A. S., Poulton, P. L., Hochman, Z., Thorburn, P. J., Gaydon, D. S., Dalglish, N. P., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp, J., Cichota, R., Vogeler, I., Li, F. Y., Wang, E., Hammer, G. L., Robertson, M. J., Dimes, J. P., Whitbread, A. M., Hunt, J., van Rees, H., McClelland, T., Carberry, P. S., Hargreaves, J. N., MacLeod, N., McDonald, C., Harsdorff, J., Wedgwood, S., & Keating, B. A. (2014). APSIM Evolution towards a new generation of agricultural systems simulation. *Environmental Modelling and Software*, 62, 327 – 350.
- Howden, S., & Crimp, S. (2005). Assessing dangerous climate change impacts on australia's wheat industry. *Modelling and Simulation Society of Australia and New Zealand*, (pp. 505–511).
- Iizumi, T., Nishimori, M., & Yokozawa, M. (2010). Diagnostics of climate model biases in summer temperature and warm-season insolation for the simulation of regional paddy rice yield in japan. *Journal of Applied Meteorology and Climatology*, 49, 574–591.
- Ingestad, T. (1977). Nitrogen and Plant Growth; Maximum Efficiency of Nitrogen Fertilizers. *Ambio*, 6, 146–151.
- Izaurralde, R., Williams, J., McGill, W., Rosenberg, N., & Quiroga Jakas, M. (2006). Simulating soil C dynamics with EPIC: Model description and testing against long-term data. *Ecological Modelling*, 192, 362–384.
- Jagtap, S. S., & Jones, J. W. (2002). Adaptation and evaluation of the CROPGRO-soybean model to predict regional yield and production. *Agriculture, Ecosystems & Environment*, 93, 73 – 85.
- Jones, J., Hoogenboom, G., Porter, C., Boote, K., Batchelor, W., Hunt, L., Wilkens, P., Singh, U., Gijsman, A., & Ritchie, J. (2003). The DSSAT cropping system model. *European Journal of Agronomy*, 18, 235 – 265.
- Jones, J. W., Antle, J. M., Basso, B., Boote, K. J., Conant, R. T., Foster, I., Godfray, H. C. J., Herrero, M., Howitt, R. E., Janssen, S., Keating, B. A., Munoz-Carpena, R., Porter, C. H., Rosenzweig, C., & Wheeler, T. R. (2017). Toward a new generation of agricultural system data, models, and knowledge products: State of agricultural systems science. *Agricultural Systems*, 155, 269 – 288.
- Keating, B., Carberry, P., Hammer, G., Probert, M., Robertson, M., Holzworth, D., Huth, N., Hargreaves, J., Meinke, H., Hochman, Z., McLean, G., Verburg, K., Snow, V., Dimes, J., Silburn, M., Wang, E., Brown, S., Bristow, K., Asseng, S., Chapman, S., McCown, R., Freebairn, D., & Smith, C. (2003). An overview of APSIM, a model designed for farming systems simulation. *European Journal of Agronomy*, 18, 267 – 288.
- Leeds, W. B., Moyer, E. J., & Stein, M. L. (2015). Simulation of future climate under changing temporal covariance structures. *Advances in Statistical Climatology, Meteorology and Oceanography*, 1, 1–14. URL: <https://www.adv-stat-clim-meteorol-oceanogr.net/1/1/2015/>. doi:10.5194/ascmo-1-1-2015.
- Lindeskog, M., Arneth, A., Bondeau, A., Waha, K., Seaquist, J., Olin, S., & Smith, B. (2013). Implications of accounting for land use in simulations of ecosystem carbon cycling in Africa. *Earth System Dynamics*, 4, 385–407.
- Liu, J., Williams, J. R., Zehnder, A. J., & Yang, H. (2007). GEPIC - modelling wheat yield and crop water productivity with high resolution on a global scale. *Agricultural Systems*, 94, 478 – 493.
- Liu, W., Yang, H., Folberth, C., Wang, X., Luo, Q., & Schulin, R. (2016a). Global investigation of impacts of PET methods on simulating crop-water relations for maize. *Agricultural and Forest Meteorology*, 221, 164 – 175.
- Liu, W., Yang, H., Liu, J., Azevedo, L. B., Wang, X., Xu, Z., Abbaspour, K. C., & Schulin, R. (2016b). Global assessment of nitrogen losses and trade-offs with yields from major crop cultivations. *Science of The Total Environment*, 572, 526 – 537.
- Lobell, D. B., & Burke, M. B. (2010). On the use of statistical models to predict crop yield responses to climate change. *Agricultural and Forest Meteorology*, 150, 1443 – 1452.
- Lobell, D. B., & Field, C. B. (2007). Global scale climate-crop yield relationships and the impacts of recent warming. *Environmental Research Letters*, 2, 014002.
- MacKay, D. (1991). Bayesian Interpolation. *Neural Computation*, 4, 415–447.
- Makowski, D., Asseng, S., Ewert, F., Bassu, S., Durand, J., Martre, P., Adam, M., Aggarwal, P., Angulo, C., Baron, C., Basso, B., Bertuzzi, P., Biernath, C., Boogaard, H., Boote, K., Brisson, N., Cammarano, D., Challinor, A., Conijn, J., & Wolf, J. (2015). Statistical analysis of large simulated yield datasets for studying climate effects. (p. 1100). doi:10.13140/RG.2.1.5173.8328.
- Mauser, W., & Bach, H. (2015). PROMET - Large scale distributed hydrological modelling to study the impact of climate change on the water flows of mountain watersheds. *Journal of Hydrology*, 376, 362 – 377.
- Mauser, W., Klepper, G., Zabel, F., Delzeit, R., Hank, T., Putzenlechner, B., & Calzadilla, A. (2009). Global biomass production potentials exceed expected future demand without the need for cropland expansion. *Nature Communications*, 6.
- McDermid, S., Dileepkumar, G., Murthy, K., Nedumaran, S., Singh, P., Srinivas, C., Gangwar, B., Subash, N., Ahmad, A., Zubair, L., & Nissanka, S. (2015). Integrated assessments of the impacts of climate change on agriculture: An overview of AgMIP regional research in South Asia. *Chapter in: Handbook of Climate Change and Agroecosystems*, (pp. 201–218).
- Mistry, M. N., Wing, I. S., & De Cian, E. (2017). Simulated vs. empirical weather responsiveness of crop yields: US evidence and implications for the agricultural impacts of climate change. *Environmental Research Letters*, 12.
- Moore, F. C., Baldos, U., Hertel, T., & Diaz, D. (2017). New science of climate change impacts on agriculture implies higher social cost of carbon. *Nature Communications*, 8.
- Müller, C., Elliott, J., Chrissanthacopoulos, J., Arneth, A., Balkovic, J., Ciais, P., Deryng, D., Folberth, C., Glotter, M., Hoek, S., Iizumi, T., Izaurralde, R. C., Jones, C., Khabarov, N., Lawrence, P., Liu, W., Olin, S., Pugh, T. A. M., Ray, D. K., Reddy, A., Rosenzweig, C., Ruane, A. C., Sakurai, G., Schmid, E., Skalsky, R., Song, C. X., Wang, X., de Wit, A., & Yang, H. (2017). Global gridded crop model evaluation: benchmarking, skills, deficiencies and implications. *Geoscientific Model Development*, 10, 1403–1422.
- Nakamura, T., Osaki, M., Koike, T., Hanba, Y. T., Wada, E., & Tadano, T. (1997). Effect of CO<sub>2</sub> enrichment on carbon and nitrogen interaction in wheat and soybean. *Soil Science and Plant Nutrition*, 43, 789–798.
- O'Hagan, A. (2006). Bayesian analysis of computer code outputs: A tutorial. *Reliability Engineering & System Safety*, 91, 1290 – 1300.
- Olin, S., Schurgers, G., Lindeskog, M., Wårldin, D., Smith, B., Bodin, P., Holmér, J., & Arneth, A. (2015). Modelling the response of yields and tissue C:N to changes in atmospheric CO<sub>2</sub> and N management in the main wheat regions of western europe. *Biogeosciences*, 12, 2489–2515. doi:10.5194/bg-12-2489-2015.
- Osaki, M., Shinano, T., & Tadano, T. (1992). Carbon-nitrogen interaction in field crop production. *Soil Science and Plant Nutrition*, 38, 553–564.
- Osborne, T., Gornall, J., Hooker, J., Williams, K., Wiltshire, A., Betts, R., & Wheeler, T. (2015). JULES-crop: a parametrisation of crops in the Joint UK Land Environment Simulator. *Geoscientific Model Development*, 8, 1139–1155.
- Ostberg, S., Schewe, J., Childers, K., & Frieler, K. (2018). Changes in crop yields and their variability at different levels of global warming. *Earth System Dynamics*, 9, 479–496.

- Oyebamiji, O. K., Edwards, N. R., Holden, P. B., Garthwaite, P. H., Schaphoff, S., & Gerten, D. (2015). Emulating global climate change impacts on crop yields. *Statistical Modelling*, 15, 499–525.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Pas-  
sos, A., Cournapeau, D., Brucher, M., Perrot, M., & Duchesnay, E. (2011). Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research*, 12, 2825–2830.
- Pirttioja, N., Carter, T., Fronzek, S., Bind, M., Hoffmann, H., Palosuo, T., Ruiz-Ramos, M., Tao, F., Trnka, M., Acutis, M., Asseng, S., Baranowski, P., Basso, B., Bodin, P., Buis, S., Cammarano, D., Deligios, P., Destain, M., Dumont, B., Ewert, F., Ferrise, R., François, L., Gaiser, T., Hlavinka, P., Jacquemin, I., Kersebaum, K., Kollas, C., Krzyszczak, J., Lorite, I., Minet, J., Minguez, M., Montesino, M., Moriando, M., Müller, C., Nendel, C., Öztürk, I., Perego, A., Rodríguez, A., Ruane, A., Ruget, F., Sanna, M., Semenov, M., Slawinski, C., Strattonovich, P., Supit, I., Waha, K., Wang, E., Wu, L., Zhao, Z., & Rötter, R. (2015). Temperature and precipitation effects on wheat yield across a European transect: a crop model ensemble analysis using impact response surfaces. *Climate Research*, 65, 87–105.
- Poppick, A., McInerney, D. J., Moyer, E. J., & Stein, M. L. (2016). Temperatures in transient climates: Improved methods for simulations with evolving temporal covariances. *Ann. Appl. Stat.*, 10, 477–505. URL: <https://doi.org/10.1214/16-AOAS903>. doi:10.1214/16-AOAS903.
- Porter et al. (IPCC) (2014). Food security and food production systems. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. In C. F. et al. (Ed.), *IPCC Fifth Assessment Report* (pp. 485–533). Cambridge, UK: Cambridge University Press.
- Portmann, F., Siebert, S., Bauer, C., & Doell, P. (2008). Global dataset of monthly growing areas of 26 irrigated crops.
- Portmann, F., Siebert, S., & Doell, P. (2010). MIRCA2000 - Global Monthly Irrigated and Rainfed crop Areas around the Year 2000: A New High-Resolution Data Set for Agricultural and Hydrological Modeling. *Global Biogeochemical Cycles*, 24, GB1011.
- Pugh, T., Müller, C., Elliott, J., Deryng, D., Folberth, C., Olin, S., Schmid, E., & Arneth, A. (2016). Climate analogues suggest limited potential for intensification of production on current croplands under climate change. *Nature Communications*, 7, 12608.
- Räisänen, J., & Ruokolainen, L. (2006). Probabilistic forecasts of near-term climate change based on a resampling ensemble technique. *Tellus A: Dynamic Meteorology and Oceanography*, 58, 461–472.
- Ratto, M., Castelletti, A., & Pagano, A. (2012). Emulation techniques for the reduction and sensitivity analysis of complex environmental models. *Environmental Modelling & Software*, 34, 1 – 4.
- Razavi, S., Tolson, B. A., & Burn, D. H. (2012). Review of surrogate modeling in water resources. *Water Resources Research*, 48.
- Roberts, M., Braun, N., R Sinclair, T., B Lobell, D., & Schlenker, W. (2017). Comparing and combining process-based crop models and statistical models with some implications for climate change. *Environmental Research Letters*, 12.
- Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A. C., Müller, C., Arneth, A., Boote, K. J., Folberth, C., Glotter, M., Khabarov, N., Neumann, K., Piontek, F., Pugh, T. A. M., Schmid, E., Stehfest, E., Yang, H., & Jones, J. W. (2014). Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the National Academy of Sciences*, 111, 3268–3273.
- Rosenzweig, C., Jones, J., Hatfield, J., Ruane, A., Boote, K., Thorburn, P., Antle, J., Nelson, G., Porter, C., Janssen, S., Asseng, S., Basso, B., Ewert, F., Wallach, D., Baigorria, G., & Winter, J. (2013). The Agricultural Model Intercomparison and Improvement Project (AgMIP): Protocols and pilot studies. *Agricultural and Forest Meteorology*, 170, 166 – 182.
- Rosenzweig, C., Ruane, A. C., Antle, J., Elliott, J., Ashfaq, M., Chatta, A. A., Ewert, F., Folberth, C., Hathie, I., Havlik, P., Hoogenboom, G., Lotze-Campen, H., MacCarthy, D. S., Mason-D'Croz, D., Contreras, E. M., Müller, C., Perez-Dominguez, I., Phillips, M., Porter, C., Raymundo, R. M., Sands, R. D., Schleussner, C.-F., Valdivia, R. O., Valin, H., & Wiebe, K. (2018). Coordinating AgMIP data and models across global and regional scales for 1.5°C and 2.0°C assessments. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 376.
- Ruane, A., I. Hudson, N., Asseng, S., Camarrano, D., Ewert, F., Martre, P., J. Boote, K., Thorburn, P., Aggarwal, P., Angulo, C., Basso, B., Bertuzzi, P., Biernath, C., Brisson, N., Challinor, A., Doltra, J., Gayler, S., Goldberg, R., Grant, R., & Wolf, J. (2016). Multi-wheat-model ensemble responses to interannual climate variability. *Environmental Modelling and Software*, 81, 86–101.
- Ruane, A. C., Antle, J., Elliott, J., Folberth, C., Hoogenboom, G., Mason-D'Croz, D., Müller, C., Porter, C., Phillips, M. M., Raymundo, R. M., Sands, R., Valdivia, R. O., White, J. W., Wiebe, K., & Rosenzweig, C. (2018). Biophysical and economic implications for agriculture of +1.5° and +2.0°C global warming using AgMIP Coordinated Global and Regional Assessments. *Climate Research*, 76, 17–39.
- Ruane, A. C., Cecil, L. D., Horton, R. M., Gordon, R., McCollum, R., Brown, D., Killough, B., Goldberg, R., Greeley, A. P., & Rosenzweig, C. (2013). Climate change impact uncertainties for maize in panama: Farm information, climate projections, and yield sensitivities. *Agricultural and Forest Meteorology*, 170, 132 – 145.
- Ruane, A. C., Goldberg, R., & Chryssanthacopoulos, J. (2015). Climate forcing datasets for agricultural modeling: Merged products for gap-filling and historical climate series estimation. *Agric. Forest Meteorol.*, 200, 233–248.
- Ruane, A. C., McDermid, S., Rosenzweig, C., Baigorria, G. A., Jones, J. W., Romero, C. C., & Cecil, L. D. (2014). Carbon-temperature-water change analysis for peanut production under climate change: A prototype for the agmip coordinated climate-crop modeling project (c3mp). *Glob. Change Biol.*, 20, 394–407. doi:10.1111/gcb.12412.
- Rubel, F., & Kottek, M. (2010). Observed and projected climate shifts 1901–2100 depicted by world maps of the Köppen-Geiger climate classification. *Meteorologische Zeitschrift*, 19, 135–141.
- Ruiz-Ramos, M., Ferrise, R., Rodriguez, A., Lorite, I., Bind, M., Carter, T., Fronzek, S., Palosuo, T., Pirttioja, N., Baranowski, P., Buis, S., Cammarano, D., Chen, Y., Dumont, B., Ewert, F., Gaiser, T., Hlavinka, P., Hoffmann, H., Hhn, J., Jurecka, F., Kersebaum, K., Krzyszczak, J., Lana, M., Mechiche-Alami, A., Minet, J., Montesino, M., Nendel, C., Porter, J., Ruget, F., Semenov, M., Steinmetz, Z., Strattonovich, P., Supit, I., Tao, F., Trnka, M., de Wit, A., & Rötter, R. (2018). Adaptation response surfaces for managing wheat under perturbed climate and co2 in a mediterranean environment. *Agricultural Systems*, 159, 260 – 274. doi:doi.org/10.1016/j.aghs.2017.01.009.
- Sacks, W. J., Deryng, D., Foley, J. A., & Ramankutty, N. (2010). Crop planting dates: an analysis of global patterns. *Global Ecology and Biogeography*, 19, 607–620.
- Schauberger, B., Archontoulis, S., Arneth, A., Balkovic, J., Caias, P., Deryng, D., Elliott, J., Folberth, C., Khabarov, N., Müller, C., A. M. Pugh, T., Rolinski, S., Schaphoff, S., Schmid, E., Wang, X., Schlenker, W., & Frieler, K. (2017). Consistent negative response of US crops to high temperatures in observations and crop models. *Nature Communications*, 8, 13931.
- Schlenker, W., & Roberts, M. J. (2009). Nonlinear temperature effects indicate severe damages to U.S. crop yields under climate change. *Proceedings of the National Academy of Sciences*, 106, 15594–15598.
- Snyder, A., Calvin, K. V., Phillips, M., & Ruane, A. C. (2018). A crop yield change emulator for use in gcam and similar models: Persephone v1.0. *Geoscientific Model Development Discussions*, 2018, 1–42.
- Storlie, C. B., Swiler, L. P., Helton, J. C., & Sallaberry, C. J. (2009). Implementation and evaluation of nonparametric regression procedures for sensitivity analysis of computationally demanding models. *Reliability Engineering & System Safety*, 94, 1735 – 1763.
- Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, 93, 485–498.
- Tebaldi, C., & Lobell, D. B. (2008). Towards probabilistic projections of climate change impacts on global crop yields. *Geophysical Research Letters*, 35.
- Valade, A., Caias, P., Vuichard, N., Viovy, N., Caubel, A., Huth, N., Marin, F., & Martin, J. F. (2014). Modeling sugarcane yield with a process-based model from site to continental scale: Uncertainties arising from model structure and parameter values. *Geoscientific Model Development*, 7, 1225–1245.
- Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., & Schewe, J. (2014). The Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP): Project framework. *Proceedings of the National Academy of Sciences*, 111, 3228–3232.
- White, J. W., Hoogenboom, G., Kimball, B. A., & Wall, G. W. (2011). Method-

- ologies for simulating impacts of climate change on crop production. *Field Crops Research*, 124, 357 – 368.
- Williams, K., Gornall, J., Harper, A., Wiltshire, A., Hemming, D., Quaife, T., Arkebauer, T., & Scoby, D. (2017). Evaluation of JULES-crop performance against site observations of irrigated maize from Mead, Nebraska. *Geoscientific Model Development*, 10, 1291–1320.
- Williams, K. E., & Falloon, P. D. (2015). Sources of interannual yield variability in JULES-crop and implications for forcing with seasonal weather forecasts. *Geoscientific Model Development*, 8, 3987–3997.
- de Wit, C. (1957). Transpiration and crop yields. *Verslagen van Landbouwkundige Onderzoeken* : 64.6.
- Wolf, J., & Oijen, M. (2002). Modelling the dependence of european potato yields on changes in climate and co<sub>2</sub>. *Agricultural and Forest Meteorology*, 112, 217 – 231.
- Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D. B., Huang, Y., Huang, M., Yao, Y., Bassu, S., Ciais, P., Durand, J. L., Elliott, J., Ewert, F., Janssens, I. A., Li, T., Lin, E., Liu, Q., Martre, P., Müller, C., Peng, S., Peuelas, J., Ruane, A. C., Wallach, D., Wang, T., Wu, D., Liu, Z., Zhu, Y., Zhu, Z., & Asseng, S. (2017). Temperature increase reduces global yields of major crops in four independent estimates. *Proc. Natl. Acad. Sci.*, 114, 9326–9331.