

The GGCMI Phase II experiment: global crop yield responses to changes in carbon dioxide, temperature, water, and nitrogen levels

James Franke^{a,b,*}, Joshua Elliott^{b,c}, Christoph Müller^d, Alexander Ruane^e, Abigail Snyder^f, Jonas Jägermeyr^{c,b,d,e}, Juraj Balkovic^{g,h}, Philippe Ciais^{i,j}, Marie Dury^k, Pete Falloon^l, Christian Folberth^g, Louis François^k, Tobias Hank^m, Munir Hoffmannⁿ, Cesar Izaurralde^{o,p}, Ingrid Jacquemin^k, Curtis Jones^o, Nikolay Khabarov^g, Marian Kochⁿ, Michelle Li^{b,l}, Wenfeng Liu^{r,i}, Stefan Olin^s, Meridel Phillips^{e,t}, Thomas Pugh^{u,v}, Ashwan Reddy^o, Xuhui Wang^{i,j}, Karina Williams^l, Florian Zabel^m, Elisabeth Moyer^{a,b}

^aDepartment of the Geophysical Sciences, University of Chicago, Chicago, IL, USA

^bCenter for Robust Decision-making on Climate and Energy Policy (RDCEP), University of Chicago, Chicago, IL, USA

^cDepartment of Computer Science, University of Chicago, Chicago, IL, USA

^dPotsdam Institute for Climate Impact Research, Leibniz Association (Member), Potsdam, Germany

^eNASA Goddard Institute for Space Studies, New York, NY, United States

^fJoint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD, USA

^gEcosystem Services and Management Program, International Institute for Applied Systems Analysis, Laxenburg, Austria

^hDepartment of Soil Science, Faculty of Natural Sciences, Comenius University in Bratislava, Bratislava, Slovak Republic

ⁱLaboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, 91191 Gif-sur-Yvette, France

^jSino-French Institute of Earth System Sciences, College of Urban and Environmental Sciences, Peking University, Beijing, China

^kUnité de Modélisation du Climat et des Cycles Biogéochimiques, UR SPHERES, Institut d'Astrophysique et de Géophysique, University of Liège, Belgium

^lMet Office Hadley Centre, Exeter, United Kingdom

^mDepartment of Geography, Ludwig-Maximilians-Universität, Munich, Germany

ⁿGeorg-August-University Göttingen, Tropical Plant Production and Agricultural Systems Modelling, Göttingen, Germany

^oDepartment of Geographical Sciences, University of Maryland, College Park, MD, USA

^pTexas AgriLife Research and Extension, Texas A&M University, Temple, TX, USA

^qDepartment of Statistics, University of Chicago, Chicago, IL, USA

^rEAWAG, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

^sDepartment of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

^tEarth Institute Center for Climate Systems Research, Columbia University, New York, NY, USA

^uKarlsruhe Institute of Technology, IMK-IFU, 82467 Garmisch-Partenkirchen, Germany.

^vSchool of Geography, Earth and Environmental Science, University of Birmingham, Birmingham, UK.

Abstract

Concerns about food security under climate change have motivated efforts to better understand the future changes in yields by using detailed process-based models in agronomic sciences. Process-based models differ on many details affecting yields and considerable uncertainty remains in future yield projections. Phase II of the Global Gridded Crop Model Intercomparison (GGCMI), an activity of the Agricultural Model Intercomparison and Improvement Project (AgMIP), consists of a large simulation set with perturbations in atmospheric CO₂ concentrations, temperature, precipitation, and applied nitrogen inputs and constitutes a data-rich basis of projected yield changes across twelve models and five crops (maize, soy, rice, spring wheat, and winter wheat) using global gridded simulations. In this paper we present the simulation output database from Phase II of the GGCMI effort, a targeted experiment aimed at understanding the sensitivity to and interaction between multiple climate variables (as well as management) on yields, and illustrate some initial summary results from the model intercomparison project. We also present the construction of a simple “emulator or statistical representation of the simulated 30-year mean climatological output in each location for each crop and model. The emulator captures the response of the process-based models in a lightweight, computationally tractable form that facilitates model comparison as well as potential applications in subsequent modeling efforts such as integrated assessment.

Keywords: climate change, food security, model emulation, AgMIP, crop model

1. Introduction

2 Projecting crop yield response to a changing climate is of
3 great importance, especially as the global food production sys-
4 tem will face pressure from increased demand over the next
5 century. Climate-related reductions in supply could therefore
6 have severe socioeconomic consequences. Multiple studies
7 with different crop or climate models predict sharp reduction in
8 yields on currently cultivated cropland under business-as-usual
9 climate scenarios, although their yield projections show consid-
10 erable spread (e.g. Porter et al. (IPCC), 2014, Rosenzweig et al.,
11 2014, Schauberger et al., 2017, and references therein). Model
12 differences are unsurprising because crop responses in models
13 can be complex, with crop growth a function of complex inter-
14 actions between climate inputs and management practices.

15 Computational Models have been used to project crop yields
16 since the 1950's, beginning with statistical models (Heady,
17 1957, Heady & Dillon, 1961) that attempt to capture the rela-
18 tionship between input factors and resultant yields. These sta-
19 tistical models were typically developed on a small scale for lo-
20 cations with extensive histories of yield data. The emergence of
21 computers allowed development of numerical models that sim-
22 ulate the process of photosynthesis and the biology and phe-
23 nology of individual crops (first proposed by de Wit (1957),
24 Duncan et al. (1967) and attempted by Duncan (1972)). For a
25 history of crop model development see the appendix of Rosen-
26 zweig et al. (2014). A half-century of improvement in both
27 models and computing resources means that researchers can
28 now run crop simulation models for many years at high spatial
29 resolution on the global scale.

30 Both types of models continue to be used, and compara-
31 tive studies have concluded that when done carefully, both ap-
32 proaches can provide similar yield estimates (e.g. Lobell &

33 Burke, 2010, Moore et al., 2017, Roberts et al., 2017, Zhao
34 et al., 2017). Models tend to agree broadly in major response
35 patterns, including a reasonable representation of the spatial
36 pattern in historical yields of major crops (e.g. Elliott et al.,
37 2015, Müller et al., 2017) and projections of decreases in yield
38 under future climate scenarios.

Process models do continue to struggle with some important details, including reproducing historical year-to-year variability (e.g. Müller et al., 2017), reproducing historical yields when driven by reanalysis weather (e.g. Glotter et al., 2014), and low sensitivity to extreme events (e.g. Glotter et al., 2015). These issues are driven in part by the diversity of new cultivars and genetic variants, which outstrips the ability of academic modeling groups to capture them (e.g. Jones et al., 2017). Models do not simulate many additional factors affecting production, including pests/diseases/weeds. For these reasons, individual studies must generally re-calibrate models to ensure that short-term predictions reflect current cultivar mixes, and long-term projections retain considerable uncertainty (Wolf & Oijen, 2002, Jagtap & Jones, 2002, Iizumi et al., 2010, Angulo et al., 2013, Asseng et al., 2013, 2015). Inter-model discrepancies can also be high in areas not yet cultivated (e.g. Challinor et al., 2014, White et al., 2011). Finally, process-based models present additional difficulties for high-resolution global studies because of their complexity and computational requirements. For economic impacts assessments, it is often impossible to integrate a set of process-based crop models directly into an integrated assessment model to estimate the potential cost of climate change to the agricultural sector.

Nevertheless, process-based models are necessary for understanding the global future yield impacts of climate change for many reasons. First, cultivation may shift to new areas, where no yield data are currently available and therefore statistical models cannot apply. Yield data are also often limited in the de-

*Corresponding author at: 4734 S Ellis, Chicago, IL 60637, United States.
email: jfranke@uchicago.edu

67 developing world, where future climate impacts may be the most
 68 critical. Second, only process-based models can capture the
 69 growth response to elevated CO₂, novel conditions that are not
 70 represented in historical data (e.g. Pugh et al., 2016, Roberts
 71 et al., 2017). Similarly process-based models can represent
 72 novel changes in management practices (e.g. fertilizer input)
 73 that may ameliorate climate-induced damages.

74 Statistical emulation of crop simulations has been used to
 75 combine advantageous features of both statistical and process-
 76 based models. The statistical representation of complicated nu-
 77 matical simulation (e.g. O'Hagan, 2006, Conti et al., 2009), in
 78 which simulation output acts as the training data for a statisti-
 79 cal model, has been of increasing interest with the growth of
 80 simulation complexity and volume of output. Such emulators
 81 or "surrogate models" have been used in a variety of fields in-
 82 cluding hydrology (e.g. Razavi et al., 2012), engineering (e.g.
 83 Storlie et al., 2009), environmental sciences (e.g. Ratto et al.,
 84 2012), and climate (e.g. Castruccio et al., 2014, Holden et al.,
 85 2014). For agricultural impacts studies, emulation of process-
 86 based models allows exploring crop yields in regions outside
 87 ranges of current cultivation and with input variables outside
 88 historical precedents, in a lightweight, flexible form that is com-
 89 patible with economic studies.

90 In the past decade, many studies have developed emulators of
 91 crop yields from process-based models. Early studies propos-
 92 ing or describing potential emulators include Howden & Crimp
 93 (2005), Räisänen & Ruokolainen (2006) and Lobell & Burke
 94 (2010). In an early application, Ferrise et al. (2011) used a Arti-
 95 ficial Neural Net trained on simulation outputs to predict wheat₁₀₁
 96 yields in the Mediterranean. Studies developing single-model₁₀₂
 97 emulators include Holzkämper et al. (2012) for the CropSyst₁₀₃
 98 model, Ruane et al. (2013) for the CERES wheat model, Oye-₁₀₄
 99 bamiji et al. (2015) for the LPJmL model (for multiple crops,₁₀₅
 100 using multiple scenarios as a training set). In recent years, emu-₁₀₆




Figure 1: Presently cultivated area for rain-fed crops in the real world. Blue contour area indicates grid cells with more than 20,000 hectares (approx. 10% of the land in a grid cell near the equator for example) of crop cultivated. Gray contour shows area with more than 10 hectares cultivated. Cultivated areas for maize, rice, and soy are taken from the MIRCA2000 ("monthly irrigated and rain-fed crop areas around the year 2000") dataset (Portmann et al., 2010). Areas for winter and spring wheat areas are adapted from MIRCA2000 data and sorted by growing season. For analogous figure of irrigated crops, see Figure S1.

lators have begun to be used in the context of multi-model inter-
 comparisons, with Blanc & Sultan (2015), Blanc (2017), Ost-
 berg et al. (2018) and Mistry et al. (2017) using them to analyze
 the five crop models of the Inter-Sectoral Impacts Model Inter-
 comparison Project (ISIMIP) (Warszawski et al., 2014) (for
 maize, soy, wheat, and rice). Approaches differ: Blanc & Sultan

107 (2015) and Blanc (2017) used local weather variables (and CO₂)₁₃₅ Intercomparison and Improvement Project (AgMIP) (Rosen-
 108 values) and yields but emulate across soil types using historical₁₃₆ zweig et al., 2013, 2014), an international effort conducted un-
 109 simulations and a future climate scenario (RCP8.5 over mul-₁₃₇ der a framework similar to the Climate Model Intercomparison
 110 tiple climate models); Ostberg et al. (2018) used global mean₁₃₈ Project (CMIP) (Taylor et al., 2012, Eyring et al., 2016). The
 111 temperature change (and CO₂) as regressors but pattern-scale₁₃₉ GGCMI protocol builds on the AgMIP Coordinated Climate-
 112 to emulate local yields using multiple climate scenarios; Mis-₁₄₀ Crop Modeling Project (C3MP) (Ruane et al., 2014, McDer-
 113 try et al. (2017) used local weather and yields and a historical₁₄₁ mid et al., 2015) and will contribute to the AgMIP Coordinated
 114 simulation and compare with data.₁₄₂ Global and Regional Assessments (CGRA) (Ruane et al., 2018,
 142 Rosenzweig et al., 2018).

115 Those studies all temporal data in climate runs and attempt₁₄₃ GGCMI Phase II is designed to allow addressing goals such
 116 to untangle dependences.₁₄₄ as understanding where highest-yield regions may shift un-
 117 Recently efforts have been made to generate datasets that al-₁₄₅ der climate change; exploring future adaptive management
 118 low more systematic sampling of the input variable space (the₁₄₆ strategies; understanding how interacting parameters affect
 119 focus of this study): Makowski et al. (2015) for temperature,₁₄₇ crop yield; quantifying uncertainties across models and major
 120 CO₂, and nitrogen, Pirttioja et al. (2015) and Snyder et al.₁₄₈ drivers; and testing strategies for producing lightweight emu-
 121 (2018) for temperature, water, and CO₂, and (Fronzek et al.,₁₄₉ lators of process-based models. In this paper, we describe the
 122 2018) for temperature and water, with all studies simulating se-₁₅₀ GGCMI Phase II experiments, summarize output and present
 123 lected sites for a limited number of crops. The use of limited₁₅₁ initial results, demonstrate that it is tractable to emulation, and
 124 input parameter space or restricted geographic scope may im-₁₅₂ present a simple climatological emulator as a potential tool for
 125 pede the ability to build future projections and to understand₁₅₃ impacts assessments.₁₅₄
 126 interaction effects in global process-based crop models.₁₅₄

The Global Gridded Crop Model Intercomparison (GGCMI)

127 Phase II experiment seeks to provide a comprehensive global₁₅₅ 2. Materials and Methods
 128 dataset to allow systematically exploring how process-based
 129 crop models for the major crop respond to the main climate₁₅₆
 130 and management drivers and their interactions. The experiment₁₅₇
 131 involves running a suite of process-based crop models across₁₅₈ 2.1. GGCMI Phase II: experiment design
 132 historical conditions perturbed by a set of defined input pa-₁₅₉ GGCMI Phase II is the continuation of a multi-model com-
 133 rameters, and was conducted as part of the Agricultural Model₁₆₀ parison exercise begun in 2014. The initial Phase I compared
 134 (1980-2010) scenario with a primary goal of model evaluation

2. Materials and Methods

2.1. GGCMI Phase II: experiment design

GGCMI Phase II is the continuation of a multi-model comparison exercise begun in 2014. The initial Phase I compared harmonized yields of 21 models for 19 crops over a historical (1980-2010) scenario with a primary goal of model evaluation

Input variable	Abbr.	Tested range	Unit
CO ₂	C	360, 510, 660, 810	ppm
Temperature	T	-1, 0, 1, 2, 3, 4, 5*, 6	°C
Precipitation	W	-50, -30, -20, -10, 0, 10, 20, 30, (and W _{inf})	%
Applied nitrogen	N	10, 60, 200	kg ha ⁻¹

Table 1: Phase II input variable test levels. Temperature and precipitation values indicate the perturbations from the historical, climatology. * Only simulated by one model. W-percentage does not apply to the irrigated (W_{inf}) simulations.

(Elliott et al., 2015, Müller et al., 2017). Phase II compares simulations of 12 models for 5 crops (maize, rice, soybean, spring wheat, and winter wheat) over hundreds of scenarios in which individual climate or management inputs are adjusted from their historical values. The reduced set of crops includes the three major global cereals and the major legume and accounts for over 50% of human calories (in 2016, nearly 3.5 billion tons or 32% of total global crop production by weight (Food and Agriculture Organization of the United Nations, 2018).

The major goals of GGCMI Phase II are to:

- Enhance understanding of how models work by characterizing their sensitivity to input climate and nitrogen drivers.
- Study the interactions between climate variables and nitrogen inputs in driving modeled yield impacts.
- Explore differences in crop response to warming across the Earth's climate regions.
- Provide a dataset that allows statistical emulation of crop model responses for downstream modelers.
- Illustrate differences in potential adaptation via growing season changes.

The guiding scientific rationale of GGCMI Phase II is to provide a comprehensive, systematic evaluation of the response of process-based crop models to different values for carbon dioxide, temperature, water, and applied nitrogen (collectively known as "CTWN"). Phase II of the GGCMI project consists of a series of simulations, each with one or more of the CTWN dimensions perturbed over the 31-year historical time series (1980-2010) used in Phase I. In most cases, historical daily climate inputs are taken from the 0.5 degree NASA AgMERRA daily gridded re-analysis product specifically designed for agricultural modeling, with satellite-corrected precipitation (Ruane et al., 2015). Two models require sub-daily input data and use

alternative sources. See Elliott et al. (2015) for additional details.

The experimental protocol consists of 9 levels for precipitation perturbations, 7 for temperature, 4 for CO₂, and 3 for applied nitrogen, for a total of 672 simulations for rain-fed agriculture and an additional 84 for irrigated (Table 1). For irrigated simulations, soil water is held at either field capacity or, for those models that include water-log damage, at maximum beneficial level. Temperature perturbations are applied as absolute offsets from the daily mean, minimum, and maximum temperature time series for each grid cell used as inputs. Precipitation perturbations are applied as fractional changes at the grid cell level, and carbon dioxide and nitrogen levels are specified as discrete values applied uniformly over all grid cells. Note that CO₂ changes are applied independently of changes in climate variables, so that higher CO₂ is not associated with higher temperatures. An additional, identical set of scenarios (at the same C, T, W, and N levels) simulate adaptive agronomy under climate change by varying the growing season for crop production. (These adaptation simulations are not shown or analyzed here.) The resulting GGCMI data set captures a distribution of crop responses over the potential space of future climate conditions.

The 12 models included in GGCMI Phase II are all mechanistic process-based crop models that are widely used in impacts assessments (Table 2). Although some of the models shares a common base (e.g. LPJmL and LPJ-GUESS and the EPIC models), they have developed independently from this shared base, for more details on the genealogy of the models see Figure S1 in Rosenzweig et al. (2014). Differences in model structure does mean that several key factors are not standardized across the experiment, including secondary soil nutrients, carry over effects across growing years including residue management and soil moisture, and extent of simulated area for

Model (Key Citations)	Maize	Soy	Rice	Winter Wheat	Spring Wheat	N Dim.	Simulations per Crop
APSIM-UGOE , Keating et al. (2003), Holzworth et al. (2014)	X	X	X	–	X	Yes	37
CARAIB , Dury et al. (2011), Pirttioja et al. (2015)	X	X	X	X	X	No	224
EPIC-IIASA , Balkovi et al. (2014)	X	X	X	X	X	Yes	35
EPIC-TAMU , Izaurrealde et al. (2006)	X	X	X	X	X	Yes	672
JULES* , Osborne et al. (2015), Williams & Falloon (2015), Williams et al. (2017)	X	X	X	–	X	No	224
GEPIC , Liu et al. (2007), Folberth et al. (2012)	X	X	X	X	X	Yes	384
LPJ-GUESS , Lindeskog et al. (2013), Olin et al. (2015)	X	–	–	X	X	Yes	672
LPJmL , von Bloh et al. (2018)	X	X	X	X	X	Yes	672
ORCHIDEE-crop , Valade et al. (2014)	X	–	X	–	X	Yes	33
pDSSAT , Elliott et al. (2014), Jones et al. (2003)	X	X	X	X	X	Yes	672
PEPIC , Liu et al. (2016a,b)	X	X	X	X	X	Yes	130
PROMET*† , Mauser & Bach (2015), Hank et al. (2015), Mauser et al. (2009)	X	X	X	X	X	Yes†	239
Totals	12	10	11	9	12	–	3993 (maize)

Table 2: Models included in the GGCMI Phase II and the number of C, T, W, and N simulations performed for rain-fed crops (“Sims per Crop”), with 672 as the maximum. “N-Dim.” indicates if simulations include varying nitrogen levels; two models omit this dimension. All models provided the same set of simulations across all modeled crops, but some omitted individual crops in some cases. (For example, APSIM did not simulate winter wheat.) Irrigated simulations are provided at the level of the other covariates for each model (for an additional 84 simulations for the fully-sampled models). Geographic extent of simulation varies to some extent within a certain model for different scenarios (672 rain-fed simulations does not necessarily equal 672 climatological yields in all areas). This geographic variance only applies for areas far outside the area of currently cultivated crops. Two models (marked with *) use non-AgMERRA climate inputs. For further details on models, see Elliott et al. (2015). †PROMET provided simulations at only two nitrogen levels so is not emulated across the nitrogen dimension.

227 different crops. Growing seasons are identical across models,²⁴⁵
 228 but vary by crop and by location on the globe. All stresses²⁴⁶
 229 except factors related to nitrogen, temperature, and water (e.g.²⁴⁷
 230 Alkalinity, salinity) are disabled. No additional nitrogen inputs,²⁴⁸
 231 such as atmospheric deposition, are considered, but some mod-²⁴⁹
 232 els have individual assumptions on soil organic matter that may²⁵⁰
 233 release additional nitrogen through mineralization. See Rosen-²⁵¹
 234 zweig et al. (2014), Elliott et al. (2015) and Müller et al. (2017)²⁵²
 235 for further details on models and underlying assumptions.²⁵³

236 Each model is run at 0.5 degree spatial resolution and covers²⁵⁴
 237 all currently cultivated areas and much of the uncultivated land²⁵⁵
 238 area. Coverage extends considerably outside currently culti-²⁵⁶
 239 vated areas because cultivation will likely shift under climate²⁵⁷
 240 change. See Figure 1 for the present-day cultivated area of²⁵⁸
 241 rain-fed crops, and Figure S1 in the supplemental material for²⁵⁹
 242 irrigated crops. Some areas such as Greenland, far-northern²⁶⁰
 243 Canada, Siberia, Antarctica, the Gobi and Sahara deserts, and²⁶¹
 244 central Australia are not simulated as they are assumed to re-²⁶²

main non-arable even under an extreme climate change. Grow-
 ing seasons are standardized across models with data adapted
 from several sources (Sacks et al., 2010, Portmann et al., 2008,
 2010).

The participating modeling groups provide simulations at
 any of four initially specified levels of participation, so the num-
 ber of simulations varies by model, with some sampling only a
 part of the experiment variable space. Most modeling groups
 simulate all five crops in the protocol, but some omitted one
 or more. Table 2 provides details of coverage for each model.
 Note that the three models that provide less than 50 simulations
 are excluded from the emulator analysis.

All models produce as output, crop yields ($\text{tons ha}^{-1} \text{ year}^{-1}$)
 for each 0.5 degree grid cell. Because both yields and yield
 changes vary substantially across models and across grid cells,
 we primarily analyze relative change from a baseline. We take
 as the baseline the scenario with historical climatology (i.e. T
 and P changes of 0), C of 360 ppm, and applied N at 200 kg

263 ha⁻¹. We show absolute yields in some cases to illustrate geo-
264 graphic differences in yields for a single model.

265 2.2. Simulation model validation approach

266 To verify the skill of the process-based models used, we re-
267 peat the validation exercises presented in Müller et al. (2017)
268 for GGCMI Phase I. Note however that the GGCMI Phase II
269 simulations are designed for evaluating changes in yield but not
270 absolute yields, and so omit the calibrations used in predict-
271 ing modeling to account for cultivar, pest loss, and manage-
272 ment differences. The Phase II simulations also do not repro-
273 duce realistic nitrogen application levels for individual coun-
274 tries, since nitrogen is one of the parameters systematically var-
275 ied. The Müller et al. (2017) validation procedure evaluates re-
276 sponse to year-to-year temperature and precipitation variations
277 in a control run driven by historical climate and compares it
278 to detrended historical yields from the FAO (Food and Agri-
279 culture Organization of the United Nations, 2018) by calculat-
280 ing the Pearson correlation coefficient. The procedure offers no
281 means of assessing CO₂ fertilization, since CO₂ has been rel-
282 atively constant over the historical data collection period. Ni-
283 trogen data are limited for many countries, and as mentioned
284 the GGCMI Phase II runs impose fixed and uniform nitrogen
285 application, introducing some uncertainty into the analysis. We
286 evaluate one or more control runs for each model, since some
287 modeling groups provide historical runs for three different ni-
288 trogen levels.

289 2.3. Climatological-mean yield emulator design

290 To demonstrate the properties of the GGCMI Phase II
291 dataset, we construct an emulator of 30-year climatological
292 mean yields, which are of most interest to impact modelers.
293 This approach differs from previous studies of crop model em-
294 ulation, which have typically emulated at the annual level. An-
295 nual emulation is required when the input training set consists




Figure 2: Example showing both climatological mean yields and distribution of yearly yields for three 30-year scenarios. Figure shows rain-fed maize for a grid cell in northern Iowa (a representative high-yield region) from the pDSSAT model, for the baseline climatology (1981–2010) and for scenarios with temperature shifted by (T) +3 and (T) +6 °C, with other variables held at baseline values. Dashed vertical lines and black dots indicate the climatological mean yield.




Figure 3: Example showing simple temperature relationship developed from year-to-year values vs. climatological mean values. Figure shows irrigated maize for four adjacent grid cell in northern Iowa (a representative high-yield region) from the pDSSAT model, for the baseline climatology (1981–2010) and for scenarios with temperature shifted (T) +6 °C, with other variables held at baseline values. Irrigated yields are shown to control for precipitation effects. Blue and red lines indicate total least squares linear regression across each temper-
ature scenario. Black dots indicate the climatological mean yield values for each climatological temperature scenario.

296 of non-stationary projections of evolving yields (such as an³¹³ efforts.

297 RCP run). Recent studies (e.g. Fronzek et al., 2018, ?) that used

298 a training set of stationary simulations with fixed variations in³¹⁴
 299 parameters allow emulating the climatological mean response³¹⁵
 300 instead. The two can differ for multiple reasons, including any³¹⁶
 301 year-to-year memory in the crop model, or if the distribution of³¹⁷
 302 growing-season daily temperatures associated with interannual³¹⁸
 303 variability is different from that associated with long-term CO₂-³¹⁹
 304 driven changes. Note that the GGCMI Phase II datasets will³²⁰
 305 not capture distributional shifts, because all simulations are run³²¹
 306 with fixed offsets from the historical climatology. (For meth-³²²
 307 ods to generate adjust historical climate data inclusive of dis-³²³
 308 tributional changes, see XX and XX). The year-over-year yield³²⁴
 309 response to individual factors in GGCMI Phase II nevertheless³²⁵
 310 often exceeds the climatological response (Figure 3). Emula-³²⁶
 311 tion approaches are an area of active ongoing study and one of³²⁷
 312 the goals of the GGCMI Phase II dataset is to facilitate these³²⁸

Emulation involves fitting individual regression models for each crop, simulation model, and 0.5 degree geographic pixel from the GGCMI Phase II data set. The regressors are the applied constant perturbations in temperature, water, nitrogen and CO₂, we aggregate the simulation outputs in the time dimension, and regress on the 30-year mean yields. (See Figure 2 for illustration). The regression therefore omits information about yield responses to year-to-year climate perturbations, which are more complex. Emulating inter-annual yield variations would likely require considering statistical details of the historical climate time series, including changes in marginal distribution and temporal dependencies. (Future work should explore this). The climatological emulation indirectly includes any yield response to geographically distributed factors such as soil type, insolation, and the baseline climate itself, because we construct separate emulators for each grid cell. The emulator parameter ma-

329




Figure 4: Illustration of the distribution of regional yield changes across the multi-model ensemble, split by Köppen-Geiger climate regions (Rubel & Kottek (2010)). We show responses of a single crop (rain-fed maize) to applied uniform temperature perturbations, for three discrete precipitation perturbation levels (rain-fed (rf) -20%, rain-fed (0), and rain-fed +20%), with CO₂ and nitrogen held constant at baseline values (360 ppm and 200 kg ha⁻¹ yr⁻¹). Y-axis is fractional change in the regional average climatological potential yield relative to the baseline. The figure shows all modeled land area; see Figure S6 in the supplemental material for only currently-cultivated land. Panel text gives the percentage of rain-fed maize presently cultivated in each climate zone (Portmann et al., 2010). Box-and-whiskers plots show distribution across models, with median marked. Box edges are first and third quartiles, i.e. box height is the interquartile range (IQR). Whiskers extend to maximum and minimum of simulations but are limited at 1.5-IQR; otherwise the outlier is shown. Models generally agree in most climate regions (other than cold continental), with projected changes larger than inter-model variance. Outliers in the tropics (strong negative impact of temperature increases) are the pDSSAT model; outliers in the high-rainfall case (strong positive impact of precipitation increases) are the JULES model. Inter-model variance increases in the case where precipitation is reduced, suggesting uncertainty in model response to water limitation. The right panel with global changes shows yield responses to an globally uniform temperature shift; note that these results are not directly comparable to simulations of more realistic climate scenarios with the same global mean change.

trices are portable and the yield computations are cheap even at³⁶⁴
 the half-degree grid cell resolution, so we do not aggregate in³⁶⁵
 space at this time.³⁶⁶

We regress climatological mean yields against a third-order³⁶⁷
 polynomial in C, T, W, and N with interaction terms. The³⁶⁸
 higher-order terms are necessary to capture any nonlinear re-³⁶⁹
 sponds, which are well-documented in observations for tem-³⁷⁰
 perature and water perturbations (e.g. Schlenker & Roberts³⁷¹
 (2009) for T and He et al. (2016) for W). We include inter-³⁷²
 action terms (both linear and higher-order) because past stud-³⁷³
 ies have shown them to be significant effects. For example,³⁷⁴
 Lobell & Field (2007) and Tebaldi & Lobell (2008) showed³⁷⁵
 that in real-world yields, the joint distribution in T and W is³⁷⁶
 needed to explain observed yield variance. (C and N are fixed³⁷⁷
 in these data.) Other observation-based studies have shown the³⁷⁸
 importance of the interaction between water and nitrogen (e.g.³⁷⁹
 Aulakh & Malhi, 2005), and between nitrogen and carbon diox-³⁸⁰
 ide (Osaki et al., 1992, Nakamura et al., 1997). We do not fo-³⁸¹
 cuse on comparing different functional forms in this study, and³⁸²
 instead choose a relatively simple parametrization that allows³⁸³
 for some interpretation of coefficients. Some prior studies have³⁸⁴
 used more complex functional forms and larger numbers of pa-³⁸⁵
 rameters, e.g. 39 in Blanc & Sultan (2015) and Blanc (2017),³⁸⁶
 who borrow information across space by fitting grid points si-
 multaneously across a large region in a panel regression. **We**
choose a simpler emulation at grid-cell level to avoid the
requirement of assuming responses are uniform across space and
to maximize interpretability.³⁸⁷

The limited GGCMI variable sample space means that use³⁵⁸
 of the full polynomial expression described above, which has³⁵⁹
 34 terms for the rain-fed case (12 for irrigated), can be prob-³⁶⁰
 lematic, and can lead to over-fitting and unstable parameter es-³⁶¹
 timations. We therefore reduce the number of terms through a³⁶²
 feature selection cross-validation process in which terms in the³⁶³

polynomial are tested for importance. In this procedure higher-³⁶⁴
 order and interaction terms are added successively to the model;³⁶⁵
 we then follow the reduction of the the aggregate mean squared³⁶⁶
 error with increasing terms and eliminate those terms that do³⁶⁷
 not contribute significant reductions. See supplemental docu-³⁶⁸
 ments for more details. We select terms by applying the feature³⁶⁹
 selection process to the three models that provided the com-³⁷⁰
 plete set of 672 rain-fed simulations (pDSSAT, EPIC-TAMU,³⁷¹
 and LPJmL); the resulting choice of terms is then applied for³⁷²
 all emulators.³⁷³

Feature importance is remarkably consistent across all three³⁷⁴
 models and across all crops (see Figure S4 in the supplemental³⁷⁵
 material). The feature selection process results in a final poly-³⁷⁶
 nomial in 23 terms, with 11 terms eliminated. We omit the N³³⁷⁷
 term, which cannot be fitted because we sample only three ni-³⁷⁸
 trogen levels. We eliminate many of the C terms: the cubic,³⁷⁹
 the CT, CTN, and CWN interaction terms, and all higher order³⁸⁰
 interaction terms in C. Finally, we eliminate two 2nd-order in-³⁸¹
 teraction terms in T and one in W. Implication of this choice³⁸²
 include that nitrogen interactions are complex and important,³⁸³
 and that water interaction effects are more nonlinear than those³⁸⁴
 in temperature. The resulting statistical model (Equation 1) is³⁸⁵
 used for all grid cells, models, and crops:³⁸⁶

$$\begin{aligned}
 Y = & K_1 & (1) \\
 & + K_2 C + K_3 T + K_4 W + K_5 N \\
 & + K_6 C^2 + K_7 T^2 + K_8 W^2 + K_9 N^2 \\
 & + K_{10} C W + K_{11} C N + K_{12} T W + K_{13} T N + K_{14} W N \\
 & + K_{15} T^3 + K_{16} W^3 + K_{17} T W N \\
 & + K_{18} T^2 W + K_{19} W^2 T + K_{20} W^2 N \\
 & + K_{21} N^2 C + K_{22} N^2 T + K_{23} N^2 W
 \end{aligned}$$

To fit the parameters K , we use a Bayesian Ridge probabilis-

tic estimator (MacKay, 1991), which reduces volatility in parameter estimates when the sampling is sparse, by weighting parameter estimates towards zero. The Bayesian Ridge method is necessary to maintain a consistent functional form across all models, and locations as the linear least squares fails to provide a stable result in many cases. In the GGCMI Phase II experiment, the most problematic fits are those for models that provided a limited number of cases or for low-yield geographic regions where some modeling groups did not run all scenarios. Because we do not attempt to emulate models that provided less than 50 simulations, the lowest number of simulations emulated across the full parameter space is 130 (for the PEPIC model). We use the implementation of the Bayesian Ridge emulator from the scikit-learn package in Python (Pedregosa et al., 2011).

The resulting parameter matrices for all crop model emulators are available on request, as are the raw simulation data and a Python application to emulate yields. The yield output for a single GGCMI model that simulates all scenarios and all five crops is ~ 12.5 GB; the emulator is ~ 100 MB, a reduction by over two orders of magnitude.

2.4. Emulator evaluation

Because no general criteria exist for defining an acceptable model emulator, we develop a metric of emulator performance specific to GGCMI. For a multi-model comparison exercise like GGCMI, a reasonable criterion is what we term the “normalized error”, which compares the fidelity of an emulator for a given model and scenario to the inter-model uncertainty. We define the normalized error e for each scenario as the difference between the fractional yield change from the emulator and that in the original simulation, divided by the standard deviation of the multi-model spread (Equations 2 and 3):

$$F_{scn.} = \frac{Y_{scn.} - Y_{baseline}}{Y_{baseline}} \quad (2)$$

$$e_{scn.} = \frac{F_{em, scn.} - F_{sim, scn.}}{\sigma_{sim, scn.}} \quad (3)$$

Here $F_{scn.}$ is the fractional change in a model’s mean emulated or simulated yield from a defined baseline, in some scenario (scn.) in C, T, W, and N space; $Y_{scn.}$ and $Y_{baseline}$ are the absolute emulated or simulated mean yields. The normalized error e is the difference between the emulated fractional change in yield and that actually simulated, normalized by $\sigma_{sim.}$, the standard deviation in simulated fractional yields $F_{sim, scn.}$ across all models. The emulator is fit across all available simulation outputs, and then the error is calculated across the simulation scenarios provided by all nine models (Figure 9 and Figures S12 and Figures S13 in supplemental documents). Note that the normalized error e for a model depends not only on the fidelity of its emulator in reproducing a given simulation but on the particular suite of models considered in the intercomparison exercise. The rationale for this choice is to relate the fidelity of the emulation to an estimate of true uncertainty, which we take as the multi-model spread.

3. Results

3.1. Simulation results

Crop models in the GGCMI ensemble show a broadly consistent responses to climate and management perturbations in most regions, with a strong negative impact of increased temperature in all but the coldest regions. We illustrate this result for rain-fed maize in Figure 4, which shows yields for the primary Köppen-Geiger climate regions (Rubel & Kottek, 2010). In warming scenarios, models show decreases in maize yield in the temperate, tropical, and arid regions that account for nearly three-quarters of global maize production. These impacts are




Figure 5: Illustration of the spatial pattern of potential yields and potential yield changes in the GGCMI Phase II ensemble, for three major crops. Left column (a) shows multi-model mean climatological yields for the baseline scenario for (top–bottom) rain-fed maize, soy, and rice. (For wheat see Figure S11 in the supplemental material.) White stippling indicates areas where these crops are not currently cultivated. Absence of cultivation aligns well with the lowest yield contour (0.2 ton ha^{-1}). Right column (b) shows the multi-model mean fractional yield change in the extreme $T + 4 \text{ }^{\circ}\text{C}$ scenario (with other inputs at baseline values). Areas without hatching or stippling are those where confidence in projections is high: the multi-model mean fractional change exceeds two standard deviations of the ensemble. ($\Delta > 2\sigma$). Hatching indicates areas of low confidence ($\Delta < 1\sigma$), and stippling areas of medium confidence ($1\sigma < \Delta < 2\sigma$). Crop model results in cold areas, where yield impacts are on average positive, also have the highest uncertainty.

robust for even moderate climate perturbations. In the temperate zone, even a 1 degree temperature rise with other variables held fixed leads to a median yield reduction that outweighs the variance across models. A 6 degree temperature rise results in median loss of $\sim 25\%$ of yields with a signal to noise of nearly three. A notable exception is the cold continental region, where models disagree strongly, extending even to the sign of impacts. Model simulations of other crops produce similar responses to warming, with robust yield losses in warmer locations and high inter-model variance in the cold continental regions (Figures S7).

The effects of rainfall changes on maize yields are also as expected and are consistent across models. Increased rainfall mit-

igates the negative effect of higher temperatures, most strongly in arid regions. Decreased rainfall amplifies yield losses and also increases inter-model variance more strongly, suggesting that models have difficulty representing crop response to water stress. We show only rain-fed maize here; see Figure S5 for the irrigated case. As expected, irrigated crops are more resilient to temperature increases in all regions, especially so where water is limiting.

Mapping the distribution of baseline yields and yield changes shows the geographic dependencies that underlie these results. Figure 5 shows baseline and changes in the $T+4$ scenario for rain-fed maize, soy, and rice in the multi-model ensemble mean, with locations of model agreement marked. Absolute yield po-

tentials are have strong spatial variation, with much of the₄₇₄
Earth's surface area unsuitable for any given crop. In general,₄₇₅
models agree most on yield response in regions where yield₄₇₆
potentials are currently high and therefore where crops are cur-₄₇₇
rently grown. Models show robust decreases in yields at low₄₇₈
latitudes, and highly uncertain median increases at most high₄₇₉
latitudes. For wheat crops see Figure S11; wheat projections₄₈₀
are both more uncertain and show fewer areas of increased yield₄₈₁
in the inter-model mean.₄₈₂

₄₈₃ 3.2. *Simulation model validation results*

Figure 6 shows the Pearson time series correlation between
the simulation model yield and FAO yield data. Figure 6 can be
compared to Figures 1,2,3,4 and 6 in Müller et al. (2017). The
results are mixed, with many regions for rice and wheat be-
ing difficult to model. No single model is dominant, with each
model providing near best-in-class performance in at least one
location-crop combination. The presence of very few vertical
dark green color bars clearly illustrates the power of a multi-
model intercomparison project like the one presented here. The
ensemble mean does not beat the best model in each case, but
shows positive correlation in over 75% of the cases presented
here. The EPIC-TAMU model performs best for soy, CARIAB,
EPIC-TAMU, and PEPIC perform best for maize, PROMET
performs best for wheat, and the EPIC family of models per-
form best for rice. Reductions in skill over the performance
illustrated in Müller et al. (2017) can be attributed to the nitro-₄₉₉
gen levels or lack of calibration in some models.₅₀₀

₅₀₁ *** or harmonization *** Christoph

Soy is qualitatively the easiest crop to represent (except in₅₀₂
Argentina), which is likely due in part to the invariance of the₅₀₃
response to nitrogen application (soy fixes atmospheric nitrogen₅₀₄
very efficiently). Comparison to the FAO data is therefore easier₅₀₅
than the other crops because the nitrogen application levels do₅₀₆
not matter. US maize has the best performance across models,₅₀₇
₅₀₈

with nearly every model representing the historical variability
to a reasonable extent. Especially good example years for US
maize are 1983, 1988, and 2004 (top left panel of Figure 6),
where every model gets the direction of the anomaly compared
to surrounding years correct. 1983 and 1988 are famously bad
years for US maize along with 2012 (not shown). US maize
is possibly both the most uniformly industrialized (in terms of
management practices) crop and the one with the best data col-
lection in the historical period of all the cases presented here.₅₁₆

The FAO data is at least one level of abstraction from ground
truth in many cases, especially in developing countries. The
failure of models to represent the year-to-year variability in rice
in some countries in southeast Asia is likely partly due to model
failure and partly due to lack of data. It is possible to speculate
that the difference in performance between Pakistan (no suc-
cessful models) and India (many successful models) for rice
may reside at least in part in the FAO data and not the mod-
els themselves. The same might apply to Bangladesh and In-
dia for rice. Partitioning of these contributions is impossible at
this stage. Additionally, there is less year-to-year variability in
rice yields (partially due to the fraction of irrigated cultivation).
Since the Pearson r metric is scale invariant, it will tend to score
the rice models more poorly than maize and soy. An example
of very poor performance can be seen with the pDSSAT model
for rice in India (top right panel of Figure 6).₅₂₄

₅₂₅ 3.3. *Emulator performance*

Emulation provides not only a computational tool but a
means of understanding and interpreting crop yield response
across the parameter space. Emulation is only possible, how-
ever, when crop yield responses are sufficiently smooth and
continuous to allow fitting with a relatively simple functional
form. In the GGCMI simulations, this condition largely but
not always holds. Responses are quite diverse across locations,
crops, and models, but in most cases local responses are reg-₅₃₄




Figure 6: Time series correlation coefficients between simulated crop yield and FAO data (Food and Agriculture Organization of the United Nations, 2018) at the country level. The top panels indicate two example cases: US maize (a good case), and rice in India (mixed case), both for the high nitrogen application case. The heatmaps illustrate the Pearson r correlation coefficient between the detrended simulation mean yield at the country level compared to the detrended FAO yield data for the top producing countries for each crop with continuous FAO data over the 1980–2010 period. Models that provided different nitrogen application levels are shown with low, med, and high label (models that did not simulate different nitrogen levels are analogous to a high nitrogen application level). The ensemble mean yield is also correlated with the FAO data (not the mean of the correlations). Wheat contains both spring wheat and winter wheat simulations.

ular enough to permit emulation. Figure 7 illustrates the geographic diversity of responses even in high-yield areas for a single crop and model (rain-fed maize in pDSSAT for various high-cultivation areas). This heterogeneity validates the choice of emulating at the grid cell level.

Each panel in Figure 7 shows model yield output from scenarios varying only along a single dimension (CO_2 , temperature, precipitation, or nitrogen addition), with other inputs held fixed at baseline levels; in all cases yields evolve smoothly across the space sampled. For reference we show the results




Figure 7: Example illustrating spatial variations in yield response and emulation ability. We show rain-fed maize in the pDSSAT model in six example locations selected to represent high-cultivation areas around the globe. Legend includes hectares cultivated in each selected grid cell. Each panel shows variation along a single variable, with others held at baseline values. Dots show climatological mean yield, solid lines a simple polynomial fit across this 1D variation, and dashed lines the results of the full emulator of Equation 1. The climatological response is sufficiently smooth to be represented by a simple polynomial. Because precipitation changes are imposed as multiplicative factors rather than additive offsets, locations with higher baseline precipitation have a larger absolute spread in inputs sampled than do drier locations (not visible here).




Figure 8: Example illustrating across-model variations in yield response. We show simulations and emulations from six models for rain-fed maize in the same Iowa location shown in Figure 7, with the same plot conventions. Models that did not simulate the nitrogen dimension are omitted for clarity. The inter-model standard deviation is larger for the perturbations in CO₂ and nitrogen than for those in temperature or precipitation illustrating that the sensitivity to weather is better constrained than to management at elevated CO₂. While most model responses can readily be emulated with a simple polynomial, some response surfaces are more complicated and lead to emulation error.

552 of the full emulation fitted across the parameter space. The 557 polynomial fit readily captures the climatological response to 558 perturbations.

555 Crop yield responses generally follow similar functional 561 forms across models, though with a spread in magnitude. Fig-

ure 8 illustrates the inter-model diversity of yield responses to the same perturbations, even for a single crop and location (rain-fed maize in northern Iowa, the same location shown in the Figure 7). The differences make it important to construct emulators separately for each individual model, and the fidelity

of emulation can also differ across models. This figure illustrates a common phenomenon, that models differ more in response to perturbations in CO₂ and nitrogen perturbations than to those in temperature or precipitation. (Compare also Figures 4 and S18.) For this location and crop, CO₂ fertilization effects can range from ~5–50%, and nitrogen responses from nearly flat to a 60% drop in the lowest-application simulation.

While the nitrogen dimension is important and uncertain, it is also the most problematic to emulate in this work because of its limited sampling. The GGCMI protocol specified only three nitrogen levels (10, 60 and 200 kg N y⁻¹ ha⁻¹), so a third-order fit would be over-determined but a second-order fit can result in potentially unphysical results. Steep and nonlinear declines in yield with lower nitrogen levels means that some regressions imply a peak in yield between the 100 and 200 kg N y⁻¹ ha⁻¹ levels. While there may be some reason to believe over-application of nitrogen at the wrong time in the growing season could lead to reduced yields, these features are almost certainly an artifact of under sampling. In addition, the polynomial fit cannot capture the well-documented saturation effect of nitrogen application (e.g. Ingestad, 1977) as accurately as would be possible with a non-parametric model.

To assess the ability of the polynomial emulation to capture the behavior of complex process-based models, we evaluate the normalized emulator error. That is, for each grid cell, model, and scenario we evaluate the difference between the model yield and its emulation, normalized by the inter-model standard deviation in yield projections. This metric implies that emulation is generally satisfactory, with several distinct exceptions. Almost all model-crop combination emulators have normalized errors less than one over nearly all currently cultivated hectares (Figure 9), but some individual model-crop combinations are problematic (e.g. PROMET for rice and soy, JULES for soy and winter wheat, Figures S14–S15). Normalized errors for soy




Figure 9: Assessment of emulator performance over currently cultivated areas based on normalized error (Equations 3, 2). We show performance of all 9 models emulated, over all crops and all sampled T and P inputs, but with CO₂ and nitrogen held fixed at baseline values. Large columns are crops and large rows models; squares within are T,P scenario pairs. Colors denote the fraction of currently cultivated hectares with $e < 1$. Of the 756 scenarios with these CO₂ and N values, we consider only those for which all 9 models submitted data. JULES did not simulate spring wheat and LPJ-GUESS did not simulate rice and soy. Emulator performance is generally satisfactory, with some exceptions. Emulator failures (significant areas of poor performance) occur for individual crop-model combinations, with performance generally degrading for hotter and wetter scenarios.




Figure 10: Illustration of our test of emulator performance, applied to the CARAIB model for the T+4 scenario for rain-fed crops. Contour colors indicate the normalized emulator error e , where $e > 1$ means that emulator error exceeds the multi-model standard deviation. White areas are those where crops are not simulated by this model. Models differ in their areas omitted, meaning the number of samples used to calculate the multi-model standard deviation is not spatially consistent in all locations. Emulator performance is generally good relative to model spread in areas where crops are currently cultivated (compare to Figure 1) and in temperate zones in general; emulation issues occur primarily in marginal areas with low yield potentials. For CARAIB, emulation of soy is more problematic, as was also shown in Figure 9.

are somewhat higher across all models not because emulator fidelity is worse but because models agree more closely on yield changes for soy than for other crops (see Figure S16, lowering the denominator. Emulator performance often degrades in geographic locations where crops are not currently cultivated. Figure 10 shows a CARAIB case as an example, where emulator performance is satisfactory over cultivated areas for all crops other than soy, but uncultivated regions show some problematic areas.

It should be noted that this assessment metric is relatively forgiving. First, each emulation is evaluated against the simulation actually used to train the emulator. Had we used a spline interpolation the error would necessarily be zero. Second, the performance metric scales emulator fidelity not by the magnitude of yield changes but by the inter-model spread in those changes. Where models differ more widely, the standard for emulators becomes less stringent. Because models disagree on the magnitude of CO₂ fertilization, this effect is readily seen when comparing assessments of emulator performance in simulations at baseline CO₂ (Figure 9) with those at higher CO₂ levels (Figure S13). Widening the inter-model spread leads to an apparent increase in emulator skill.

3.4. Emulator applications

Because the emulator or “surrogate model” transforms the discrete simulation sample space into a continuous response surface at any geographic scale, it can be used for a variety of applications. Emulators provide an easy way to compare a ensemble of climate or impacts projections. They also provide a means for generating continuous damage functions. As an example, we show a damage function constructed from 4D emulations for aggregated yield at the global scale, for maize on currently cultivated land, with simulated values shown for comparison. (Figure 11; see Figures S16- S19 in the supplemental material for other crops and dimensions.) The emu-

630 lated values closely match simulations even at this aggregation level. Note that these functions are presented only as
 631 examples and do not represent true global projections, because they are developed from simulation data with a uniform
 632 temperature shift while increases in global mean temperature should manifest non-uniformly. The global coverage of the
 633 GGCMI simulations allows impacts modelers to apply arbitrary geographically-varying climate projections, as well as arbitrary
 634 aggregation mask, to develop damage functions for any climate scenario and any geopolitical or geographic level.
 635
 636
 637
 638
 639




Figure 11: Global emulated damages for maize on currently cultivated lands for the GGCMI models emulated, for uniform temperature shifts with other inputs held at baseline. (The damage function is created from aggregating up emulated values at the grid cell level, not from a regression of global mean yields.) Lines are emulations for rain-fed (solid) and irrigated (dashed) crops; for comparison, dots are the simulated values for the rain-fed case. For most models, irrigated crops show a sharper reduction than do rain-fed because of the locations of cultivated areas: irrigated crops tend to be grown in warmer areas where impacts are more severe for a given temperature shift. (The exceptions are PROMET, JULES, and LPJmL.) For other crops and scenarios see Figures S16–S19 in the supplemental material.

4. Conclusions and discussion

The GGCMI Phase II experiment provides a database targeted to allow detailed study of crop yields from process-based models under climate change. The experiment is designed to facilitate not only comparing the sensitivities of process-based crop yield models to changing climate and management inputs but also evaluating the complex interactions between driving factors (CO_2 , temperature, precipitation, and applied nitrogen). Its global nature also allows identifying geographic shifts in high yield potential locations. We expect that the simulations will yield multiple insights in future studies, and show here a selection of preliminary results to illustrate their potential uses.

First, the GGCMI Phase II simulations allow identifying major areas of uncertainty. Across the major crops, inter-model uncertainty is greatest for wheat and least for soy. Across factors impacting yields, inter-model uncertainty is largest for CO_2 fertilization and nitrogen response effects. Across geographic regions, projections are most uncertain in the high latitudes where yields may increase, and most robust in low latitudes where yield impacts are largest.

Second, the GGCMI Phase II simulations allow understanding the way that climate-driven changes and locations of cultivated land combine to produce yield impacts. One counterintuitive result immediate apparent is that irrigated maize shows steeper yield reductions under warming than does rain-fed maize when considered only over currently cultivated land. The effect results from geographic differences in cultivation. In any given location, irrigation increases crop resiliency to temperature increase, but irrigated maize is grown in warmer locations where the impacts of warming are more severe (Figures S5–S6). The same behavior holds for rice and winter wheat, but not for soy or spring wheat (Figures S8–S10). Irrigated wheat and maize are also more sensitive to nitrogen fertilization levels than are analogous non-irrigated crops, presumably

674 because those rain-fed crops are limited by water as well as⁷⁰⁸
675 nitrogen availability (Figure S19). (Soy as an efficient atmo-⁷⁰⁹
676 spheric nitrogen-fixing crop is relatively insensitive to nitrogen, and⁷¹⁰
677 rice is not generally grown in water-limited conditions).⁷¹¹

678 Third, we show that even the relatively limited GGCMI⁷¹²
679 Phase II sampling space allows emulation of the climatological⁷¹³
680 response of crop models with a relatively simple reduced-form⁷¹⁴
681 statistical model. The systematic parameter sampling in the⁷¹⁵
682 GGCMI Phase II procedure provides information on the influ-⁷¹⁶
683 ence of multiple interacting factors in a way that single projec-⁷¹⁷
684 tions cannot, and emulating the resulting response surface then⁷¹⁸
685 produces a tool that can aid in both physical interpretation of⁷¹⁹
686 the process-based models and in assessment of agricultural im-⁷²⁰
687 pacts under arbitrary climate scenarios. Emulating the climato-⁷²¹
688 logical response isolates long-term impacts from any confound-⁷²²
689 ing factors that complicate year-over-year changes, and the use⁷²³
690 of simple functional forms offer the possibility of physical in-⁷²⁴
691 terpretation of parameter values. We anticipate that systematic⁷²⁵
692 parameter sampling will become the norm in future model in-⁷²⁶
693 tercomparison exercise.⁷²⁷

694 While the GGCMI Phase II database should offer the foun-⁷²⁹
695 dation for multiple future studies, several cautions need to be
696 noted. Because the simulation protocol was designed to fo-⁷³⁰
697 cus on change in yield under climate perturbations and not
698 on replicating real-world yields, the models are not formally
699 calibrated so cannot be used for impacts projections unless in
700 used in conjunction with historical data (or data products). Be-
701 cause the GGCMI simulations apply uniform perturbations to
702 historical climate inputs, they do not sample changes in higher
703 order moments, and cannot address the additional crop yield⁷³⁶
704 impacts of potential changes in climate variability. Although⁷³⁷
705 distributional changes in model projections are fairly uncertain⁷³⁸
706 at present, follow-on experiments may wish to consider them.⁷³⁹
707 Several recent studies have described procedures for generating⁷⁴⁰

simulations that combine historical data with model projections
of not only mean changes in temperature and precipitation but
changes in their marginal distributions (cite) or temporal depen-
dence ().

— NEEDS EDIT – still theres lots of future work can be
done even with this database—

Detailed examination of interaction terms between the ma-
jor input drivers. More robust quantification of the sensitiv-
ity of different models to the input drivers. Comparison with
field-level experimental data can then aid in model evaluation.
As mentioned previously, database allows study of geographic
shifts in optimal growing regions for different crops. The out-
put dataset also contains other runs and variables not analyzed
or shown here. Runs include several which allowed adaptation
to climate changes by altering growing seasons, and additional
variables include above ground biomass, LAI, and root biomass
(as many as 25 output variables for some models). Emulation
studies that are possible include study of year-over-year vs cli-
matological emulation, and more systematic evaluation of dif-
ferent statistical model specifications and formal calculation of
uncertainties in derived parameters.

The future of food security is one of the larger challenges
facing humanity at present. The development (and emulation)
of multi-model ensembles such as GGCMI Phase II provides
a way to begin to quantify uncertainties in crop responses to a
range of potential climate inputs and explore the potential ben-
efits of adaptive responses.

5. Acknowledgments

We thank Michael Stein and Kevin Schwarzwald, who pro-
vided helpful suggestions that contributed to this work. This re-
search was performed as part of the Center for Robust Decision-
Making on Climate and Energy Policy (RDCEP) at the Univer-
sity of Chicago, and was supported through a variety of sources.

741 RDCEP is funded by NSF grant #SES-1463644 through the
 742 Decision Making Under Uncertainty program. J.F. was sup-
 743 ported by the NSF NRT program, grant #DGE-1735359. C.M.
 744 was supported by the MACMIT project (01LN1317A) funded
 745 through the German Federal Ministry of Education and Re-
 746 search (BMBF). C.F. was supported by the European Research
 747 Council Synergy grant #ERC-2013-SynG-610028 Imbalance-
 748 P. P.F. and K.W. were supported by the Newton Fund through
 749 the Met Office Climate Science for Service Partnership Brazil
 750 (CSSP Brazil). A.S. was supported by the Office of Science
 751 of the U.S. Department of Energy as part of the Multi-sector
 752 Dynamics Research Program Area. Computing resources were
 753 provided by the University of Chicago Research Computing
 754 Center (RCC). S.O. acknowledges support from the Swedish
 755 strong research areas BECC and MERGE together with sup-
 756 port from LUCCI (Lund University Centre for studies of Car-
 757 bon Cycle and Climate Interactions).

758 **6. References**

759 Angulo, C., Ritter, R., Lock, R., Enders, A., Fronzek, S., & Ewert, F. (2013).
 760 Implication of crop model calibration strategies for assessing regional im-
 761 pacts of climate change in europe. *Agric. For. Meteorol.*, 170, 32 – 46.

762 Asseng, S., Ewert, F., Martre, P., Ritter, R. P., B. Lobell, D., Cammarano, D.,
 763 A. Kimball, B., Ottman, M., W. Wall, G., White, J., Reynolds, M., D. Al-
 764 derman, P., Prasad, P. V. V., Aggarwal, P., Anothai, J., Basso, B., Bier-
 765 nath, C., Challinor, A., De Sanctis, G., & Zhu, Y. (2015). Rising tempera-
 766 tures reduce global wheat production. *Nature Climate Change*, 5, 143–147.
 767 doi:10.1038/nclimate2470.

768 Asseng, S., Ewert, F., Rosenzweig, C., Jones, J., Hatfield, J., Ruane, A.,
 769 J. Boote, K., Thorburn, P., Ritter, R. P., Cammarano, D., Brisson, N.,
 770 Basso, B., Martre, P., Aggarwal, P., Angulo, C., Bertuzzi, P., Biernath,
 771 C., Challinor, A., Doltra, J., & Wolf, J. (2013). Uncertainty in simulat-
 772 ing wheat yields under climate change. *Nature Climate Change*, 3, 827832.
 773 doi:10.1038/nclimate1916.

774 Aulakh, M. S., & Malhi, S. S. (2005). Interactions of Nitrogen with Other
 775 Nutrients and Water: Effect on Crop Yield and Quality, Nutrient Use Ef-
 776 ficiency, Carbon Sequestration, and Environmental Pollution. *Advances in*
 777 *Agronomy*, 86, 341 – 409.

778 Balkovi, J., van der Velde, M., Skalsk, R., Xiong, W., Folberth, C., Khabarov,
 779 N., Smirnov, A., Mueller, N. D., & Obersteiner, M. (2014). Global wheat
 780 production potentials and management flexibility under the representative
 781 concentration pathways. *Global and Planetary Change*, 122, 107 – 121.

782 Blanc, E. (2017). Statistical emulators of maize, rice, soybean and wheat yields
 783 from global gridded crop models. *Agricultural and Forest Meteorology*, 236,
 784 145 – 161.

785 Blanc, E., & Sultan, B. (2015). Emulating maize yields from global gridded
 786 crop models using statistical estimates. *Agricultural and Forest Meteorol-*
 787 *ogy*, 214-215, 134 – 147.

788 von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., & Zaehle, S.
 789 (2018). Implementing the Nitrogen cycle into the dynamic global vegeta-
 790 tion, hydrology and crop growth model LPJmL (version 5.0). *Geoscientific*
 791 *Model Development*, 11, 2789–2812.

792 Castruccio, S., McInerney, D. J., Stein, M. L., Liu Crouch, F., Jacob, R. L.,
 793 & Moyer, E. J. (2014). Statistical Emulation of Climate Model Projections
 794 Based on Precomputed GCM Runs. *Journal of Climate*, 27, 1829–1844.

795 Challinor, A., Watson, J., Lobell, D., Howden, S., Smith, D., & Chhetri, N.
 796 (2014). A meta-analysis of crop yield under climate change and adaptation.
 797 *Nature Climate Change*, 4, 287 – 291.

798 Conti, S., Gosling, J. P., Oakley, J. E., & O'Hagan, A. (2009). Gaussian process
 799 emulation of dynamic computer codes. *Biometrika*, 96, 663–676.

800 Duncan, W. (1972). SIMCOT: a simulation of cotton growth and yield. In
 801 C. Murphy (Ed.), *Proceedings of a Workshop for Modeling Tree Growth,*
 802 *Duke University, Durham, North Carolina* (pp. 115–118). Durham, North
 803 Carolina.

804 Duncan, W., Loomis, R., Williams, W., & Hanau, R. (1967). A model for
 805 simulating photosynthesis in plant communities. *Hilgardia*, (pp. 181–205).

806 Dury, M., Hambuckers, A., Warnant, P., Henrot, A., Favre, E., Ouberrous,
 807 M., & François, L. (2011). Responses of European forest ecosystems to
 808 21st century climate: assessing changes in interannual variability and fire
 809 intensity. *iForest - Biogeosciences and Forestry*, (pp. 82–99).

810 Elliott, J., Kelly, D., Chryssanthacopoulos, J., Glotter, M., Jhunjhnuwala, K.,
 811 Best, N., Wilde, M., & Foster, I. (2014). The parallel system for integrating
 812 impact models and sectors (pSIMS). *Environmental Modelling and Soft-*
 813 *ware*, 62, 509–516.

814 Elliott, J., Müller, C., Deryng, D., Chryssanthacopoulos, J., Boote, K. J.,
 815 Büchner, M., Foster, I., Glotter, M., Heinke, J., Iizumi, T., Izaurrealde, R. C.,
 816 Mueller, N. D., Ray, D. K., Rosenzweig, C., Ruane, A. C., & Sheffield, J.
 817 (2015). The Global Gridded Crop Model Intercomparison: data and mod-
 818 eling protocols for Phase 1 (v1.0). *Geoscientific Model Development*, 8,
 819 261–277.

820 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J.,
 821

- & Taylor, K. E. (2016). Overview of the coupled model intercomparison project phase 6 (cmip6) experimental design and organization. *Geoscientific Model Development*, 9, 1937–1958.
- Ferrise, R., Moriondo, M., & Bindi, M. (2011). Probabilistic assessments of climate change impacts on durum wheat in the mediterranean region. *Natural Hazards and Earth System Sciences*, 11, 1293–1302.
- Folberth, C., Gaiser, T., Abbaspour, K. C., Schulin, R., & Yang, H. (2012). Regionalization of a large-scale crop growth model for sub-Saharan Africa: Model setup, evaluation, and estimation of maize yields. *Agriculture, Ecosystems & Environment*, 151, 21 – 33.
- Food and Agriculture Organization of the United Nations (2018). FAOSTAT database. URL: <http://www.fao.org/faostat/en/home>.
- Fronzek, S., Pirttioja, N., Carter, T. R., Bindi, M., Hoffmann, H., Palosuo, T., Ruiz-Ramos, M., Tao, F., Trnka, M., Acutis, M., Asseng, S., Baranowski, P., Basso, B., Bodin, P., Buis, S., Cammarano, D., Deligios, P., Destain, M.-F., Dumont, B., Ewert, F., Ferrise, R., François, L., Gaiser, T., Hlavinka, P., Jacquemin, I., Kersebaum, K. C., Kollas, C., Krzyszczak, J., Lorite, I. J., Minet, J., Minguez, M. I., Montesino, M., Moriondo, M., Müller, C., Nen-del, C., Öztürk, I., Perego, A., Rodríguez, A., Ruane, A. C., Ruget, F., Sanna, M., Semenov, M. A., Slawinski, C., Stratonovitch, P., Supit, I., Waha, K., Wang, E., Wu, L., Zhao, Z., & Rötter, R. P. (2018). Classifying multi-model wheat yield impact response surfaces showing sensitivity to temperature and precipitation change. *Agricultural Systems*, 159, 209–224.
- Glotter, M., Elliott, J., McInerney, D., Best, N., Foster, I., & Moyer, E. J. (2014). Evaluating the utility of dynamical downscaling in agricultural impacts projections. *Proceedings of the National Academy of Sciences*, 111, 8776–8781.
- Glotter, M., Moyer, E., Ruane, A., & Elliott, J. (2015). Evaluating the Sensitivity of Agricultural Model Performance to Different Climate Inputs. *Journal of Applied Meteorology and Climatology*, 55, 151113145618001.
- Hank, T., Bach, H., & Mauser, W. (2015). Using a Remote Sensing-Supported Hydro-Agroecological Model for Field-Scale Simulation of Heterogeneous Crop Growth and Yield: Application for Wheat in Central Europe. *Remote Sensing*, 7, 3934–3965.
- He, W., Yang, J., Zhou, W., Drury, C., Yang, X., D. Reynolds, W., Wang, H., He, P., & Li, Z.-T. (2016). Sensitivity analysis of crop yields, soil water contents and nitrogen leaching to precipitation, management practices and soil hydraulic properties in semi-arid and humid regions of Canada using the DSSAT model. *Nutrient Cycling in Agroecosystems*, 106, 201–215.
- Heady, E. O. (1957). An Econometric Investigation of the Technology of Agricultural Production Functions. *Econometrica*, 25, 249–268.
- Heady, E. O., & Dillon, J. L. (1961). *Agricultural production functions*. Iowa State University Press.
- M., Kanudia, A., & F. B. (2014). Plsim-entsem v1.0: A spatiotemporal emulator of future climate change for impacts assessment. *Geoscientific Model Development*, 7, 433–451. doi:10.5194/gmd-7-433-2014.
- Holzkämper, A., Calanca, P., & Fuhrer, J. (2012). Statistical crop models: Predicting the effects of temperature and precipitation changes. *Climate Research*, 51, 11–21.
- Holzworth, D. P., Huth, N. I., deVoil, P. G., Zurcher, E. J., Herrmann, N. I., McLean, G., Chenu, K., van Oosterom, E. J., Snow, V., Murphy, C., Moore, A. D., Brown, H., Whish, J. P., Verrall, S., Fainges, J., Bell, L. W., Peake, A. S., Poulton, P. L., Hochman, Z., Thorburn, P. J., Gaydon, D. S., Dalgliesh, N. P., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp, J., Cichota, R., Vogeler, I., Li, F. Y., Wang, E., Hammer, G. L., Robertson, M. J., Dimes, J. P., Whitbread, A. M., Hunt, J., van Rees, H., McClelland, T., Carberry, P. S., Hargreaves, J. N., MacLeod, N., McDonald, C., Harsdorf, J., Wedgwood, S., & Keating, B. A. (2014). APSIM Evolution towards a new generation of agricultural systems simulation. *Environmental Modelling and Software*, 62, 327 – 350.
- Howden, S., & Crimp, S. (2005). Assessing dangerous climate change impacts on australia's wheat industry. *Modelling and Simulation Society of Australia and New Zealand*, (pp. 505–511).
- Iizumi, T., Nishimori, M., & Yokozawa, M. (2010). Diagnostics of climate model biases in summer temperature and warm-season insolation for the simulation of regional paddy rice yield in japan. *Journal of Applied Meteorology and Climatology*, 49, 574–591.
- Ingestad, T. (1977). Nitrogen and Plant Growth; Maximum Efficiency of Nitrogen Fertilizers. *Ambio*, 6, 146–151.
- Izaurrealde, R., Williams, J., McGill, W., Rosenberg, N., & Quiroga Jakas, M. (2006). Simulating soil C dynamics with EPIC: Model description and testing against long-term data. *Ecological Modelling*, 192, 362–384.
- Jagtap, S. S., & Jones, J. W. (2002). Adaptation and evaluation of the CROPGRO-soybean model to predict regional yield and production. *Agriculture, Ecosystems & Environment*, 93, 73 – 85.
- Jones, J., Hoogenboom, G., Porter, C., Boote, K., Batchelor, W., Hunt, L., Wilkens, P., Singh, U., Gijsman, A., & Ritchie, J. (2003). The DSSAT cropping system model. *European Journal of Agronomy*, 18, 235 – 265.
- Jones, J. W., Antle, J. M., Basso, B., Boote, K. J., Conant, R. T., Foster, I., Godfray, H. C. J., Herrero, M., Howitt, R. E., Janssen, S., Keating, B. A., Munoz-Carpena, R., Porter, C. H., Rosenzweig, C., & Wheeler, T. R. (2017). Toward a new generation of agricultural system data, models, and knowledge products: State of agricultural systems science. *Agricultural Systems*, 155, 269 – 288.
- Keating, B., Carberry, P., Hammer, G., Probert, M., Robertson, M., Holzworth, D., Huth, N., Hargreaves, J., Meinke, H., Hochman, Z., McLean, G., Ver-

- 907 burg, K., Snow, V., Dimes, J., Silburn, M., Wang, E., Brown, S., Bristow, K.,⁹⁵⁰
 908 Asseng, S., Chapman, S., McCown, R., Freebairn, D., & Smith, C. (2003).⁹⁵¹
 909 An overview of APSIM, a model designed for farming systems simulation.⁹⁵²
 910 *European Journal of Agronomy*, 18, 267 – 288. ⁹⁵³
- 911 Lindeskog, M., Arneth, A., Bondeau, A., Waha, K., Seaquist, J., Olin, S., &⁹⁵⁴
 912 Smith, B. (2013). Implications of accounting for land use in simulations of⁹⁵⁵
 913 ecosystem carbon cycling in Africa. *Earth System Dynamics*, 4, 385–407. ⁹⁵⁶
- 914 Liu, J., Williams, J. R., Zehnder, A. J., & Yang, H. (2007). GEPIC - modelling⁹⁵⁷
 915 wheat yield and crop water productivity with high resolution on a global⁹⁵⁸
 916 scale. *Agricultural Systems*, 94, 478 – 493. ⁹⁵⁹
- 917 Liu, W., Yang, H., Folberth, C., Wang, X., Luo, Q., & Schulin, R. (2016a).⁹⁶⁰
 918 Global investigation of impacts of PET methods on simulating crop-water⁹⁶¹
 919 relations for maize. *Agricultural and Forest Meteorology*, 221, 164 – 175. ⁹⁶²
- 920 Liu, W., Yang, H., Liu, J., Azevedo, L. B., Wang, X., Xu, Z., Abbaspour, K. C.,⁹⁶³
 921 & Schulin, R. (2016b). Global assessment of nitrogen losses and trade-offs⁹⁶⁴
 922 with yields from major crop cultivations. *Science of The Total Environment*,⁹⁶⁵
 923 572, 526 – 537. ⁹⁶⁶
- 924 Lobell, D. B., & Burke, M. B. (2010). On the use of statistical models to predict⁹⁶⁷
 925 crop yield responses to climate change. *Agricultural and Forest Meteorol-*⁹⁶⁸
 926 *ogy*, 150, 1443 – 1452. ⁹⁶⁹
- 927 Lobell, D. B., & Field, C. B. (2007). Global scale climate-crop yield relation-⁹⁷⁰
 928 ships and the impacts of recent warming. *Environmental Research Letters*,⁹⁷¹
 929 2, 014002. ⁹⁷²
- 930 MacKay, D. (1991). Bayesian Interpolation. *Neural Computation*, 4, 415–447.⁹⁷³
- 931 Makowski, D., Asseng, S., Ewert, F., Bassu, S., Durand, J., Martre, P.,⁹⁷⁴
 932 Adam, M., Aggarwal, P., Angulo, C., Baron, C., Basso, B., Bertuzzi,⁹⁷⁵
 933 P., Biernath, C., Boogaard, H., Boote, K., Brisson, N., Cammarano,⁹⁷⁶
 934 D., Challinor, A., Conijn, J., & Wolf, J. (2015). Statistical analysis of⁹⁷⁷
 935 large simulated yield datasets for studying climate effects. (p. 1100).⁹⁷⁸
 936 doi:10.13140/RG.2.1.5173.8328. ⁹⁷⁹
- 937 Mauser, W., & Bach, H. (2015). PROMET - Large scale distributed hydrolog-⁹⁸⁰
 938 ical modelling to study the impact of climate change on the water flows of⁹⁸¹
 939 mountain watersheds. *Journal of Hydrology*, 376, 362 – 377. ⁹⁸²
- 940 Mauser, W., Klepper, G., Zabel, F., Delzeit, R., Hank, T., Putzenlechner, B.,⁹⁸³
 941 & Calzadilla, A. (2009). Global biomass production potentials exceed ex-⁹⁸⁴
 942 pected future demand without the need for cropland expansion. *Nature Com-*⁹⁸⁵
 943 *munications*, 6. ⁹⁸⁶
- 944 McDermid, S., Dilleepkumar, G., Murthy, K., Nedumaran, S., Singh, P., Srinivasan,⁹⁸⁷
 945 C., Gangwar, B., Subash, N., Ahmad, A., Zubair, L., & Nissanka, S.⁹⁸⁸
 946 (2015). Integrated assessments of the impacts of climate change on agricul-⁹⁸⁹
 947 ture: An overview of AgMIP regional research in South Asia. *Chapter in:*⁹⁹⁰
 948 *Handbook of Climate Change and Agroecosystems*, (pp. 201–218). ⁹⁹¹
- 949 Mistry, M. N., Wing, I. S., & De Cian, E. (2017). Simulated vs. empirical⁹⁹²
 weather responsiveness of crop yields: US evidence and implications for
 the agricultural impacts of climate change. *Environmental Research Letters*,
 12.
- Moore, F. C., Baldos, U., Hertel, T., & Diaz, D. (2017). New science of climate
 change impacts on agriculture implies higher social cost of carbon. *Nature
 Communications*, 8.
- Müller, C., Elliott, J., Chryssanthacopoulos, J., Arneth, A., Balkovic, J., Ciais,
 P., Deryng, D., Folberth, C., Glotter, M., Hoek, S., Iizumi, T., Izaurrealde,
 R. C., Jones, C., Khabarov, N., Lawrence, P., Liu, W., Olin, S., Pugh, T.
 A. M., Ray, D. K., Reddy, A., Rosenzweig, C., Ruane, A. C., Sakurai, G.,
 Schmid, E., Skalsky, R., Song, C. X., Wang, X., de Wit, A., & Yang, H.
 (2017). Global gridded crop model evaluation: benchmarking, skills, de-
 ficiencies and implications. *Geoscientific Model Development*, 10, 1403–
 1422.
- Nakamura, T., Osaki, M., Koike, T., Hanba, Y. T., Wada, E., & Tadano, T.
 (1997). Effect of CO₂ enrichment on carbon and nitrogen interaction in
 wheat and soybean. *Soil Science and Plant Nutrition*, 43, 789–798.
- O'Hagan, A. (2006). Bayesian analysis of computer code outputs: A tutorial.
Reliability Engineering & System Safety, 91, 1290 – 1300.
- Olin, S., Schurgers, G., Lindeskog, M., Wårliind, D., Smith, B., Bodin, P.,
 Holmér, J., & Arneth, A. (2015). Modelling the response of yields and tissue
 C:N to changes in atmospheric CO₂ and N management in the main wheat
 regions of western europe. *Biogeosciences*, 12, 2489–2515. doi:10.5194/bg-
 12-2489-2015.
- Osaki, M., Shinano, T., & Tadano, T. (1992). Carbon-nitrogen interaction in
 field crop production. *Soil Science and Plant Nutrition*, 38, 553–564.
- Osborne, T., Gornall, J., Hooker, J., Williams, K., Wiltshire, A., Betts, R., &
 Wheeler, T. (2015). JULES-crop: a parametrisation of crops in the Joint UK
 Land Environment Simulator. *Geoscientific Model Development*, 8, 1139–
 1155.
- Ostberg, S., Schewe, J., Childers, K., & Frieler, K. (2018). Changes in crop
 yields and their variability at different levels of global warming. *Earth Sys-
 tem Dynamics*, 9, 479–496.
- Oyebamiji, O. K., Edwards, N. R., Holden, P. B., Garthwaite, P. H., Schaphoff,
 S., & Gerten, D. (2015). Emulating global climate change impacts on crop
 yields. *Statistical Modelling*, 15, 499–525.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O.,
 Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Pas-
 sos, A., Cournapeau, D., Brucher, M., Perrot, M., & Duchesnay, E. (2011).
 Scikit-learn: Machine Learning in Python. *Journal of Machine Learning
 Research*, 12, 2825–2830.
- Pirttioja, N., Carter, T., Fronzek, S., Bindi, M., Hoffmann, H., Palosuo, T.,
 Ruiz-Ramos, M., Tao, F., Trnka, M., Acutis, M., Asseng, S., Baranowski,

- 993 P., Basso, B., Bodin, P., Buis, S., Cammarano, D., Deligios, P., Destain, M., Dumont, B., Ewert, F., Ferrise, R., François, L., Gaiser, T., Hlavinka, P., Jacquemin, I., Kersebaum, K., Kollas, C., Krzyszczak, J., Lorite, I., Minet, J., Minguez, M., Montesino, M., Moriondo, M., Müller, C., Nendel, C., Öztürk, I., Perego, A., Rodríguez, A., Ruane, A., Ruget, F., Sanna, M., Semenov, M., Slawinski, C., Stratonovich, P., Supit, I., Waha, K., Wang, E., Wu, L., Zhao, Z., & Rötter, R. (2015). Temperature and precipitation effects on wheat yield across a European transect: a crop model ensemble analysis using impact response surfaces. *Climate Research*, 65, 87–105.

1000 Porter et al. (IPCC) (2014). Food security and food production systems. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. In C. Fi et al. (Ed.), *IPCC Fifth Assessment Report* (pp. 485–533). Cambridge, UK: Cambridge University Press.

1008 Portmann, F., Siebert, S., Bauer, C., & Doell, P. (2008). Global dataset of monthly growing areas of 26 irrigated crops.

1010 Portmann, F., Siebert, S., & Doell, P. (2010). MIRCA2000 - Global Monthly Irrigated and Rainfed crop Areas around the Year 2000: A New High Resolution Data Set for Agricultural and Hydrological Modeling. *Global Biogeochemical Cycles*, 24, GB1011.

1014 Pugh, T., Müller, C., Elliott, J., Deryng, D., Folberth, C., Olin, S., Schmid, E., & Arneth, A. (2016). Climate analogues suggest limited potential for intensification of production on current croplands under climate change. *Nature Communications*, 7, 12608.

1018 Räisänen, J., & Ruokolainen, L. (2006). Probabilistic forecasts of near-term climate change based on a resampling ensemble technique. *Tellus A: Dynamical Meteorology and Oceanography*, 58, 461–472.

1021 Ratto, M., Castelletti, A., & Pagano, A. (2012). Emulation techniques for the reduction and sensitivity analysis of complex environmental models. *Environmental Modelling & Software*, 34, 1 – 4.

1024 Razavi, S., Tolson, B. A., & Burn, D. H. (2012). Review of surrogate modeling in water resources. *Water Resources Research*, 48.

1026 Roberts, M., Braun, N., R Sinclair, T., B Lobell, D., & Schlenker, W. (2017). Comparing and combining process-based crop models and statistical models with some implications for climate change. *Environmental Research Letters*, 12.

1030 Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A. C., Müller, C., Arneth, A., Boote, K. J., Folberth, C., Glotter, M., Khabarov, N., Neumann, K., Piontek, F., Pugh, T. A. M., Schmid, E., Stehfest, E., Yang, H., & Jones, J. W. (2014). Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the National Academy of Sciences*, 111, 3268–3273.

1034 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1209 1210 1211 1212 1213 1214 1215 1216 1217 1217 1218 1219 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1259 1260 1261 1262 1263 1264 1265 1266 1267 1267 1268 1269 1269 1270 1271 1272 1273 1274 1275 1276 1277 1277 1278 1279 1279 1280 1281 1282 1283 1284 1285 1286 1287 1287 1288 1289 1289 1290 1291 1292 1293 1294 1295 1296 1297 1297 1298 1299 1299 1300 1301 1302 1303 1304 1305 1306 1307 1307 1308 1309 1309 1310 1311 1312 1313 1314 1315 1316 1316 1317 1318 1318 1319 1320 1321 1322 1323 1324 1325 1325 1326 1327 1327 1328 1329 1329 1330 1331 1332 1333 1334 1335 1336 1337 1337 1338 1339 1339 1340 1341 1341 1342 1343 1343 1344 1345 1345 1346 1347 1347 1348 1349 1349 1350 1351 1352 1353 1354 1355 1356 1356 1357 1358 1358 1359 1360 1360 1361 1362 1362 1363 1364 1364 1365 1366 1366 1367 1368 1368 1369 1370 1370 1371 1372 1372 1373 1374 1374 1375 1376 1376 1377 1378 1378 1379 1380 1380 1381 1382 1382 1383 1384 1384 1385 1386 1386 1387 1388 1388 1389 1390 1390 1391 1392 1392 1393 1394 1394 1395 1396 1396 1397 1398 1398 1399 1399 1399 1400 1400 1401 1402 1402 1403 1404 1404 1405 1406 1406 1407 1408 1408 1409 1410 1410 1411 1412 1412 1413 1414 1414 1415 1416 1416 1417 1418 1418 1419 1420 1420 1421 1422 1422 1423 1424 1424 1425 1426 1426 1427 1428 1428 1429 1430 1430 1431 1432 1432 1433 1434 1434 1435 1436 1436 1437 1438 1438 1439 1440 1440 1441 1442 1442 1443 1444 1444 1445 1446 1446 1447 1448 1448 1449 1450 1450 1451 1452 1452 1453 1454 1454 1455 1456 1456 1457 1458 1458 1459 1460 1460 1461 1462 1462 1463 1464 1464 1465 1466 1466 1467 1468 1468 1469 1470 1470 1471 1472 1472 1473 1474 1474 1475 1476 1476 1477 1478 1478 1479 1480 1480 1481 1482 1482 1483 1484 1484 1485 1486 1486 1487 1488 1488 1489 1490 1490 1491 1492 1492 1493 1494 1494 1495 1496 1496 1497 1498 1498 1499 1499 1499 1500 1500 1501 1502 1502 1503 1504 1504 1505 1506 1506 1507 1508 1508 1509 1509 1509 1510 1510 1511 1512 1512 1513 1514 1514 1515 1516 1516 1517 1518 1518 1519 1520 1520 1521 1522 1522 1523 1524 1524 1525 1526 1526 1527 1528 1528 1529 1530 1530 1531 1532 1532 1533 1534 1534 1535 1536 1536 1537 1538 1538 1539 1540 1540 1541 1542 1542 1543 1544 1544 1545 1546 1546 1547 1548 1548 1549 1550 1550 1551 1552 1552 1553 1554 1554 1555 1556 1556 1557 1558 1558 1559 1560 1560 1561 1562 1562 1563 1564 1564 1565 1566 1566 1567 1568 1568 1569 1570 1570 1571 1572 1572 1573 1574 1574 1575 1576 1576 1577 1578 1578 1579 1580 1580 1581 1582 1582 1583 1584 1584 1585 1586 1586 1587 1588 1588 1589 1590 1590 1591 1592 1592 1593 1594 1594 1595 1596 1596 1597 1598 1598 1599 1599 1599 1600 1600 1601 1602 1602 1603 1604 1604 1605 1606 1606 1607 1608 1608 1609 1610 1610 1611 1612 1612 1613 1614 1614 1615 1616 1616 1617 1618 1618 1619 1620 1620 1621 1622 1622 1623 1624 1624 1625 1626 1626 1627 1628 1628 1629 1630 1630 1631 1632 1632 1633 1634 1634 1635 1636 1636 1637 1638 1638 1639 1640 1640 1641 1642 1642 1643 1644 1644 1645 1646 1646 1647 1648 1648 1649 1650 1650 1651 1652 1652 1653 1654 1654 1655 1656 1656 1657 1658 1658 1659 1660 1660 1661 1662 1662 1663 1664 1664 1665 1666 1666 1667 1668 1668 1669 1670 1670 1671 1672 1672 1673 1674 1674 1675 1676 1676 1677 1678 1678 1679 1680 1680 1681 1682 1682 1683 1684 1684 1685 1686 1686 1687 1688 1688 1689 1690 1690 1691 1692 1692 1693 1694 1694 1695 1696 1696 1697 1698 1698 1699 1699 1699 1700 1700 1701 1702 1702 1703 1704 1704 1705 1706 1706 1707 1708 1708 1709 1710 1710 1711 1712 1712 1713 1714 1714 1715 1716 1716 1717 1718 1718 1719 1720 1720 1721 1722 1722 1723 1724 1724 1725 1726 1726 1727 1728 1728 1729 1729 1729 1730 1730 1731 1732 1732 1733 1734 1734 1735 1736 1736 1737 1738 1738 1739 1739 1739 1740 1740 1741 1742 1742 1743 1744 1744 1745 1746 1746 1747 1748 1748 1749 1749 1749 1750 1750 1751 1752 1752 1753 1754 1754 1755 1756 1756 1757 1758 1758 1759 1759 1759 1760 1760 1761 1762 1762 1763 1764 1764 1765 1766 1766 1767 1768 1768 1769 1770 1770 1771 1772 1772 1773 1774 1774 1775 1776 1776 1777 1778 1778 1779 1779 1779 1780 1780 1781 1782 1782 1783 1784 1784 1785 1786 1786 1787 1788 1788 1789 1789 1789 1790 1790 1791 1792 1792 1793 1794 1794 1795 1796 1796 1797 1798 1798 1799 1799 1799 1800 1800 1801 1802 1802 1803 1804 1804 1805 1806 1806 1807 1808 1808 1809 1810 1810 1811 1812 1812 1813 1814 1814 1815 1816 1816 1817 1818 1818 1819 1819 1819 1820 1820 1821 1822 1822 1823 1824 1824 1825 1826 1826 1827 1828 1828 1829 1829 1829 1830 1830 1831 1832 1832 1833 1834 1834 1835 1836 1836 1837 1838 1838 1839 1839 1839 1840 1840 1841 1842 1842 1843 1844 1844 1845 1846 1846 1847 1848 1848 1849 1849 1849 1850 1850 1851 1852 1852 1853 1854 1854 1855 1856 1856 1857 1858 1858 1859 1859 1859 1860 1860 1861 1862 1862 1863 1864 1864 1865 1866 1866 1867 1868 1868 1869 1869 1869 1870 1870 1871 1872 1872 1873 1874 1874 1875 1876 1876 1877 1878 1878 1879 1879 1879 1880 1880 1881 1882 1882 1883 1884 1884 1885 1886 1886 1887 1888 1888 1889 1889 1889 1890 1890 1891 1892 1892 1893 1894 1894 1895 1896 1896 1897 1898 1898 1899 1899 1899 1900 1900 1901 1902 1902 1903 1904 1904 1905 1906 1906 1907 1908 1908 1909 1909 1909 1910 1910 1911 1912 1912 1913 1914 1914 1915 1916 1916 1917 1918 1918 1919 1919 1919 1920 1920 1921 1922 1922 1923 1924 1924 1925 1926 1926 1927 1928 1928 1929 1929 1929 1930 1930 1931 1932 1932 1933 1934 1934 1935 1936 1936 1937 1938 1938 1939 1939 1939 1940 1940 1941 1942 1942 1943 1944 1944 1945 1946 1946 1947 1948 1948 1949 1949 1949 1950 1950 1951 1952 1952 1953 1954 1954 1955 1956 1956 1957 1958 1958 1959 1959 1959 1960 1960 1961 1962 1962 1963 1964 1964 1965 1966 1966 1967 1968 1968 1969 1969 1969 1970 1970 1971 1972 1972 1973 1974 1974 1975 1976 1976 1977 1978 1978 1979 1979 1979 1980 1980 1981 1982 1982 1983 1984 1984 1985 1986 1986 1987 1988 1988 1989 1989 1989 1990 1990 1991 1992 1992 1993 1994 1994 1995 1996 1996 1997 1998 1998 1999 1999 1999 2000 2000 2001 2002 2002 2003 2004 2004 2005 2006 2006 2007 2008 2008 2009 2009 2009 2010 2010 2011 2012 2012 2013 2014 2014 2015 2016 2016 2017 2018 2018 2019 2019 2019 2020 2020 2021 2022 2022 2023 2024 2024 2025 2026 2026 2027 2028 2028 2029 2029 2029 2030 2030 2031 2032 2032 2033 2034 2034 2035 2036 2036 2037 2038 2038 2039 2039 2039 2040 2040 2041 2042 2042 2043 2044 2044 2045 2046 2046 2047 2048 2048 2049 2049 2049 2050 2050 2051 2052 2052 2053 2054 2054 2055 2056 2056 2057 2058 2058 2059 2059 2059 2060 2060 2061 2062 2062 2063 2064 2064 2065 2066 2066 2067 2068 2068 2069 2069 2069 2070 2070 2071 2072 2072 2073 2074 2074 2075 2076 2076 2077 2078 2078 2079 2079 2079 2080 2080 2081 2082 2082 2083 2084 2084 2085 2086 2086 2087 2088 2088 2089 2089 2089 2090 2090 2091 2092 2092 2093 2094 2094 2095 2096 2096 2097 2098 2098 2099 2099 2099 2100 2100 2101 2102 2102 2103 2104 2104 2105 2106 2106 2107 2108 2108 2109 2109 2109 2110 2110 2111 2112 2112 2113 2114 2114 2115 2116 2116 2117 2118 2118 2119 2119 2119 2120 2120 2121 2122 2122 2123 2124 2124 2125 2126 2126 2127 2128 2128 2129 2129 2129 2130 2130 2131 2132 2132 2133 2134 2134 2135 2136 2136 2137 2138 2138 2139 2139 2139 2140 2140 2141 2142 2142 2143 2144 2144 2145 2146 2146 2147 2148 2148 2149 2149 2149 2150 2150 2151 2152 2152 2153 2154 2154 2155 2156 2156 2157 2158 2158 2159 2159 2159 2160 2160 2161 2162 2162 2163 2164 2164 2165 2166 2166 2167 2168 2168 2169 2169 2169 2170 2170 2171 2172 2172 2173 2174 2174 2175 2176 2176 2177 2178 2178 2179 2179 2179 2180 2180 2181 2182 2182 2183 2184 2184 2185 2186 2186 2187 2188 2188 2189 2189 2189 2190 2190 2191 2192 2192 2193 2194 2194 2195 2196 2196 2197 2198 2198 2199 2199 2199 2200 2200 2201 2202 2202 2203 2204 2204 2205 2206 2206 2207 2208 2208 2209 2209 2209 2210 2210 2211 2212 2212 2213 2214 2214 2215 2216 2216 2217 2218 2218 2219 2219 2219 2220 2220 2221 2222 2222 2223 2224 2224 2225 2226 2226 2227 2228 2228 2229 2229 2229 2230 2230 2231 2232 2232 2233 2234 2234 2235 2236 2236 2237 2238 2238 2239 2239 2239 2240 2240 2241 2242 2242 2243 2244 2244 2245 2246 2246 2247 2248 2248 2249 2249 2249 2250 2250 2251 2252 2252 2253 2254 2254 2255 2256 2256 2257 2258 2258 2259 2259 2259 2260 2260 2261 2262 2262 2263 2264 2264 2265 2266 2266 2267 2268 2268 2269 2269 2269 2270 2270 2271 2272 2272 2273 2274 2274 2275 2276 2276 2277 2278 2278 2279 2279 2279 2280 2280 2281 2282 2282 2283 2284 2284 2285 2286 2286 2287 2288 2288 2289 2289 2289 2290 2290 2291 2292 2292 2293 2294 2294 2295 2296 2296 2297 2298 2298 2299 2299 2299 2300 2300 2301 2302 2302 2303 2304 2304 2305 2306 2306 2307 2308 2308 2309 2309 2309 2310 2310 2311 2312 2312 2313 2314 2314 2315 2316 2316 2317 2318 2318 2319 2319 2319 2320 2320 2321 2322 2322 2323 2324 2324 2325 2326 2326 2327 2328 2328 2329 2329 2329 2330 2330 2331 2332 2332 2333 2334 2334 2335 2336 2336 2337 2338 2338 2339 2339 2339 2340 2340 2341 2342 2342 2343 2344 2344 2345 2346 2346 2347 2348 2348 2349 2349 2349 2350 2350 2351 2352 2352 2353 2354 2354 2355 2356 2356 2357 2358 2358 2359 2359 2359 2360 2360 2361 2362 2362 2363 2364 2364 2365 2366 2366 2367 2368 2368 2369 2369 2369 2370 2370 2371 2372 2372 2373 2374 2374 2375 2376 2376 2377 2378 2378 2379 2379 2379 2380 2380 2381 2382 2382 2383 2384 2384 2385 2386 2386 2387 2388 2388 2389 2389 2389 2390 2390 2391 2392 2392 2393 2394 2394 2395 2396 2396 2397 2398 2398 2399 2399 2399 2400 2400 2401 2402 2402 2403 2404 2404 2405 2406 2406 2407 2408 2408 2409 2409 2409 2410 2410 2411 2412 2412 2413 2414 2414 2415 2416 2416 2417 2418 2418 2419 2419 2419 2420 2420 2421 2422 2422 2423 2424 2424 2425 2426 2426 2427 2428 2428 2429 2429 2429 2430 2430 2431 2432 2432 2433 2434 2434 2435 2436 2436 2437 2438 2438 2439 2439 2439 2440 2440 2441 2442 2442 2443 2444 2444 2445 2446 2446 2447 2448 2448 2449 2449 2449 2450 2450 2451 2452 2452 2453 2454 2454 2455 2456 2456 2457 2458 2458 2459 2459 2459 2460 2460 2461 2462 2462 2463 2464 2464 2465 2466 2466 2467 2468 2468 2469 2469 2469 2470 2470 2471 2472 2472 2473 2474 2474 2475 2476 2476 2477 2478 2478 2479 2479 2479 2480 2480 2481 2482 2482 2483 2484 2484 2485 2486 2486 2487 2488 2488 2489

- 1079 observations and crop models. *Nature Communications*, 8, 13931. 1122
- 1080 Schlenker, W., & Roberts, M. J. (2009). Nonlinear temperature effects indicate 123
1081 severe damages to U.S. crop yields under climate change. *Proceedings of the National Academy of Sciences*, 106, 15594–15598. 124
- 1082
- 1083 Snyder, A., Calvin, K. V., Phillips, M., & Ruane, A. C. (2018). A crop yield 1122
1084 change emulator for use in gcam and similar models: Persephone v1.0. *Geoscientific Model Development Discussions*, 2018, 1–42.
- 1085
- 1086 Storlie, C. B., Swiler, L. P., Helton, J. C., & Sallaberry, C. J. (2009). Implementation 1122
1087 and evaluation of nonparametric regression procedures for sensitivity
1088 analysis of computationally demanding models. *Reliability Engineering & System Safety*, 94, 1735 – 1763.
- 1089
- 1090 Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 1122
1091 and the experiment design. *Bulletin of the American Meteorological Society*,
93, 485–498.
- 1092
- 1093 Tebaldi, C., & Lobell, D. B. (2008). Towards probabilistic projections of cli- 1122
1094 mate change impacts on global crop yields. *Geophysical Research Letters*,
35.
- 1095
- 1096 Valade, A., Ciais, P., Vuichard, N., Viovy, N., Caubel, A., Huth, N., Marin, F., & 1122
1097 Martin, J. F. (2014). Modeling sugarcane yield with a process-based model
1098 from site to continental scale: Uncertainties arising from model structure
1099 and parameter values. *Geoscientific Model Development*, 7, 1225–1245.
- 1100 Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., & Schewe, 1122
1101 J. (2014). The Inter-Sectoral Impact Model Intercomparison Project
1102 (ISI-MIP): Project framework. *Proceedings of the National Academy of Sciences*, 111, 3228–3232.
- 1103
- 1104 White, J. W., Hoogenboom, G., Kimball, B. A., & Wall, G. W. (2011). Method- 1122
1105 ologies for simulating impacts of climate change on crop production. *Field Crops Research*, 124, 357 – 368.
- 1106
- 1107 Williams, K., Gornall, J., Harper, A., Wiltshire, A., Hemming, D., Quaife, T., 1122
1108 Arkebauer, T., & Scoby, D. (2017). Evaluation of JULES-crop performance
1109 against site observations of irrigated maize from Mead, Nebraska. *Geoscientific Model Development*, 10, 1291–1320.
- 1110
- 1111 Williams, K. E., & Falloon, P. D. (2015). Sources of interannual yield vari- 1122
1112 ability in JULES-crop and implications for forcing with seasonal weather
1113 forecasts. *Geoscientific Model Development*, 8, 3987–3997.
- 1114 de Wit, C. (1957). Transpiration and crop yields. *Verslagen van Land-*
1115 *bouwkundige Onderzoeken* : 64.6, .
- 1116
- 1117 Wolf, J., & Oijen, M. (2002). Modelling the dependence of european potato 1122
1118 yields on changes in climate and co2. *Agricultural and Forest Meteorology*,
112, 217 – 231.
- 1119 Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D. B., Huang, Y., Huang, M., Yao, 1122
1120 Y., Bassu, S., Ciais, P., Durand, J. L., Elliott, J., Ewert, F., Janssens, I. A.,
1121 Li, T., Lin, E., Liu, Q., Martre, P., Miller, C., Peng, S., Peuelas, J., Ruane,