

Impact of Centre Vortices on the Gluon Propagator



THE UNIVERSITY

of ADELAIDE

James Biddle

Supervisors: Prof. Derek B. Leinweber

Dr. Waseem Kamleh

Department of Physics

University of Adelaide

This dissertation is submitted for the degree of

Master of Philosophy

August 2018

Table of contents

List of figures	xiii
List of tables	xv
1 Introduction	1
2 Lattice QCD	3
2.1 QCD in the Continuum	3
2.2 Pure Gauge Action	3
2.3 Gauge Fixing	4
2.3.1 Landau Gauge	4
2.3.2 Maximal Centre Gauge	4
3 Topology of the Lattice	5
3.1 Confinement	5
3.2 Centre Vortices	5
3.2.1 Centre Projection	5
3.3 Instantons	5
3.3.1 Topological Charge	5
4 Landau Gauge Gluon Propagator	7
4.1 Lattice Definition of the Gluon Propagator	7
4.2 Momentum Variables	9
5 Smoothing	11
5.1 Smoothing Methods	11
5.1.1 Cooling	11
5.1.2 Over-Improved Smearing	11
5.2 Results from the Gluon Propagator	11

6	Gluon Propagator on Vortex-Modified Backgrounds	13
6.1	Results	13
6.2	Cooling and the Average Action	13
6.3	Summary	13
7	Centre Vortex Visualisations	15
7.1	3D Models	15
7.1.1	Time Slices	15
7.1.2	Time-Oriented Links	15
7.1.3	Topological Charge	15
7.2	Centre Vortices and Topological Charge	15
8	Conclusion	17
	References	19

I would like to dedicate this thesis to

Declaration

I certify that this work contains no material which has been accepted for the award of any other degree or diploma in my name, in any university or other tertiary institution and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text. In addition, I certify that no part of this work will, in the future, be used in a submission in my name, for any other degree or diploma in any university or other tertiary institution without the prior approval of the University of Adelaide and where applicable, any partner institution responsible for the joint-award of this degree.

I give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library Search and also through web search engines, unless permission has been granted by the University to restrict access for a period of time.

I acknowledge the support I have received for my research through the provision of an Australian Government Research Training Program Scholarship.

James Biddle
August 2018

Acknowledgements

And I would like to acknowledge ...

Abstract

This is where you write your abstract ...

List of figures

List of tables

Chapter 1

Introduction

Chapter 2

Lattice QCD

Since the first efforts to construct a non-perturbative approach to QCD in 1974[11], lattice QCD has developed over the past 40 years into a powerful tool used to probe the low-energy behaviour of the strong nuclear force. Rather than treat spacetime as a set of continuous axes, it is instead discretised into a finite set of points on a four-dimensional hypercube. This prescription allows for the explicit calculation of path integrals present in QCD, at the cost of introducing finite-spacing errors that must be systematically accounted for.

2.1 QCD in the Continuum

2.2 Pure Gauge Action

For the purpose of this research, we are interested in the behaviour of gluons, and as such we need to develop a description of pure gauge fields. In the continuum, a pure gauge field has the Lagrangian[9]

$$\mathcal{L} = \frac{1}{2} F_{\mu\nu} F^{\mu\nu} \quad (2.1)$$

and corresponding action

$$\mathcal{S} = \int d^4x \mathcal{L}, \quad (2.2)$$

where $F_{\mu\nu}$ is the field-strength tensor, and can be written in terms of the traceless, Hermitian, gauge potential A_μ as

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - ig[A_\mu, A_\nu] \quad (2.3)$$

When considering the path integral formulation of a gauge field theory, integrals such as the generating functional,

$$\mathcal{Z} = \int \mathcal{D}A_\mu \exp(i\mathcal{S}[A_\mu(x)]), \quad (2.4)$$

and others of a similar form appear frequently. This integral closely resembles the partition function found in statistical mechanics, $Z_{\text{classical}} = \int d^3x d^3p \exp(-\beta H(x, p))$, with the notable exception of the factor of i in the exponential. This factor leads to an oscillatory weight term, rendering numerical simulations impossible. To ensure that the weight factor is purely real, it is necessary to perform a Wick rotation into Euclidean space[10, 11]

$$x_0 \rightarrow -ix_0$$

The generating functional now becomes

$$\mathcal{Z}_{\text{Eucl}} = \int \mathcal{D}A_\mu \exp(-\mathcal{S}_{\text{Eucl}}[A_\mu(x)]). \quad (2.5)$$

The Wick rotation also has the consequence of reducing the metric $g_{\mu\nu}$ to the identity, meaning that there is no longer any differentiation between covariant and contravariant tensors.

Within this framework, we can now consider discretising spacetime into a finite lattice, with each lattice site separated by a spacing a . When spacetime is discretised, it becomes necessary to consider derivatives as finite differences and integrals as finite sums. For example, we can construct the lattice form of Eq. 2.3 as

$$F_{\text{Lat}}^{\mu\nu}(x) = \frac{A_\nu(x + a\hat{\mu}) - A_\nu(x)}{a} - \frac{A_\mu(x + a\hat{\nu}) - A_\mu(x)}{a} - ig[A_\mu(x), A_\nu(x)]. \quad (2.6)$$

The notation $A_\nu(x + a\hat{\mu})$ denotes the field A_ν located at the site one lattice spacing in the $\hat{\mu}$ direction from x .

2.3 Gauge Fixing

2.3.1 Landau Gauge

2.3.2 Maximal Centre Gauge

Chapter 3

Topology of the Lattice

3.1 Confinement

3.2 Centre Vortices

3.2.1 Centre Projection

3.3 Instantons

3.3.1 Topological Charge

Chapter 4

Landau Gauge Gluon Propagator

In a gauge field theory, the propagator of the gauge boson is the two-point correlation function

4.1 Lattice Definition of the Gluon Propagator

We begin with the definition of the coordinate space propagator [12, 4, 6].

$$D_{\mu\nu}^{ab}(x) = \langle A_\mu^a(x) A_\nu^b(0) \rangle. \quad (4.1)$$

The propagator in momentum space is simply related by the discrete Fourier transform,

$$D_{\mu\nu}^{ab}(p) = \sum_x e^{-ip \cdot x} \langle A_\mu^a(x) A_\nu^b(0) \rangle. \quad (4.2)$$

Noting that the coordinate space propagator $D_{\mu\nu}^{ab}(x - y)$ only depends on the difference $x - y$, we can make use of translational invariance to average over the four-dimensional volume to obtain the form for the momentum space propagator.

$$\begin{aligned} D_{\mu\nu}^{ab}(p) &= \frac{1}{V} \sum_{x,y} e^{-ip \cdot x} \langle A_\mu^a(x + y) A_\nu^b(y) \rangle \\ &= \frac{1}{V} \sum_{x,y} \langle e^{-ip \cdot (x+y)} A_\mu^a(x + y) e^{+ip \cdot y} A_\nu^b(y) \rangle \\ &= \frac{1}{V} \langle A_\mu^a(p) A_\nu^b(-p) \rangle. \end{aligned} \quad (4.3)$$

Hence we find that the momentum space gluon propagator on a finite lattice with four-dimensional volume V is given by

$$D_{\mu\nu}^{ab}(p) \equiv \frac{1}{V} \langle A_\mu^a(p) A_\nu^b(-p) \rangle. \quad (4.4)$$

In the continuum, the Landau-gauge momentum-space gluon propagator has the following form [7, 3]

$$D_{\mu\nu}^{ab}(q) = \left(\delta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) \delta^{ab} D(q^2), \quad (4.5)$$

where $D(q^2)$ is the scalar gluon propagator. Contracting Gell-Mann index b with a and Lorentz index ν with μ one has

$$D_{\mu\mu}^{aa}(q) = (4 - 1) (n_c^2 - 1) D(q^2), \quad (4.6)$$

such that the scalar function can be obtained from the gluon propagator via

$$D(q^2) = \frac{1}{3(n_c^2 - 1)} D_{\mu\mu}^{aa}(q), \quad (4.7)$$

where $n_c = 3$ is the number of colours.

As the lattice gauge links $U_\mu(x)$ naturally reside in the fundamental representation of $SU(3)$, it is convenient to work with the corresponding 3×3 matrix representation of the gauge potential $A_\mu = A_\mu^a (\lambda_a/2)$, where λ_a are the eight Gell-Mann matrices. Using the orthogonality relation $\text{Tr}(\lambda_a \lambda_b) = \delta_{ab}$ for the Gell-Mann matrices, it is straightforward to see that

$$2 \text{Tr}(A_\mu A_\mu) = A_\mu^a A_\mu^a, \quad (4.8)$$

which can be substituted into equation 4.7 to obtain the final expression for the lattice scalar gluon propagator,

$$D(p^2) = \frac{2}{3(n_c^2 - 1)V} \langle \text{Tr} A_\mu(p) A_\mu(-p) \rangle. \quad (4.9)$$

Following the formalism of Ref. [7], we calculate the lattice gluon propagator using the mid-point definition of the gauge potential in terms of the lattice link variables [1],

$$A_\mu(x + \hat{\mu}/2) = \frac{1}{2ig_0} (U_\mu(x) - U_\mu^\dagger(x)) - \frac{1}{6ig_0} \text{Tr} (U_\mu(x) - U_\mu^\dagger(x)) + \mathcal{O}(a^2). \quad (4.10)$$

The gluon fields $U_\mu(x)$ are first gauge-fixed by maximizing an $\mathcal{O}(a^2)$ -improved functional using a Fourier-accelerated algorithm [5, 2, 8]. The gauge potential in momentum space is then obtained by taking the discrete Fourier transform,

$$A_\mu(p) = \sum_x e^{-ip \cdot (x + \hat{\mu}/2)} A_\mu(x + \hat{\mu}/2). \quad (4.11)$$

4.2 Momentum Variables

Chapter 5

Smoothing

5.1 Smoothing Methods

5.1.1 Cooling

5.1.2 Over-Improved Smearing

5.2 Results from the Gluon Propagator

Chapter 6

Gluon Propagator on Vortex-Modified Backgrounds

6.1 Results

6.2 Cooling and the Average Action

6.3 Summary

Chapter 7

Centre Vortex Visualisations

7.1 3D Models

7.1.1 Time Slices

7.1.2 Time-Oriented Links

7.1.3 Topological Charge

7.2 Centre Vortices and Topological Charge

Chapter 8

Conclusion

References

- [1] Alles, B., Henty, D., Panagopoulos, H., Parrinello, C., Pittori, C., and Richards, D. G. (1997). α_s from the nonperturbatively renormalised lattice three gluon vertex. *Nucl. Phys.*, B502:325–342.
- [2] Bonnet, F. D. R., Bowman, P. O., Leinweber, D. B., Williams, A. G., and Richards, D. G. (1999). Discretization errors in Landau gauge on the lattice. *Austral. J. Phys.*, 52:939–948.
- [3] Bonnet, F. D. R., Bowman, P. O., Leinweber, D. B., Williams, A. G., and Zanotti, J. M. (2001). Infinite volume and continuum limits of the Landau gauge gluon propagator. *Phys. Rev.*, D64:034501.
- [4] Cucchieri, A. (1999). Infrared behavior of the gluon propagator in lattice Landau gauge: The Three-dimensional case. *Phys. Rev.*, D60:034508.
- [5] Davies, C. T. H., Batrouni, G. G., Katz, G. R., Kronfeld, A. S., Lepage, G. P., Wilson, K. G., Rossi, P., and Svetitsky, B. (1988). Fourier Acceleration in Lattice Gauge Theories. 1. Landau Gauge Fixing. *Phys. Rev.*, D37:1581.
- [6] Langfeld, K., Reinhardt, H., and Gattnar, J. (2002). Gluon propagators and quark confinement. *Nucl. Phys.*, B621:131–156.
- [7] Leinweber, D. B., Skullerud, J. I., Williams, A. G., and Parrinello, C. (1998). Gluon propagator in the infrared region. *Phys. Rev.*, D58:031501.
- [8] Roberts, D. S., Bowman, P. O., Kamleh, W., and Leinweber, D. B. (2011). Wave Functions of the Proton Ground State in the Presence of a Uniform Background Magnetic Field in Lattice QCD. *Phys. Rev.*, D83:094504.
- [9] Ryder, L. H. (1996). *Quantum field theory*. Cambridge university press.
- [10] Schäfer, T. and Shuryak, E. V. (1998). Instantons in QCD. *Rev. Mod. Phys.*, 70:323–426.
- [11] Wilson, K. G. (1974). Confinement of Quarks. *Phys. Rev.*, D10:2445–2459. [319(1974)].
- [12] Zwanziger, D. (1991). Vanishing of zero momentum lattice gluon propagator and color confinement. *Nucl. Phys.*, B364:127–161.

