

## RWorksheet\_Cautivar#4a.Rmd

James Clark Cautivar

2024-10-16

1. The table below shows the data about shoe size and height. Create a data frame.

```
RespondentsData <- data.frame(
  Shoe_Size = c(6.5, 9.0, 8.5, 8.5, 10.5, 7.0, 9.5, 9.0, 13.0, 7.5, 10.5, 8.5, 12.0, 10.5, 13.0, 11.5, 8.5),
  Height = c(66.0, 68.0, 64.5, 65.0, 70.0, 64.0, 70.0, 71.0, 72.0, 64.0, 74.5, 67.0, 71.0, 71.0, 77.0, 77.0, 77.0),
  Gender = c("F", "F", "F", "F", "M", "F", "F", "F", "M", "F", "M", "F", "M", "M", "M", "M", "F", "F", "F")
)
RespondentsData
```

##	Shoe_Size	Height	Gender
## 1	6.5	66.0	F
## 2	9.0	68.0	F
## 3	8.5	64.5	F
## 4	8.5	65.0	F
## 5	10.5	70.0	M
## 6	7.0	64.0	F
## 7	9.5	70.0	F
## 8	9.0	71.0	F
## 9	13.0	72.0	M
## 10	7.5	64.0	F
## 11	10.5	74.5	M
## 12	8.5	67.0	F
## 13	12.0	71.0	M
## 14	10.5	71.0	M
## 15	13.0	77.0	M
## 16	11.5	72.0	M
## 17	8.5	59.0	F
## 18	5.0	62.0	F
## 19	10.0	72.0	M
## 20	6.5	66.0	F
## 21	7.5	64.0	F
## 22	8.5	67.0	M
## 23	10.5	73.0	M
## 24	8.5	69.0	F
## 25	10.5	72.0	M
## 26	11.0	70.0	M
## 27	9.0	69.0	M
## 28	13.0	70.0	M

- Describe the data. The data shows informations about respondent's shoe size, height, and their gender.
- Create a subset by males and females with their corresponding shoe size and height. What its result? Show the R scripts.

```
Males <- subset(RespondentsData, Gender == "M", select = c(Gender, Shoe_Size, Height))
Males
```

```
##      Gender Shoe_Size Height
## 5      M      10.5    70.0
## 9      M      13.0    72.0
## 11     M      10.5    74.5
## 13     M      12.0    71.0
## 14     M      10.5    71.0
## 15     M      13.0    77.0
## 16     M      11.5    72.0
## 19     M      10.0    72.0
## 22     M       8.5    67.0
## 23     M      10.5    73.0
## 25     M      10.5    72.0
## 26     M      11.0    70.0
## 27     M       9.0    69.0
## 28     M      13.0    70.0
```

```
Females <- subset(RespondentsData, Gender == "F", select = c(Gender, Shoe_Size, Height))
Females
```

```
##      Gender Shoe_Size Height
## 1      F       6.5    66.0
## 2      F       9.0    68.0
## 3      F       8.5    64.5
## 4      F       8.5    65.0
## 6      F       7.0    64.0
## 7      F       9.5    70.0
## 8      F       9.0    71.0
## 10     F       7.5    64.0
## 12     F       8.5    67.0
## 17     F       8.5    59.0
## 18     F       5.0    62.0
## 20     F       6.5    66.0
## 21     F       7.5    64.0
## 24     F       8.5    69.0
```

- c. Find the mean of shoe size and height of the respondents. Write the R scripts and its result.

```
AllShoeSize <- RespondentsData$Shoe_Size
mean(AllShoeSize)
```

```
## [1] 9.410714
```

```
AllHeight <- RespondentsData$Height
mean(AllHeight)
```

```
## [1] 68.57143
```

- d. Is there a relationship between shoe size and height? Why? I would say that there is a relationship between shoe size and height. Because taller people tend to have bigger shoe size while shorter people have smaller shoe size.
2. Construct character vector months to a factor with factor() and assign the result to factor\_months\_vector. Print out factor\_months\_vector and assert that R prints out the factor levels below the actual values.

```
months_vector <- c("March", "April", "January", "November", "January", "September", "October", "September", "November", "April")
```

```
factor_months_vector <- factor(months_vector)
factor_months_vector
```

```
## [1] March      April      January   November  January   September October
## [8] September November August     January   November  November  February
## [15] May        August     July      December  August    August    September
## [22] November  February   April
## 11 Levels: April August December February January July March May ... September
```

3. Then check the `summary()` of the `months_vector` and `factor_months_vector`. Interpret the results of both vectors. Are they both equally useful in this case?

```
summary(months_vector)
```

```
##      Length      Class      Mode
##         24 character character
```

```
summary(factor_months_vector)
```

```
##      April      August  December  February   January      July      March      May
##         2         4         1         2         3         1         1         1
## November  October September
##         5         1         3
```

Personally, i think that they're both useful because they show different results which are helpful if you need a specific information about a vector.

4. Create a vector and factor for the table below.

```
Directions <- c("East", "West", "North")
Frequency <- c(1, 4, 3)
```

```
new_order_data <- factor(Directions, levels = c("East", "West", "North"))
print(new_order_data)
```

```
## [1] East West North
## Levels: East West North
```

5. Enter the data below in Excel with file name = `import_march.csv`

- a. Import the excel file into the Environment Pane using `read.table()` function. Write the code.

```
file <- read.table("import_march.csv", header = TRUE, sep = ",")
```

- b. View the dataset. Write the R scripts and its result.

```
print(file)
```

```
##      Students Strategy.1 Strategy.2 Strategy3
## 1      Male           8          10          8
## 2              4           8           6
## 3              0           6           4
## 4      Female        14           4          15
## 5              10           2          12
## 6              6           0           9
```