

Baryonic Matter Organisation on Planetary, Stellar and Galactic Scales: The Λ CDM (M, λ) Parametrization as a Scale-Free Inheritance Law

JAMIESON G. DAHL¹

¹*Independent Researcher, Safety Bay, Australia*

ABSTRACT

Λ CDM galaxy formation describes a galaxy with two numbers, a mass budget and a spin draw, and three decades of semi-analytic success show that the pair is enough. Nothing in the theory says why. This paper says why: (M, λ) is the inheritance pair of dissipative baryonic organisation, the budget and the collapse-depth record that every parent system hands every child, processed by one universal machine at every scale. The galaxy recipe cosmology found by simulation is the top rung of a law this paper calibrates four rungs down. Jupiter’s satellite system closes at 0.9% on the pair inherited from Sol’s natal disc at Jupiter’s formation slot. The same law, inverted, reconstructs Hyperion, the star whose disc produced Sol. Run forward at full scale, it generates the Milky Way. Dark matter cannot dissipate and lies outside the theory; Λ CDM’s cosmological core is imported intact.

The machine. A natal disc’s formation seats, its natal slots, follow $r_n = R_{\text{disc}} \cdot \rho^n$ with $\rho = 1 - \frac{\sqrt{\ln 2}}{2} \approx 0.584$, a geometric constant with no fitted parameter. Two boundaries hold the disc in a vice: the magnetospheric Alfvén barrier inside, and outside the Davis Dam, the pressure wall against the ambient cloud, at $R_{\text{disc}} = 30.07 (M_{\star}/M_{\text{prim}}) D^{-1/3}$ AU with the nebula density D equal to one for Sol. Each slot’s allocation sums rock, snow-line ice, pebble drift, and envelope capture. In primordial solar units Sol sits at exactly (1, 1, 1) with its dam at Neptune; ten clean calibrations and 23 fitted systems follow. One dynamical event, Jupiter’s destabilisation of the embryo swarm, joins the Mercury–Vulcan merger to account for every Solar System mass anomaly.

The hierarchy. A daughter system inherits both defining inputs from its seat in the parent’s natal disc: the slot allocation budgets her inventory, and the parent’s local pressure sets her dam through $D_d/D_p = 0.39 (r_{\text{site}}/R_{\text{disc}})^{-2.12}$, an exponent that matches the disc’s own density profile. Three generations check one another. The Galilean disc prefers Jupiter’s formation site over its present orbit by a factor of 35; all four moons sit in situ in their observed order, Callisto on the dam, the inner trio displaced outward in the Laplace tidal pattern. Inverted, the recursion calibrates Sol’s

A-class progenitor and predicts its remnant: a white dwarf of $0.69 \pm 0.04 M_{\odot}$, cooling age near 3.4 Gyr, on a thin-disc orbit consistent with co-natal dispersal, travelling with a main-sequence sibling that carries Sol’s isotopic fingerprint.

The factory. The dam is also a production line. The T Tauri firehose drove $660 M_{\oplus}$ of solids through it, and the Kuiper belt is the residue that stuck. The factory is a moving shoreline whose deposits are dated cohorts, and the trans-Neptunian census confirms that the dam’s exterior carries structure in time, not in space: the resonant populations are the earliest cohort swept into Neptune’s comb, the cold classicals are the mid-firehose deposition band, and the Kuiper cliff is the radius where production shut down. The boundary still operates. Its stocked shelf is observed as the heliopause hydrogen wall, its counterparts as the astrospheres of other stars, and its galactic instance is already named in the literature: the galactopause, where thermal instability condenses cloud-mass products that rain back inward. A dense-cloud transit 2–3 Myr ago, independently proposed to explain the deep-sea ^{60}Fe layers, predicts young volatile-fresh bodies at 40–150 AU carrying live radionuclides.

The galaxy. A galaxy forms the way a planetary system does, one rung up. A seed condenses in a collapsing protogalactic cloud, and its outflow against the ambient medium sets a Davis Dam, the pressure boundary where the factory runs. This is the same production line that minted the Kuiper belt at Sol’s dam, one rung up, and here its products are stars. The vice is also why the cloud does not become one gigantic black hole: the collapse runs just deep enough to make the seed, the seed’s outflow stands up the two dams, and from then on further infall is shut off and the fuel is minted into stars instead. The dam is the system’s outer edge at every epoch. While the cloud is still collapsing, ambient pressure is at its peak and pins the dam to a small radius, so the factory makes its stars close in, building the bulge and feeding the seed together, which is why black-hole mass tracks classical-bulge mass ever after. With that the seed’s part is played. It templates, it does not labor: the factory’s output is limited only by the fuel arriving at the boundary, so the budget, not the seed, decides how much galaxy gets built, and the finished structure outweighs its founder ten thousand to one. As ambient pressure falls the dam advances, and the disc is its deposition track: every annulus of stars is minted at the dam as it passes, so the disc ages from the centre outward, the inside-out growth observers measure, at constant mass per radius, which is the Mestel profile (Mestel 1963) and the origin of the flat rotation curve. Each stellar cohort the factory mints lands at a golden-angle offset of 137.5° from the last, the division of the circle at which successive batches least overlap. The arms are the manufacturing trail: cohorts one lattice step apart trace a logarithmic spiral of pitch 12.65° , two steps apart 24.2° , both angles fixed by ρ and the golden angle alone; the measured pitch distribution across 4,378 disk galaxies is bimodal at $12.0^{\circ} \pm 3.4$ and $23^{\circ} \pm 4.3$. Once the deposited mass outweighs the seed, dynamics pass from Keplerian to self-gravitating, and the templating epoch survives as fossils: the founder coupling, the deposition profile, the pitch lattice,

and a bulge fraction set by collapse depth. A generative run of this sequence with one tuned normalisation reproduces the Milky Way’s gross statistics eight for eight: stellar mass, star count, white-dwarf census, star-formation rate, half-mass epoch, old-mass fraction, bulge fraction, and the inside-out age gradient. Slot-determined stellar metallicity dissolves the Sun’s metal-rich anomaly along the way.

The paper closes with a dated predictions register. Every mechanism is listed with the observation that would kill it.

Keywords: planet formation, protoplanetary disks, Solar System formation, Kuiper belt, Oort cloud, satellite systems, spiral galaxies, Galaxy structure, stellar abundances

1. INTRODUCTION

Λ CDM galaxy formation rests on a remarkably economical recipe. To first order a galaxy is two numbers, its halo mass and its spin parameter λ (Peebles 1969), plus an assembly history. Three decades of semi-analytic modelling attest that the pair suffices, yet within Λ CDM the pair is phenomenology; nothing in the theory says why two numbers, or why these two. This paper proposes the why, and the evidence for the proposal lives far below the galactic scale. Our thesis is that (M, λ) is the universal inheritance pair of dissipative baryonic organisation. Every bounded baryonic system is determined by the mass budget and the collapse-depth record handed down by its parent, processed by one universal machine. Λ CDM’s galaxy recipe is the top rung of this law, operating in the self-gravitating dynamical regime, discovered top-down by simulation at the one scale where the parent is the environment itself. The law’s lower rungs are not conjecture. They are calibrated in this paper, at the sub-percent level where the hierarchy is resolvable (Table 1).

The machine operates in two dynamical regimes. In the Keplerian regime, where the seed dominates its natal disc, it is fully calibrated: a ladder of natal slots at a fixed geometric ratio, held between two pressure boundaries, with a closed-form mass allocation. In the self-gravitating regime the disc has come to outweigh the seed that organised it. The Keplerian formulas for period structure and resonant locking do not transfer there, and they are not needed: the templating epoch writes its record into the structure itself, and Section 9 reads that record. The boundary grammar crosses the transition whole. Dissipative matter settles into flat ordered components, dissipationless matter occupies spheroidal holding reservoirs, and boundaries between media operate as factories at every scale we examine. Where dissipation itself fails the theory ends: dark matter is the permanently unorganisable remainder, and Λ CDM’s cosmological core enters this paper as infrastructure, imported rather than contested.

The machine tier is developed first and carries the paper’s calibrations; its strongest instance is the Solar System. The architecture of the Solar System is unusually well

Table 1. The inheritance pair across the hierarchy. Each system is determined by a mass budget and a collapse-depth (spin) draw handed down by its parent. The machine processing the pair is universal; its dynamical regime changes once, at the top, where the disc outweighs its seed. Throughout this paper the dissipative disc from which a bound system’s members condense is called the *natal disc*, and its discrete formation seats are the *natal slots*. Protogalactic, protoplanetary, circumplanetary, and protolunar discs are scale-specific instances of one object, the realisation of the inheritance pair.

System	Parent	Budget (M)	Spin draw (λ)	Status
Galaxy	environment	$f_b M_{\text{halo}}$	λ_{halo} (tidal torques)	generative facsimile (§9)
Star + disc	cloud / parent disc	core / slot allocation	recorded $\Omega = D^{2/3}$ (jaw-lock)	10 clean calibrations (§3)
Sol	A-class progenitor	slot-0 allocation	$D = 1$ at the progenitor’s dam	reconstruction (§7)
Giant-planet sub-cascade	host star’s disc	slot allocation σ_{AAF}	$D_d = 0.39 x^{-2.12} D_p$	measured at 0.9% (§7)

constrained by both dynamical and cosmochemical evidence, yet no single theoretical framework simultaneously accounts for: (i) the exact masses of the eight planets, (ii) Mars’s anomalously small mass relative to extrapolated rocky surface density, (iii) the Moon-forming impactor’s existence, and (iv) the near-5:2 mean-motion resonance between Saturn and Jupiter (their orbital periods sitting close to a 5:2 integer ratio).

Existing frameworks address subsets. The Grand Tack hypothesis (Walsh et al. 2011) postulates that Jupiter migrated inward to ~ 1.5 AU and then outward, dragging Saturn behind it; this scenario explains Mars’s mass deficit but is in tension with isotopic evidence (Kruijer et al. 2017) indicating that Jupiter’s core formed in < 1 Myr and separated the non-carbonaceous and carbonaceous reservoirs without subsequent excursions. The Nice Model (Tsiganis et al. 2005) postulates a large-scale post-formation outward migration of Saturn through Neptune. This paper requires no such reorganization: all four giants form at or within an AU of their present orbits, and Neptune sits in situ at the dam to four significant figures, where it anchors the entire calibration. Core accretion (Pollack et al. 1996) and pebble accretion (Lambrechts & Johansen 2012; Bitsch et al. 2015) provide mechanisms for individual planet growth but do not produce a closed-form mass-allocation rule across the entire disc.

We propose a complementary approach: a *local-capacity* rule that allocates planetary mass to each orbital radius r on the basis of the viscous-disc surface density profile, snow-line ice partition, pebble drift, and gas-envelope accretion timescales. The rule is closed-form and admits direct comparison against observed planetary masses. The principal advantage is parsimony: three system-level parameters plus per-planet formation radius and time suffice to account for ten Solar System bodies (including the proposed impactors Theia and Vulcan), the two anchor planets fit exactly, and every other body’s deviation from its cascade allocation is attributed to one of two dynamical events.

The model places Jupiter and Saturn essentially in-situ at their geometric cascade slots ($r_3 = 5.98$ AU and $r_2 = 10.24$ AU respectively, with both observed positions within ~ 0.8 AU of slot center). This eliminates the conventional invocation of Grand Tack-style migration sequences (Walsh et al. 2011) or Nice-Model-style outer-giant reorganization (Tsiganis et al. 2005). The inner-system perturbations that previously required migration explanations are instead supplied by a single dynamical event: Jupiter’s standard gravitational scattering of the pre-consolidation embryo swarm at the snow-line pile-up zone (slot 4-5 region, 2–3.5 AU).

Evidence structure.—The law is calibrated where the hierarchy can be resolved and predictive where it cannot yet be. Sections 2 through 7 carry the calibrations: ten clean system fits, the Solar System at exactly (1, 1, 1) in primordial units, and the three-generation closure at 0.9%. Sections 8 and 9 carry the same law into territory where its tests are still running, and state for each prediction the observation that would kill it; the register in Section 10 consolidates these with dates. Two matters lie outside the theory by its own structure and are said so once: dark matter, which cannot dissipate and therefore cannot be organised by any dissipative machine, and late-stage system evolution.

Scope of this paper.—The paper proceeds as follows. Section 2 introduces the local-capacity machine: the geometric cascade, the two-jawed vice (Alfvén boundary and Davis Dam), the nebula-density variable, and Tanigawa-Ikoma gas-envelope physics. Section 3 presents the Solar System calibration, including the Vulcan-Mercury merger. Section 4 establishes Venus and Neptune as the ISU primordial-plane reference and quantifies impact signatures via inclination and eccentricity excesses. Section 5 demonstrates Saturn and Jupiter in-situ at slots 2 and 3 respectively, with the slot 4-5 embryo-swarm dispersal mechanic as the framework’s single dynamical event. Section 6 demonstrates scale-invariance of the cascade at multi-stellar and cluster scales. Section 7 formalises the hierarchical recursion by which each generation budgets and confines the next, and carries its three-generation calibration. Section 8 extends the boundary physics outward, deriving the Kuiper belt and Oort cloud as reservoirs at the system’s magnetic and gravitational outer boundaries, with the dam as a moving production shoreline. Section 9 presents the galactic tier: the regime transition, the inheritance pair under its Λ CDM names, the signatures of the geometry surviving self-gravity, and the boundary-factory programme. Section 10 enumerates testable predictions. Section 11 discusses novelty, the empirical f_{disc} stochasticity finding, model limitations, and relation to prior frameworks. We conclude in Section 12. A browser-based interactive implementation of the model, with all calibrated systems pre-loaded, is available at <https://jamesgdahl.github.io/HYDROS-Planet-Formation-Model/>.

2. THE LOCAL-CAPACITY ALLOCATION MODEL

2.1. *Physical foundation*

The model rests on three established disc-physics results:

Viscous disc surface density.—For a steady-state viscous accretion disc with viscosity $\nu \propto r$, self-similar solutions of the Lynden-Bell–Pringle (hereafter LBP) equations yield surface density

$$\Sigma(r) \propto r^{-1}, \quad (1)$$

giving constant mass-per-annulus $2\pi r\Sigma(r)dr = \text{const}$ (Lynden-Bell & Pringle 1974; Pringle 1981; Shakura & Sunyaev 1973). This is observationally confirmed for discs around T Tauri stars (young pre-main-sequence stars still accreting from their discs), with measured power-law indices $\gamma \approx 0.9\text{--}1.1$ (Hartmann et al. 1998; Andrews et al. 2009, 2010; Williams & Cieza 2011; Andrews 2020; Long et al. 2018; Trapman et al. 2019). The Hayashi minimum-mass solar nebula (MMSN) (Hayashi 1981) adopts a steeper $\Sigma \propto r^{-1.5}$ for solid material, but modern observations with the Atacama Large Millimeter/submillimeter Array (ALMA) favor the LBP scaling. We adopt $\Sigma \propto r^{-1}$ here.

The Davis Dam (outer-edge pressure barrier).—We name the outer boundary for Leverett Davis Jr., who identified its present-day descendant before the solar wind itself was accepted: Davis (1955) proposed that the Sun’s corpuscular emission carves a cavity in the interstellar medium whose edge is set by *pressure balance against the interstellar field*, the heliopause as a confinement surface, three years before Parker (1958) supplied the outflow that inflates it. (Earlier drafts of this framework called this boundary the “Anti-Alfvén Dam”; we retire that coinage in favour of the discoverer.) The two boundaries form a *vice* with distinct owners: the inner jaw is stellar (spin and field, through disc-locking) while the outer jaw is environmental, the ambient pressure differential of the birth site. The single spin parameter reads both jaws at once because they correlate through their common driver, the birth-environment density; the extreme effective spins of the inverted regime are the outer jaw’s signature, not stellar rotation. The dam’s local masonry is magnetic: through the bulk of the disc the magnetorotational instability (MRI), the magnetic stirring that makes disc gas effectively viscous, is *suppressed* by non-ideal effects (ambipolar diffusion, low ionization; Bai & Stone 2013; Turner et al. 2014); only where ionization rises does the midplane re-couple to the magnetic field. At the outer disc boundary, cosmic-ray and stellar ionization penetrate the thinning column and revive the MRI (Gammie 1996; Flock et al. 2015), producing a sharp rise in effective viscosity: a step, because the bulk on one side is magnetically dead and the edge on the other is alive. Solid material drifting inward from the outer disc pools against this viscosity step, and the associated pressure gradient halts further drift (Pinilla et al. 2012, 2016; Lyra et al. 2009; Regály et al. 2012). Observed disc outer radii behave like such transport-set

boundaries: gas-disc sizes track angular-momentum transport rather than free viscous spreading (Najita & Bergin 2018), and measured dust radii do not grow with age over 1–10 Myr (Hendler et al. 2020), edges are boundary-controlled, not spreading fronts. We name this boundary the *Davis Dam*, $R_{\bar{A}}$, in explicit parallel to the inner *Alfvén Dam*, R_A (next paragraph). The dam metaphor is deliberate: both boundaries are *viscous* barriers in a flowing fluid, not gravitational traps, material accumulates because the downstream channel narrows. The two dams bracket the planet-forming annulus $[R_A, R_{\bar{A}}]$, each backed by a reservoir that fills to a baseline mass set by the balance of inward drift flux against the dam’s holding capacity: the inner reservoir is the model’s intercept a (Eq. 8); the outer reservoir is the edge pile-up of Eq. 12. The literature’s “dead zone” label for the quiescent interior is a misnomer on both counts: the zone still accretes via layered MRI in its upper layers (Gammie 1996), and the dynamically important structure is the outer boundary where material pools, not the interior. In the model we identify $R_{\bar{A}} \equiv R_{\text{disc}}$.

The inner-reservoir mass a (Eq. 8) corresponds physically to what Chatterjee & Tan (2014) term the “Inside-Out Planet Formation reservoir”: solids drift inward via aerodynamic drag and pool against the inner edge, supplying the close-in planet zone (Boley & Ford 2013; Hansen & Murray 2013; Liu, Ormel & Lin 2017; Romanova & Lovelace 2006). Indeed, in-situ assembly of close-in super-Earths from a smooth disc fails without exactly such an inner solid pile-up (Ogihara et al. 2015): the reservoir is required, not optional. Our intercept a is the steady-state baseline fill of that reservoir; the scaling $a \propto M_\star \cdot \Omega^{2/7}$ reflects the dam-pressure dependence on stellar magnetic moment and spin via $R_A \propto \Omega^{4/7}$.

The Alfvén Dam (magnetospheric inner barrier).—Stellar magnetic-field pressure halts disc gas at the Alfvén radius,

$$R_A = \left(\frac{\mu^4}{2GM_\star \dot{M}} \right)^{1/7}, \quad (2)$$

where μ is the stellar magnetic moment and $\dot{M}$ is the accretion rate (Ghosh & Lamb 1979; Königl 1991; Shu et al. 1994; Long et al. 2005; Romanova et al. 2008); the underlying magnetic-pressure concept traces to Alfvén (1942). The truncation formula itself carries no explicit spin dependence: the $R_A \propto \Omega^{4/7}$ scaling we use enters through the disc-locking equilibrium, in which the star’s rotation is regulated to corotation with the truncation radius throughout the accretion phase (Königl 1991; Bouvier et al. 2007), dynamically coupling spin to the inner boundary. We refer to this boundary as the *Alfvén Dam*: the magnetic field obstructs the inward-flowing disc, and solids pool against it, filling the inner reservoir a described above. For solar-mass stars during the protoplanetary phase, $R_A \sim 0.05\text{--}0.2$ AU, scaling with stellar mass and spin.

Snow line and ice condensation.—The snow line r_{snow} is the orbital radius beyond which the disc is cold enough for water to condense onto grains as ice, enhancing the local

solid surface density by a factor $\sim 3\text{--}4$ (Hayashi 1981; Pollack & Bodenheimer 1989; Mulders et al. 2017). Its position is set by the disc’s thermal structure, which depends on dust opacity and therefore on the size distribution of the suspended dust (Min et al. 2011; Mulders, Ciesla & Min 2015; Bitsch et al. 2015). In a viscously heated disc, opacity-rich ISM-like (small-grain-dominated) dust blankets the midplane, keeping it warm and pushing the snow line outward; grain growth lowers the opacity and draws the snow line up to a factor ~ 2 closer to the star (Mulders, Ciesla & Min 2015). We parameterize this through the dimensionless *grain-opacity parameter* $g \in [0, 1]$ (Eq. 6), interpolating between the grain-grown, opacity-poor limit $r_{\text{snow}}(g=0) = 1.6$ AU and the ISM-like, opacity-rich limit $r_{\text{snow}}(g=1) = 3.3$ AU at solar mass and $f_{\text{disc}} = 0.01$, matching the Mulders-Ciesla viscous-disc range.

Pebbles drifting inward across the snow line additionally create a local pile-up: they lose their volatiles at the ice-sublimation boundary and accumulate there as a pressure-trap concentration of refractory and residual-ice material (Mulders et al. 2017; Schoonenberg & Ormel 2017; Drażkowska & Alibert 2017). In the two-population picture of Birnstiel et al. (2012), the small grains that carry the opacity (and so set r_{snow}) coexist with the grown pebbles that carry the inward mass flux (and so feed the trap). In the model the trap’s amplitude and position both follow r_{snow} (Eq. 11), giving g a dual role: it sets where the snow-line bump sits and how much mass it concentrates.

What a slot is.—The slot is geometric in origin, magnetic in operation, and fossil in survival. Its position is fixed by the projection constant ρ , with no field in the derivation; its effects are delivered entirely through the magnetized gas. Every component of this embodiment is published. Long-lived axisymmetric pressure bumps arise spontaneously in magnetorotational turbulence as zonal flows, generated by magnetic-pressure variations (Johansen et al. 2009), driven by magnetic flux concentration (Bai & Stone 2014); solids drift into them and form planetesimals there (Dittrich et al. 2013), and dust trapping in such rings is now directly observed (Dullemond et al. 2018). Migration torque reverses across density structure, so the same bumps trap passing planets (Masset et al. 2006), with the magnetospheric cavity rim as the family’s innermost member (Lin et al. 1996). One structure therefore performs both of the lattice’s jobs: it builds occupants and it catches them. What the literature lacks, and this paper supplies, is the organisation: the bumps stand on the ρ lattice anchored at the dam, and they precede their occupants. The living lattice has been photographed: HL Tau’s ladder of rings at an age younger than any planet-formation timescale (ALMA Partnership 2015), and the ringed-disc census of the DSHARP survey (Huang et al. 2018). The mainstream reading of those rings runs the causal arrow the other way, as gaps carved by already-formed planets; HL Tau’s youth is this paper’s witness, and the adjacent-ring ratio distribution of the DSHARP catalogue is the registered discriminator (§10): lattice occupancy predicts ratio peaks at 1.31 and 1.71 and their multiples, planet carving predicts none. When the gas disperses

the enforcer dies but the law does not; the addresses persist as fossils, which is why they remain fittable gigayears later.

2.2. Fundamental inputs

The model has three system-level inputs and one per-planet input:

$M_\star$, **primordial stellar mass:** (in $M_\odot$, before pre-main-sequence mass loss).

Ω , **primordial spin:** dimensionless, with $\Omega = 1$ defined as the Sun’s primordial value.

f_{disc} , **disc-to-star mass ratio:** $\in [0.001, 0.3]$, the integrated gas-disc mass divided by the stellar mass.

r , **per-planet orbital radius:** (in AU).

Per-planet formation time t_{form} is derived from the cascade rule (§2.5) rather than fit independently. Numerically the model works in *primordial-solar units* (stellar mass, spin, and nebula density all equal 1 for Sol, with one mass unit $\equiv 1.14$ present-day $M_\odot$) so Sol is the exact baseline $(M_\star, \Omega, D) = (1, 1, 1)$ and the calibration constants carry no residual 1.14 factors. Catalogue host masses, quoted in present-day solar masses, are adopted *directly* as primordial-unit values: under uniform fractional early mass loss (each star, like Sol, retaining 1/1.14 of its birth mass through the T Tauri wind era) the 1.14 cancels in the ratio. Young hosts that have not yet shed their T Tauri mass are the exception, β Pic (~ 20 Myr) and HR 8799 (~ 30 – 40 Myr) enter at literature mass /1.14. Spectral classes follow the same convention as primordial identities: a slot of X units is the star the literature quotes at $X M_\odot$ after uniform loss, so letter-class thresholds scale by 1.14 while ignition thresholds (deuterium, hydrogen) remain physical at birth mass. Sol is born, lives, and is classified G, and the convention validates against the sky: Alpha Centauri B’s 0.80-unit slot classifies K, matching its observed K1V.

The nebula density D : spin’s physical twin.—Under the vice reading of the two dams (§2), the second input may be stated equivalently as the *nebula density* D (Sol $\equiv 1$): the formation cloud’s ambient confinement establishes the Davis Dam at

$$R_{\text{disc}} = 30.07 \cdot \frac{M_\star}{1.14 M_\odot} \cdot D^{-1/3} \text{ AU}, \quad \Omega = D^{2/3}, \quad (3)$$

so that $D = 1$ places Sol’s dam at 30.07 AU, Neptune’s exact slot, with spin the angular-momentum *record* of the same squeeze (the jaw-lock: both jaws driven by birth-site density). The calibration machinery that anchors R_{disc} on the outermost planet is therefore a bisection of D expressed through the lock, and fitted spins double as density readouts. Inverting Sol’s $D = 1$ through the collapse centrifugal radius (at the measured rotational-energy fraction $\beta \approx 0.02$; Burkert & Bodenheimer 2000) yields a birth core of ~ 0.015 pc at $n(\text{H}_2) \approx 10^6 \text{ cm}^{-3}$: a *clustered* prestellar core,

independently consistent with the meteoritic birth-cluster evidence and the cluster-era boundaries of §8. In the inverted regime the fitted D_{eff} exceeds any static cloud (TRAPPIST-1: $\sim 5 \times 10^4$), there the outer jaw involves more than birth pressure, and the lock reads as effective compression rather than literal cloud density.

The grain-opacity constant g .—A fourth quantity, the dimensionless grain-opacity parameter $g \in [0, 1]$ (from fully grain-grown, opacity-poor dust ($g = 0$) to ISM-like, small-grain-dominated dust ($g = 1$)) sets the snow-line position (Eq. 6). It is treated as a *universal constant*, $g = 0.82$ (Table 2), not a per-system input: calibrated once on Sol, the same value serves every calibrated system, since the mass-position data that drive the calibrations constrain M_* , Ω , and f_{disc} but give little independent leverage on g , whose snow-line effect is partially degenerate with both (Eq. 6). The value sits close to the ISM-like, opacity-rich limit, consistent with the opacity-bearing small-grain population being inherited from the interstellar medium’s grain-size distribution (Mathis et al. 1977) with only modest processing by the epoch of planet formation. We note the calibration sample is confined almost entirely to the solar neighbourhood, so the apparent constancy of g may be a property of our local galactic region rather than truly universal; per-system constraints await compositional or resolved-snow-line observations (§11).

Rotational disc compression.—Primordial stellar rotation sets the centrifugal radius for in-falling envelope material; rapidly-rotating young stars truncate their gas discs at correspondingly smaller radii. Modern ALMA surveys reveal a wide spread in protoplanetary disc sizes, with characteristic radii ranging from ~ 3 AU (compact discs around young M-dwarfs and rapidly-rotating solar analogs) to $\gtrsim 200$ AU (extended T Tauri systems) (Andrews et al. 2010; Long et al. 2018; Trapman et al. 2019; Andrews 2020; Manara et al. 2023; Hendler et al. 2020; Najita & Bergin 2018). The compression scaling $R_{\text{disc}} \propto \Omega^{-1/2}$ adopted here (Eq. 4) captures this trend: compact super-Earth chains (e.g., TRAPPIST-1, Kepler-90) require primordial $\Omega \gtrsim 5$ –20 to compress their formation discs to match observed planet positions, while wide-architecture systems like Sol sit near $\Omega = 1$.

The scaling has a deeper reading. $R \propto \Omega^{-1/2}$ is equivalent to $\Omega \propto R^{-2}$: precisely the signature of angular-momentum conservation ($\Omega \propto J/MR^2$ at fixed J). The relation therefore does not require spin to *cause* compression; it records that disc size and spin are two readings of one conserved quantity, and the causal driver can be external: ambient pressure confines the disc–star system to radius R , and spin follows by conservation, communicated to the star by disc-locking (Königl 1991; Bouvier et al. 2007). On this reading “primordial spin” is the recorded compression of the birth environment, fast rotators are stars born squeezed, with the natal draw of specific angular momentum (Burkert & Bodenheimer 2000) setting the family constant (30 AU at solar values) and environmental pressure selecting the system’s position along the (R, Ω) hyperbola. The measured spread of that natal draw is itself a predic-

tion in waiting: propagated through $R \propto \Omega^{-1/2}$, the [Burkert & Bodenheimer \(2000\)](#) angular-momentum distribution fixes not just the scaling but the *width* of the disc-size distribution at fixed stellar mass, directly comparable to the ALMA spread cited above.

2.3. Derived disc properties

$$R_{\text{disc}} = 30.07 \cdot \frac{M_{\star}}{M_{\odot, \text{prim}}} \cdot \Omega^{-1/2} \text{ AU}, \quad (4)$$

$$R_A = 0.20 \cdot \frac{M_{\star}}{M_{\odot, \text{prim}}} \cdot \Omega^{4/7} \text{ AU}, \quad (5)$$

$$r_{\text{snow}} = [1.6 + 1.7 g^{2.2}] \cdot \left(\frac{M_{\star}}{M_{\odot, \text{prim}}} \right)^2 \sqrt{\frac{f_{\text{disc}}}{0.01}} \text{ AU}, \quad (6)$$

$$\sigma \equiv \frac{M_{\star} \cdot Z \cdot f_{\text{rock}} \cdot f_{\text{disc}} \cdot \eta_{\text{rock}}}{R_{\text{disc}}} \quad [M_{\oplus}/\text{AU}], \quad (7)$$

$$a = 0.596 \cdot \frac{M_{\star}}{M_{\odot, \text{prim}}} \cdot \Omega^{2/7} \quad [M_{\oplus}], \quad (8)$$

where $M_{\odot, \text{prim}} = 1.14 M_{\odot}$ (Sol calibration anchor) and the universal physical constants are listed in Table 2. The $M_{\star}^2$ exponent in Eq. 6 is a model choice calibrated within the framework; accretional snow-line treatments give shallower stellar-mass scalings ([Kennedy & Kenyon 2008](#); [Mulders, Ciesla & Min 2015](#)), and the difference is testable across the M-dwarf catalogue systems. The quantity σ (Eq. 7) we name the *Annulus Allocation Factor*: it is the intensive rate at which rocky mass is allocated per unit radius along the disc. For an LBP $\Sigma \propto r^{-1}$ profile, $\sigma = dM/dr = 2\pi r \Sigma(r)$ is a radial constant, which is what makes the linear allocation rule (Eq. 10) well-defined as a function of r alone. The 1970s disc-dynamics literature ([Lynden-Bell & Pringle 1974](#); [Pringle 1981](#)) arrived at this constant implicitly via the self-similar viscous solution but did not name it.

Disc compression regime.—The dimensionless compression $C \equiv R_A/R_{\text{disc}}$ classifies the disc: $C < 1$ is the normal regime; $C \geq 1$ is the *inverted* regime, where rocky allocation concentrates at the magnetospheric inner edge (relevant for compact M-dwarf systems such as TRAPPIST-1).

Half-step occupancy in the inverted regime.—Inversion changes how viscous matter behaves, and thereby which rungs of the ladder are fed. With the entire disc inside the magnetosphere, magnetic stresses stiffen the gas toward corotation (reducing the Keplerian shear that cross-feeds adjacent annuli) and the compressed reservoir’s gas density crushes the random velocities of solids. In the normal regime a rung body’s feeding zone runs at its geometric (Gaussian-HWHM) width: adjacent zones at full

Table 2. Universal physical constants.

Symbol	Value	Meaning
g	0.82	Grain opacity (0 grain-grown $\rightarrow$ 1 ISM-like)
Z	0.014	Solar metallicity
f_{rock}	0.22	Rocky fraction of condensables
η_{rock}	0.78	Rock retention factor
$\eta_{\text{ice/rock}}$	3.5	Ice/rock ratio past snow line
ϵ_{pebble}	0.40	Pebble capture efficiency
M_{thresh}	$3.0 M_{\oplus}$	Gas-accretion threshold
A_0	$4.45 M_{\oplus}^{-1}$	Tanigawa-Ikoma H/He accretion coefficient
$k_{\text{H/He}}$	0.691 Myr^{-1}	Envelope decay rate
t_{disc}	5 Myr	Disc dispersal time
$M_{\odot}/M_{\oplus}$	332946	Solar mass in $M_{\oplus}$

step touch at half amplitude, so the midpoint material is consumed during consolidation and full-step occupancy is enforced by starvation, not exclusion. In the inverted regime, low-mass bodies' zones become *drag-limited*, narrower than geometric: midpoint material survives, and interstitial bodies consolidate at the half-step sites $r_n\sqrt{\rho}$ wherever the resulting half-step lattice is Hill-stable for the occupant ($\text{gap}/R_{H,\text{mut}} \geq 7$). We name this half-step occupancy state the *Luger lattice*, after the work that exhibited its complete form (the TRAPPIST-1 resonant chain, whose member h was predicted from the occupied rungs before detection (Luger et al. 2017)) and we note the eponym is earned twice over: the lattice's natural period ratio $\rho^{-3/4} = 1.497$ sits within 0.2% of the 3:2 commensurability that dominates that chain. Giants are Hill-limited rather than drag-limited, their reach spans the midpoint regardless of damping, so the gate excludes them naturally. The two occupancy harmonics map onto the two observed resonant-chain families: full-ladder chains (adjacent period ratio $\rho^{-3/2} = 2.24$) relax into 2:1 locks (GJ 876, HR 8799) while Luger-lattice chains are born essentially *on* 3:2: TRAPPIST-1's seven planets occupy seven consecutive lattice sites in the calibrated fit (§11.2), with its outermost member h anchored directly on the dam (slot 0 = R_{disc}), and compact Kepler chains follow the same pattern. Mixed occupancy follows the same gate: in Kepler-90 the two giants hold full rungs while the six small inner planets occupy the lattice. The inverted ladder's inward projection is terminated not by geometry but by the star itself: during the disc era the host sits on its Hayashi track, bloated to $R_{\text{HT}} \approx 2.3 R_{\odot} (M_{\star}/M_{\odot})^{2/3} (t/\text{Myr})^{-1/3}$, and rungs interior to that photosphere are *consumed*: their allocation is real, its destination is the star, and the predicted planet mass at such sites is zero (TRAPPIST-1's slots 6–10; 55 Cancri's slots 9–10). The inner edge of a compact system thereby tracks the host's protostellar radius: 55 Cancri e sits at the first rung outside its formation-era photosphere.

2.4. Mass-allocation function

The total planet mass at r is

$$M(r) = M_{\text{rock}}(r) + M_{\text{ice}}(r) + M_{\text{peb}}(r) + M_{\text{H/He}}(r) + \delta M, \quad (9)$$

where δM accommodates per-planet post-formation modifications such as giant-impact mass loss.

Rock allocation.—For the normal regime ($C < 1$),

$$M_{\text{rock}}(r) = \begin{cases} \sigma \cdot \left[\left(r + \frac{5}{6} R_A \right) - R_A \right] \cdot f_{\text{trunc}}(r) & R_A \leq r < 2R_A \\ [a + \sigma \cdot r] \cdot f_{\text{trunc}}(r) & 2R_A \leq r \leq R_{\text{disc}} \\ 0 & \text{otherwise} \end{cases} \quad (10)$$

plus a snow-line pile-up bump (active when $r \leq r_{\text{snow}}$):

$$M_{\text{bump}}(r) = \frac{1}{2} \sigma r_{\text{snow}} \exp \left[-\frac{(r - r_{\text{snow}})^2}{2(0.15 r_{\text{snow}})^2} \right] \quad (11)$$

and an outer-edge contribution:

$$M_{\text{outer}}(r) = \sigma R_{\text{disc}} C \exp \left[-\frac{(r - R_{\text{disc}})^2}{2(0.3 R_{\text{disc}})^2} \right]. \quad (12)$$

Edge truncation of the feeding zone: the hexagonal void.—A planet near a disc edge loses part of its feeding zone to the void beyond the dam, and the fraction lost follows from packing geometry alone. In two dimensions the densest arrangement of equal bodies is hexagonal close-packing, in which every interior body has exactly six nearest neighbours spaced at 60° intervals: six being the two-dimensional “kissing number” (Thue 1910; Conway & Sloane 1999). A growing embryo embedded in a drifting pebble field competes for material along the six sectors separating it from those six neighbours, so its feeding zone (in the sense of Safronov 1972) divides naturally into six equal shares. For a body in the disc interior all six sectors are populated with solids. For a body sitting on the outer dam ($r = R_{\text{disc}}$), the outward-facing sector lies beyond the dam, in the void where no solid material exists: the neighbour that would have occupied that sector is absent, and the share of material that sector would have supplied is simply never delivered. The body therefore accretes from five of its six sectors (retention = $5/6 \approx 0.83$, loss = $1/6 \approx 0.17$) a geometric constant with no fitted parameter. Equivalently, the feeding zone can be written as an asymmetric radial interval with inward range L and outward range S in the ratio $L:S = 5:1$; smoothing that interval across the edge gives

$$f_{\text{trunc}}(r) = \max \left[0, 1 - \frac{(r + S) - R_{\text{disc}}}{L + S} \right], \quad S = 0.025 R_{\text{disc}}, \quad L = 5S, \quad (13)$$

equal to 1 well inside the disc (all six sectors populated), exactly $5/6$ at $r = R_{\text{disc}}$, and 0 beyond $1.125 R_{\text{disc}}$. Neptune, sitting on Sol’s outer dam, carries precisely this one-sixth ($\sim 17\%$) reduction in its rock and ice allocation. The same one-sixth loss recurs at the inner edge, where the missing sector faces the magnetospheric void inside R_A ; it enters Eq. 10 through the ramp offset $\frac{5}{6}R_A$. The outer-reservoir Gaussian (Eq. 12) is the dam pile-up itself, not a feeding zone, and is not truncated.

Ice allocation.—Full condensation past the snow line gives an ice-to-rock mass ratio of $\eta_{\text{ice/rock}} \approx 3.5$, set by solar condensable-element abundances (Lodders 2003; Asplund et al. 2009), which yield a cosmochemical ice/rock partition of $\sim 3\text{--}4$ depending on whether organics and CO/CO₂ ices are included (Pollack et al. 1994; Ciesla & Cuzzi 2006; Oka, Nakamoto & Ida 2011).

Only a fraction of this ice is retained as planet-building material, and the fraction falls with radius: slower accretion at large r allows more ice to drift inward and sublimate (Ciesla & Cuzzi 2006; Garaud & Lin 2007; Kennedy & Kenyon 2008); outer-disc solids are more easily lost to scattering and photoevaporation (Owen 2017; Picogna et al. 2019); and late-forming bodies lose ice as the disc photosphere recedes (Krijt et al. 2016, 2018). We capture these combined effects through an empirical exponential ice-retention factor:

$$\eta_{\text{ice}}(r) = \exp\left[-\frac{r - r_{\text{snow}}}{0.8R_{\text{disc}}}\right], \quad (14)$$

$$M_{\text{ice}}(r) = \sigma (r - r_{\text{snow}}) \cdot \eta_{\text{ice/rock}} \cdot \eta_{\text{ice}}(r) \cdot f_{\text{trunc}}(r) + M_{\text{bump}}(r). \quad (15)$$

The decay length $0.8 R_{\text{disc}}$ is calibrated to Uranus’s and Neptune’s ice fractions and is consistent with the depletion scaling expected from Ciesla & Cuzzi (2006)’s viscous transport modeling. The snow-line bump M_{bump} (Eq. 11) is predominantly ice past r_{snow} , since the pressure trap concentrates condensable material at the ice-condensation front (Schoonenberg & Ormel 2017; Drażkowska & Alibert 2017).

Pebble flux allocation.—Drifting solids in a protoplanetary disc are aerodynamically transported inward as a disc-wide *pebble flux* (Birnstiel et al. 2012; Lambrechts & Johansen 2012, 2014; Bitsch et al. 2015). Gas-eligible cores along the drift path capture a share of this flux via pebble accretion. We allocate the total budget across eligible cores according to per-planet weights set by their radial distance from the snow line (the dominant pebble source region). The total pebble-flux budget is

$$M_{\text{peb,total}} = f_{\text{disc}} \cdot Z \cdot (1 - f_{\text{rock}}) \cdot M_{\star} \cdot \epsilon_{\text{pebble}}, \quad (16)$$

distributed among gas-eligible planets according to weights $w_i = (r_i - r_{\text{snow}})^{-1/2}$ (zero if $r_i \leq r_{\text{snow}}$):

$$M_{\text{peb,i}} = M_{\text{peb,total}} \cdot \frac{w_i}{\sum_{j \in \text{eligible}} w_j}. \quad (17)$$

Gas-accretion threshold.—A planet is *gas-eligible* if its core mass (rock + ice) exceeds $M_{\text{thresh}} = 3 M_{\oplus}$ and its formation time $t_{\text{form}} < t_{\text{disc}} = 5$ Myr. The threshold reflects the critical core mass at which the envelope’s self-gravity overwhelms its thermal pressure support, triggering runaway gas accretion (Mizuno 1980; Pollack et al. 1996; Ikoma, Nakazawa & Emori 2000; Rafikov 2006; Piso & Youdin 2014; Lee & Chiang 2015). The classical Mizuno value ($\sim 10 M_{\oplus}$) assumed solar-metallicity opacity-rich envelopes; modern revisions accounting for grain-size evolution and reduced opacity in young protoplanetary discs place the critical core mass closer to $3\text{--}5 M_{\oplus}$ (Ikoma, Nakazawa & Emori 2000; Hori & Ikoma 2010; Piso & Youdin 2014). We adopt $3 M_{\oplus}$ consistent with the low-opacity end of this range, appropriate for the high-accretion-rate protoplanetary phase. The disc-dispersal timescale of 5 Myr is the observationally-determined median gas-disc lifetime for solar-mass stars (Haisch et al. 2001; Hartmann et al. 1998; Mamajek 2009; Williams & Cieza 2011).

Hydrogen-helium envelope.—For gas-eligible cores, the H/He envelope grows by Kelvin-Helmholtz contraction (the envelope radiates away heat, shrinks, and admits more gas) until either the core reaches its runaway-accretion mass or the gas disc disperses (Pollack et al. 1996; Ida & Lin 2004; Mordasini 2014; Lissauer et al. 2009; Tanigawa & Ikoma 2007). The Tanigawa-Ikoma 2007 formulation gives a quadratic dependence of gas-envelope mass on core mass during the runaway-accretion phase, integrated over the gas-disc dispersal window:

$$A(t_{\text{form}}) = A_0 \exp[-k_{\text{H/He}} t_{\text{form}}], \quad (18)$$

$$w_{\text{wind}}(r) = \frac{1}{1 + (\Omega/30)(0.5/r)^2}, \quad (19)$$

$$M_{\text{H/He}} = A_0 \cdot M_{\text{core}}^2 \cdot \exp[-k_{\text{H/He}} t_{\text{form}}] \cdot w_{\text{wind}}(r), \quad (20)$$

where $M_{\text{core}} = M_{\text{rock}} + M_{\text{ice}} + M_{\text{peb}}$, and t_{form} is the formation time inferred from the Annulus Allocation Factor via $t_{\text{form}} = 0.10 \cdot r/\sigma$ (Eq. 22, §2.5).

The accretion coefficient $A_0 = 4.45 M_{\oplus}^{-1}$ is the most empirically-tuned parameter in the model. Its physical origin is the integrated runaway gas accretion rate (Tanigawa-Ikoma 2007), which scales as

$$A_0 \sim \frac{\Sigma_{\text{gas},0} \cdot \Omega_K \cdot \tau_{\text{disc}}}{M_{\star}}, \quad (21)$$

where $\Sigma_{\text{gas},0}$ is the local gas surface density at formation, Ω_K is the Keplerian angular velocity, and τ_{disc} is the gas-dispersal timescale. For Sol’s minimum-mass solar nebula values at Jupiter’s slot-3 position ($\Sigma_{\text{gas},0} \approx 8 \text{ g/cm}^2$ at 5.2 AU, $\Omega_K \approx 0.53 \text{ rad/yr}$, $\tau_{\text{disc}} \approx 3 \text{ Myr}$), substituting into Eq. 21 reproduces $A_0 \approx 4\text{--}6$ in the correct order of magnitude. The framework currently treats A_0 as a calibration constant tuned to Jupiter’s bulk composition; a first-principles derivation from disc evolution models (Pollack-Bitsch-Mordasini class) is identified as future work.

The exponential decay rate $k_{\text{H/He}} = 0.691 \text{ Myr}^{-1}$ (half-life $\sim 1 \text{ Myr}$) reflects the rapid loss of available gas as the disc disperses: protoplanetary disc gas surface density declines exponentially over $\sim 1\text{--}3 \text{ Myr}$ (Haisch et al. 2001; Hartmann et al. 1998; Mamajek 2009), suppressing envelope accretion for late-forming cores.

Inner gas-envelope boundary (solar-wind suppression).—The wind-suppression factor $w_{\text{wind}}(r)$ encodes the well-established extreme-ultraviolet (XUV) photoevaporation of close-in gas envelopes by stellar irradiation (Murray-Clay et al. 2009; Lopez, Fortney & Miller 2012; Owen & Wu 2013; Jin et al. 2014; Lopez & Fortney 2013). Young stars exhibit XUV luminosities $\sim 10^3$ times the modern Sun’s value (Tu et al. 2015; Ribas et al. 2005), and this flux drives hydrodynamic escape from H/He envelopes at small orbital radii. The Ω dependence in w_{wind} captures the correlation between primordial stellar rotation and XUV luminosity: rapidly-rotating young stars emit substantially higher XUV flux (Wright et al. 2011; Tu et al. 2015), stripping envelopes more efficiently. The r^{-2} scaling is the canonical inverse-square dependence of incident flux. Together, $w_{\text{wind}}(r)$ establishes an effective *inner boundary* for gas-envelope retention: close-in planets in fast-rotating systems retain little to no H/He, consistent with the observed bimodal radius distribution of Kepler super-Earths and sub-Neptunes (Fulton et al. 2017; Owen & Wu 2017).

2.5. Formation-time cascade

The formation time t_{form} for a planet at radius r is inferred directly from the Annulus Allocation Factor σ (Eq. 7) via a linear pebble-drift-timescale relation:

$$t_{\text{form}}(r) = 0.10 \cdot \frac{r}{\sigma} \quad [\text{Myr}]. \quad (22)$$

This is the only place in the model where t_{form} enters as an independent quantity, and it feeds directly into the H/He envelope accretion (Eq. 18) and the gas-window eligibility ($t_{\text{form}} < t_{\text{disc}}$, §2 paragraph on gas-accretion threshold). The formula’s physical content is:

Scaling with σ (the Annulus Allocation Factor).—Higher σ means more local solid mass per AU available to a growing core. For pebble accretion, the core-assembly rate scales linearly with the local solid surface density and therefore with σ : doubling σ roughly halves the time to reach the critical core mass. The $1/\sigma$ dependence in Eq. 22 captures this directly. Across systems with very different disc masses (e.g., HR 8799’s $f_{\text{disc}} \sim 30\%$ versus TRAPPIST-1’s $f_{\text{disc}} \sim 0.5\%$), this single scaling reproduces the observed formation-timescale spread without per-system tuning.

Scaling with r .—At fixed σ , formation time grows linearly with orbital radius. The dominant physical effect is the radial pebble-flux integration length: cores at larger r must wait longer for inward-drifting pebbles to traverse the disc and reach them (Ormel & Klahr 2010; Lambrechts & Johansen 2012, 2014; Bitsch et al. 2015, 2018;

Johansen & Lambrechts 2017). The linear r dependence (as opposed to the steeper $r^{5/2}$ that pure local-density arguments would suggest for a $\Sigma \propto r^{-1}$ LBP profile) reflects the gas-drag-mediated accretion regime in which core growth is rate-limited by the pebble-flux supply rather than by the local solid reservoir alone, a result robust across the Ormel & Klahr (2010), Lambrechts & Johansen (2012), and Bitsch et al. (2015) pebble-accretion frameworks.

Cascade structure and calibration.—Together, these two scalings produce an outward-sweeping formation cascade in which inner planets form first and outer planets form later, in a single disc. The proportionality constant 0.10 is calibrated so that Sol’s Jupiter (at $r = 5.20$ AU, $\sigma = 0.304 M_{\oplus}/\text{AU}$) lands at $t_{\text{form}} = 1.71$ Myr, consistent with Kruijer et al. (2017)’s isotopic constraint of < 1 Myr for Jupiter’s core formation followed by envelope growth. Once calibrated, the same constant reproduces all eight Solar System bodies’ formation timings (Table 3) and generalizes across the 22 further catalogued systems.

Sensitivity.—For cores below $M_{\text{thresh}} = 3 M_{\oplus}$ the envelope physics is gated off and t_{form} has no further effect. For gas-eligible cores, t_{form} sets the envelope mass through Eq. 18; across the calibrated systems the observed envelope masses correspond to formation times within a factor of ~ 2 of the cascade value, indicating that the r/σ scaling captures the dominant physics.

Mass-driven vs. radius-driven cascades.—In Sol, mass and radius correlate (heavier planets are further out), so the r/σ cascade produces a clean monotonic t_{form} -by-radius sequence. In compact systems such as 55 Cnc or Kepler-90, where inner gas giants exist at $r < 1$ AU and the system has been compressed by primordial rotation (Eq. 23), t_{form} is driven primarily by core mass (through the envelope term’s exponential sensitivity to t_{form}) rather than by formation radius. This dual behavior, natural radius-cascade in spread architectures, mass-cascade in compact architectures, is an emergent property of the model, not an imposed feature.

2.6. Boundary conditions

The allocation vanishes outside the two viscous dams. At the inner edge the ramp in Eq. 10 clamps to zero (for $r < R_A/6$, inside the magnetospheric void); at the outer edge f_{trunc} (Eq. 13) fades the allocation smoothly to zero by $1.125 R_{\text{disc}}$. The linear rule $a + \sigma r$ is never extrapolated beyond the disc. In the present model $R_{\bar{A}} \equiv R_{\text{disc}}$.

2.7. Anchoring the cascade to the outermost planet

The outermost observed planet anchors the cascade: slot 0 sits at the Davis Dam, so the outermost planet’s orbital radius marks R_{disc} itself (or an inner slot of it, $r_{\text{out}} = \rho^k R_{\text{disc}}$, when the outermost surviving body formed at slot $k > 0$). The primordial spin then follows from Eq. 4:

$$\Omega = \left(\frac{30 M_{\star}/M_{\odot, \text{prim}}}{R_{\text{disc}}} \right)^2. \quad (23)$$

Table 3. Solar System fit. Inputs: $M_\star = 1.14 M_\odot$, $\Omega = 0.995$, $f_{\text{disc}} = 0.01$. Venus and Neptune (ISU anchors) fit exactly; all other deviations are attributed to the two dynamical events (footnotes).

Body	Slot	r_{slot} (AU)	r_{obs} (AU)	t_{form} (Myr)	$M_{\text{pred}} (M_\oplus)$	$M_{\text{obs}} (M_\oplus)$
Mercury	8	0.41	0.387	0.13	0.108	0.055 [†]
Venus	7	0.69	0.723	0.24	0.815	0.815
Earth	6	1.19	1.000	0.33	0.900	1.000 [‡]
Mars	5	2.04	1.524	0.67	1.323	0.107 [*]
Jupiter	3	5.98	5.203	1.71	318.9	317.83
Saturn	2	10.25	9.537	3.13	133.7	95.16 ^{***}
Uranus	1	17.55	19.189	6.31	28.7	14.54 ^{††}
Neptune	0	30.07	30.070	9.94	17.15	17.15
<i>Inferred bodies (not observed)</i>						
Vulcan (merger partner)	9	0.24	...	0.08	0.062	merged → Mercury
Theia swarm members ^{**}	4,5	2–3.5	...	...	$3.9 \sum$	scattered ^{***}

[†] Mercury’s mass is the iron-enriched merger remnant of slot 8 + slot 9 Vulcan ($\eta_{\text{retain}} = 0.30$ at $\Delta v/v_{\text{esc}} \approx 2.5$; §3.1).

[‡] Earth’s +11% excess attributed to inner-scattered Theia swarm embryo impact (Moon-forming event; §5.6).

^{*} Mars’s −92% deficit attributed to Jupiter-driven slot 4-5 swarm dispersal: 1-2 inner-scattered embryos survive as Mars at $r < r_5$ (§5.6).

^{**} Slot 4 (snow-line pile-up zone) entirely dispersed by Jupiter scattering, the “Theia swarm”.

^{***} Saturn’s −29% deficit attributed to cumulative envelope-stripping by outer-scattered swarm impactors via multi-pass Jupiter acceleration (§5.6).

^{††} Uranus’s −49% deficit attributed to outer-scattered swarm impactor event producing the 98° axial tilt + envelope strip (§5.6).

Compact systems (e.g., TRAPPIST-1, Kepler-90) thereby resolve to fast primordial rotators with strongly compressed discs, while wide architectures like Sol sit near $\Omega = 1$. In rare cases the outermost planet lies beyond the dam entirely: a gas giant that formed at slot 0 and was destabilized outward (HD 219134’s outermost giant is the catalogue’s clearest example).

3. CALIBRATION TO THE SOLAR SYSTEM

We apply the model to the Solar System using inputs $M_\star = 1.14 M_\odot$, $\Omega = 0.995$ (from Eq. 23 with R_{disc} at Neptune’s orbit), and $f_{\text{disc}} = 0.01$. These three parameters, with the universal constants of Table 2, fix the disc properties ($R_{\text{disc}} = 30.07$ AU, $R_A = 0.20$ AU, $r_{\text{snow}} = 2.70$ AU, $\sigma = 0.304 M_\oplus/\text{AU}$, $a = 0.596 M_\oplus$).

Table 3 reports the resulting fit for the ten bodies of the Solar System, including the inferred Theia swarm at the snow line and Vulcan as Mercury’s merger partner. The ISU anchors Venus and Neptune fit exactly by construction; every other body’s deviation is attributed, with independent observational support, to one of the two dynamical events (table footnotes).

The deviations between predicted and observed masses are all attributable to the framework’s two dynamical events: the Mercury–Vulcan adjacent-slot merger (§3.1)

and Jupiter’s scattering of the slot 4–5 embryo swarm (§5.6), whose asymmetric inner/outer split accounts for the Theia impact on Earth, the Mars deficit, the Saturn and Uranus envelope losses and tilts, and the near-empty asteroid belt.

3.1. *The Vulcan merger and Mercury’s iron-enriched composition*

Mercury’s anomalously high core mass fraction ($\sim 70\%$ Fe by mass) (Nittler et al. 2011; Hauck et al. 2013) is the most prominent compositional anomaly in the inner Solar System. The mainstream interpretation has been that Mercury was originally a chondritic-composition body that subsequently lost a substantial fraction of its silicate mantle. Three mechanism classes have been proposed: single giant impact (Benz et al. 1988, 2007), hit-and-run sequences (Asphaug & Reufer 2014; Chau et al. 2018), and high-temperature nebular processing (Cameron 1985; Fegley & Cameron 1987; Ebel & Alexander 2011). Each has tensions with observations (Peplowski et al. 2011), and none has emerged as a consensus mechanism. The sharpest discriminator is volatile abundance: high-temperature nebular processing and vaporizing giant impacts predict strong depletion of moderately volatile elements, yet MESSENGER found Mercury’s surface volatile-*rich* (high S and K; Peplowski et al. 2011; Nittler et al. 2011): falsifying the high-temperature class outright, while remaining fully consistent with a *cold* merger of two icy-rocky inner-cascade bodies whose energy went into mantle ejection rather than global vaporization.

Our framework reverses the direction of inference. Mercury is not a stripped chondritic body; it is the *merger remnant* of two adjacent inner-cascade slot bodies, with the iron enrichment arising naturally from the energetics of the merger event itself.

The local-capacity cascade allocates planet-forming material to geometrically-spaced slots inward from the outer Davis Dam at R_{disc} . For Sol, the geometric ratio $\rho \approx 0.584$ (the half-amplitude-at-45° projection of the Gaussian accretion-zone HWHM; see §2) generates slot positions $r_n = R_{\text{disc}}\rho^n$. Slot 8 sits at $r_8 \approx 0.41$ AU (Mercury’s current position), and slot 9 sits at $r_9 \approx 0.24$ AU. The framework’s slot 9 body, which we name *Vulcan* after Le Verrier’s 19th-century hypothesis of a sub-Mercury planet (Le Verrier 1859), grew during the protoplanetary phase to $\sim 0.06 M_{\oplus}$ in primordial composition (rocky core).

Hill spacing.—Orbital separations in this paper are expressed in Hill radii. The Hill radius of a body of mass m orbiting at semi-major axis a ,

$$R_H = a \left(\frac{m}{3M_{\star}} \right)^{1/3}, \quad (24)$$

is the radius of the region in which the body’s gravity dominates over the star’s tidal field (Hill 1878; Murray & Dermott 1999). For a planet pair the *mutual* Hill radius, $R_{H,\text{mut}} = \frac{1}{2}(a_1 + a_2) [(m_1 + m_2)/3M_{\star}]^{1/3}$, is the natural unit of orbital spacing: numerical experiments show that similar-mass pairs separated by less than $\sim 10 R_{H,\text{mut}}$ are dynamically unstable, encountering each other on timescales short compared to

the system age (Gladman 1993; Chambers et al. 1996), with ~ 7 mutual Hill radii the empirical stability boundary for adjacent bodies in multi-planet chains (Smith & Lissauer 2009).

The Mercury–Vulcan merger.—The two bodies are dynamically packed at ~ 4 mutual Hill radii of separation: well below the stability threshold. They are guaranteed to encounter each other gravitationally; the only question is the encounter outcome.

The relevant encounter energy is set by the orbital velocity difference between the two slots:

$$\Delta v_{\text{orb}} = |v_{\text{orb}}(r_9) - v_{\text{orb}}(r_8)| \approx 15.4 \text{ km/s}, \quad (25)$$

where $v_{\text{orb}}(r) = 29.785\sqrt{M_\star/r}$ km/s at Sol. The escape velocity of the combined merger body ($M_{\text{combined}} \approx 0.17 M_\oplus$ of rocky material) is

$$v_{\text{esc}} = 11.186 M_{\text{combined}}^{1/3} \approx 6.2 \text{ km/s}. \quad (26)$$

The energy ratio $\Delta v_{\text{orb}}/v_{\text{esc}} \approx 2.5$ places the encounter deep in the *catastrophic-stripping regime*: the kinetic energy delivered substantially exceeds the gravitational binding energy of the merged body’s silicate mantle. Standard impact-erosion physics (Asphaug 2010; Leinhardt & Stewart 2012) predicts that under these conditions, the silicate mantle of both bodies is preferentially ejected while the dense iron cores merge and are retained. The controlling parameter $\Delta v/v_{\text{esc}}$ is classical: it is the inverse of the Safronov number that has governed accrete-versus-erode outcomes since Safronov (1972), and the quantitative accretion-efficiency curves of Leinhardt & Stewart (2012) and the hit-and-run mapping of Asphaug & Reufer (2014) supply the impact-physics basis for the retention law adopted below: mechanically, our adjacent-slot merger is a similar-mass, modest- Δv collision of exactly the class those works parameterize.

We adopt the empirical retention function

$$\eta_{\text{retain}} = \max[0.3, 1 - 0.37 (\Delta v_{\text{orb}}/v_{\text{esc}})] \quad (27)$$

calibrated against (a) Mercury’s observed retention fraction (~ 0.30 relative to combined Vulcan+Mercury primordial mass) and (b) the milder Tau Ceti e merger event (~ 0.81 retention at $\Delta v_{\text{orb}}/v_{\text{esc}} \approx 0.53$). The lower-bound 0.30 floor corresponds to the iron-core-only residual when the entire silicate envelope is ejected.

Composition prediction.—Combined primordial Vulcan + Mercury rocky core mass is $\sim 0.17 M_\oplus$. At $\eta_{\text{retain}} = 0.30$, the residual post-merger body has mass $\sim 0.051 M_\oplus$, within 8% of Mercury’s observed $0.055 M_\oplus$. The retained composition is iron-dominant because (i) iron is centrally concentrated in each pre-merger body’s differentiated core, (ii) the combined post-encounter core sinks inward as denser material, and (iii) silicate mantle is preferentially ejected as outer-layer material. Mercury’s $\sim 70\%$ Fe bulk composition is thus the natural consequence of the merger geometry, not the result of post-formation impact stripping of an originally chondritic body.

This resolves the Mercury anomaly without requiring (i) a separate giant impactor of specific kinematics, (ii) extreme nebular processing, or (iii) hit-and-run with larger embryos. The Mercury–Vulcan pair were always packed at unstable mutual Hill spacing; their merger was energetically constrained to produce an iron-enriched remnant.

Comparison to other merger systems.—The retention function’s mild end is anchored by the Tau Ceti e partial-merger interpretation: at $\Delta v_{\text{orb}}/v_{\text{esc}} \approx 0.53$ the formula gives $\eta_{\text{retain}} \approx 0.81$, a low-energy merger retaining most of the combined mass. The two anchors thus calibrate opposite ends of the merger-energy spectrum. We predict that mergers at other heliocentric distances and stellar masses follow the same scaling, with retention set by local orbital and escape velocities rather than ad hoc impactor kinematics.

The same model has been applied to the 22 further catalogued systems (Table 9), spanning spectral classes A–M and architectures from compact super-Earth chains to wide-orbit super-Jupiters and multi-star systems; the calibrated f_{disc} distribution across that catalogue is analysed in §11.2.

4. ORBITAL SIGNATURES OF IMPACT EVENTS:

VENUS AND NEPTUNE AS THE PRIMORDIAL-PLANE REFERENCE

The standard practice of using Earth’s orbit (the ecliptic) as the reference plane for Solar System inclinations is dynamically biased. Earth’s orbital plane was perturbed by the Theia impact (Canup & Asphaug 2001; Canup 2012); using it as zero-reference contaminates the inclination of every other planet by a contribution that is itself part of the impact-event signature we seek to identify. We resolve this by using *Venus and Neptune as the primordial-plane reference*. This choice is supported by four independent lines of evidence presented in §4.1.

4.1. *Venus and Neptune as ISU: a four-line proof*

The framework’s identification of Venus and Neptune as the Solar System’s only in-situ unchanged (ISU) bodies is not an arbitrary or convenience-based assignment. Four observationally independent lines of evidence converge on this same identification.

Line 1: Venus and Neptune are the most circular orbits.—The eccentricities of the eight Solar System planets, ordered from most to least circular:

Venus	$e = 0.0068$
Neptune	$e = 0.0086$
Earth	$e = 0.0167$
Uranus	$e = 0.0457$
Jupiter	$e = 0.0489$
Saturn	$e = 0.0565$
Mars	$e = 0.0934$
Mercury	$e = 0.2056$

Venus and Neptune occupy the top two positions with a clean factor-of-two gap separating them from Earth and all other planets. No other planetary pairing in the Solar System shows this kind of clean ranking separation. We note that osculating eccentricities oscillate over secular cycles (Earth’s, for instance, will fall to ~ 0.003 – 0.005 over the next ~ 30 kyr), so the snapshot ranking is suggestive rather than decisive on its own; the durable content of this diagnostic is carried by the free amplitudes and angular-momentum-deficit argument of Line 2.

Line 2: Eccentricity is a one-way integral of post-formation dynamical history.—Gas-disc dynamical friction and gas-drag damping during the disc-active phase drive orbits toward a near-circular *floor* ($e \lesssim 0.01$): damping is efficient but finite, leaving a small residual rather than $e = 0$ (Goldreich & Tremaine 1980; Ida 1990; Kominami & Ida 2002). Once the gas disc disperses (~ 5 – 10 Myr post-formation), no efficient eccentricity-damping mechanism exists for planetary-mass bodies, and Kominami & Ida (2002) show explicitly that post-gas dynamics pump eccentricity *up* from that floor. Subsequent dynamical events, migration (Walsh et al. 2011; Pierens & Raymond 2011), scattering (Tsiganis et al. 2005; Morbidelli et al. 2007), mean-motion resonance crossings, giant-impact events (Canup & Asphaug 2001), can only *add* eccentricity, not remove it: a body observed today *at* the gas-era floor has experienced none of them. Individual eccentricities do oscillate reversibly through secular angular-momentum exchange between the planets, but the system-level budget of that exchange, the angular momentum deficit (AMD; Laskar 1997), can only grow once the gas is gone, so a planet whose free eccentricity amplitude is near zero can never have been significantly excited.

Venus’s and Neptune’s $e \approx 0.008$ is therefore not just small, it is small in an irreversible sense. Whatever has happened to these two bodies since gas-disc dispersal ~ 4.5 Gyr ago, it has not been substantial enough to perturb them from their primordial near-circular state. The same statement cannot be made for any other planet in the system: Earth has been perturbed by Theia, Mars by the Borealis impactor, Mercury by Vulcan, the gas giants by mutual resonance evolution. This reading has support independent of the present framework: N -body reconstructions of the pre-excitation terrestrial system require Venus’s primordial free eccentricity amplitude to have been near zero, with the terrestrial orbits dynamically colder than today (Brasser et al. 2013).

Line 3: ISU diagnostics are the inverse of established gravitational-capture diagnostics.—The orbital signatures used to identify gravitational capture are well established in the satellite-dynamics literature (Jewitt & Haghighipour 2007; Nesvorný et al. 2007; Sheppard & Jewitt 2003):

- High orbital inclination relative to the host’s reference plane (Neptune’s Triton at $i \approx 157^\circ$; Saturn’s Phoebe at $i \approx 175^\circ$);
- High eccentricity (most Jupiter and Saturn irregular satellites have $e \gtrsim 0.1$);

- Retrograde orbital motion (the strongest single capture indicator);
- Position far from the host’s natural formation/cascade zone.

In-situ formation is, by definition, the absence of these signatures. A body that formed and remained in its current orbit shows: prograde orbit, low inclination relative to the system’s formation plane, low eccentricity, and position consistent with the local formation cascade. Applying the established capture-diagnostic methodology in the inverse direction identifies in-situ bodies by their lack of capture signatures.

For Venus: $e = 0.0068$, $i = 3.39^\circ$ (ecliptic-referenced; only 1.99° from the VN baseline), prograde orbit, position consistent with slot 7 of the cascade.

For Neptune: $e = 0.0086$, $i = 1.77^\circ$ (ecliptic-referenced; only 1.99° from the VN baseline), prograde orbit, position consistent with slot 0 of the cascade.

Both bodies score positive on every in-situ indicator and negative on every capture indicator. By the same methodology that identifies Triton as captured, Venus and Neptune are identified as in-situ.

Line 4: Mass-measurement precision.—Independent of orbital considerations, Venus and Neptune have masses known to part-per-million precision from spacecraft tracking (Magellan, Cassini-Huygens, Voyager 2). Their cascade-allocated mass predictions ($M_{\text{pred}}^{\text{Venus}} = 0.815 M_\oplus$ from the slot 7 rocky allocation; $M_{\text{pred}}^{\text{Neptune}} = 17.15 M_\oplus$ from the slot 0 cascade) match observed values to better than $0.001 M_\oplus$. No other Solar System body achieves this combination of precision measurement *and* exact cascade match. The other rocky planets either have observed masses substantially different from cascade predictions (Mercury -49% , Mars -92% , both attributable to the framework’s two dynamical events) or have δM contributions from late delivery (Earth $+11\%$ from inner-scattered swarm embryo). The non-anchor gas giants show impactor-event-induced deficits: Saturn -29% from cumulative outer-scattered swarm impacts; Uranus -49% from a single high-energy swarm impactor producing both the envelope strip and the 98° axial tilt.

Convergence.—The four lines, most circular orbit (Line 1), one-way eccentricity preservation (Line 2), inverse-of-capture diagnostics (Line 3), exact mass-cascade match (Line 4), converge on the same identification. Venus and Neptune are uniquely ISU among Solar System bodies. The designation is not a convenience choice but a physically grounded observational conclusion.

Relation to prior art.—For Venus, the equivalent concept has precedent under different nomenclature: [Jacobson et al. \(2017\)](#) attribute Venus’s absent magnetic dynamo to a primordially stratified core that was never mechanically mixed (that is, to Venus never suffering a late energetic giant impact) and [Brasser et al. \(2013\)](#) require its primordial free eccentricity to be near zero. The [Jacobson et al. \(2017\)](#) mechanism constrains more than the absence of one late impact: a stratified, never-stirred core requires Venus’s *entire* accretion to have proceeded through many small, low- Δv impactors, a

geophysical witness, independent of the orbital evidence, that Venus sits on the quiet branch of the impact-energetics ledger this framework assigns it. For Neptune, the mainstream holds the opposite: the resonant Kuiper belt populations (Pluto and the Plutinos in Neptune’s 3:2 mean-motion resonance, with pumped eccentricities) are interpreted as bodies swept up and captured during an outward migration of Neptune by several AU (Malhotra 1993, 1995). The framework reads the same populations differently. In the cascade picture the Kuiper belt is not cascade material at all: it forms *outside* the Davis Dam ($R_{\text{disc}} = 30.07$ AU, Neptune’s slot), by slow accretion in the exterior reservoir that pools against the dam from outside (§2), a different formation mechanism from the slot cascade. Neptune’s exterior mean-motion resonances (3:2 at 39.4 AU, 2:1 at 47.7 AU) all lie within that exterior region, and bodies drifting inward toward the dam are captured into the resonances of a *stationary* Neptune by drag-induced convergent drift, their eccentricities pumped to equilibrium during continued drift: the classic exterior-resonance trapping mechanism of Weidenschilling & Davis (1985). Adiabatic resonance capture requires only convergent relative drift: the standard model moves the planet, the framework moves the particles.

Drag trapping cannot hold dwarf-planet-mass bodies (drag acceleration falls with size), but those do not need it. Bodies that condense at the dam itself are born sharing Neptune’s orbital radius (the dam is two-sided, producing Neptune from the interior cascade allocation and Pluto-class bodies from the exterior pool) and a body born co-orbital with a $17 M_{\oplus}$ giant has no non-resonant survival path: non-resonant Neptune-crossing orbits are cleared on 10–100 Myr timescales, so after 4.5 Gyr the only survivors are those in protecting mean-motion resonances, whose libration geometry forbids close approaches. Pluto’s eccentricity is then the scar of the scattering that preceded capture, not the pump of a migrating resonance. Its deep libration and Kozai state, historically read as fingerprints of slow adiabatic capture, are equally fingerprints of *survival*: erosion preferentially removes large-amplitude librators (Tiscareno & Malhotra 2009), so any initial resonant population converges to deep, Kozai-rich microstates, and resonance sticking continually feeds the islands from the scattered population (Lykawka & Mukai 2007). Two further observations favour this channel: the distant 5:2 resonance is anomalously well-populated, a standing puzzle for smooth migration (Gladman et al. 2012) but the natural outcome of sticking from an eccentricity-pumped scattered source, and Triton (retrograde, Pluto-like) is the expected prize of binary-exchange capture (Agnor & Hamilton 2006) acting on the factory’s natively-binary output. The two claims interlock: binary-exchange capture *requires* a substantially binary source population, and gravitational collapse at the dam natively produces exactly the wide, near-equal-mass binary fraction observed in the cold classicals (Nesvorný et al. 2010): the factory supplies the prerequisite that makes Triton’s capture mechanism available. The competing channels predict different joint distributions of resonant occupancy, eccentricity, and libration amplitude (§10), making the resonant Kuiper belt a sharp observational discriminator between

the framework and migration-based models rather than a refutation of Neptune’s ISU status.

4.2. *The Venus-Neptune mean plane*

Each planet’s orbital plane is specified by its inclination i (tilt from a reference plane) and longitude of ascending node Ω . The orbital plane normal vector (in ecliptic-referenced Cartesian coordinates with $\hat{z}$ along the ecliptic pole) is

$$\hat{n}(i, \Omega) = (\sin i \sin \Omega, -\sin i \cos \Omega, \cos i). \quad (28)$$

For Venus ($i_V = 3.39^\circ$, $\Omega_V = 76.68^\circ$) (Williams 2024a; Standish 1998):

$$\hat{n}_V = (0.0575, -0.0136, 0.9982). \quad (29)$$

For Neptune ($i_N = 1.77^\circ$, $\Omega_N = 131.78^\circ$) (Williams 2024b; Standish 1998):

$$\hat{n}_N = (0.0230, 0.0206, 0.9995). \quad (30)$$

The Venus-Neptune mean plane normal is the average:

$$\hat{n}_{VN} = \frac{\hat{n}_V + \hat{n}_N}{|\hat{n}_V + \hat{n}_N|} \approx (0.0403, 0.0035, 0.9989). \quad (31)$$

The VN plane is offset from the ecliptic by $i_{VN} \approx 2.71^\circ$ at ascending node $\Omega_{VN} \approx 95^\circ$. By construction, both Venus and Neptune sit at 1.99° from this plane (in opposite directions).

The angle of any planet’s orbital plane from the VN plane is

$$i_{VN} = \arccos(\hat{n}_{\text{planet}} \cdot \hat{n}_{VN}). \quad (32)$$

4.3. *Inclinations relative to the VN baseline*

Computing Eq. 32 for all eight planets, then subtracting the 1.99° baseline so that Venus and Neptune sit at zero:

The pattern is consistent with the framework’s predicted impact history. Bodies identified as impact-modified (Mercury via Vulcan, Earth via inner-scattered swarm embryo, Mars via swarm dispersal, Uranus via outer-scattered swarm impactor) all show positive inclination excess relative to the VN baseline. Bodies identified as in-situ formation products (Venus, Neptune) sit at zero by construction. Jupiter and Saturn show small negative offsets, reflecting the current invariable plane (dominated by their mass-weighted angular momentum) sitting slightly below the VN primordial plane. The $\sim 0.3^\circ$ offset between the current invariable plane and the VN primordial plane is the cumulative signature of impactor-induced perturbations on Jupiter and Saturn during the slot 4-5 swarm dispersal event, not of any large-scale migration.

Table 4. Solar System inclinations relative to the Venus-Neptune primordial plane. Negative values indicate orbits below the VN plane (toward the invariable plane).

Planet	i_{VN} (°)	Δi_{VN} (°)	Status
Saturn	1.62	−0.37	Theia outer-swarm envelope stripped anchor body
Jupiter	1.81	−0.18	
Venus	1.99	$\equiv 0$	ISU
Neptune	1.99	$\equiv 0$	ISU
Mars	2.14	+0.15	Borealis impact
Uranus	2.13	+0.14	Outer-scattered swarm impactor
Earth	2.71	+0.72	Theia impact
Mercury	5.94	+3.95	Vulcan merger

Table 5. Eccentricity excess relative to the Venus-Neptune baseline ($\bar{e}_{VN} = 0.008$).

Planet	e	Δe	$e/\bar{e}_{VN}$
Venus	0.0068	−0.001	~ 1
Neptune	0.0086	+0.001	~ 1
Earth	0.0167	+0.009	2
Uranus	0.0457	+0.038	6
Jupiter	0.0489	+0.041	6
Saturn	0.0565	+0.049	7
Mars	0.0934	+0.085	12
Mercury	0.2056	+0.198	26

4.4. Eccentricity excess

Having established Venus and Neptune as the ISU reference (§4.1), we use their near-circular orbits ($e_V = 0.0068$, $e_N = 0.0086$) as the baseline ($\bar{e}_{VN} \approx 0.008$) and compute the eccentricity excess of each remaining planet relative to this baseline:

Mercury’s eccentricity ($e \approx 0.21$) is dramatically elevated, a factor ~ 26 above the VN baseline. This is consistent with a high- $\Delta v/v_{\text{esc}}$ merger event delivering substantial orbital momentum exchange in addition to the catastrophic mass-loss signature. Mars’s $e \approx 0.09$ ($12\times$ baseline) is similarly consistent with its scattering history and the Borealis impact. Saturn and Uranus show $\sim 6\text{--}7\times$ baseline eccentricity, consistent with the swarm bombardment overlaid on mutual gas-giant secular evolution; Jupiter’s comparable excess reflects its role as the scattering source.

4.5. Coupled signatures: four mass-change events, four orbital records

The combination of inclination excess and eccentricity excess provides a coupled diagnostic for the four mass-changed planets. For each event, three or four independent signatures should appear: mass anomaly (§3), inclination excess (§4.3), eccentricity excess (§4.4), and (for some) axial tilt:

Table 6. Coupled orbital signatures of the framework’s four mass-change events. Each attribution is supported by *multiple* independent orbital observables on its target body.

Target	Mass anomaly	Δi_{VN}	Δe	Tilt / spin
Mercury (Vulcan merger)	70% Fe	+3.95°	+0.198	0° tilt (locked)
Mars (swarm scattering + Borealis)	−1.22 $M_{\oplus}$	+0.15°	+0.085	25.2° tilt
Saturn (cumulative swarm impacts)	−39 $M_{\oplus}$	−0.37°	+0.049	26.7° tilt
Uranus (outer-scattered swarm impactor)	−14 $M_{\oplus}$	+0.14°	+0.038	98° tilt, envelope strip

Mercury (Vulcan merger).—Four independent signatures from the same high- $\Delta v/v_{\text{esc}}$ encounter event: composition, mass, inclination, and eccentricity all extreme. The Vulcan body (slot 9 sub-Mercury) merger explains all four through the impact retention formula calibrated on this event.

Mars (inner-scattered swarm survivor).—Hemispheric dichotomy (Borealis basin), severe mass deficit (−92%), mild inclination perturbation, substantial eccentricity excess. Mars is one of ~ 1 –2 inner-scattered embryos from the slot 4–5 swarm that survived at $r < r_5$ (below Jupiter’s chaotic zone); the Borealis-basin impact was a smaller sibling-embryo arriving at high velocity.

Saturn (cumulative outer-swarm impacts).—29% envelope deficit, 26.7° axial tilt, $7\times$ baseline eccentricity, and a small negative Δi_{VN} (the invariable-plane offset of §4.6). A multiple-impact origin predicts exactly this combination: intermediate tilt and distributed, rather than single-event, orbital perturbation.

Uranus (outer-scattered swarm impactor).—98° axial tilt, 49% envelope loss, small inclination perturbation, moderate eccentricity excess. The small Δi_{VN} despite massive impactor energy indicates an *in-plane impact trajectory*, consistent with the impactor having been scattered through the Jupiter-Saturn ecliptic plane to reach Uranus’s orbit. The impact transferred angular momentum to spin (tilting the planet) without significantly displacing the orbital plane.

4.6. The original disc plane and the invariable-plane offset

The Venus-Neptune plane’s 1.99° offset from both Venus and Neptune reflects the natural inclination dispersion of bodies forming in a protoplanetary disc of finite vertical scale height ($H/r \approx 0.03$ – 0.05 at typical disc temperatures and stellar masses; Armitage 2018). Bodies forming within such a disc inherit small random inclinations on the order of $\arctan(H/r) \approx 2$ – 3° . Venus and Neptune happen to sit on opposite sides of the disc midplane at characteristic disc-thickness offsets.

The *current* invariable plane of the Solar System, defined by the total angular momentum vector (dominated by Jupiter and Saturn), sits at $i \approx 1.58^\circ$ relative to the ecliptic (Souami & Souhay 2012), or equivalently $\approx 1.6^\circ$ from the VN primordial plane. This offset is the **cumulative impactor-perturbation signature**: Jupiter’s

and Saturn’s orbital planes shifted by $\sim 0.3^\circ$ under the swarm-impactor bombardment, dragging the system’s angular-momentum-weighted invariable plane below the primordial disc plane.

4.7. Refined ISU criterion

The orbital analysis suggests a refined criterion for assigning the ISU flag in the framework:

ISU-eligible: when a planet has *both*:

- Precise, well-characterized observed mass and orbit
- Orbital state (i, e) consistent with undisturbed formation (low Δi , low Δe relative to a system-mean baseline)

Not ISU-eligible: when orbital signatures indicate post-formation perturbation, even if mass is precisely measured. Such planets show distinct Δi or Δe excursions whose origin is identifiable as an impact, scattering, or migration event.

For Sol, only Venus and Neptune satisfy both conditions, making them the natural ISU anchors: they are the only bodies whose mass and position can be used as absolute constraints on the disc parameters without contamination from post-formation events.

Operational diagnostic: minimum-eccentricity selection.—In practice, the planets with the lowest observed eccentricities are the strong ISU candidates, with the number selected set by how cleanly an eccentricity gap separates near-circular bodies from the rest of the system. For Sol, the factor-of-two gap below Earth’s $e = 0.0167$ selects exactly Venus and Neptune. Because eccentricity integrates post-formation dynamical history one way (Line 2 above), preserved near-circularity directly identifies undisturbed bodies. Systems with no clean gap may have no undisturbed bodies; the disc parameters are then inferred jointly from all observed planets rather than pinned to ISU anchors, Beta Pictoris’s directly-imaged super-Jupiters, whose orbital states are poorly constrained, are handled this way.

5. SATURN AND JUPITER: IN-SITU OUTER ARCHITECTURE

The geometric cascade $\rho = 1 - \sqrt{\ln 2}/2 \approx 0.584$ places slot 2 at $r_2 = 10.24$ AU and slot 3 at $r_3 = 5.98$ AU for Sol. Saturn (observed 9.54 AU) and Jupiter (observed 5.20 AU) sit ~ 0.7 AU inside their cascade positions, a minor inward drift, not a 4-AU or 8-AU migration. **The Solar System’s outer architecture is essentially in-situ.** The framework reproduces Saturn’s and Jupiter’s positions and masses with the standard cascade allocation plus a single envelope-loss event for Saturn (§5.6); no Grand Tack or Nice Model is invoked.

5.1. *Jupiter as the in-situ anchor*

Jupiter at slot 3 ($r_3 = 5.98$ AU) with cascade-default $t_{\text{form}} = 1.71$ Myr predicts $M_{\text{pred}} = 318.9 M_{\oplus}$, matching the observed $317.83 M_{\oplus}$ to 0.3%. The Tanigawa-Ikoma $A_0 = 4.45$ ($1/M_{\oplus}$) and $k = 0.691 \text{ Myr}^{-1}$ calibration is anchored on Jupiter’s mass-position match, and the same constants reproduce Neptune at slot 0. Jupiter’s ~ 0.78 AU inward drift from slot 3 to its observed position is consistent with gas-disc torque damping during the gas-active phase, and is a factor of five smaller than the Grand Tack’s postulated ~ 4 AU inner excursion.

Jupiter’s stable position is consistent with the isotopic constraint of [Kruijer et al. \(2017\)](#): Jupiter formed early and remained stably between the NC and CC reservoirs without major inner excursions. The constraint is in fact stronger than a formation clock: the non-carbonaceous and carbonaceous meteorite reservoirs remained isotopically *separated* from ~ 1 to $\sim 3\text{--}4$ Myr, which requires a sustained transport barrier between inner and outer disc for the entire interval: a barrier a stationary slot-3 Jupiter atop the framework’s snow-line pile-up supplies naturally, and that any large radial excursion would have breached. The same cosmochemistry thereby witnesses both of this section’s claims at once: the giant stayed put, and the disc was partitioned by standing dams.

5.2. *Saturn in-situ at slot 2*

Saturn at slot 2 ($r_2 = 10.25$ AU) with default $t_{\text{form}} = 3.13$ Myr predicts $M_{\text{pred}} = 134 M_{\oplus}$. Saturn’s observed mass is $95.16 M_{\oplus}$, a $-39 M_{\oplus}$ (-29%) deficit relative to the cascade prediction. This deficit is attributed to envelope-stripping events during the Jupiter-driven slot 4-5 swarm dispersal (§5.6). Cumulative impactor kinetic energy from outer-scattered swarm members reaches $\sim 75\%$ – 100% of Saturn’s $\sim 2 \times 10^{35}$ J envelope binding energy over multiple Jupiter-accelerated transits, consistent with the observed deficit.

Saturn’s observed position at 9.54 AU sits 0.70 AU inside slot 2, consistent with combined disc-gas torque damping and cumulative gravitational interactions during impactor transits.

5.3. *Saturn’s axial tilt as an impact signature*

Saturn’s 26.7° axial tilt is intermediate between Jupiter’s 3.1° (anchor, no impacts) and Uranus’s 97.8° (extreme direct impact). The progressive tilt magnitudes (Jupiter $\rightarrow$ Saturn $\rightarrow$ Uranus) reflect cumulative angular momentum delivery from the slot 4-5 outer-swarm impactor population: Jupiter is the scattering source and receives no swarm impacts; Saturn receives multiple direct impacts that tilt its axis and strip $\sim 29\%$ of its primordial gas envelope; Uranus receives the highest-velocity direct impact, producing both its extreme axial tilt and substantial envelope loss. The progression is a natural prediction of the framework’s swarm-dispersal mechanism, with impact severity increasing with heliocentric distance from the Jupiter scattering source as

the swarm acquires successively higher orbital eccentricities through repeated Jupiter encounters.

The standard explanation for Saturn’s obliquity is instead a secular spin-orbit resonance with Neptune, entered as Kuiper-belt mass depletion slowed the precession of Neptune’s orbit plane (Ward & Hamilton 2004). The two accounts are not exclusive here: the framework’s trans-Neptunian evolution (§8) supplies the required belt-mass depletion without planetary migration, leaving the resonance pathway open. Saturn’s envelope-mass deficit, however, requires impacts in either case.

5.4. *Contrast with conventional outer-system narratives*

Grand Tack (Walsh et al. 2011): requires Jupiter to migrate inward to ~ 1.5 AU and back, dragging Saturn behind it. Our model has Jupiter at 5.20 AU, essentially in-situ at slot 3, with only ~ 0.8 AU inward drift.

Nice Model (Tsiganis et al. 2005): requires the outer giants to form in a compact configuration (~ 15 – 30 AU) and reorganize via specific mean-motion-resonance crossings. Our model has all four outer giants within ~ 1.6 AU of their geometric cascade slot positions; no post-formation reorganization is invoked. We note a convergence rather than only a conflict: the compact, near-resonant initial chain the Nice family must *assume* (Morbidei et al. 2007) is precisely the configuration the cascade *produces* (adjacent slots at period ratio 2.24, just wide of 2:1), the frameworks disagree on what happened next, not on how the giants started. Removing the instability does leave the Nice Model’s auxiliary successes (the putative Late Heavy Bombardment spike; Trojan capture) in need of separate accounts; irregular-satellite capture, at least, ties naturally to the framework’s own scattering encounters (§8.8; Nesvorný et al. 2007).

The near-5:2 Jupiter-Saturn period ratio, $P_S/P_J = (9.54/5.20)^{3/2} \approx 2.48$, emerges as a natural consequence of the cascade’s geometric structure: adjacent slots with ratio $\rho \approx 0.584$ have a period ratio $\rho^{-3/2} \approx 2.24$, within 10% of the observed 2.48. Both planets occupy positions close to their geometric slots without requiring ad-hoc resonance trapping or migration sequences.

5.5. *The framework’s parsimony claim*

Sol’s outer-giant architecture (Jupiter, Saturn, Uranus, Neptune all within ~ 1.6 AU of cascade slots), the Jupiter-Saturn 5:2 mean-motion resonance, Saturn’s -29% mass deficit, Saturn’s modest axial tilt, Uranus’s extreme axial tilt and envelope loss, the asteroid belt’s near-empty mass, and Earth’s Moon are all reproduced from: (i) the geometric cascade allocation; (ii) the Mercury-Vulcan adjacent-slot merger event (§3.1); (iii) Jupiter’s standard gravitational scattering of the slot 4-5 embryo swarm (§5.6).

Nothing extraordinary happened. The Solar System’s observed architecture is the natural outcome of cascade formation plus one dynamical event in the inner-mid

region. The conventional narrative’s elaborate migration sequences, resonance crossings, and reorganization events are not required by the observed data once the cascade structure is correctly identified.

5.6. *Embryo-swarm dispersal as a general mechanic*

A predictive criterion follows: for any cascade slot N with mass M_N , if there exists a perturber slot P with $M_P \gg M_N$ within $\sim 5\text{--}7$ mutual Hill radii of N ’s position, then N ’s pre-consolidation embryo swarm is dispersed by Jupiter-style asymmetric scattering. The dispersal produces:

- **Inner fraction** ($\sim 5\%$): scattered to lower AU, manifests as either an embryo impact on inner planets or a small surviving body at $r < r_N$.
- **Outer fraction** ($\sim 95\%$): scattered outward via multi-pass perturber acceleration, manifests as cumulative impacts on outer-slot bodies.

The scattering mechanism itself is established: N -body simulations show that a giant’s runaway gas accretion destabilizes and scatters nearby planetesimals in all directions, with a fraction deposited inward (Raymond & Izidoro 2017). The framework differs in elevating this from one ingredient among migration scenarios to the Solar System’s *single* dynamical event, and in tying the outward-scattered majority to the Saturn and Uranus impact record.

For Sol, Jupiter at slot 3 has Hill radius $R_{H,J} = 0.34$ AU (Eq. 24). First-order mean-motion-resonance overlap (Wisdom 1980) makes orbits formally chaotic within $\Delta a/a \approx 1.3(M_J/M_\odot)^{2/7} \approx 0.18$ of Jupiter (~ 0.9 AU on each side), and repeated distant encounters destabilize loosely-bound pre-consolidation embryo swarms out to the framework’s $\sim 10 R_H$ dispersal criterion (~ 3.4 AU). Slots 4 and 5 of the cascade ($r_4 = 3.49$ AU, $r_5 = 2.04$ AU) both lie within this reach. The framework predicts:

Table 7. Predicted dispersal of cascade slots under Jupiter’s gravitational influence for Sol.

Slot	Predicted ($M_\oplus$)	Status	Outcome
4	2.5	Fully dispersed	Theia swarm
5	1.3	Partially dispersed + survivor	Mars (0.107) + outer scatter
6	0.9	In-situ + Theia chunk	Earth + Moon
7–8	rocky	In-situ	Venus, Mercury

This dispersal mechanic is the framework’s single dynamical event beyond cascade formation. It is generalizable to other systems: any cascade slot within scattering range of a sufficiently massive perturber is predicted to be either unobservable (fully dispersed) or displaced to a position *below and outside* the perturber’s chaotic zone with surviving mass at $\sim 5\text{--}10\%$ of the slot’s primordial allocation. The settling radius is the inner stability edge of the chaotic zone, $r_{\text{survivor}} \approx r_{\text{perturber}} - 11 R_H$. For

Sol this predicts a survivor at $5.20 - 11 \times 0.34 \approx 1.46$ AU carrying $0.07\text{--}0.13 M_{\oplus}$; Mars is observed at 1.524 AU with $0.107 M_{\oplus}$ (8.1% of slot 5’s predicted $1.323 M_{\oplus}$): a position match within 5%. The position-plus-mass-fraction signature is the characteristic outcome of partial dispersal.

5.7. *Migration is gravitational, not gas-torque*

Conventional gas-disc migration mechanisms (Type-I, gas torque on sub-thermal-mass cores; Type-II, viscous transport of gap-opening giants) are not invoked in this framework. Planets in the calibrated catalogue certainly migrate, and some migrate far: Mu Arae c steps off its dam outward, the wrecking pairs of HD 60532 and HD 142 fall from their birth seats to the magnetospheric rim, and 55 Cancri b transits its entire inner system. But the driver in every case is gravitational: angular-momentum exchange with more massive bodies. A body that surrenders angular momentum to a heavier neighbour drops to a lower orbit; one that extracts angular momentum rises to a higher one. Luna’s tidal recession is the same bookkeeping with Earth’s spin as the reservoir.

The same interaction obeys a mass-ratio law that splits its outcomes. When the participants are comparable, or the migrator is itself large, the body transports intact and arrives whole at its destination. When the mass ratio is extreme, the smaller body does not migrate; it is taken apart, and the catalogue documents the spectrum: swarm dispersal with the asymmetric inner/outer split (§5.6), Martian-type remnants at 5–10% of the parent, mantle-grazed survivors, and Roche deliveries ground into rings. Large bodies migrate; small bodies at extreme ratios are ripped apart. One grammar, one threshold.

The gas disc still matters, but its role is passive and architectural: it traps rather than torques. The slot lattice is a ladder of pressure bumps at which migration torque cancels (§2), so a gravitationally driven migrator in a living disc is caught at the next vacant seat, and the magnetospheric cavity rim is the terminal trap (Lin et al. 1996); the wrecking pairs park bracketing their fitted Alfvén radii, and HD 142 c sits on the exact seat its companion vacated. Parking position therefore dates the move: a migrator seated on the lattice moved while the gas lived; one parked off-lattice moved after the gas was gone, as Jupiter’s small post-swarm recoil and Luna’s tidal march both illustrate.

This is a substantive claim that extends beyond the framework’s calibration: the observational evidence for Type-I and Type-II migration as continuous gas-torque processes may be substantially weaker than commonly attributed. What has been called migration is better described as gravitational scattering settling into trap-stabilised orbits. The distinguishing tests are the asymmetric inner/outer mass fraction of scattering (the $\sim 5\%/95\%$ split) against continuous migration’s smoothly varying distributions, and the parking statistics: gas-torque migration predicts no preference

for fossil seat positions among migrated planets, while trap-terminated gravitational migration predicts that wet-era migrators sit on them.

5.8. *Brown-dwarf vs gas-giant mutual eviction in other systems*

Eviction is the migration grammar of §5.7 carried past its boundaries. The same gravitational angular-momentum exchange that lifts a body to a higher orbit, given enough exchange, lifts it past escape velocity and out of the system entirely; the same exchange that drops a body to a lower orbit, continued, drops it through the rim and into the star. The catalogue holds both endpoints: Proxima left Alpha Centauri with an energy surplus of barely ten percent, the marginal escape of a body nudged just past the threshold, while the protostellar-consumption channel records the opposite exit, and Sol’s own eviction from Hyperion sits in between as the gentlest recorded departure. Migration, ejection, and consumption are one process read at three depths of the exchange; shredding (§5.7) is what the same process does to a body too small to survive it. Which depth each participant reaches depends on the encounter. The pair must balance its own books only when no third party pays: in a full relaxation the destroyed slots and consumed bodies carry part of the angular-momentum bill, so the menu includes one partner up and one down, both partners lifted together (Alpha Centauri’s relaxation raised B from slot 1 to the vacated dam seat while ejecting Proxima from it, both bills paid by the obliterated interior slots), mutual ejection when both clear escape, and ejection paired with a plunge into the star.

Sol’s outer giants do not undergo system ejection: encounters between $\sim 100 M_{\oplus}$ -scale bodies do not deliver hyperbolic excess velocity (speed beyond what is needed to escape the system), so mass is redistributed within the bound system rather than ejected. Genuine mutual eviction requires bodies above roughly the deuterium-burning threshold ($13 M_J \approx 4000 M_{\oplus}$, the mass at which an object briefly fuses deuterium and is classed a brown dwarf; Spiegel et al. 2011), where mutual encounter velocities can exceed the system escape velocity. The exit assignments then follow the encounter, not a rule: one ejected and one consumed, both ejected, or both displaced outward with the deficit charged to the rest of the cascade. The framework flags empty adjacent slots whose predicted masses both exceed this threshold as mutually evicted. The distinction matters for Sol: the literature’s “ejected fifth giant” hypotheses (Nesvorný 2011; Nesvorný & Morbidelli 2012) demonstrate that Jupiter encounters *can* eject an ice-giant-mass body, their five-planet simulations succeed at it, but the framework’s point is parsimony, not impossibility: no fifth body is needed here, because the outer architecture is reproduced by Jupiter’s scattering of the slot 4–5 swarm alone, and the cascade allocates no slot for one.

Jupiter’s recoil from swarm scattering.—Jupiter’s ~ 0.8 AU inward drift from slot 3 ($r_3 = 5.98$ AU) to its observed position (5.20 AU) is consistent with momentum-conservation recoil from scattering $\sim 3.9 M_{\oplus}$ of the slot 4–5 swarm material outward. For typical scattering velocity boost $\Delta v_{\text{swarm}} \approx 30$ km/s, momentum conservation

gives $\Delta v_J = -(M_{\text{swarm}}/M_J) \Delta v_{\text{swarm}} \approx -0.37 \text{ km/s}$, producing an inward drift $\Delta r_J \approx 0.4 \text{ AU}$. This accounts for roughly half of Jupiter’s observed inward drift from slot 3; the remainder is consistent with disc-gas dynamical friction during the gas-active phase. No large-scale Type-II migration is required.

6. CASCADE FRAMEWORK AT STELLAR SCALES

The cascade framework’s geometric ratio $\rho = 0.5837$ and mass-allocation formula are dimensionless. Their natural extension to multi-star systems and stellar nurseries follows once the host mass and primordial f_{disc} are scaled appropriately. This section presents three demonstrations: (i) Alpha Centauri as binary/triple formation in a single protoplanetary disc; (ii) the Crab Nebula’s supernova progenitor as a scaled-up version of the same physics producing a multi-star cluster; (iii) the predicted compositional bimodality between cascade bodies forming inside vs outside their parent’s snow line.

6.1. *Alpha Centauri as in-disc multi-star formation*

Applied to Alpha Centauri with input mass $M_\star = 1.08 M_\odot$, spin = 1.29 (Sol-primordial units), and a viable $f_{\text{disc}} = 0.25$ (the all-stellar architecture leaves f_{disc} formally unconstrained; §11.2), the framework’s cascade allocation produces a 9-slot system with $R_{\text{disc}} = 25.0 \text{ AU}$, the same configuration the framework fits in GJ 667, at mass ratio 7.5 instead of 2.1:

- Slot 0 (25.0 AU): Proxima Centauri ($0.122 M_\odot$ current = $40,625 M_\oplus$), the pair’s runt, born on the dam seat and ejected from it during the primordial relaxation to its current marginally-bound $\sim 8,700 \text{ AU}$ orbit.
- Slot 1 (14.6 AU): Alpha Cen B ($0.91 M_\odot$ current = $302,770 M_\oplus$), risen in the same relaxation to 23.5 AU, parked just under the seat Proxima vacated: the catalogue’s third seat capture, at stellar mass.
- Slots 2–3 (8.5–5.0 AU): brown dwarfs ($\sim 18,000$ and $\sim 6,400 M_\oplus$); mutually evicted and dispersed.
- Slots 4–7 (2.9–0.6 AU): gas giants of decreasing mass; all obliterated by multi-perturber scattering among B, Proxima, and the evicted middleweights.
- Slot 8 (0.34 AU): rocky body; obliterated.

The cascade fits Proxima (slot 0) and Cen B (slot 1) exactly. The exchange operates as everywhere else: the heavyweight rose one seat ($14.6 \rightarrow 23.5 \text{ AU}$), the lightweight took the ejection, and B is the Alpha Cen system’s GJ 667 B, Proxima its C, flung farther because the relaxation ran deeper. Proxima’s prograde, plane-aligned, moderately eccentric orbit identifies it as a primordial cascade body that was internally scattered rather than externally captured. Quantitatively, B and Proxima formed

10.4 AU apart with a mutual Hill radius of 13 AU: a packing index of 0.8 (§8.8): their Hill spheres overlapped at birth, making the relaxation compulsory rather than contingent.

The relaxation’s ledger balances once the casualties pay. Lifting Proxima from its dam-seat birth orbit (circular at 25.0 AU) to its present $a \approx 8,700$ AU orbit (Kervella et al. 2017) requires a specific energy $GM_A \cdot 0.020 \text{ AU}^{-1}$; the pass at the fitted periapsis $q = 21$ AU (§8.9) supplies $\sim GM_B/q$, about twice the requirement: a comfortable final kick rather than a fine-tuned one. The larger line item is B’s own climb from slot 1 to the vacated seat, which costs roughly five Proxima-ejections, and the payers are the casualties: the brown-dwarf and giant slots below, sunk and shredded, released the binding energy that lifted the survivor and expelled the runt. Proxima still did not quite unbind, stalling on a marginally-bound orbit, and a partially radial final kick of exactly this kind leaves the moderately eccentric ($e \approx 0.5$), prograde, plane-aligned orbit that is observed.

The same encounter edited Proxima’s own planetary system, and the edits remain legible in its fit residuals. A stellar flyby truncates a protoplanetary disc at $r_t \approx 0.28 q (M_{\text{pert}}/M_*)^{-0.32}$ (Breslau et al. 2014); the fitted encounter ($q = 21$ AU, §8.9) gives $r_t = 3.1$ AU, beheading the disc’s outer rungs (the would-be slot-0 gas giant (2.7 AU, in the stirred zone) lost 43% of its inventory and is unobserved) while every surviving planet closes essentially exactly: c at +0.1% via a merger absorbing slot 2 (the same mechanism as 55 Cancri’s c), b at −0.3% and d at −2.3% with late delivery from the inward debris. The encounter energetics forbid $q \gtrsim 45$ AU and the truncation requires the pass to reach the planet region; the fitted $q = 21$ AU sits inside the window, at twice the early manual estimate ($q \approx 10$ AU, which pinned r_t at c’s orbit: a coarser reading the joint fit supersedes). This is also why Proxima’s system resists unperturbed-cascade calibration: no anchor family closes its per-planet masses, because the allocation curve was overwritten by the encounter. The framework’s one resistant system is the one it independently identifies as an ejected, disc-truncated star.

Slot-determined metallicity, tested on the triple.—Because B and Proxima are slot children of A’s natal disc, their compositions are A’s cloud plus their seats’ solid enrichment, and the framework computes the offsets exactly: +0.11 dex for B at slot 1 and +0.08 dex for Proxima at the dam seat, with the ordering $A < \text{Proxima} < B$. For co-natal systems this arithmetic is immune to galactic chemical evolution; every internal difference is seat arithmetic. The B–A prediction tests against a known anomaly: differential abundance analyses consistently measure B slightly more metal-rich than A, by +0.02 to +0.05 dex (Morel 2018), a difference co-natal formation does not explain and the seat structure requires; the predicted sign and order are observed, with the magnitude high by a factor of two to three. Proxima’s measured metallicity carries M-dwarf systematics of ± 0.1 dex and does not yet discriminate; the sharp test is the ordering, plus a compositional tag the snow line supplies for free: B’s seat lay inside

A’s disc’s ice line (14.6 against 15.7 AU) while Proxima’s lay outside, so B’s solids were dry (computed 0% ice against Proxima’s 18%) and B should be volatile-poor relative to Proxima, the inverse arrangement of Sol and Eos.

6.2. Thought experiment: the Crab Nebula’s primordial cluster

This subsection is a thought experiment, not a claim or proof: it depends on assumed (not measured) primordial parameters and illustrates how the framework extrapolates to massive-star scales.

The Crab Pulsar (PSR B0531+21, $\sim 1.4 M_{\odot}$ neutron star, formed in SN 1054 AD) had a B-class progenitor of approximately $10 M_{\odot}$. Applied to this progenitor with $M_{\star} = 10 M_{\odot}$, spin = 2.58 (typical B-class primordial fast rotation), and $f_{\text{disc}} = 0.25$ (binary-fragmentation regime, identical to Alpha Centauri), the framework allocates an 8-slot cascade with $R_{\text{disc}} = 164$ AU:

- Slot 0 (164 AU): B-class star ($\sim 3.4 M_{\odot}$)
- Slot 1 (96 AU): B-class star ($\sim 3.1 M_{\odot}$)
- Slot 2 (56 AU): A-class star ($\sim 1.5 M_{\odot}$)
- Slot 3 (33 AU): K-class star ($\sim 0.65 M_{\odot}$)
- Slot 4 (19 AU): M-class star ($\sim 0.26 M_{\odot}$)
- Slot 5 (11 AU): M-class star ($\sim 0.10 M_{\odot}$)
- Slot 6 (6.5 AU): brown dwarf ($\sim 0.04 M_{\odot}$)
- Slot 7 (3.8 AU): gas giant ($\sim 15 M_{\oplus}$)

This is a septuple star system: one $10 M_{\odot}$ central host plus six cascade-allocated stellar companions, with one brown dwarf and one gas giant. The total cascade mass is approximately $9 M_{\odot}$, producing a primordial system mass of $\sim 19 M_{\odot}$ in stars alone. The architecture is comparable to compact young clusters such as θ^1 Orionis (Trapezium core, 4–5 confirmed massive stars within ~ 0.02 pc) or AR Cassiopeiae and ν Scorpii (both confirmed septuple systems).

The framework’s mutual-eviction detector flags every cascade member as gravitationally unstable. Within the ~ 30 Myr main-sequence lifetime of the central B-class star, mutual scattering would have dispersed most cascade members. By the time of supernova in 1054 AD, only 1–2 surviving close companions likely remained; SN 1054 then destroyed any remaining envelope and inner companions while leaving the surviving cluster siblings moving outward at ~ 10 – 30 km/s peculiar velocities.

6.2.1. *Stellar nurseries as cascade fragmentation events*

The Crab progenitor’s primordial 7-star configuration is identical in structure to observed compact young clusters: θ^1 Orionis, the Arches cluster, R136 in 30 Doradus, Westerlund 1. Each has at its dynamical core a high-mass primary plus a small number of massive companions in tight orbits. The cascade framework identifies these as fragmentation patterns of massive primordial discs, not stochastic Jeans-instability events. Disc fragmentation is an established binary-formation channel: gravitational instability in massive discs forms stellar and brown-dwarf companions beyond ~ 50 AU (Kratte & Lodato 2016); the cascade adds the prescription of *where* (the slot positions) and *how much* (the slot allocations).

The key parameters distinguishing single-star vs multi-star formation outcomes are: host mass $M_\star$, primordial f_{disc} , and cascade slot mass at slot 0 relative to the stellar formation threshold, the hydrogen-burning minimum mass of $0.08 M_\odot \approx 25,400 M_\oplus$ (Chabrier & Baraffe 2000). For low-mass primaries ($M_\star < 1.4 M_\odot$, G/K/M class), slot 0 mass falls below the stellar threshold unless $f_{\text{disc}} > 0.3$; the cascade produces planets only. For intermediate-mass primaries (A/B class), slot 0 readily crosses the stellar threshold at $f_{\text{disc}} \gtrsim 0.05$; cascade fragmentation yields binary, triple, or higher-multiplicity systems. For massive primaries (O class), the slot 0 stellar threshold is crossed at any reasonable f_{disc} ; cascade fragmentation inevitably produces multi-star clusters.

This is consistent with observation: the stellar multiplicity fraction increases monotonically with primary mass, from $\sim 25\%$ for M-dwarfs to $\sim 80\%$ for B-class to $\sim 100\%$ for O-class (Duchêne & Kraus 2013). The cascade framework predicts this scaling directly from slot 0 mass allocation versus the deuterium-burning and hydrogen-burning thresholds.

6.3. *Compositional bimodality across the parent snow line*

A direct compositional consequence of the cascade framework at stellar scales: cascade bodies forming inside their parent’s snow line have qualitatively different ice budgets from those forming beyond it. The framework’s snow line is a function of parent luminosity (and hence parent mass), scaling roughly as $r_{\text{snow}} \propto L_\star^{1/2}$. For a $M_\star = 2.5 M_\odot$ A-class parent, the snow line falls at ~ 40 AU; for a $M_\star = 10 M_\odot$ B-class parent, the snow line is at ~ 650 AU, well outside the entire cascade.

Cascade bodies forming beyond the parent’s snow line incorporate condensed ice during slot allocation. The cascade rock/ice/pebble allocation formula gives explicit ice fractions for such bodies. Cascade bodies forming inside the snow line acquire essentially no condensed ice; their composition is dominated by rock, pebble, and the H/He envelope (with dissolved water vapor only at ISM gas levels).

This compositional bimodality affects multiple observables:

- **D/H ratio** (the deuterium-to-hydrogen isotope ratio): ice-condensed water at low temperatures is enhanced in deuterium relative to gas-phase water (Tielens

1983). Stars formed beyond the parent’s snow line should have slightly elevated D/H, though the signature is diluted by H/He envelope mass.

- **$^{18}\text{O}/^{16}\text{O}$ ratio:** cold gas-grain chemistry selectively enriches ice in ^{18}O . Outer-slot stars carry this enrichment in their composition.
- **Total volatile budget:** outer-slot stars have $\sim 10\text{--}500 M_{\oplus}$ of cascade-condensed ice contributing to their composition; inner-slot stars have effectively zero cascade-ice budget.
- **Subsurface ice retention in moons/planets:** the sub-cascade of a star inherits its parent’s primordial volatile content. Stars formed dry (inside parent snow line) host planets with reduced primordial water budgets compared to wet (outer-slot) stars.

For the Crab progenitor cluster (slot 1–7 all inside the 650 AU snow line), all cascade-formed companions were dry stars and sub-stellar bodies. By contrast, the Alpha Centauri pair splits across A’s disc ice line (15.7 AU): Proxima, born on the dam seat at 25 AU, condensed ice-enriched solids, while Cen B’s slot-1 seat at 14.6 AU lay just inside the line and built dry.

Observationally, this bimodality should appear in stellar compositional surveys as two compositional families: ice-rich outer-slot stars and ice-poor inner-slot stars. The ice-poor inner-slot population should be statistically associated with cascade companions to high-mass primaries; the ice-rich outer-slot population should be statistically isolated or associated with low-mass primaries (where the snow line falls within the cascade range).

A direct prediction for the Sun follows from the calibrated reconstruction (§7.6): Sol’s seat lay beyond Hyperion’s snow line, so its $\sim 411 M_{\oplus}$ of seat solids were 45% ice, water that contributed a few percent of Sol’s primordial oxygen budget, with the remainder of the heavy inventory arriving as envelope-dissolved disc gas.

7. THE HIERARCHICAL RECURSION: EACH GENERATION BUDGETS THE NEXT

Parents template; they never labor. A daughter system’s two defining inputs are both read off the parent’s structure at the daughter’s seat, and nothing else about the parent matters. The budget is the slot allocation: the daughter’s inventory is the mass the parent’s natal disc assigned to her slot (§2). The boundary is the parent’s local pressure. A daughter forming at radius r_{site} inside her parent’s disc inherits the nebula density

$$\frac{D_d}{D_p} = 0.39 \left(\frac{r_{\text{site}}}{R_{\text{disc}}} \right)^{-2.12}, \quad (33)$$

and the exponent is not free in practice. It reproduces the disc’s own density profile ($\rho \propto \Sigma/H \propto r^{-2.1}$), so the law says simply: record the ambient compression at

your seat. The daughter’s dam follows from her pair through $R_{\text{disc}} = 30.07 M D^{-1/3}$, her cascade runs on the universal machinery, and her slots budget granddaughters. Two numbers enter at any rung and a complete subtree comes out. This is the (M, λ) inheritance pair of Table 1, with λ realised at these scales as the recorded compression D through the jaw-lock (§2). Two daughters of two different parents fix the constants of Eq. 33: Proxima against Alpha Centauri A’s disc, and Jupiter against Sol’s. That the resulting exponent independently matches the disc profile is the law’s first self-consistency check.

7.1. *The measured rung: Jupiter’s satellite system*

The Galilean system supplies the clean test, because the daughter’s pair can be obtained two independent ways. Fitted from the moons alone, Jupiter’s disc requires $D_J = 11.87$. Predicted from Jupiter’s seat in Sol’s disc via Eq. 33, the answer depends on which radius is used. Jupiter’s formation slot ($r_3 = 5.98$ AU) gives $D = 11.97$, agreement to 0.9%. Jupiter’s current orbit (5.20 AU) gives 16.1 and misses by 35%. The satellite disc remembers where Jupiter formed, not where it is. A disc 2,400 times smaller than the orbit it remembers thereby confirms, independently, the ~ 0.8 AU inward recoil that the interior fit attributes to the embryo-swarm scattering event (§5.6).

With the inherited pair imposed, all four Galileans sit in situ in their observed order. Callisto is the dam resident, at slot 0 (0.012586 AU against an observed 0.012585), and is fittingly the one undifferentiated, geologically primitive Galilean. The position residuals of the inner three decline outward in exactly the Laplace-resonance tidal pattern: Io +13.0%, Europa +4.9%, Ganymede −2.3%, Callisto 0.0%. The framework’s geometry plus four gigayears of known tidal drift accounts for the system without any orbital rearrangement.

The Roche parameter completes Jupiter’s interior census. Slots 5 and 6 fall inside the fluid Roche limit (1.36×10^5 km): their allocations could never consolidate and persist as the ring system, whose main edge sits on the line. Slot 4, at 1.6 times the limit, consolidated and was lost the way satellite-formation theory has long required moons to be lost: dragged inward across the line by gas drag: satellite-formation theory has long required generations of moons to be lost inward in exactly this way (Canup & Ward 2006), and the machine itemises one. Its bulk fed Jupiter. Its highest shards survive as the Amalthea group, stranded from the seat down to the line: Thebe on slot 4 to 1.5%, Amalthea between, at a measured density of 0.86 g cm^{-3} a rubble pile sitting inside the Roche limit for its own density and held together by friction, and Metis and Adrastea on the line feeding the main ring. Each fragment trails its own gossamer ring. The debris field of the torn slot-4 body is not inferred; it is photographed.

7.2. *The diagnostic rung: Saturn*

Saturn’s seat (slot 2, 10.25 AU) fixes its inherited pair, and the pair refuses to fit the present mid-sized moons at any input. Mimas through Rhea total ~ 0.7 milli- $M_{\oplus}$, two orders of magnitude below any cascade allocation for a disc of Saturn’s budget. The refusal is the result. Cassini’s gravity and ring-pollution measurements date the rings to $\sim 10^7$ – 10^8 yr (Iess et al. 2019), and dynamical arguments date the mid-sized moons comparably (Čuk et al. 2016). The recursion reads the same story from the masses alone. Saturn’s interior satellite cascade was destroyed recently; the present mid-moons are re-accreted debris; the rings are the destruction ledger, visible because fresh. The Roche parameter sharpens the ring reading into a law. Saturn’s forbidden zone (ice-Roche near 1.37×10^5 km) swallows slot 4 and everything below, and the ring system spans that zone edge to edge: the zone is primordial, a permanent trap in which no slot could ever consolidate, while the ring material is fresh, the latest refill from the recent wreck. This resolves why rings exist at all and why Cassini finds them young in a single statement, and it generalises four for four: every giant’s ring system sits inside its primary’s Roche zone, the standing allocation of forbidden slots, refilled by each destruction epoch. Slot 3, marginal at 1.05 times the line, repeats Jupiter’s slot-4 fate one seat deeper: the body just above the line is the first casualty of inward drag. Titan and Iapetus formed beyond the dam at the exterior boundary (the factory of §8 at sub-cascade scale) and rode out the violence untouched. They are the system’s Kuiper analog, and Titan is the dam-exterior dominant for the same reason the dam’s outer face is everywhere the best-fed seat in a system.

7.3. *The adopted rung: Neptune*

Neptune adds a third reading of the same machinery. Its inherited pair puts its dam near 3.2×10^5 km: the inner regulars, Naiad through Proteus, sit inside as interior cascade products, and Triton sits just beyond the wall. Triton’s retrograde orbit rules out condensation in place and marks it as adopted, and the framework names the source. Neptune is Sol’s slot-0 resident, so Neptune’s trans-dam territory and Sol’s dam-exterior factory floor are the same place. Triton is compositionally Pluto’s twin, and Pluto is a Sol-dam cohort object: Triton came off the same production line and was bound by the planet parked beside it, during the firehose era when the newborn cohort was dense and binary-exchange capture easy. The adoption is also the system’s destruction event, the third instance of the wrecked-interior pattern after HD 69830 and Saturn: the original regulars were destroyed by the arriving heavyweight, the present inner moons are re-accreted debris, Neptune’s faint rings are the remaining ledger, and Nereid rides its $e = 0.75$ orbit as the scattering relic. The trans-dam adoptee outweighs the interior survivors five hundred to one, the same exterior dominance Titan shows, for the same reason: the dam’s far side is the best-fed seat in any system. The test is isotopic: Triton and Pluto should match in D/H

and nitrogen isotopes as production-line siblings, and the re-accreted inner moons should show young surfaces.

7.4. *The rebuilt rung: Uranus*

Uranus completes the four-giant gallery. Its inherited pair ($D = 1.22$ from Sol's slot-1 seat) puts its dam near 1.6×10^5 km, and the census splits cleanly across the wall: the ring system, the small-moon swarm, and Miranda inside; Ariel, Umbriel, Titania, and Oberon outside, prograde and equatorial, native trans-dam products outweighing the interior ~ 125 to one. The architecture is the receipt for a catastrophe this paper has already charged to Uranus. The Solar System fit attributes Uranus's 98° tilt and envelope loss to the outer share of the slot-4–5 embryo swarm (§5.6), and the satellite system records the same event: the major moons orbit in the tilted equatorial plane, so they formed in a disc rebuilt after the knockdown; the rings and inner swarm are the wreck ledger; Miranda, sitting near the interior half-step site, is the re-accreted body, and its patchwork coronae geology has read as shattering and reassembly since Voyager; and the dam seat itself is empty, its original resident a casualty. Across the four giants one variable now organises the trans-dam yield: Jupiter completed its gas capture and drank its exterior pool dry (its only trans-dam natives are the small prograde Himalia-group candidates), while Saturn, Uranus, and Neptune left pools that condensed into Titan, the Uranian big four, and the cohort Neptune adopted Triton from. The emptiest larder belongs to the biggest eater.

7.5. *The kinetic rung: Earth and Luna*

The recursion's smallest rung begins with a prohibition. Run Earth's inherited pair from Sol's slot-6 seat and the gas-era dam lands at 1,700 km, inside the planet: the nebula's ambient pressure at 1.2 AU crushes any terrestrial circumplanetary disc to nothing. Terrestrial planets cannot run the machine during the gas era, which is why none of them carries a primordial satellite system. Earth's pair arrived another way. The Theia impact, the inner share of the same slot-4–5 swarm event this paper charges for Mars, Saturn, and Uranus, delivered it kinetically: spin (the ~ 5 -hour post-impact day is the λ -record, written by collision rather than collapse), a stirred core and dynamo whose field is the Alfvén boundary, a debris disc as the natal disc, and a Davis Dam where the new magnetosphere meets the post-gas solar wind. With the impact-era field and wind, that dam stands at $5\text{--}7 R_\oplus$, and the cascade under it has exactly two seats outside the Roche limit: slot 0 near $6.6 R_\oplus$ and slot 1 at $3.8 R_\oplus$, which is Luna's accretion radius to one percent. Slots 2 and below are Roche-forbidden and feed the open seats. The machine therefore predicts one large moon and at most one small companion, and it predicts the companion's fate. The two-seat configuration is born packed: $\Delta r = 2.73 R_\oplus$ against a mutual Hill radius of $0.84 R_\oplus$ gives $P \approx 3.2$, far below the Lissauer threshold, so relaxation is compulsory by the same criterion that convicts 55 Cancri and HD 69830. Earth's spin-down sets the direction: tidal recession scales as $m/a^{11/2}$, so Luna receded some 600 times faster

than its thirty-times-lighter outer sibling and ran it down, caught it into the co-orbital points, released it as the migrating geometry destabilised, and absorbed it at roughly escape velocity, the one collision speed that pancakes rather than craters. Lunar science met both bodies and the ending from the other direction: Luna at slot 1, and the companion whose slow farside merger explains the highland asymmetry (Jutzi & Asphaug 2011). We name the companion *Endymion*: Selene’s beloved, granted eternal sleep and kept by her forever, which is the object’s literal fate. (The historical “Lilith” attached to claimed second moons belongs to an unrelated and discredited 1898 observational episode.) The Moon’s two faces are the fossil of slot 0’s eviction. Tidal recession carried Luna from 3.8 to $60.3 R_{\oplus}$ at a rate still measured today, the post-formation clock running in plain sight.

The configuration was the machine’s most extreme interior. The vice stood nearly closed, with the corotation boundary at $2.3 R_{\oplus}$ against the dam at 6.6 : compression $C = 0.35$, fifty times Sol’s, a hair short of inversion, which is why the debris disc was compact. Of the cascade’s fifteen slots, eleven lay inside Earth itself and two inside the Roche limit, leaving the two seats. The census maps one-to-one onto what impact simulations have always found and never categorised: the in-planet slots’ allocations are the 50–70% fallback fraction, the Roche-forbidden slots’ allocation is the Roche-interior disc that viscously feeds the moon, and the disc’s much-noted compactness is the kinetically delivered λ -record itself, capped by the magnetopause dam.

The slot, not the Roche limit, is Luna’s address. The conventional account reaches “just outside the Roche limit” and stops; Roche is a floor where accretion becomes possible, not a reason the Moon sits at 3.8 rather than 3.0 or $4.5 R_{\oplus}$. The machine supplies the address: slot 1 of the impact disc’s cascade. Roche explains why not closer; the slot explains why exactly there, and the Moon becomes an ordinary slot occupant, the same kind of object as Venus at Sol’s slot 7 or Callisto at Jupiter’s slot 0. The agreement also hides a discriminator. Mainstream accretion simulations recover $\sim 3.8 R_{\oplus}$ as ~ 1.3 Roche radii, a density-anchored number; the machine derives the same value as $\rho \times R_{\text{dam}}$, a field-and-wind-anchored number. For Earth the two coincide. In any other impact-built satellite system where they separate, Pluto–Charon among the candidates and resolved exomoon systems eventually, the moon’s address decides whether tides or the machine set it. The test is entered in the register (§10).

The control case clinches it. Venus took no late giant impact and is the paper’s in-situ-unchanged anchor, and its three famous absences, no spin, no magnetic field, no moon, are one absence: the kinetic delivery never arrived. Mars runs the same sequence at one-tenth scale and supplies receipts Luna cannot. Its gas-era dam computes to 260 km, subsurface as the prohibition requires; the Borealis impact, the proposed origin of the hemispheric dichotomy (Andrews-Hanna et al. 2008), delivered the pair, and the dynamo’s receipt survives as the crustal magnetization, the field dying ~ 4.0 Gyr ago, after the formation epoch it enabled. The impact-era mag-

netopause dam brackets at $4.5\text{--}6.9 R_{\text{M}}$, and the cascade beneath it holds exactly two seats above the rocky Roche limit ($2.67 R_{\text{M}}$): slot 0 at $6.92 R_{\text{M}}$, where Deimos sits in situ to this day, and slot 1 at $4.04 R_{\text{M}}$, Phobos’s formation seat, from which measured tidal decay carried it to its present 2.77, a gentler start than near-synchronous origins demand of the tidal quality factor. The corotation radius ($6.03 R_{\text{M}}$) falls between the moons, and both obey it: Deimos recedes, Phobos falls. Why Mars kept both children while Earth ate one is a single number: the pair packs at $P = 285$ against Luna–Endymion’s 3.2; same architecture, tiny occupants, relaxation impossible. The mass ledger reads differently from Luna’s, and informatively so: even the engine’s minimum disc fraction allocates $\sim 8 \times 10^{19}$ kg per seat, the canonical Borealis disc scale, while the moons total $\sim 10^{16}$ kg; the deficit is the grinding ledger of the ring–moon cycle (Hesselbrock & Minton 2017), which is running now: Phobos sits inside the fluid Roche limit for its own density (2.77 against $3.13 R_{\text{M}}$), strength-held, ring-bound within ~ 50 Myr. The Roche zone is permanent and its fill recurrent: the ring law, live, one planet over. Across the terrestrial zone the machine is off by default and switched on only by collision, and every switch position matches.

7.6. The recursion run upward: Sol’s progenitor

Nothing restricts the recursion to descent. Sol’s own pair, $(M, D) = (1, 1)$, is itself an inheritance, and inverting Eq. 33 reconstructs the parent: a star of ≈ 2.2 primordial solar masses with $D \approx 0.165$, $f_{\text{disc}} \approx 0.13$, and a dam near 121 AU, whose slot-0 allocation makes one solar mass. We name it *Hyperion*, the Titan father of Helios, completing the mythological lineage the field began when it named the Moon-forming impactor Theia, mother of Selene; the name is shared with Saturn VII, where no confusion can arise. Sol is the dam child, evicted before its own disc assembled (its pristine 30 AU disc requires this), by the same marginal-energy mechanism the catalogue documents for Proxima’s ejection from Alpha Centauri. The reconstruction also answers the meteoritic short-lived radionuclides (^{26}Al , ^{60}Fe , ^{53}Mn), which require a supernova within parsecs of Sol’s formation: Hyperion’s birth cluster supplies the neighbour.

The reconstruction ends in a falsifiable package. Hyperion’s mass and epoch fix its remnant: a carbon–oxygen white dwarf of $0.69 \pm 0.04 M_{\odot}$ with cooling age ~ 3.4 Gyr ($T_{\text{eff}} \approx 6,200\text{--}6,600$ K), on a thin-disc orbit displaced from Sol’s by the tens of km s^{-1} that 4.6 Gyr of independent disc heating imposes on any co-natal pair. A candidate already exists in the catalogue, fourteen light-years away: van Maanen’s star matches the package in mass ($0.68 M_{\odot}$), temperature (6,100–6,200 K), and cooling age (~ 3.2 Gyr), and the literature’s own progenitor reconstruction for it ($2.6 M_{\odot}$, 0.9 Gyr main sequence) agrees with the inheritance-law inversion to within 4% in mass. It is moreover the archetypal metal-polluted white dwarf, currently accreting debris from its old planetary system, which supplies the decisive archival test: its photospheric metal ratios sample the planets condensed from Hyperion’s disc, the

reservoir that made Sol, and should be solar-rock to high precision if the identification holds, and *dry*: the inheritance run places Hyperion’s snow line at 51 AU, beyond every planet-mass seat, so the predicted diet is waterless rock, which is what the star’s hydrogen-free DZ pollution already shows. The candidate’s single status is itself a prediction kept: Hyperion’s asymptotic-giant mass loss shed over half the system’s mass and unbound its companions. It does not travel alone in the package, however. The progenitor’s slot-1 allocation is stellar ($0.4\text{--}1.6\text{ M}_{\odot}$), so a main-sequence sibling exists in the field, released at the AGB unbinding ~ 3.2 Gyr ago (and therefore less phase-mixed than Sol), carrying Sol’s isotopic fingerprint offset by a computed $+0.04$ dex in $[\text{M}/\text{H}]$ and a volatile-poor pattern (her solids were 18% ice against Sol’s 45%, formed inside the snow line), an abundance signature that is effectively unforgeable; if van Maanen’s star is Hyperion, the sibling search anchors on its orbit, not Sol’s. The same composition run reproduces Sol’s bulk metallicity at $Z = 0.0155$ against the protosolar 0.0153, matching the known photosphere-versus-bulk offset unprompted.

A conditional chain follows, and we state it as the speculation it is, with its dependencies explicit. *If* van Maanen’s star is Hyperion, then the inheritance run anchored on its measured progenitor mass places an M-class companion at the next seat: a $0.33\text{ M}_{\odot}$ red dwarf, ejected at birth in the same relaxation that evicted Sol, today essentially unaged (an M dwarf spends 10^{-4} of its mass in 4.6 Gyr of burning). The archival search then finds exactly one star consistent with that companion on every measured axis: GJ 4040, at 46 light-years, with $[\text{Fe}/\text{H}] = +0.04$ against the predicted $+0.043$ (Mann et al. 2015), $\text{H}\alpha$ in absorption and a 30-day rotation period (Díez Alonso et al. 2019), both marks of an old star, kinematics at the median of the co-natal heating distribution (stellar velocity dispersions grow with age as field stars are scattered by clouds and arms; by 4.6 Gyr a co-natal pair typically differs by $\sim 50\text{ km s}^{-1}$; Holmberg et al. 2009) with respect to both Sol and the white dwarf (the expected signature of an at-birth ejection, which predicts no residual coherence with either), and a position nearly equidistant from both (46 and 49 light-years), as independent dispersal requires. GJ 4040 could be Eos *only if* van Maanen’s star is Hyperion; the chain shares the fate of its first link, and both links face the same two tests already stated, the oxygen deficit and the Gaia astrometric architecture. An isotope match at a predicted metallicity offset is a far sharper sieve than isotopes alone. Both predictions are searchable in Gaia-era archives and are entered in the register (§10).

7.7. The root of the recursion

The tree has a measured top. Across the fitted catalogue, systems at essentially one galactocentric radius span 5.6 decades in D . The parent that sets a system’s pair is always the local structure, a disc or a core or a cloud, never the Galaxy at large. At the root, where no parent disc exists, the pair becomes an environmental draw, and this paper measures the draw’s statistics twice: as the stochasticity of f_{disc} across the planetary catalogue (§11), and one tier up as the λ -distribution of galaxy formation. They are the same measurement, made at two scales, under two nomenclatures (§9).

8. THE EXTERIOR RESERVOIR: KUIPER BELT, OORT CLOUD, AND THE BOUNDARY CENSUS

Standard theory supplies *mechanisms* for the trans-Neptunian reservoirs (the Oort cloud as scattered planetesimals lifted by galactic tides (Oort 1950; Duncan et al. 1987), the Kuiper belt as the remnant outer planetesimal disc) but their locations, masses, and mutual structure enter as initial conditions, and each carries a standing anomaly (the belt’s two-orders-of-magnitude “missing mass,” the unexplained Kuiper cliff, the Oort cloud’s $\sim 5\%$ trapping efficiency). In the cascade framework both reservoirs are instead *necessary consequences* of the two outer boundaries every star possesses: the magnetic boundary (the Davis Dam) and the gravitational boundary (the star’s Hill radius in the galactic potential, Eq. 24 applied one level up the hierarchy). Material pools at a boundary while the boundary has working fluid, and fossilizes in place when it dries. The belt and the cloud are the deposit sequences of the two boundaries.

8.1. The dam’s exterior side

The dam traps from both directions. In steady accretion $\dot{M} = 3\pi\nu\Sigma$ is constant in radius, so surface density scales as $\Sigma \propto 1/\alpha$: gas flowing inward through the MRI-active exterior ($\alpha \sim 10^{-2}$) meets the dead-zone interior ($\alpha \sim 10^{-3}$ – 10^{-4}) and queues against the transport-capacity step, a pile-up of order the viscosity contrast (10–100 $\times$) over a width of order the scale height. Solids respond aerodynamically: drag delivers pebbles to the resulting pressure maximum, where the concentrated ring crosses the streaming instability threshold (Youdin & Goodman 2005) and collapses directly into ~ 100 -km bodies: natively producing the like-coloured binaries observed throughout the cold classical population (Nesvorný et al. 2010). The cascade itself cannot operate exterior to the dam (the slot geometry is anchored to the dam from inside), so growth out there stops at Kuiper-belt-object scale: *the exterior pool makes a belt because it cannot make a planet.*

This dissolves the classical mass problem. In-situ coagulation required a 10–30 $M_{\oplus}$ primordial belt eroded by 99.5% to today’s $\sim 0.02 M_{\oplus}$ (Pitjeva & Pitjev 2018); streaming-instability assembly at the dam bump requires only ~ 0.1 – $1 M_{\oplus}$ of exterior solids, comfortably supplied by a percent-level residue of envelope infall that never crossed the dam, and needs no disposal mechanism. Li & Chiang (2025) object

that pressure-bump trapping generically over-feeds the belt, and favour late-nebula streaming instability (at 2–5% residual gas) for the cold classicals. Their objection applies to bumps fed by the full disc’s drifting dust; the dam-edge trap is fed only by the exterior pool, the interior pebble flux having been consumed by the cascade slots upstream, so the trapped mass is capped by the exterior supply. The late-nebula timing they favour is moreover natural here: the dam persists as the gas wanes, and the exterior pool is the last reservoir still feeding it.

Assembly at boundary interfaces is observed at far larger scales: star formation occurs *within* galactic outflows (Maiolino et al. 2017) and is widespread in them (Gallagher et al. 2019), flow–medium frontiers, environments naively expected only to disperse material, instead pooling, compressing, and collapsing gas into bound objects. This is not an analogy but the same physics: a wind–medium pressure boundary in magnetized flow, pooling what arrives at it until the local collapse threshold is crossed, with only the participants scaled by twelve orders of magnitude; the dam-edge factory pooling exterior solids into Kuiper belt objects and the outflow frontier condensing gas into stars are one boundary class (§8.6), and the product mass in each case is set by the local capture physics, not by the mechanism.

8.2. Belt architecture from a stationary Neptune

Neptune is the dam’s interior product (cascade slot 0, gas-fed); the dwarf planets are its exterior products (ice-fed, streaming-instability-born). Both sides deliver to the same orbital radius, with the consequences developed in §4.1: exterior products born while the dam lay within Neptune’s $\sim 10 R_H$ reach ($\lesssim 38$ AU) are milled (scattered, eccentricity-pumped, surviving only in protecting resonances) while products born after the dam moved beyond that reach are never disturbed. The observed hot/cold dichotomy follows from a *moving factory and a stationary planet*: the hot population (Plutinos, scattered disc, high inclinations) is early output born inside the reach; the cold classicals (low- e , low- i , binary-rich) are later output born beyond it. The belt’s inner edge (~ 39 AU) is Neptune’s clearing reach plus the 3:2 resonance at 39.4 AU; the Kuiper cliff coincides with the 2:1 at 47.7 AU, the outermost first-order trap for inward-drifting exterior material (§4.1). The cliff is doubly determined: the dam was transiting the ~ 48 AU station just as the gas-era supply died (§8.3), so the 2:1 trap and the factory’s shutdown radius coincide, the belt ends where production ended, with the resonance as the last shelf stocked.

8.3. The migrating dam: generational deposition

The dam’s position is set by the pressure balance between the wind and the ambient medium, and the stellar spin is that balance’s recorded history: compression causes spin through the jaw-lock ($\Omega = D^{2/3}$, §2), so as the ambient pressure relaxes and the dam marches outward, the braking star writes the march into its rotation (Skumanich 1972), and Eq. 4 reads the station back from the record. With Sol’s primordial rotation period of 1.55 d and today’s 25.4 d, the recorded stations are approximately:

30 AU through the gas era (disc-locked spin); ~ 48 AU by ~ 0.2 Gyr; ~ 80 AU by ~ 1 Gyr; and $30 \cdot (0.061)^{-1/2} = 121.4$ AU today, within 0.2% of the heliopause measured by Voyager 1 at 121.6 AU (Stone et al. 2013). We identify the present-day dam with the heliopause: the surface where the Sun’s magnetized outflow meets the interstellar medium is the same boundary, holding nothing because it has no working fluid. The boundary today is a balance of two near-vacuums (wind ram pressure at 121 AU is $\sim 10^{-4}$ of its 1-AU value (~ 0.1 pPa), matched against a comparably tenuous interstellar medium) roughly twelve orders of magnitude below the disc-era working-fluid density at the same boundary. A pressure-balance surface is exactly as strong as the medium it holds, so the dam’s clientele has shrunk in proportion: from pebble pile-ups in the gas era to the deflection of sub- $0.1 \mu\text{m}$ charged grains (the only particles whose charge-to-mass ratio still couples them to so feeble a structure) today. (The identification carries caveats: the agreement uses the primordial-mass convention and equatorial rotation period, the heliopause breathes with the solar cycle, and a derivation showing the wind–interstellar-medium standoff scales as $\Omega^{-1/2}$ remains outstanding. The formation-era law itself has a candidate derivation: $R \propto \Omega^{-1/2}$ is the angular-momentum-conservation locus of a pressure-confined system (§2), with environmental compression as cause and spin as record. The identification then requires the same causality to persist after the disc: the boundary sits where ambient pressure dictates, while spin, braked by the wind that mediates that very balance, continues to track the boundary’s secular expansion. Spin is the trend-line record; the instantaneous environment is weather around it, with dense-cloud crossings as the ring-depositing excursions and the spin-read station as the baseline the boundary relaxes back to.)

The present-day dam, though feeble, visibly carries its pile-up: the *hydrogen wall* (the density enhancement of interstellar neutrals stacked against the heliopause, detected in Lyman- α absorption toward nearby stars and crossed by the Voyagers (Wood 2004)) together with the measured filtration of small charged interstellar grains at the boundary. The factory’s shelf is stocked and observed; what it lacks today is throughput. The same absorption technique detects *astrospheres*, hydrogen walls, around other nearby stars (Wood 2004), establishing the dam as a universal, directly observed stellar structure rather than a solar peculiarity.

Cementing requires both working fluid and supply, so deposition is episodic: the gas era builds the main belt, and later cohorts form only when the Sun traverses dense interstellar material, which both re-immerses the dam in a dissipative medium and supplies fresh solids. The radial map of the belt is then a *time log* of the Sun’s environmental history read against the spin-down clock. The newest observations show exactly such banded structure: dust fluxes above all model predictions approaching 60 AU (Doner et al. 2024), and a populated ring at 70–90 AU separated from the known belt by a near-empty gap at 55–70 AU (Fraser et al. 2024). In this reading the 70–90 AU ring is a wet episode at the ~ 1 Gyr dam station, and the gap is a quiet

transit. The present-day silence is overdetermined: the Sun currently resides inside the Local Bubble, a supernova-blown cavity of hot, $\sim 100\times$ underdense gas (Frisch et al. 2011), which it entered ~ 5 Myr ago (Zucker et al. 2022), the factory today lacks fluid *and* feedstock. Kuiper-belt-object formation is a property of local interstellar conditions, switched on only when a star’s environment supplies both. (Interstellar processing of Oort-cloud comet *surfaces* (erosion by interstellar grains, cosmic-ray irradiation) is established (Stern 2003); the framework extends the star–interstellar-medium interaction from surface weathering to episodic accretion and formation.)

The spin-down clock makes deposition radius a birthdate: a cohort at radius R was cemented when the dam stood there, at the epoch when $\Omega(t) = (30 \text{ AU}/R)^2$ in primordial units. This yields a sharp prediction. The youngest possible Kuiper belt objects date to the last wet episode before Local Bubble entry, and because the dam has moved less than 1% over the past ~ 100 Myr, *any recently-formed cohort must lie within a few AU of the present heliopause radius* ($\sim 105\text{--}121$ AU): dynamically cold (born too recently to be stirred) and spectrally fresh (surfaces younger than the space-weathering timescale, in contrast to every known trans-Neptunian population). The clock carries one stated exception, and the exception is itself dated. A sufficiently dense interstellar cloud drives the dam far *inward* for the duration of the transit (the ram pressure of the Sun’s ~ 25 km/s motion through the cloud dominating the balance), and a production flash at the compressed stations deposits a thin cohort of small bodies at radii the baseline clock forbids. Independent evidence indicates exactly one recent such event: the live ^{60}Fe and ^{244}Pu layers in deep-sea archives, which Opher et al. (2024) attribute to a cold-cloud passage $\sim 2\text{--}3$ Myr ago that compressed the heliosphere drastically. The framework therefore predicts a sparse population of *young, volatile-fresh, kilometre-class bodies at $\sim 40\text{--}150$ AU carrying live ^{60}Fe* (half-life 2.6 Myr : still $\sim 50\%$ extant): self-labelling, lab-verifiable, and tied to an independently dated event. The two-sided test is then clean: a fresh-surfaced cohort whose location and radionuclide stamp match a dated transit *confirms* the excursion clause, while a fresh-surfaced population at small radii with *no* corresponding transit record falsifies the clock outright. (Distant comets active beyond conventional sublimation radii, usually attributed to supervolatile ices, are noted as candidate transit-cohort members without being claimed as such.)

The firehose budget.—The primordial-mass convention adopted in §2 fixes the factory’s total throughput with no new assumptions: Sol shed $\sim 0.14 M_\odot$ across the T Tauri epoch, and every gram of it crossed the Davis Dam. At disc metallicity that is $\sim 660 M_\oplus$ of solids delivered *through* the dam wall over ~ 10 Myr: of order $65 M_\oplus$ per Myr at peak. The belt’s assembly then requires only that a fractional residue stall and stick at the pressure boundary: a capture efficiency of a few $\times 10^{-3}$ supplies the $\sim 0.1\text{--}1 M_\oplus$ exterior pool the streaming-instability reading needs (§8.1), while even the classical massive-primordial-belt requirement ($10\text{--}30 M_\oplus$) would demand only $\sim 4\%$. The budget is overdetermined in the framework’s favour: the factory was supply-rich

by two to three orders of magnitude, and the belt’s final mass is set by the capture physics at the wall, not by the availability of material. The same convention names the epochs: the *firehose* era (gas-age, $\sim 65 M_{\oplus}/\text{Myr}$ of solids at the wall) built the main belt; everything since is the *trickle*.

Production-rate budget.—The belt’s census constrains the shape of the production history. The main belt’s $\sim 10^5$ objects larger than 100 km overwhelmingly date to the gas era, when the factory ran at rates of order 10^4 objects per Myr; the post-gas factory is supply-limited and episodic, delivering clutches of order 10^2 – 10^4 objects per wet episode (streaming instability collapses clumps in bursts, not singly) across a few tens of episodes over solar history. Averaged, that is roughly one new Kuiper belt object per Myr-scale interval: but the delivery must be bursty rather than continuous: a steady drip would smear deposits along the dam’s entire outward track, and the observed emptiness of the 55–70 AU gap (Fraser et al. 2024) is itself direct evidence of the episodic duty cycle.

Compositional variance structure.—Cohort deposition predicts a distinctive variance pattern: *low scatter within a cohort, high scatter between cohorts*. The gas-era population formed from a single well-mixed reservoir over ~ 10 Myr and should be internally consistent in composition, isotopic ratios, and age: as the cold classicals conspicuously are, constituting the most homogeneous (uniformly very red) colour group in the trans-Neptunian region (Tegler & Romanishin 2000). The same survey’s colour *bimodality*, a very-red class and a neutral class, maps onto the framework’s two source populations: dam-born cohort material versus milled inner-disc debris mixed into the dynamically hot population. Each later ring instead samples whatever interstellar cloud powered its wet episode, so successive rings should differ from one another (in ice inventory, dust mineralogy, isotopic vintage, and surface age) while remaining uniform internally. The vintage prediction has a thermometer: deuterium enrichment in ice is set by low-temperature grain-surface chemistry (Tielens 1983), and successive dam stations sit at progressively colder radii, so the cohort ladder predicts a *monotonic D/H gradient* with formation radius across the rings: a measurable ordering, not merely a qualitative variance claim. The dynamically hot population, in this picture a mixture of milled dam-born bodies and inner-disc-born swarm debris scattered outward (§4.1, §5.6), should by contrast be compositionally bimodal at any radius. Radius, in short, predicts composition class: discrete rings, discrete vintages.

The age inversion.—This reading inverts the classical understanding of the trans-Neptunian region. Classically, Kuiper belt objects and comets are the Solar System’s *oldest* accessible material: pristine relics of the solar nebula, the rationale for cometary sample missions. In this framework the belt and cloud are by definition its *youngest* components: the only parts of the Solar System still (episodically) under construction, with assembly ages spanning from the gas era down to the last wet episode before Local Bubble entry. The planets finished forming within ~ 10 Myr; the belt never

stopped. Later cohorts are built from interstellar matter that postdates the Sun: including condensates from stars that died *after* the Solar System formed: *post-solar grains*, the inverse of the presolar grains treasured in cometary samples. This yields the framework’s most direct sample-return test: a radiometric or isotopic assembly age younger than 4.56 Gyr in any trans-Neptunian solid is inexplicable classically and expected here, with the youngest material residing in the outermost rings.

Existing cometary samples do not yet test this (every comet sampled or closely visited is a Jupiter-family comet, drained from the *old* hot population) but they already test the mixture prediction: comet 81P/Wild 2’s classically-surprising content of inner-solar-system igneous material, including high-temperature crystalline silicates and a CAI-like refractory inclusion, alongside an anomalously low presolar-grain abundance (Brownlee et al. 2006) is the expected positive signature of inner-disc-born swarm debris rather than of pristine outer-nebula ice: the refractory content is diagnostic of inner-disc origin, not merely consistent with it. We note a paradigm blind spot: anomalous-isotope stardust in samples is classified as *presolar* by assumption, so post-solar grains would be misfiled; cometary grain inventories have not been screened with the age direction treated as open. Encounters with dynamically-new Oort-cloud comets (e.g., ESA’s *Comet Interceptor*) are the first opportunity to sample populations where young material is predicted.

8.4. *The gravitational boundary: the Oort cloud*

The second boundary is gravitational: the Sun’s Hill radius in the galactic potential, $R_{H,\odot} \approx 1.3 \text{ pc} \approx 2.7 \times 10^5 \text{ AU}$, with long-term-stable orbits confined (as for any satellite system) to roughly half that: matching the outer Oort cloud’s observed extent of $\sim 10^5 \text{ AU}$. This boundary is also epoch-dependent: in the birth cluster the tidal radius was only $\sim 2 \times 10^4 \text{ AU}$, and the inner Oort cloud and Sednoid region plausibly fossilize that cluster-era station (Brasser et al. 2006), while exchange within the cluster commons stocked the cloud with sibling debris: indeed Levison et al. (2010) conclude the Sun *captured* a large fraction of its Oort cloud from other cluster members, which predicts a compositional split the framework inherits gladly: long-period comets (partly sibling material) should be isotopically heterogeneous relative to the native, dam-pooled Kuiper cohorts. The Kuiper belt and Oort cloud are thus the same phenomenon at the two boundaries: drag pools in the disc plane at the magnetic boundary (a flat belt); gravity pools in three dimensions at the tidal boundary, isotropized by galactic tides (a sphere). The flat-versus-spherical dichotomy, explained in standard theory by two unrelated histories, follows here from the holding mechanism alone.

8.5. *The desert, and exterior slots*

The two-boundary census makes a hard negative prediction: neither boundary ever occupied $\sim 150\text{--}2,000 \text{ AU}$ (the dam will reach only $\sim 160\text{--}180 \text{ AU}$ by the end of the main sequence; the cluster-era tidal radius never compressed below a few thousand

Table 8. The Solar System’s boundary census. Every boundary the system has had either persists today or left a fossil population at its old station.

Boundary	Physics	Primordial	Today
Alfvén Dam	magnetic (stellar)	0.20 AU	~ 0.04 AU ($\sim 9 R_{\odot}$)
Davis Dam	pressure (interstellar)	30.07 AU	~ 121 AU (heliopause)
Resonance brackets	drag fossil	39.4–47.7 AU	frozen
Cluster tidal radius	gravitational	$\sim 2 \times 10^4$ AU	inner Oort fossil
Galactic Hill radius	gravitational	,	2.7×10^5 AU

AU), so *no in-situ population should exist in that range*: only high-eccentricity lifted bodies passing through, consistent with the known Sednoids (Sedna, $e = 0.84$).

8.6. The factory class

The full Minor Planet Center census (6,034 trans-Neptunian objects) shows the dam’s exterior carries no standing geometry of its own: structure outside the dam is *temporal*. The interior, a held dissipative disc, writes space-structure: slots. The exterior has one organizing element only, the boundary itself: a single moving shoreline whose deposits record *when* it stood at each radius (§8.3), post-processed by the nearest giant’s resonances. And the shoreline is not a solar peculiarity. It is the local instance of a factory class operating at every wind–medium interface, and every component of that class is already in the literature under its own name: at the galactic scale first, because that is the only scale at which one can photograph someone else’s boundary.

At stellar scale the boundary and its stocked shelf are directly observed: the hydrogen wall at our heliopause and the equivalent *astrospheres* of nearby stars (Wood 2004). At galactic scale the boundary is named: the *galactopause*, where Shull & Moss (2020) show that during strong outflows thermal instability precipitates kpc-scale clouds at the wind–IGM interface (30–65 kpc), which then lose pressure support and rain back inward on ~ 200 Myr timescales. Condensation at the pressure boundary, followed by inward delivery of the product: the same two factory steps as the dam’s, eleven orders of magnitude up, with the product mass set in each case by the local capture physics, pebble pile-ups cement km-to-100-km ice bodies at a stellar dam; thermally unstable coronal gas condenses cloud-mass objects at a galactic one, which fall inward as the disc’s fuel. Star formation at flow boundaries is likewise established componentwise: sequential triggering at shock fronts (Elmegreen & Lada 1977), star formation *within* outflows predicted as positive feedback and then observed (Maiolino et al. 2017; Gallagher et al. 2019), young clusters manufactured at ram-pressure interfaces, and globular clusters condensing from the protogalactic boundary medium (Fall & Rees 1985). The factory manufactures belts here and fuel there; the grammar is identical.

What has not previously been assembled is the identification: heliopause, astrospheres, and galactopause as *one class of production boundary whose output is dated cohorts*. Each boundary’s deposit sequence is a time log, radius (or infall phase) maps to epoch through the boundary’s own motion, so every boundary population is an archive of its system’s environmental history, and the archives are mutually calibrating: the Solar System supplies the resolved, sample-accessible instance (cohort vintages, the D/H ladder, the transit-cohort radionuclide stamp above), while external systems supply the statistics. At the galactic tier the same clock logic predicts age–radius *stratigraphy* in nuclear stellar populations rather than spatial lattices; Section 9 develops this against the Galactic Center’s measured, strongly episodic star-formation history.

The deep desert prediction of the two-boundary census stands as stated: a Rubin-class survey finding a substantial low-eccentricity in-situ population at 150–2,000 AU would falsify the census; sparse late-cohort material near the dam’s historical stations ($\lesssim 121$ AU) is expected; the deep desert is not populated in situ.

8.7. Universality

Every magnetized accreting star has a dam, and every star has a galactic Hill radius; the framework therefore predicts that exo-Kuiper belts are universal, located at $R \propto M_\star \Omega_{\text{prim}}^{-1/2}$, with comet clouds filling each star’s Hill sphere to its stability edge. Cold debris belts are indeed detected around $\gtrsim 20\%$ of FGK stars (sensitivity-limited, so the true fraction is higher), as rings at tens-of-AU radii, and their measured radii already show a systematic increase with host luminosity (Matrà et al. 2018). Regressing the resolved-belt sample against the dam scaling is an archival test requiring no new observations: if it holds, every resolved debris ring is a measurement of its host’s primordial rotation.

β Pictoris already provides a sharper, system-level instance. Fitting the cascade to its two imaged giants alone yields $R_{\text{disc}} = 28.4$ AU (spin 2.64), with b and c as inward migrants from slots 1 and 0 and the inner cascade (slots 2–7, $\sim 700 M_\oplus$ of predicted inventory) shredded by their passage: supplying the archetypal debris disc that in-situ readings leave unexplained. The fit is then checked out-of-sample by the system’s resolved planetesimal belts: mid-infrared imaging locates dust-replenishing rings at ~ 6 , 16, and 30 AU (Okamoto et al. 2004), coincident with cascade slots 3, 1, and 0 (5.6, 16.5, 28.4 AU) to within 3–6%, three independently observed belts on three slots of a ladder fitted only to the planet positions, reading naturally as collisional debris at the giants’ formation slots and at the destroyed slot-3 body.

HR 8799 is a second instance of the same mechanic: its four wide super-Jupiters leave every inner slot obliterated by simultaneous scattering in the fit, and the system indeed hosts a warm inner dust belt while no inner planets are known. Two of two wide multi-super-Jupiter systems in the catalogue thus carry shredded inner cascades, generalizing to a population prediction with a mass threshold: outer giants near

Jupiter’s mass *preserve* their inner systems, consistent with the observed positive correlation between cold Jupiters and inner super-Earths (Zhu & Wu 2018), while multiple wide super-Jupiters *destroy* them, leaving inner-planet deserts filled with multi-ring debris at slot radii. The known excess of directly-imaged giant planets around debris-disc hosts (Meshkat et al. 2017) reads here not as a shared massive-disc origin but as cause and wreckage; the discriminators are the dust’s location and structure (interior to the giants, slot-laddered) and the absence of surviving inner planets specifically above the super-Jupiter threshold.

8.8. *Born-hot chains: the packing index at formation*

The cascade fixes where bodies form and how massive they are; Hill stability (Eq. 24) then determines whether the configuration can persist. For each adjacent pair of slots define the *packing index* $P = \Delta r / R_{H,\text{mut}}$, the slot separation in units of the pair’s mutual Hill radius, evaluated at the formation positions, with destroyed slots carried at their predicted masses (stability acts on the at-birth configuration, not the surviving one). Numerical integrations of multi-planet chains require $P \gtrsim 7$ for long-term stability (Smith & Lissauer 2009); a system fitted below that limit was born dynamically hot and *had* to relax.

The fitted catalogue splits cleanly at the boundary. Quiet systems sit far above it (Sol’s terrestrials at $P = 41\text{--}56$, TRAPPIST-1 at 8–10 on its half-step lattice (§2), the Galilean satellites at 15–18) and none carries scattering, migration, or remnant signatures. Sol’s one marginal pair is Jupiter–Saturn at $P = 7.2$, grazing the limit: precisely the pair implicated in the framework’s single dynamical event (§5.6). Every system fitted below the boundary shows independent evidence of violence, and the outcomes sort into recognizable fates. *Resonance lock*: adjacent slots sit at a period ratio $\rho^{-3/2} = 2.24$, just wide of 2:1, so modest convergent drift captures a packed chain into the neighbouring integer commensurability rather than chaos. We emphasize the distinction: 2.24 is the *birth* ratio, and the observed integer locks are the *relaxed* states, GJ 876 ($P = 3.8$) settled into the canonical Laplace 4:2:1 chain (Rivera et al. 2010), and HR 8799’s four giants ($P = 3.7$) into a proposed 8:4:2:1 protection (Goździewski & Migaszewski 2014). The cascade explains why capture lands specifically in first-order 2:1 chains, the resonances convergent disc migration also selects (Pierens & Raymond 2011): the ladder is born adjacent to them. The Luger lattice of the inverted regime (§2) supplies the second observed family by the same logic, $\rho^{-3/4} = 1.497$ sits on 3:2 essentially at birth, which is why compact chains like TRAPPIST-1’s barely needed to move to lock (Luger et al. 2017), and why its observed pairs sit within $\sim 1\%$ of exact commensurability while the giant chains had to converge 11% to reach theirs. An occupancy prediction follows directly: HD 7924 (the catalogue’s deepest inversion ($C = 241$), whose known planets occupy lattice sites while the ladder’s entire lower half (six rungs, $\sim 10 M_\oplus$ of allocation) formed inside the Hayashi photosphere and was consumed) carries vacant full rungs at $2.5\text{--}6.8 M_\oplus$

(0.02–0.09 AU; radial-velocity semi-amplitudes 1–2.5 ms^{-1}): either detectable with modest precision gains, or absorbed in mergers like its slot 2, which the fit already credits to planet b. The HR 8799 lock model carries a bonus that doubles as a slot test: Goździewski & Migaszewski (2014) predict a fifth, innermost planet (HR 8799f) at ~ 7.5 – 9.7 AU to extend the chain, and the cascade fitted to the four known giants independently places its next rung, slot 4, at 7.9 AU, inside their predicted window. Two unrelated formalisms, a resonance-dynamics fit and a geometric ladder, point at the same empty orbit. *Outside-in shredding*: β Pictoris ($P = 3.5$) and HR 8799, whose packed outer giants destroyed their inner cascades (above). *Inside-out wrecking*: HD 134987 ($P = 3.3$ at the fitted parameterization) places c in situ at the dam edge (5.8 AU) and b as an inward migrant from slot 2 ($2.0 \rightarrow 0.81$ AU), with slot 1’s occupant, the chain’s dominant member, now absent despite decades of radial-velocity coverage: ejected during the exchange. The survivor is a lone warm Jupiter over a swept interior, parked where magnetospheric-cavity simulations show inward-migrating giants halt and accumulate (Romanova & Lovelace 2006): the configuration whose population-level signature is the well-documented loneliness of close-in giants (Steffen et al. 2012), whose differential form (warm Jupiters and hot Neptunes retain companions; hot Jupiters specifically do not) maps the launch-energy threshold of the wrecking fate. 55 Cancri repeats the pattern in compressed form ($P = 5.1$; eleven slots inside 1.3 AU): b swept inward from slot 2 ($0.46 \rightarrow 0.11$ AU), capturing c near the observed 3:1 commensurability on the way past, while d was driven outward ($1.3 \rightarrow 5.5$ AU) and the innermost survivor e was ground to a bare core. Notably, wreckage *visibility* is age-gated: β Pic (~ 20 Myr) and HR 8799 (~ 30 – 40 Myr) still display their debris, while ancient wrecks like 55 Cancri (~ 10 Gyr) retain only dynamical scars, consistent with the absence of any confirmed debris disc there after gigayears of collisional grinding and drag removal. *Marginal rearrangement*: Mu Arae’s three giants pack at $P = 6.0$ – 6.1 , just under the limit, and show precisely a marginal outcome, the lightest giant displaced inward 2.1 AU, and a $10.5 M_{\oplus}$ body parked deep inside the Alfvén void, reading as the 5–10% Martian-band survivor of a 105 – $210 M_{\oplus}$ casualty of the relaxation.

Across every packed chain the *lightest* member absorbs the relaxation, and the rule continues unbroken to stellar masses. Alpha Centauri B and Proxima formed at adjacent slots 10.4 AU apart with a mutual Hill radius of 13 AU: $P = 0.8$, overlapping Hill spheres at birth (the Hill formalism is only indicative at stellar mass ratios, but an index below unity needs no precision). Coexistence as born was impossible; the $7.5\times$ lighter Proxima took the kick, and its wide, bound, plane-aligned orbit (Kervella et al. 2017) is the relaxation product (§6.1). GJ 667 repeats the pattern at $P = 0.7$ with the kick rule executed exactly: B formed at the dam edge (slot 0, 21.6 AU) and settled inward 9 AU to the observed 12.6 AU A–B separation, while C, the runt of the triple, was flung from slot 1 to its observed ~ 230 AU. The same fit predicts a $0.46 M_{\odot}$ K-class sibling at slot 2 and a $17.5 M_J$ brown dwarf at slot 3, mutually evicted

and ejected outright, GJ 667 was born a quadruple plus a brown dwarf and fired two runaways. Packed-chain ejecta (stars, brown dwarfs, giants like HD 134987’s, and planetesimal debris) are thus a standing output of cascade formation, stocking the interstellar inventory that the exterior factories of §8 later sample.

8.9. Gravitational stripping: encounter-edited cascades

Two catalogue systems carry violence their own packing indices forbid. HD 20794’s chain packs at $P = 14\text{--}22$ with no giant anywhere, yet its ledger shows a destroyed dam-edge body, a half-stripped outer member, and inner planets carrying mass surpluses an order of magnitude beyond Sol-style late delivery; Proxima Centauri’s planets defeat every unperturbed anchor family outright. Born-quiet systems do not wreck themselves: the editing was external, and for Proxima the framework independently names the editor (§6.1). We therefore promote encounter editing from forensic narrative to a fitted mechanism. A flagged system gains one parameter (the encounter periapsis q , with the perturber mass M_{pert} supplied ($0.91 M_{\odot}$ for Proxima’s named assailant; $0.5 M_{\odot}$ for an anonymous cluster-era flyby)) from which the *stripping factor* follows as the tidal truncation radius $r_t = 0.28 q (M_{\text{pert}}/M_{\star})^{-0.32}$ (Breslau et al. 2014). The transform edits the baseline allocation in three zones: slots exterior to r_t are *destroyed* (predicted planet mass zero; inventory freed), the band $0.5 r_t\text{--}r_t$ is *stirred* (graduated loss, up to 60% at r_t), and the interior is *enriched*, 15% of the freed inventory rains inward and is swept up outermost-first. q is then bisected against the mean per-planet $|\log(\text{pred}/\text{obs})|$: precisely the quantity that consensus mass-closure cannot see, and the quantity encounter editing exists to close.

Both systems close. HD 20794 fits at $q = 15.8$ AU ($r_t = 5.2$ AU): the dam-edge slot is destroyed, g sits in the stirred zone at -23% , f’s raw $+310\%$ surplus collapses to -10% (the late-delivery band) once the inward debris is credited, and b ($2.7 M_{\oplus}$ parked beneath the Alfvén dam) is the *encounter fragment*: 8% of the $34 M_{\oplus}$ freed inventory, the inward share of the same 5%/95% asymmetric split that delivers Theia and the Mars survivor in Sol (§5.6), here operating on an encounter’s spoils rather than a swarm’s. Proxima fits at $q = 21$ AU, inside the energetically allowed window (§6.1): d, b, and c close at -2.3% , -0.3% , and $+0.1\%$ respectively, the cleanest per-planet closure in the catalogue, from the system that was previously unfittable at any compression. The two fits agree on the transport physics: the inward-capture fractions implied independently are 10–15% (Proxima) and $\sim 18\%$ (HD 20794), bracketing the calibrated 15%. One parameter, physically derived from stellar mass and periapsis, converts the framework’s two resistant systems into its two best-documented crime scenes.

9. THE GALACTIC TIER

A galaxy forms the way a planetary system does, one rung up. This section states the generation sequence, identifies its fossils in the present-day Galaxy, and gives the tests that are running now. The law driving the sequence is the same inheritance pair measured at 0.9% three rungs down (§7.1); at this tier the pair already has names.

9.1. *The generation sequence*

A seed condenses in a collapsing protogalactic cloud. Its outflow against the ambient medium sets a pressure boundary, the galactopause, which is the Davis Dam of the seed’s natal disc (§9.4), and the core is then constructed exactly the way Sol’s planets were. While the collapse holds the dam deep, the natal disc runs the machine of §2 unchanged: cascade seats at the fixed ratio ρ beneath the dam, each allocating by the same rule, with only the products rescaled by local capture physics. At galactic seat allocations of 10^5 – $10^6 M_\odot$ no single body can hold together as a star, so the seats fill with direct-collapse objects and massive clusters where Sol’s seats filled with planets. The interior seats are consumed by the seed precisely as protostellar consumption eats a compact system’s inner slots: slot children dragged across the inner boundary are part of how the seed grows. The surviving deposit of this deep-dam epoch is the bulge and nuclear star cluster, and black-hole mass tracks classical-bulge mass ever after because seed and bulge are the same epoch’s co-products: one cascade, two ledger columns. As ambient pressure falls, the boundary sweeps outward and lays the disc down inside-out at constant mass per radius, with successive cohorts deposited at golden-angle offsets along the slot ladder. Once the deposited mass outweighs the seed, dynamics pass from Keplerian to self-gravitating, the sweep ends at the galactopause, and the templating epoch survives only in what it built. A generative integration of this sequence, the inheritance pair driven forward 13.6 Gyr under a pressure-driven factory law with one tuned normalisation, reproduces the Milky Way’s gross statistics eight for eight: stellar mass by construction, and star count, white-dwarf census, star-formation rate, half-mass epoch, old-mass fraction, unconverted-baryon fraction, bulge fraction, and the inside-out age gradient emergent, with the bulge entering through the compression dial below.

9.2. *The inheritance pair under its Λ CDM names*

Λ CDM galaxy formation parameterises a galaxy by halo mass and spin (Peebles 1969), and the correspondence to the framework’s pair is identity of content, not analogy. The budget is the same number outright: $B = f_b M_{\text{halo}}$. The spin draw is the same physical variable read at opposite ends of the system. λ measures the system-wide angular momentum acquired by tidal torques; the framework’s D records the central compression the collapse achieved; a shallower angular-momentum barrier means deeper collapse and higher central compression, so the two are anticorrelated readings of one underlying draw, collapse depth. Λ CDM discovered the pair top-down, by simulation, at the one scale where the parent is the environment itself, and

within Λ CDM the pair is phenomenology. The recursion supplies the missing why: budget and boundary are what any parent hands any child, and the same two-number law fits Saturn’s moons, which no semi-analytic model can attempt with $(M_{\text{halo}}, \lambda)$. The theories tile rather than compete. This paper imports Λ CDM’s cosmological scaffold wholesale, and Λ CDM’s galaxy recipe acquires lower rungs and a derivation. Each becomes evidence for the other: the stochasticity of f_{disc} across the planetary catalogue (§11) and the λ -distribution of haloes are the same measurement, made at two scales, under two nomenclatures.

9.3. Fossils of the templating epoch

The seed’s present dynamical irrelevance does not erase what it templated. Sgr A* rests at the dynamical centre to less than 1 km s^{-1} (Reid & Brunthaler 2004), the founder still at the barycentre of everything it organised, and the templating epoch survives in four fossils.

The deposition profile.—The allocation rule lays down constant mass per radius, which is the Mestel profile, the one surface density whose rotation curve is flat (Mestel 1963). The shape of the Galaxy’s baryonic deposition is thereby a direct print of the allocation law. The dark component that dominates the full rotation curve is Λ CDM’s, imported with the rest of the scaffold (§9.5).

The spiral pitch.—Cohorts deposited at the most-irrational (golden) angular offset along the slot ladder trace a logarithmic spiral of pitch $\arctan[\ln(1/\rho)/2.400] = 12.65^\circ$, with a doubled-step harmonic at 24.2° and a forced ratio $\tan \varphi_2 / \tan \varphi_1 = 2$. The measured pitch distribution across 4,378 disk galaxies is bimodal at $12.0^\circ \pm 3.4$ and $23^\circ \pm 4.3$ (Yu & Ho 2020), and $\tan 23^\circ / \tan 12^\circ = 1.997$. One constant, imported unchanged from the planetary cascade, places both peaks; the mainstream alternatives require two unrelated mechanisms to do the same. The theory further predicts a third harmonic at 33.9° , suppressed density near 18° , and pitch independent of shear rate within each peak (§10).

The founder coupling.—The machine makes the organised inventory linear in the seed ($\sigma_{\text{AAF}} \propto M_{\text{seed}}$), so a templated structure stays mass-correlated with its founder long after the founder is dynamically negligible. This is the black-hole–bulge relation and its famous asymmetry: near-linear ($M_{\text{BH}} \propto M_{\text{bulge}}^{1.17 \pm 0.08}$), tight for classical bulges, absent for pseudobulges and discs (Kormendy & Ho 2013). The asymmetry is structural. Classical bulges are regime-1 deposits, co-eval with seed feeding; pseudobulges are later secular rearrangement, causally disconnected from the templating epoch, and therefore uncorrelated. The zero-zero corner exists: M33, bulgeless, carries a nuclear black-hole limit of order $10^3 M_\odot$ (Gebhardt et al. 2001). The relation’s ~ 0.3 dex scatter is the environmental draw again, the same stochasticity the planetary catalogue measures in f_{disc} .

The bulge family.—The generative sequence has one structural dial, the early-compression history $\gamma = d \ln R_{\text{dam}} / d \ln(1 + z)$, and the dial is the spin draw under another name. Setting $\gamma = 2.1$, which corresponds to protogalactic overdensity growth modest beside virialisation, reproduces the Milky Way’s 25% bulge and ends central deposition at $z \approx 1.7$, matching both the measured ages of bulge stars and the observed “blue nugget” compaction epoch, the compact star-forming phase through which massive galaxies are seen to pass at $z \approx 2\text{--}4$ before growing inside-out (Zolotov et al. 2015). Bulge-to-total ratio is therefore a one-parameter family in collapse depth, and the family’s corollary has been in the data for decades: the morphology–density relation (Dressler 1980). Ambient density is the dial, so morphology tracks environment because the dial is the address.

9.4. *The boundary factory at galactic scale*

The galactopause *is* the Davis Dam, one rung up. The same wind–ambient pressure boundary that stands at 121 AU around Sol, holding the hydrogen wall, stands at tens of kiloparsecs around the Galaxy, holding the precipitating halo clouds: one object, two addresses, and the galactic instance was named and its boundary condensation modelled before this paper asked for it (Shull & Moss 2020). The rest of the factory class (§8.6) is likewise already published piecewise, star formation at flow boundaries established from triggered shells (Elmegreen & Lada 1977) to outflow interiors (Maiolino et al. 2017; Gallagher et al. 2019). What the framework adds is the identification across scales, and the clock. Boundary deposits are dated cohorts, so nuclear populations carry age–radius stratigraphy rather than spatial structure, and the Galactic Centre already reads this way. The nuclear stellar disc’s star-formation history is strongly episodic: $\sim 80\%$ of its mass is older than 8 Gyr, gigayears of near-quiescence follow, and a discrete burst arrives ~ 1 Gyr ago (Nogueras-Lara et al. 2020). That is the duty cycle in fossil form. The current boundary components are individually observed: the quiescent inner rim at the circumnuclear disc, front jitter in X-ray light echoes, and the escaped polar front of the last major episode in the Fermi bubbles (Su et al. 2010). The “paradox of youth,” stars at 0.04–0.5 pc with ages of 4–6 Myr (Paumard et al. 2006) where standard star formation is forbidden, coincides in epoch with that last episode. It is the factory’s most recent firing, deposited exactly where and when the front last stood. The stratigraphic tests are running: the clockwise disc’s stellar age will equal the bubble-launch epoch as both datings tighten; the distinct disc features will show measurably different ages, where a single fragmentation event predicts one; and the ~ 1 Gyr nuclear burst will prove radially annular. Externally the programme is statistical. Astrospheres establish the boundary as universal (Wood 2004); cavity-edge star-forming rings in face-on hosts test cohort age against independently dated activity fossils, with the bar-resonance alternative discriminated by unbarred hosts and by radial rather than azimuthal age

structure; and ram-pressure fronts manufacturing young clusters supply the running existence proof.

9.5. *Scope: the dissipation gate*

Everything above concerns the dissipative fraction. Dark matter cannot cool, therefore cannot pool, therefore cannot enter any dam, slot, or shoreline. It is the permanently unorganisable remainder: present everywhere, including the Solar System, where some five Jupiter masses of it stream unbound through the Sun’s Oort volume at any instant, and organised nowhere. The theory makes no claim on its dynamics and imports Λ CDM’s cosmological core intact. The claim of this paper is the baryonic layer: the visible fraction, organised by one inheritance law from the Galaxy down to the moons.

10. PREDICTIONS AND OBSERVATIONAL TESTS

The model makes several falsifiable predictions:

Asymmetric scattering split (5%/95%).—N-body simulations of Jupiter’s gravitational scattering of the slot 4-5 embryo swarm should reproduce the framework’s predicted asymmetric inner/outer mass partition: roughly 5% of the swarm mass delivered inward (yielding the Mars survivor and the Earth-impacting body) and roughly 95% delivered outward (producing cumulative impacts on Saturn and Uranus).

Cascade-slot signatures in exoplanet populations.—Surveys of compact exoplanet systems should show the geometric ratio $\rho \approx 0.584$ in their orbital spacings, with deviations attributable to scattering events or migration through the chaotic zones of dominant perturbers. The ~ 5 –10% surviving-mass-fraction predicted for partial swarm dispersal (§5.6) is testable against the observed mass-deficit-at-position patterns in multi-planet systems.

Packed-chain ejecta.—Chains fitted below the Lissauer limit shed their lightest members (§8.8). The framework therefore predicts a standing galactic population of runaway stars, brown dwarfs, and free-floating giants as routine cascade output, and that wide, eccentric, plane-aligned low-mass companions (Proxima, GJ 667 C) are relaxation products of their own systems rather than captures: testable through composition, since a scattered sibling shares its system’s abundances while a capture need not.

Multi-star formation thresholds.—The framework predicts multi-star formation outcomes wherever the slot-0 mass allocation exceeds the deuterium-burning threshold ($\sim 25,400 M_{\oplus} = 0.08 M_{\odot}$). The calibrated f_{disc} catalog (§11.2) provides direct test cases. Future ALMA-era disc-mass surveys combined with companion-search programs around the catalogued systems can validate the slot-0-threshold criterion.

Resonant Kuiper belt census.—The co-orbital survivorship channel (§4.1) predicts resonance-island populations loaded by survival statistics from a source at Neptune’s own orbit: strong 3:2, weak 2:1, strong 5:2, with deep-libration, Kozai-rich microstates after 4.5 Gyr of erosion. Migration sweeping loads the 2:1 most heavily. The observed census (3:2 strong, 2:1 weak, 5:2 anomalously strong; Gladman et al. 2012) already leans toward survivorship; a dedicated N -body comparison (Pluto-mass bodies injected co-orbital with a stationary Neptune, integrated over the age of the Solar System, counted against the observed resonant census) is a decisive and currently feasible test. Two further discriminators separate the channels: migration sweeping ties a captured body’s eccentricity to the migration distance (Malhotra 1995), predicting a tight e -resonance relation, whereas scatter-then-survive predicts a broad survival distribution with *no* such correlation; and drag-driven trapping at a stationary Neptune (Weidenschilling & Davis 1985) sets population ratios by drift speed alone, with no migration timescale anywhere in the prediction.

Exterior-reservoir tests.—Three follow from §8: (i) *exo-belt scaling*, resolved debris-belt radii should regress onto $R \propto M_{\star} \Omega_{\text{prim}}^{-1/2}$ (archival ALMA data). The existing empirical relation is $R \propto L_{\star}^{0.19}$ (Matrà et al. 2018) (roughly $M_{\star}^{0.8}$ on the main sequence, shallower than the dam law’s linear mass term) with scatter that dominates the trend; the framework’s distinctive content is precisely that the scatter is not noise but the $\Omega_{\text{prim}}^{-1/2}$ term, so the test is whether independently-inferred spin orders the residuals; (ii) *the desert* (no in-situ (low-eccentricity) population at 150–2,000 AU under the two-boundary census, versus low- e clustering at the exterior-slot radii (151, 259, 444 AU) under the outward ladder) a clean three-way verdict for Rubin-class surveys; (iii) *cohort stratification* (members of the 70–90 AU ring (Fraser et al. 2024) should be compositionally and surface-age distinct from the cold classicals if they are a later, dam-station generation; (iv) *the youngest cohort*) any objects formed in the last wet episode before Local Bubble entry must sit at ~ 105 –121 AU, just inside the modern heliopause, dynamically cold and spectrally fresh; the largest such members fall within the reach of Rubin-class coadded photometry, and the New Horizons dust counter should see sustained or rising fluxes if its trajectory approaches such a ring.

11. DISCUSSION AND FUTURE WORK

11.1. *Novelty statement: which claims are new*

The framework presented here synthesises established disc-physics and impact-dynamics results into a coherent allocation rule for planetary mass and orbital position. Many individual ingredients are well-established in the literature; others are proposed here for the first time. We list each major claim with its provenance, distinguishing rigorously-derived from synthesised from hypothesised content. We hedge speculative claims and emphasise that several core proposals are testable predictions rather than proven results.

Existing-literature foundations (well-established).—The viscous-disc surface density profile $\Sigma \propto r^{-1}$ (Lynden-Bell & Pringle 1974; Pringle 1981), the magnetospheric Alfvén truncation (Alfvén 1942; Königl 1991), ALMA-era observations of disc sizes (Andrews 2020; Long et al. 2018; Manara et al. 2023), the canonical pebble-accretion framework (Lambrechts & Johansen 2012; Bitsch et al. 2015), the Tanigawa-Ikoma runaway gas accretion ($M_{\text{gas}} \propto M_{\text{core}}^2$; Tanigawa & Ikoma 2007), and the Chatterjee-Tan inside-out planet formation reservoir (Chatterjee & Tan 2014) are adopted directly from the published literature.

Novel framework claims.—The following claims, to our knowledge, do not appear in the existing literature and are proposed here:

1. *The geometric cascade ratio* $\rho = 1 - \frac{\sqrt{\ln 2}}{2} \approx 0.584$ (§2). Derived from the half-amplitude-at-45° projection of the Gaussian accretion-zone HWHM; a pure geometric constant with no empirical fit parameter. The cascade slot positions $r_n = R_{\text{disc}} \cdot \rho^n$ fit Sol’s planets and the 22 further catalogued systems. Geometric orbital ladders have an empirical pedigree (the Titius–Bode tradition and its exoplanet generalizations, which fit per-system ladders $a_n = a_0 e^{bn}$ with system-specific spacing (Bovaird & Lineweaver 2013)) but the cascade ratio differs in kind: a single, derived, universal constant anchored to a physical boundary (R_{disc}), not a per-system fit.
2. *Jupiter and Saturn essentially in-situ at slots 3 and 2* (§5); the slot 4-5 embryo-swarm dispersal as the Solar System’s single dynamical event beyond cascade formation. Contradicts the Grand Tack and Nice Model migration sequences; testable via N-body verification of the asymmetric $\sim 5\%/95\%$ inner/outer scattering split.
3. *The Borealis impactor and Theia share a common Jupiter-driven slot 4-5 swarm origin.* Hypothesis; the literature establishes both impacts independently but does not link them dynamically. Testable via Borealis-ejecta cosmochemistry vs. Theia-snow-line-origin signatures.
4. *Alfvén Dam and Davis Dam* as paired terms for the inner (magnetospheric) and outer (pressure-confinement) disc truncations (§2). Terminological proposal honouring the discoverers of each boundary’s physics (Alfvén for the magnetospheric truncation concept, Davis for the pressure-confined cavity (Davis 1955)) rather than claiming coinage: the underlying physics is established; what is new is the dam-and-reservoir framing, the identification of the outer boundary as the planet-forming edge, and its present-day identification with the heliopause.
5. *The Annulus Allocation Factor* σ (Eq. 7) as a named intensive quantity. The mathematical quantity ($dM/dr = 2\pi r \Sigma$) follows directly from Lynden-Bell & Pringle (1974) but was not previously named.

6. *Mercury as a slot-8/slot-9 merger remnant* with iron enrichment from the high-energy encounter retention function (§3.1). The slot-9 Vulcan body is a framework prediction; the merger mechanism resolves Mercury’s mantle anomaly without invoking post-formation impact stripping, hit-and-run sequences, or extreme nebular processing. The retention formula $\eta = \max(0.3, 1 - 0.37\Delta v/v_{\text{esc}})$ is calibrated against both Mercury and Tau Ceti e.
7. *Venus and Neptune as the ISU primordial-plane anchors of the Solar System* (§4.1). Identification via four convergent diagnostics (most-circular orbits, one-way-integral eccentricity, inverse-capture diagnostics, precise mass-cascade match). The component facts are long catalogued, Venus and Neptune hold the two most circular planetary orbits, the invariable plane is the standard dynamical reference (Souami & Souchay 2012), secular reconstructions find Venus primordially cold (Brasser et al. 2013), and Jacobson et al. (2017) infer that Venus never suffered a late energetic giant impact, but Neptune’s in-situ status requires a non-migratory reading of the resonant Kuiper belt, small bodies drag-trapped at a stationary Neptune’s exterior resonances (Weidenschilling & Davis 1985), dwarf planets surviving in protecting resonances after condensing at the dam alongside Neptune : rather than swept up by a migrating planet (Malhotra 1995). The joint use of the pair as a primordial-plane reference for reading impact history is, to our knowledge, new (§4.1, “Relation to prior art”).
8. *Scale-invariant cascade ratio* across stellar, sub-stellar, and multi-star scales (§6). Demonstrated for Jupiter’s Galilean moon system (all four moons in situ on the inherited pair, closing at 0.9%; §7.1) and Alpha Centauri’s triple-star architecture. The Crab Nebula progenitor reconstruction (§6.2) is presented as an illustrative thought experiment only; Sol’s own progenitor is treated as a calibrated reconstruction with named falsifiers (§7.6).
9. *The trans-Neptunian reservoirs as the Solar System’s youngest components* (§8). The Kuiper belt and Oort cloud are derived as necessary deposits at the system’s two outer boundaries, assembled episodically across the entire age of the Solar System rather than only at its formation: inverting the pristine-relic paradigm and predicting cohort-stratified ages, compositions, and *post-solar* grain content in the outer rings.
10. *Stochastic f_{disc} across stellar class, and its identifiability structure* (§11.2). Across the eight cleanly machine-calibrated systems, $R^2 \leq 0.03$ vs $M_\star$ with f_{disc} spanning a factor ~ 21 over three decades of host mass; and the model itself delimits where f_{disc} is measurable at all: all-giant, no-ISU architectures leave it formally unconstrained. Consistent with the cross-cluster ALMA disc-mass compilation of Manara et al. (2023).

Mathematically derivable vs. empirically calibrated.—The local-capacity allocation function $M(r)$ is mathematically defined by the closed-form expressions of §2. Given the three system inputs $(M_\star, \Omega, f_{\text{disc}})$ and the universal constants of Table 2, the planet masses are uniquely determined, this is a derived (not fitted) result. The Jupiter-driven slot 4-5 swarm dispersal hypothesis and the inner/outer asymmetric scattering split are physically motivated but *not* mathematically derivable from the allocation function alone. We present these as hypotheses consistent with the observational record, testable by the specific predictions of §10, rather than as theorems flowing from the underlying disc-physics framework.

What our framework cannot prove.—We explicitly do not claim to have proven: (i) that the asymmetric scattering split has the precise 5%/95% ratio predicted (requires N-body verification with disc-gas torques); (ii) that the Borealis impactor was a slot 4-5 outer-swarm sibling of Theia (requires ejecta cosmochemistry to test); (iii) that the Vulcan slot-9 body existed as a distinct embryo or remains as buried iron mass within Mercury (requires high-precision gravity-gradient or interior-structure measurements distinguishing merger-remnant from primordially-mantle-stripped chondritic body); Each of these is a candidate explanation consistent with observation, not a proof. Additionally, the Crab-progenitor reconstruction (§6.2) is *not* a claim but an illustrative thought experiment of how the framework behaves under a speculative scenario; it should not be cited as a testable hypothesis, prediction, or evidence for the framework. Sol’s progenitor reconstruction (§7.6) is, by contrast, calibrated, and carries its falsifiers with it.

11.2. Empirical finding: disc-to-star mass ratio is stochastic across stellar classes

A non-trivial empirical finding emerges from machine calibration of the catalogue. Calibration is restricted to systems whose architectures actually constrain the disc mass: f_{disc} has leverage only through sub-threshold rocky planets (whose masses scale directly with the disc budget), mantle-stripped remnants, or ISU anchors. In a system composed entirely of gas-eligible giants with no ISU anchor, the per-planet formation-time fit absorbs any disc mass, and f_{disc} is *formally unconstrained*: a structural identifiability property of the model, not a fitting deficiency. Planets observed deep inside the Alfvén Dam ($r < 0.5 R_A$; e.g. hot Jupiters) occupy no slot at all: they are delivered-inward bodies whose formation slot is indeterminate, and they are likewise excluded from calibration targets. Fifteen of the 22 catalogued systems fall in the unconstrained class (viable parameterizations exist for all of them, but their disc masses are consistency conditions, not measurements). Table 9 reports the eight systems with clean, identifiable machine calibrations.

Across the eight cleanly-calibrated systems the disc-to-star mass ratio is *stochastic* with respect to stellar class: f_{disc} spans a factor of ~ 16 (0.010–0.16); systems of nearly identical host mass differ by a factor of sixteen (HD 20794 at 0.010 versus 55 Cancri at 0.16, both late-G hosts); and a regression of $\log f_{\text{disc}}$ on $\log M_\star$ yields

Table 9. Machine-calibrated primordial spin and f_{disc} for the ten systems whose architectures constrain the disc mass (sub-threshold rocky planets, stripped remnants, or ISU anchors among the calibration targets), ordered by host mass. Systems composed entirely of gas-eligible giants are formally unconstrained in f_{disc} (see text) and excluded from statistics. Stellar masses are in primordial-solar units ($1 \equiv 1.14$ present-day $M_{\odot}$). [†]Fitted with one encounter parameter (gravitational stripping, §8.9): HD 20794’s innermost planet is the encounter fragment, the inward share of the 5%/95% split, parked beneath the Alfvén dam, and Proxima closes only under its Cen B encounter ($q = 21$ AU; §6.1); both encounter fits close to $< 0.01\%$.

System	$M_{\star} (M_{\odot})$	Ω	f_{disc}
Jupiter (sub-cascade)	0.00095	5.18	0.031
TRAPPIST-1 (M8V)	0.089	1861	0.048
Proxima Centauri (M5.5V) [†]	0.122	1.90	0.025
Tau Ceti (G8V)	0.78	308	0.022
HD 20794 (G8V) [†]	0.81	10.3	0.010
HD 219134 (K3V)	0.81	487	0.045
55 Cancri (G8V)	0.95	454	0.157
Mu Arae (G3V)	1.10	165	0.090
Kepler-90 (G0V)	1.13	1127	0.032
Sol (G2V)	1.00	0.995	0.010

$R^2 < 0.001$ for the nine stellar hosts ($R^2 < 0.001$ including the circumplanetary sub-cascade, which extends the host-mass baseline by three decades). The Jupiter sub-cascade calibrates to $f_{\text{disc}} = 0.032$: a planetary-scale disc drawing its ratio from within the stellar-host range, consistent with scale-invariant boundary physics.

Multi-star formation outcomes require the slot-0 allocation to cross the deuterium- or hydrogen-burning thresholds, which demands a high- f_{disc} disc; the catalogued multi-star systems (Alpha Centauri, GJ 667) admit viable parameterizations at $f_{\text{disc}} \sim 0.2$ – 0.3 , but their all-stellar architectures place them in the unconstrained class, so those disc masses are consistency conditions rather than measurements.

This empirical finding is independently supported by ALMA-era protoplanetary disc surveys, which measure $M_{\text{disc}}/M_{\star}$ directly via millimetre continuum emission. Surveys of the Lupus (Ansdell et al. 2016), Chamaeleon (Pascucci et al. 2016), Ophiuchus (Williams et al. 2019), and ODISEA (Cieza et al. 2019) star-forming regions consistently find ratios spanning 0.1%–30% at fixed stellar mass. The mean trend is real and in fact super-linear (Pascucci et al. (2016) report $M_{\text{disc}} \propto M_{\star}^{1.7-1.9}$, steepening with cluster age) but the stochasticity claim rests on the *scatter*, not the slope: 0.5–0.9 dex at fixed stellar mass (Ansdell et al. 2016), dwarfing any deterministic prediction for an individual system (Manara et al. 2023). (A super-linear mean is itself congenial to the cloud-density conjecture of §11: denser cores yield both more massive stars and proportionally heavier, faster-spinning discs.)

The combined model + observational finding is: f_{disc} is set primarily by stochastic local conditions during cloud collapse and disc assembly (turbulent fragmentation,

infall asymmetries, magnetic flux trapping), *not by the deterministic properties of the resulting star*. This is consistent with the conclusion of [Manara et al. \(2023\)](#) in the Protostars and Planets VII (PPVII) review that disc-mass diversity reflects formation-environment variability more than stellar-mass scaling.

Theoretical expectation.—The stochasticity is not only observed but predicted. The angular momentum that seeds a disc is set by the turbulent velocity field of its natal core: effectively a random draw with a large intrinsic spread at fixed core mass ([Burkert & Bodenheimer 2000](#)). Zoom-in simulations following collapse from giant-molecular-cloud scales find the accretion process heterogeneous “in time, in space, and among protostars of otherwise similar mass” ([Kuffmeier et al. 2017](#)), and disc population synthesis from radiation-hydrodynamical cluster simulations produces an enormous diversity of disc masses and sizes from the chaotic dynamics of star formation itself ([Bate 2018](#)).

The cascade’s structural explanation.—The framework supplies its own reason for the same decoupling. When star formation occurs within discs (§6), a star of a given mass can emerge from progenitor discs of almost any mass: the same stellar mass can be allocated at slot 0 of a modest disc, at an inner slot of a massive one, or assembled by independent cloud collapse. The new star’s own disc budget is in turn inherited from its local share of the *progenitor’s* disc, not set by the star’s final mass. Stellar mass and f_{disc} are therefore decoupled by construction: f_{disc} records a system’s formation lineage rather than any property of the star it orbits. The catalogue’s $R^2 < 0.1$ is the planet-architecture-level imprint of that decoupling.

A single-variable hypothesis (unproven).—The two per-system inputs may not be independent of each other. Spin, on the pressure-confinement reading of §2, is the recorded *compression* of the birth site; f_{disc} , on the structural reading above, is the recorded *mass supply* of the birth site. Both would then be functions of a single underlying variable, the density of the formation environment, and planet formation would reduce to a two-parameter theory: stellar mass and birth-site density, plus universal constants. We emphasize that this is a hypothesis with no proof at present. Across the ten cleanly machine-calibrated systems the rank correlation between fitted spin and f_{disc} is $\rho \approx 0.60$ at $n = 10$: positive, as a common density driver would produce, but not significant ($p \approx 0.07$). We report it with its full volatility history: development revisions of the catalogue moved the value between $\rho \approx 0.6$ and 0.81 as calibrations improved and the constrained sample grew from eight to ten, rank statistics this volatile should be treated as motivation for a larger sample, not as an established correlation. Density also enters the two quantities through different moments (confining pressure versus local surface density). Establishing or refuting the hypothesis requires a larger identifiable sample, birth-environment tagging of calibrated systems, and rotation–density statistics in young clusters. The other piece of internal evidence already pointing this way is the quasi-universality of the specific-

angular-momentum constant (30 AU at solar values) across the cleanly-calibrated set.

Implications.—This stochasticity has direct consequences for the framework’s generalisability:

- The cascade slot mass budget at each radius scales with f_{disc} ; systems with high f_{disc} have more massive cascade bodies at every slot, and the slot-0 mass allocation crosses the stellar formation threshold at lower f_{disc} for massive primaries (driving the multi-star outcome statistics).
- Two systems with identical $M_{\star}$ but different f_{disc} produce qualitatively different cascade architectures (planet-only vs. multi-star), even at fixed stellar class.
- Population-level statistical inferences about planetary architectures cannot rely on stellar class alone; the system-specific f_{disc} is an independent, observable, dominant input.

11.3. Model limitations

The cascade framework is closed-form and deterministic given its three system inputs ($M_{\star}$, Ω , f_{disc}) plus per-planet (r , t_{form}) via the AAF cascade rule (Eq. 22).

The grain-opacity constant is calibrated only on Sol.—All catalogue fits use $g = 0.82$ (§2); present mass-position data cannot constrain it per system. Resolved snow-line imaging or compositional measurements of individual systems would lift the degeneracy and test whether g is genuinely universal: or merely uniform across the solar neighbourhood that the calibration sample occupies.

Observational anchors carry tracer and calibration systematics.—Two soft spots are inherited from the disc-observation literature. ALMA “disc sizes” are tracer-dependent, CO radii are chemically depleted relative to the true gas extent (Trapman et al. 2019), so identifying a measured radius with the physical dam edge carries a tracer correction. And absolute gas-disc masses remain systematically uncertain (the disc mass-budget problem; Manara et al. 2023), so the catalogue’s *absolute* f_{disc} values are reliable only to that systematic floor; the stochasticity conclusion and all cross-system comparisons are differential and robust to it.

Linear rocky allocation neglects pebble-drift physics.—The linear σr rocky allocation neglects detailed pebble-drift dynamics in favor of an aggregate empirical scaling. A more physics-rich treatment would couple Birnstiel-style two-population dust evolution (Birnstiel et al. 2012) to the local-capacity framework, particularly for the pebble accumulation at outer cascade slots that contributes to gas-giant core masses.

Empirical A_0 accretion coefficient.—The Tanigawa-Ikoma H/He accretion coefficient ($A_0 = 4.45 M_{\oplus}^{-1}$, $k = 0.691 \text{ Myr}^{-1}$) appears as an empirically-tuned constant. Its order of magnitude is reproduced from MMSN disc parameters at Jupiter’s slot-3 position (Eq. 21), but a fully first-principles derivation from disc viscous evolution and gas-accretion timescales remains outstanding.

δM post-formation modifications.—The δM post-formation modification term is a structural feature of the model that absorbs collisional and dynamical events following primary cascade assembly. We emphasize that most δM assignments have independent physical interpretation: Jupiter-induced dispersal of the slot 4-5 embryo swarm is constrained by dynamical and isotopic evidence; impact-retention energetics for Mercury follow from the geometric cascade and $\Delta v/v_{\text{esc}}$ encounter physics (§3.1) without ad hoc mass-stripping mechanisms. The remaining flexibility is bounded by the cascade’s slot-position predictions and the four-line ISU criterion (§4.1); arbitrary δM assignments would violate the inclination/eccentricity baseline established by the Venus-Neptune primordial plane.

Sub-cascade scale-invariance is only partially validated.—The cascade ratio $\rho = 0.5837$ generalises successfully from stellar to sub-stellar scales (Jupiter’s circumplanetary cascade, §6). However, the framework’s R_{disc} formula uses stellar primordial-spin scaling (Eq. 4); for circumplanetary discs around gas giants, Hill-sphere truncation by the host star may impose additional boundary conditions not captured by the spin-only formula. Saturn’s moon system, where Titan dominates the mass budget by $75\times$ over the next-largest body, was identified as a non-cascade “runaway-growth + fragments” regime where the geometric cascade does not apply.

f_{disc} is an input only at the recursion’s root.—For a system whose parent is identified, the framework now *computes* the pair rather than fitting it: the budget is the parent’s seat allocation and the boundary follows from the inheritance law (§7), as measured at 0.9% for Jupiter’s disc and run upward to calibrate Hyperion’s. The limitation that remains is confined to root systems, those with no parent disc, where f_{disc} is a genuine environmental draw taken as input: its distribution across stellar classes is documented but not predicted, and deriving it from cloud-collapse physics (turbulent fragmentation, infall asymmetries, magnetic flux trapping) would require coupling the cascade framework to larger-scale star-formation theory.

11.4. Extensions

Future work should: (i) implement N-body simulations of Jupiter-driven slot 4-5 swarm dispersal to verify the predicted $\sim 5\%/95\%$ inner/outer scattering asymmetry and the cumulative impact-energy delivery to Saturn and Uranus; (ii) extend the embryo-swarm dispersal criterion to other multi-planet systems to predict scattering signatures across the exoplanet catalog; (iii) derive the A_0 and k envelope-accretion constants from disc physics rather than from their present calibration against Jupiter

and Neptune; the slot’s physical embodiment as a zonal-flow pressure bump (§2) names the route, since A_0 is physically the bump-fed accretion efficiency, accessible to the magnetized-disc simulations already cited, with Pollack–Bitsch–Mordasini-class disc-evolution models constraining the k depletion clock; (iv) test the prediction that gas-disc Type-I and Type-II migration are minor or absent dynamical effects, with observed planet positions explained by cascade-default formation plus scattering events alone; and (v) test the single-variable hypothesis that Ω and f_{disc} both derive from birth-site density (§11.2), beginning with per-system spin refits and rotation–density statistics in young clusters.

11.5. *Relation to existing frameworks: the LBP track that was disregarded*

The present model adopts viscous-disc surface-density scaling from the gas-dynamics tradition (Lynden-Bell & Pringle 1974; Shakura & Sunyaev 1973; Pringle 1981) and applies it to planet mass allocation, a perspective rarely combined in the planet-formation literature, which has predominantly used Hayashi (1981)’s MMSN $\Sigma \propto r^{-1.5}$ scaling. Modern ALMA observations (Andrews 2020; Andrews et al. 2010; Long et al. 2018; Trapman et al. 2019; Manara et al. 2023) robustly favour the LBP $\Sigma \propto r^{-1}$ profile, with measured power-law indices $\gamma \approx 0.9\text{--}1.1$. The MMSN framework, which underweights the disc’s outer mass reservoir, can be considered superseded.

The Annulus Allocation Factor was implicit in the 1970s viscous-disc literature. —A historical observation deserves explicit acknowledgement. For the LBP $\nu \propto r$ self-similar solution, the mass per unit radius $dM/dR = 2\pi R\Sigma(R) = \text{constant}$ emerges as a *trivial mathematical consequence*, the quantity we name the Annulus Allocation Factor sits one line below their derivation. Lynden-Bell & Pringle (1974) proved it; Pringle (1981) consolidated it; Shakura & Sunyaev (1973) established the α -disc viscosity prescription that makes $\nu \propto r$ natural. The framework was set aside not because the math was wrong, but because:

1. Hayashi (1981) adopted a steeper $\Sigma \propto r^{-1.5}$ for solid material in the MMSN, partly because it produced the correct rough orders of magnitude for Solar System planetary masses under contemporary observational constraints;
2. the planet-formation community subsequently built on the MMSN framework for four decades, treating viscous-disc gas-dynamics work as relevant to stars (accretion onto compact objects, AGN, X-ray binaries) rather than to planet allocation;
3. the closed-form constant dM/dR that emerges in the LBP framework was never crystallised into a named planet-formation quantity, because no one was looking for it within the MMSN-shaped planet-formation landscape.

Safronov (1972), Cameron (1978), and Weidenschilling (1977) all worked the relevant territory in the 1970s, Safronov with his “feeding zone” concept (planet mass = integrated annular mass), and Weidenschilling with the inverse reconstruction of disc

surface density from observed planetary masses (the empirical demonstration that planet masses read out a smooth annular mass law, of which our forward allocation is the mirror) but none crystallised the intensive radial constant. The load-bearing departure is the profile: Weidenschilling’s reconstruction assumed the MMSN’s $r^{-3/2}$, where the LBP r^{-1} adopted here is what renders the per-annulus allocation a radial constant at all.

11.6. *Component provenance: who found the pieces*

The same acknowledgement extends beyond the allocation constant. Nearly every component mechanism of this framework was first identified (as a phenomenon, if not in its final mechanistic role) by earlier workers, each within the context where their methods encountered it. We record that provenance explicitly, because the framework’s claim is the *assembly* and its universality, not the parts:

- *The allocation constant*: Lynden-Bell & Pringle (1974) proved it, Pringle (1981) consolidated it, Shakura & Sunyaev (1973) supplied the viscosity that makes it natural; Weidenschilling (1977) demonstrated empirically that planet masses invert to a smooth annular mass law, and Safronov (1972) supplied the feeding-zone calculus and the $\Delta v/v_{\text{esc}}$ accrete-or-erode parameter.
- *The inner dam*: magnetospheric truncation by Ghosh & Lamb (1979) (the magnetic-pressure concept tracing to Alfvén 1942); its coupling to stellar spin by disc-locking, Königl (1991) and Bouvier et al. (2007); the trapping of solids against it as a planet-forming reservoir by Chatterjee & Tan (2014); the parking of inward-migrating giants at it by Romanova & Lovelace (2006).
- *The outer dam*: the pressure-confined cavity (the boundary itself, in its present-day form) by Davis (1955), inflated by Parker’s wind (Parker 1958); the ionization-bounded viscosity transition (the dam’s disc-era masonry) by Gammie (1996), sharpened by the demonstration that the disc bulk is magnetically dead (Bai & Stone 2013); solid trapping at pressure maxima by Pinilla et al. (2012); the transport-set character of observed disc edges by Najita & Bergin (2018).
- *The packing physics*: the Hill-spacing stability threshold by Smith & Lissauer (2009); chain relaxation at gas dispersal by Liu, Ormel & Lin (2017); the observation that real chains assemble at marginal spacing by Hansen & Murray (2013).
- *Rung inference itself*: geometric spacing regressions in exoplanet systems by Bovaird & Lineweaver (2013); the prediction of an unseen member from the occupied rungs of a chain by Goździewski & Migaszewski (2014) for HR 8799f, and its confirmed precedent, TRAPPIST-1h, predicted from the inner chain’s resonance relations and found at the predicted period (Luger et al. 2017); the half-step occupancy state of §2 is named the *Luger lattice* accordingly.

- *The exterior factory's parts*: collapse at pressure maxima by Youdin & Goodman (2005); its binary signature by Nesvorný et al. (2010); drag-trapping into resonances at a stationary planet by Weidenschilling & Davis (1985); the cluster-commons exchange by Levison et al. (2010); boundary mass assembly at galactic scale by Maiolino et al. (2017).
- *The standing barrier in cosmochemistry*: the sustained NC/CC reservoir separation by Kruijer et al. (2017).

Each of these works contains a piece of the present mechanism, observed where its discoverers' instruments and questions happened to point. What this paper adds is the identification: that they are one structure (two dams, a geometric ladder between them, and the dynamics of its occupants) operating identically from circumplanetary discs to star-forming clouds.

12. CONCLUSION

Λ CDM galaxy formation describes a galaxy with two numbers and does not say why two suffice. This paper has proposed the answer and calibrated it where the hierarchy can be resolved: (M, λ) is the inheritance pair of dissipative baryonic organisation, the mass budget and the collapse-depth record that every parent system hands every child, processed by one universal machine whose formation seats, the natal slots, stand at the fixed geometric ratio $\rho = 1 - \sqrt{\ln 2}/2 \approx 0.584$ between two pressure boundaries.

At planetary scale the machine is calibrated, not conjectured. In primordial solar units Sol sits at exactly $(1, 1, 1)$ with its dam at Neptune; ten clean calibrations and 23 fitted systems follow across spectral classes A through M and both dynamical regimes of the disc. The Solar System retains its radical parsimony: Jupiter and Saturn essentially in situ, one dynamical event, the slot 4–5 swarm dispersal, paying out through its asymmetric split the Moon-forming impactor, the Mars survivor, and the envelope deficits and tilts of Saturn and Uranus, with the Mercury–Vulcan merger as the only other invocation. Migration throughout the catalogue is gravitational, angular-momentum exchange with heavier bodies, with the gas disc trapping rather than torquing: migrators park on seats and rims, and the parking position dates the move.

The hierarchy closes in both directions. Jupiter's satellite disc prefers Jupiter's formation seat over its present orbit by a factor of 35 and closes at 0.9% on the inherited pair; Saturn's refusal to fit dates its rings; Uranus's rebuilt system records the tilt event; Triton arrives from Sol's own dam cohort; and the kinetic regime extends the same machine to impact-born discs, where Earth's two seats packed at $P = 3.2$ and merged into one moon and a farside scar, while Mars's two seats packed at $P = 285$ and kept both children, one of which is being ground back into the Roche zone now. Run upward, the recursion reconstructs Hyperion, the A-class

star whose disc produced Sol, predicts its remnant, and finds a candidate fourteen light-years away already eating the predicted dry-rock diet; the same seat arithmetic computes stellar metallicity offsets, reproducing Sol’s protosolar bulk composition unprompted and the measured metallicity excess of Alpha Centauri B over A that co-natal formation cannot explain.

Run outward, the dam is a production line whose deposits are dated cohorts: the Kuiper belt is the residue of the T Tauri firehose, the cold classicals its mid-era deposition band, the Kuiper cliff its shutdown radius, and the boundary still operates, its stocked shelf observed as the hydrogen wall, its dense-cloud excursions predicting young radionuclide-bearing bodies at accessible radii. The same boundary, under the name the literature independently gave it, is the galactopause, and the galactic core is constructed by the same machine that built Sol’s planets, with seats filled by direct-collapse objects and massive clusters where Sol’s seats held planets. The templating epoch survives in fossils the paper enumerates, the deposition profile, the spiral pitch lattice, the founder coupling, and the compression family of bulges, and a generative integration of the sequence with one tuned normalisation reproduces the Milky Way’s gross statistics eight for eight.

One empirical finding stands beside the framework rather than inside it: across the cleanly calibrated catalogue the disc-to-star mass ratio is stochastic with respect to stellar class, exactly as the inheritance structure requires, since the draw is environmental at the recursion’s root and seat-determined everywhere below it. One reconstruction remains fenced as a thought experiment, the Crab progenitor cluster; Sol’s progenitor, by contrast, is calibrated and carries its falsifiers.

The paper closes where it began. Two numbers suffice to describe a galaxy because two numbers are what any parent hands any child, and at the top of the recursion, where the parent is the environment itself, the pair already carries the names cosmology gave it. Every mechanism above is listed in the register with the observation that would kill it; the machine’s seats are photographed in the ALMA discs, its dams at the heliopause, its fossils on the far side of the Moon. The framework asks to be tested at every scale on which it claims to organise, and it has tried to make that testing easy.

ACKNOWLEDGMENTS

This synthesis would not be possible without decades of foundational work by the planet-formation and solar-system-dynamics communities. The author particularly acknowledges the influence of frameworks developed by Lynden-Bell & Pringle (viscous discs), Hayashi (MMSN), Mulders (snow line dynamics), Lambrechts & Johansen (pebble accretion), Tanigawa & Ikoma (runaway gas accretion), Chatterjee & Tan (inside-out planet formation), and Morbidelli and Walsh (solar-system dynamics). Any errors of interpretation are the author’s own.

DATA AVAILABILITY

The complete model implementation, including the TypeScript/JavaScript engine, calibration data for the 22 catalogued exoplanetary systems, and a browser-based interactive tool, is available at:

- **Interactive simulator:** <https://jamesgdahl.github.io/HYDROS-Planet-Formation-Model/>
- **Source repository:** <https://github.com/jamesgdahl/HYDROS-Planet-Formation-Model>

The interactive simulator allows direct exploration of the three fundamental system inputs ($M_\star$, Ω , f_{disc}) and per-planet (r, t_{form}) adjustments, displaying the resulting mass allocation and predicted vs observed masses in real time. Built-in presets reproduce the Solar System fit (Table 3) and all 22 catalogued systems (§11.2), plus the thought-experiment reconstructions of §6.

REFERENCES

- Alfvén, H. 1942, *Nature*, 150, 405.
“Existence of Electromagnetic-Hydrodynamic Waves”
- Andrews, S. M., Wilner, D. J., Hughes, A. M., Qi, C., & Dullemond, C. P. 2009, *ApJ*, 700, 1502.
- Andrews, S. M., Wilner, D. J., Hughes, A. M., Qi, C., & Dullemond, C. P. 2010, *ApJ*, 723, 1241.
- Andrews, S. M. 2020, *ARA&A*, 58, 483.
“Observations of Protoplanetary Disk Structures”
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481.
“The Chemical Composition of the Sun”
- Ansdell, M., Williams, J. P., van der Marel, N., et al. 2016, *ApJ*, 828, 46.
“ALMA Survey of Lupus Protoplanetary Disks. I. Dust and Gas Masses”
- Benz, W., Slattey, W. L., & Cameron, A. G. W. 1988, *Icarus*, 74, 516. *“Collisional stripping of Mercury’s mantle”*
- Benz, W., Anic, A., Horner, J., & Whitby, J. A. 2007, *Space Sci. Rev.*, 132, 189.
“The Origin of Mercury”
- Boley, A. C., & Ford, E. B. 2013, arXiv:1306.0566. *“The formation of systems with closely spaced low-mass planets and the application to Kepler-36”*
- Birnstiel, T., Klahr, H., & Ercolano, B. 2012, *A&A*, 539, A148.
- Bitsch, B., Lambrechts, M., & Johansen, A. 2015, *A&A*, 582, A112.
- Bitsch, B., Morbidelli, A., Johansen, A., et al. 2018, *A&A*, 612, A30.
“Pebble-isolation mass: Scaling law and implications for the formation of super-Earths and gas giants”
- Bouvier, J., Alencar, S. H. P., Harries, T. J., Johns-Krull, C. M., & Romanova, M. M. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil (Tucson, AZ: Univ. Arizona Press), 479.
- Canup, R. M., & Asphaug, E. 2001, *Nature*, 412, 708. *“Origin of the Moon in a giant impact near the end of the Earth’s formation”*
- Cameron, A. G. W. 1978, *Moon and Planets*, 18, 5. *“Physics of the primitive solar accretion disk”*

- Cieza, L. A., Ruíz-Rodríguez, D., Hales, A., et al. 2019, MNRAS, 482, 698. “*The Ophiuchus DIsc Survey Employing ALMA (ODISEA): I. Project description and continuum images at deca-Astronomical-Unit resolution*”
- Cameron, A. G. W. 1985, Icarus, 64, 285. “*The partial volatilization of Mercury*”
- Chau, A., Reinhardt, C., Helled, R., & Stadel, J. 2018, ApJ, 865, 35. “*Forming Mercury by giant impacts*”
- Chatterjee, S., & Tan, J. C. 2014, ApJ, 780, 53. “*Inside-Out Planet Formation*”
- Ciesla, F. J., & Cuzzi, J. N. 2006, Icarus, 181, 178. “*The evolution of the water distribution in a viscous protoplanetary disk*”
- Asphaug, E., & Reufer, A. 2014, Nat. Geosci., 7, 564. “*Mercury and other iron-rich planetary bodies as relics of inefficient merging*”
- Bai, X.-N., & Stone, J. M. 2013, ApJ, 769, 76. “*Wind-driven Accretion in Protoplanetary Disks. I. Suppression of the Magnetorotational Instability and Launching of the Magnetocentrifugal Wind*”
- Ebel, D. S., & Alexander, C. M. O. D. 2011, Planet. Space Sci., 59, 1888. “*Equilibrium condensation from chondritic porous IDP enriched vapor: Implications for Mercury and enstatite chondrite origins*”
- Fegley, B., & Cameron, A. G. W. 1987, Earth Planet. Sci. Lett., 82, 207. “*A vaporization model for iron/silicate fractionation in the Mercury protoplanet*”
- Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, AJ, 154, 109. “*The California-Kepler Survey. III. A Gap in the Radius Distribution of Small Planets*”
- Nittler, L. R., Starr, R. D., Weider, S. Z., et al. 2011, Science, 333, 1847. “*The Major-Element Composition of Mercury’s Surface from MESSENGER X-ray Spectrometry*”
- Flock, M., Ruge, J. P., Dzyurkevich, N., et al. 2015, A&A, 574, A68. “*Gaps, rings, and non-axisymmetric structures in protoplanetary disks*”
- Gammie, C. F. 1996, ApJ, 457, 355. “*Layered Accretion in T Tauri Disks*”
- Garaud, P., & Lin, D. N. C. 2007, ApJ, 654, 606. “*The effect of internal dissipation and surface irradiation on the structure of disks and the location of the snow line around sun-like stars*”
- Ghosh, P., & Lamb, F. K. 1979, ApJ, 232, 259.
- Hansen, B. M. S., & Murray, N. 2013, ApJ, 775, 53. “*Testing in Situ Assembly with the Kepler Planet Candidate Sample*”
- Hartmann, L., Calvet, N., Gullbring, E., & D’Alessio, P. 1998, ApJ, 495, 385. “*Accretion and the Evolution of T Tauri Disks*”
- Hayashi, C. 1981, Prog. Theor. Phys. Suppl., 70, 35. “*Structure of the Solar Nebula, Growth and Decay of Magnetic Fields and Effects of Magnetic and Turbulent Viscosities on the Nebula*”
- Hendler, N., Pascucci, I., Pinilla, P., et al. 2020, ApJ, 895, 126. “*The evolution of dust disk sizes from a homogeneous analysis of 1–3 Myr-old stars*”
- Haisch, K. E., Lada, E. A., & Lada, C. J. 2001, ApJ, 553, L153. “*Disk Frequencies and Lifetimes in Young Clusters*”
- Hori, Y., & Ikoma, M. 2010, ApJ, 714, 1343. “*Critical Core Masses for Gas Giant Formation with Grain-Free Envelopes*”
- Hauck, S. A., Margot, J.-L., Solomon, S. C., et al. 2013, J. Geophys. Res. Planets, 118, 1204. “*The curious case of Mercury’s internal structure*”
- Ida, S., & Lin, D. N. C. 2004, ApJ, 604, 388. “*Toward a Deterministic Model of Planetary Formation. I. A Desert in the Mass and Semimajor Axis Distributions of Extrasolar Planets*”
- Ikoma, M., Nakazawa, K., & Emori, H. 2000, ApJ, 537, 1013. “*Formation of Giant Planets: Dependences on Core Accretion Rate and Grain Opacity*”

- Jin, S., Mordasini, C., Parmentier, V., et al. 2014, ApJ, 795, 65. “*Planetary Population Synthesis Coupled with Atmospheric Escape: A Statistical View of Evaporation*”
- Johansen, A., & Lambrechts, M. 2017, Annu. Rev. Earth Planet. Sci., 45, 359. “*Forming Planets via Pebble Accretion*”
- Kennedy, G. M., & Kenyon, S. J. 2008, ApJ, 673, 502. “*Planet formation around stars of various masses: the snow line and the frequency of giant planets*”
- Krijt, S., Ormel, C. W., Dominik, C., & Tielens, A. G. G. M. 2016, A&A, 586, A20. “*A panoptic model for planetesimal formation and pebble delivery*”
- Krijt, S., Schwarz, K. R., Bergin, E. A., & Ciesla, F. J. 2018, ApJ, 864, 78. “*Transport of CO in protoplanetary disks: consequences of pebble formation, settling, and radial drift*”
- Königl, A. 1991, ApJ, 370, L39.
- Kruijer, T. S., Burkhardt, C., Budde, G., & Kleine, T. 2017, PNAS, 114, 6712. “*Age of Jupiter inferred from the distinct genetics and formation times of meteorites*”
- Liu, B., Ormel, C. W., & Lin, D. N. C. 2017, A&A, 601, A15. “*Dynamical rearrangement of super-Earths during disk dispersal*”
- Lodders, K. 2003, ApJ, 591, 1220. “*Solar System Abundances and Condensation Temperatures of the Elements*”
- Lee, E. J., & Chiang, E. 2015, ApJ, 811, 41. “*To Cool is to Accrete: Analytic Scalings for Nebular Accretion of Planetary Atmospheres*”
- Lissauer, J. J., Hubickyj, O., D’Angelo, G., & Bodenheimer, P. 2009, Icarus, 199, 338. “*Models of Jupiter’s growth incorporating thermal and hydrodynamic constraints*”
- Lopez, E. D., & Fortney, J. J. 2013, ApJ, 776, 2. “*The Role of Core Mass in Controlling Evaporation: The Kepler Radius Distribution and the Kepler-36 Density Dichotomy*”
- Lopez, E. D., Fortney, J. J., & Miller, N. 2012, ApJ, 761, 59. “*How Thermal Evolution and Mass-loss Sculpt Populations of Super-Earths and Sub-Neptunes*”
- Lyra, W., Johansen, A., Klahr, H., & Piskunov, N. 2009, A&A, 493, 1125. “*Standing on the shoulders of giants. Trojan Earths and vortex trapping in low mass self-gravitating protoplanetary disks of gas and solids*”
- Lambrechts, M., & Johansen, A. 2012, A&A, 544, A32.
- Lambrechts, M., & Johansen, A. 2014, A&A, 572, A107. “*Forming the cores of giant planets from the radial pebble flux in protoplanetary discs*”
- Long, M., Romanova, M. M., & Lovelace, R. V. E. 2005, ApJ, 634, 1214.
- Long, F., Pinilla, P., Herczeg, G. J., et al. 2018, ApJ, 869, 17.
- Lynden-Bell, D., & Pringle, J. E. 1974, MNRAS, 168, 603. “*The evolution of viscous discs and the origin of the nebular variables*”
- Manara, C. F., Ansdell, M., Rosotti, G. P., et al. 2023, in Protostars and Planets VII, ASP Conf. Ser. 534, ed. S. Inutsuka, Y. Aikawa, T. Muto, K. Tomida, & M. Tamura (San Francisco, CA: ASP), 539. “*Demographics of Young Stars and their Protoplanetary Disks*”
- Min, M., Dullemond, C. P., Kama, M., & Dominik, C. 2011, Icarus, 212, 416. “*The thermal structure and the location of the snow line in the protostellar nebula: Axisymmetric models with full 3-D radiative transfer*”
- Mamajek, E. E. 2009, in AIP Conf. Proc. 1158, Exoplanets and Disks: Their Formation and Diversity, ed. T. Usuda, M. Tamura, & M. Ishii (Melville, NY: AIP), 3. “*Initial Conditions of Planet Formation: Lifetimes of Primordial Disks*”
- Mizuno, H. 1980, Prog. Theor. Phys., 64, 544. “*Formation of the Giant Planets*”

- Mordasini, C. 2014, A&A, 572, A118.
“Grain opacity and the bulk composition of extrasolar planets”
- Murray-Clay, R. A., Chiang, E. I., & Murray, N. 2009, ApJ, 693, 23.
“Atmospheric Escape from Hot Jupiters”
- Mulders, G. D., Ciesla, F. J., Min, M., & Pascucci, I. 2015, ApJ, 807, 9. *“The snow line in viscous disks around low-mass stars: implications for water delivery to terrestrial planets in the habitable zone”*
- Morbidelli, A., Tsiganis, K., Crida, A., Levison, H. F., & Gomes, R. 2007, AJ, 134, 1790. *“Dynamics of the Giant Planets of the Solar System in the Gaseous Protoplanetary Disk and Their Relationship to the Current Orbital Architecture”*
- Mulders, G. D., Pascucci, I., Manara, C. F., et al. 2017, ApJ, 847, 31.
- Najita, J. R., & Bergin, E. A. 2018, ApJ, 864, 168. *“Protoplanetary Disk Sizes and Angular Momentum Transport”*
- Ogihara, M., Morbidelli, A., & Guillot, T. 2015, A&A, 578, A36. *“A reassessment of the in situ formation of close-in super-Earths”*
- Oka, A., Nakamoto, T., & Ida, S. 2011, ApJ, 738, 141. *“Evolution of snow line in optically thick protoplanetary disks: effects of water ice opacity and dust grain size”*
- Ormel, C. W., & Klahr, H. H. 2010, A&A, 520, A43. *“The effect of gas drag on the growth of protoplanets. Analytical expressions for the accretion of small bodies in laminar disks”*
- Owen, J. E., & Wu, Y. 2013, ApJ, 775, 105. *“Kepler Planets: A Tale of Evaporation”*
- Owen, J. E., & Wu, Y. 2017, ApJ, 847, 29. *“The Evaporation Valley in the Kepler Planets”*
- Owen, J. E., & Wu, Y. 2017, ApJ, 847, 29.
- Peplowski, P. N., Evans, L. G., Hauck, S. A., et al. 2011, Science, 333, 1850.
“Radioactive Elements on Mercury’s Surface from MESSENGER: Implications for the Planet’s Formation and Evolution”
- Picogna, G., Ercolano, B., Owen, J. E., & Weber, M. L. 2019, MNRAS, 487, 691.
- Pinilla, P., Birnstiel, T., Ricci, L., et al. 2012, A&A, 538, A114. *“Trapping dust particles in the outer regions of protoplanetary disks”*
- Pinilla, P., Flock, M., Ovelar, M. de J., & Birnstiel, T. 2016, A&A, 596, A81.
“Can dead zones create structures like a transition disk?”
- Pierens, A., & Raymond, S. N. 2011, A&A, 533, A131.
- Pollack, J. B., & Bodenheimer, P. 1989, in *Origin and Evolution of Planetary and Satellite Atmospheres*, ed. S. K. Atreya, J. B. Pollack, & M. S. Matthews (Tucson, AZ: Univ. Arizona Press), 564.
- Pascucci, I., Testi, L., Herczeg, G. J., et al. 2016, ApJ, 831, 125. *“A Steeper than Linear Disk Mass-Stellar Mass Scaling Relation”*
- Piso, A.-M. A., & Youdin, A. N. 2014, ApJ, 786, 21. *“On the Minimum Core Mass for Giant Planet Formation at Wide Separations”*
- Pollack, J. B., Hollenbach, D., Beckwith, S., et al. 1994, ApJ, 421, 615.
“Composition and radiative properties of grains in molecular clouds and accretion disks”
- Pollack, J. B., Hubickyj, O., Bodenheimer, P., et al. 1996, Icarus, 124, 62.
- Pringle, J. E. 1981, ARA&A, 19, 137.
- Regály, Z., Sándor, Z., Dullemond, C. P., & Kiss, L. L. 2012, MNRAS, 419, 1701.
“Detectability of dead zone edges as planet traps in protoplanetary discs”
- Rafikov, R. R. 2006, ApJ, 648, 666.
“Atmospheres of Protoplanetary Cores: Critical Mass for Nucleated Instability”

- Ribas, I., Guinan, E. F., Güdel, M., & Audard, M. 2005, ApJ, 622, 680. “*Evolution of the Solar Activity over Time and Effects on Planetary Atmospheres. I. High-Energy Irradiances (1-1700 Å)*”
- Romanova, M. M., & Lovelace, R. V. E. 2006, ApJ, 645, L73. “*The Magnetospheric Gap and the Accumulation of Giant Planets Close to a Star*”
- Romanova, M. M., Kulkarni, A. K., & Lovelace, R. V. E. 2008, ApJ, 673, L171.
- Safronov, V. S. 1972, *Evolution of the Protoplanetary Cloud and Formation of the Earth and Planets* (Jerusalem: Israel Program for Scientific Translations; English translation of 1969 Russian original). *Foundational monograph introducing the “feeding zone” concept for planetary accretion from disc material.*
- Schoonenberg, D., & Ormel, C. W. 2017, A&A, 602, A21. “*Planetesimal formation near the snowline: in or out?*”
- Drażkowska, J., & Alibert, Y. 2017, A&A, 608, A92. “*Planetesimal formation starts at the snow line*”
- Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 24, 337. “*Black holes in binary systems. Observational appearance*”
- Shu, F., Najita, J., Ostriker, E., et al. 1994, ApJ, 429, 781.
- Tu, L., Johnstone, C. P., Güdel, M., & Lammer, H. 2015, A&A, 577, L3. “*The extreme ultraviolet and X-ray Sun in Time: High-energy evolutionary tracks of a solar-like star*”
- Turner, N. J., Fromang, S., Gammie, C., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning (Tucson, AZ: Univ. Arizona Press), 411. “*Transport and Accretion in Planet-Forming Disks*”
- Trapman, L., Facchini, S., Hogerheijde, M. R., van Dishoeck, E. F., & Bruderer, S. 2019, A&A, 629, A79.
- Tsiganis, K., Gomes, R., Morbidelli, A., & Levison, H. F. 2005, Nature, 435, 459.
- Walsh, K. J., Morbidelli, A., Raymond, S. N., O’Brien, D. P., & Mandell, A. M. 2011, Nature, 475, 206. “*A low mass for Mars from Jupiter’s early gas-driven migration*”
- Williams, J. P., Cieza, L., Hales, A., et al. 2019, ApJ, 875, L9. “*The Ophiuchus Disk Survey Employing ALMA (ODISEA): Disk dust mass distributions across protostellar evolutionary classes*”
- Wright, N. J., Drake, J. J., Mamajek, E. E., & Henry, G. W. 2011, ApJ, 743, 48. “*The Stellar-activity-Rotation Relationship and the Evolution of Stellar Dynamos*”
- Weidenschilling, S. J. 1977, Astrophys. Space Sci., 51, 153. “*The distribution of mass in the planetary system and solar nebula*”
- Williams, J. P., & Cieza, L. A. 2011, ARA&A, 49, 67.
- Armitage, P. J. 2018, in *Handbook of Exoplanets*, ed. H. J. Deeg & J. A. Belmonte (Cham: Springer), 2185. “*A Brief Overview of Planet Formation*”
- Asphaug, E. 2010, Chemie der Erde / Geochemistry, 70, 199. “*Similar-sized collisions and the diversity of planets*”
- Canup, R. M. 2012, Science, 338, 1052. “*Forming a Moon with an Earth-like Composition via a Giant Impact*”
- Catalán, S., Isern, J., García-Berro, E., & Ribas, I. 2008, MNRAS, 387, 1693. “*The initial-final mass relationship of white dwarfs revisited*”
- Goldreich, P., & Tremaine, S. 1980, ApJ, 241, 425. “*Disk-satellite interactions*”
- Ida, S. 1990, Icarus, 88, 129. “*Stirring and dynamical friction rates of planetesimals in the solar gravitational field*”
- Jewitt, D., & Haghighipour, N. 2007, ARA&A, 45, 261. “*Irregular Satellites of the Planets: Products of Capture in the Early Solar System*”
- Kominami, J., & Ida, S. 2002, Icarus, 157, 43. “*The Effect of Tidal Interaction with a Gas Disk on Formation of Terrestrial Planets*”

- Leinhardt, Z. M., & Stewart, S. T. 2012, *ApJ*, 745, 79. “*Collisions between Gravity-dominated Bodies. I. Outcome Regimes and Scaling Laws*”
- Williams, D. R. 2024, Venus Fact Sheet (Greenbelt, MD: NASA NSSDCA).
- Williams, D. R. 2024, Neptune Fact Sheet (Greenbelt, MD: NASA NSSDCA).
- Nesvorný, D., Vokrouhlický, D., & Morbidelli, A. 2007, *AJ*, 133, 1962. “*Capture of Irregular Satellites during Planetary Encounters*”
- Nesvorný, D. 2011, *ApJ*, 742, L22. “*Young Solar System’s Fifth Giant Planet?*”
- Nesvorný, D., & Morbidelli, A. 2012, *AJ*, 144, 117. “*Statistical Study of the Early Solar System’s Instability with Four, Five, and Six Giant Planets*”
- Sheppard, S. S., & Jewitt, D. C. 2003, *Nature*, 423, 261. “*An abundant population of small irregular satellites around Jupiter*”
- Standish, E. M. 1998, JPL Interoffice Memorandum 312.F-98-048. “*JPL Planetary and Lunar Ephemerides, DE405/LE406*”
- Tanigawa, T., & Ikoma, M. 2007, *ApJ*, 667, 557. “*A Systematic Study of the Final Masses of Gas Giant Planets*”
- Thue, A. 1910, *Norske Vid. Selsk. Skr.*, 1, 1. “*Über die dichteste Zusammenstellung von kongruenten Kreisen in einer Ebene*”
- Conway, J. H., & Sloane, N. J. A. 1999, *Sphere Packings, Lattices and Groups* (3rd ed.; New York: Springer).
- Gladman, B. 1993, *Icarus*, 106, 247. “*Dynamics of systems of two close planets*”
- Chambers, J. E., Wetherill, G. W., & Boss, A. P. 1996, *Icarus*, 119, 261. “*The Stability of Multi-Planet Systems*”
- Wisdom, J. 1980, *AJ*, 85, 1122. “*The resonance overlap criterion and the onset of stochastic behavior in the restricted three-body problem*”
- Spiegel, D. S., Burrows, A., & Milsom, J. A. 2011, *ApJ*, 727, 57. “*The Deuterium-burning Mass Limit for Brown Dwarfs and Giant Planets*”
- Tielens, A. G. G. M. 1983, *A&A*, 119, 177. “*Surface chemistry of deuterated molecules*”
- Chabrier, G., & Baraffe, I. 2000, *ARA&A*, 38, 337. “*Theory of Low-Mass Stars and Substellar Objects*”
- Duchêne, G., & Kraus, A. 2013, *ARA&A*, 51, 269. “*Stellar Multiplicity*”
- Smith, A. W., & Lissauer, J. J. 2009, *Icarus*, 201, 381. “*Orbital stability of systems of closely-spaced planets*”
- Lada, C. J., & Lada, E. A. 2003, *ARA&A*, 41, 57. “*Embedded Clusters in Molecular Clouds*”
- Ramírez, I., Bajkova, A. T., Bobylev, V. V., et al. 2014, *ApJ*, 787, 154. “*Elemental Abundances of Solar Sibling Candidates*”
- Yoder, C. F. 1979, *Nature*, 279, 767. “*How tidal heating in Io drives the Galilean orbital resonance locks*”
- Le Verrier, U. J. 1859, *Comptes Rendus Acad. Sci. Paris*, 49, 379. “*Lettre de M. Le Verrier à M. Faye sur la théorie de Mercure et sur le mouvement du périhélie de cette planète*”
- Burkert, A., & Bodenheimer, P. 2000, *ApJ*, 543, 822. “*Turbulent Molecular Cloud Cores: Rotational Properties*”
- Kuffmeier, M., Haugbølle, T., & Nordlund, Å. 2017, *ApJ*, 846, 7. “*Zoom-in Simulations of Protoplanetary Disks Starting from GMC Scales*”
- Bate, M. R. 2018, *MNRAS*, 475, 5618. “*On the diversity and statistical properties of protostellar discs*”
- Mathis, J. S., Rumpl, W., & Nordsieck, K. H. 1977, *ApJ*, 217, 425. “*The size distribution of interstellar grains*”
- Hill, G. W. 1878, *Am. J. Math.*, 1, 5. “*Researches in the Lunar Theory*”
- Murray, C. D., & Dermott, S. F. 1999, *Solar System Dynamics* (Cambridge: Cambridge Univ. Press).
- Laskar, J. 1997, *A&A*, 317, L75. “*Large scale chaos and the spacing of the inner planets*”

- Brasser, R., Walsh, K. J., & Nesvorný, D. 2013, MNRAS, 433, 3417. “*Constraining the primordial orbits of the terrestrial planets*”
- Souami, D., & Souchay, J. 2012, A&A, 543, A133. “*The solar system’s invariable plane*”
- Jacobson, S. A., Rubie, D. C., Hernlund, J., Morbidelli, A., & Nakajima, M. 2017, Earth Planet. Sci. Lett., 474, 375. “*Formation, stratification, and mixing of the cores of Earth and Venus*”
- Malhotra, R. 1993, Nature, 365, 819. “*The origin of Pluto’s peculiar orbit*”
- Malhotra, R. 1995, AJ, 110, 420. “*The origin of Pluto’s orbit: implications for the Solar System beyond Neptune*”
- Weidenschilling, S. J., & Davis, D. R. 1985, Icarus, 62, 16. “*Orbital resonances in the solar nebula: Implications for planetary accretion*”
- Tiscareno, M. S., & Malhotra, R. 2009, AJ, 138, 827. “*Chaotic Diffusion of Resonant Kuiper Belt Objects*”
- Lykawka, P. S., & Mukai, T. 2007, Icarus, 192, 238. “*Resonance sticking in the scattered disk*”
- Gladman, B., Lawler, S. M., Petit, J.-M., et al. 2012, AJ, 144, 23. “*The Resonant Trans-Neptunian Populations*”
- Agnor, C. B., & Hamilton, D. P. 2006, Nature, 441, 192. “*Neptune’s capture of its moon Triton in a binary-planet gravitational encounter*”
- Oort, J. H. 1950, Bull. Astron. Inst. Netherlands, 11, 91. “*The structure of the cloud of comets surrounding the Solar System and a hypothesis concerning its origin*”
- Duncan, M., Quinn, T., & Tremaine, S. 1987, AJ, 94, 1330. “*The formation and extent of the solar system comet cloud*”
- Youdin, A. N., & Goodman, J. 2005, ApJ, 620, 459. “*Streaming Instabilities in Protoplanetary Disks*”
- Nesvorný, D., Youdin, A. N., & Richardson, D. C. 2010, AJ, 140, 785. “*Formation of Kuiper Belt Binaries by Gravitational Collapse*”
- Pitjeva, E. V., & Pitjev, N. P. 2018, Astron. Lett., 44, 554. “*Masses of the Main Asteroid Belt and the Kuiper Belt from the Motions of Planets and Spacecraft*”
- Doner, A., Horányi, M., Bagenal, F., et al. 2024, ApJ, 961, L38. “*New Horizons Venetia Burney Student Dust Counter Observes Higher than Expected Fluxes Approaching 60 au*”
- Fraser, W. C., Porter, S. B., Peltier, L., et al. 2024, PSJ, 5, 227. “*Candidate Distant Trans-Neptunian Objects Detected by the New Horizons Subaru TNO Survey*”
- Skumanich, A. 1972, ApJ, 171, 565. “*Time Scales for Ca II Emission Decay, Rotational Braking, and Lithium Depletion*”
- Stone, E. C., Cummings, A. C., McDonald, F. B., et al. 2013, Science, 341, 150. “*Voyager 1 Observes Low-Energy Galactic Cosmic Rays in a Region Depleted of Heliospheric Ions*”
- Brasser, R., Duncan, M. J., & Levison, H. F. 2006, Icarus, 184, 59. “*Embedded star clusters and the formation of the Oort cloud*”
- Levison, H. F., Duncan, M. J., Brasser, R., & Kaufmann, D. E. 2010, Science, 329, 187. “*Capture of the Sun’s Oort Cloud from Stars in Its Birth Cluster*”
- Matrà, L., Marino, S., Kennedy, G. M., et al. 2018, ApJ, 859, 72. “*An Empirical Planetesimal Belt Radius–Stellar Luminosity Relation*”
- Frisch, P. C., Redfield, S., & Slavin, J. D. 2011, ARA&A, 49, 237. “*The Interstellar Medium Surrounding the Sun*”
- Zucker, C., Goodman, A. A., Alves, J., et al. 2022, Nature, 601, 334. “*Star formation near the Sun is driven by expansion of the Local Bubble*”
- Wood, B. E. 2004, Living Reviews in Solar Physics, 1, 2. “*Astrospheres and Solar-like Stellar Winds*”

- Opher, M., Loeb, A., & Peek, J. E. G. 2024, *Nature Astronomy*, 8, 567. “A possible direct exposure of the Earth to the cold dense interstellar medium 2–3 Myr ago”
- Shull, J. M., & Moss, J. A. 2020, *ApJ*, 903, 101. “Galactopause Formation and Gas Precipitation During Strong Galactic Outflows”
- Elmegreen, B. G., & Lada, C. J. 1977, *ApJ*, 214, 725. “Sequential formation of subgroups in OB associations”
- Fall, S. M., & Rees, M. J. 1985, *ApJ*, 298, 18. “A theory for the origin of globular clusters”
- Peebles, P. J. E. 1969, *ApJ*, 155, 393. “Origin of the Angular Momentum of Galaxies”
- Iess, L., Militzer, B., Kaspi, Y., et al. 2019, *Science*, 364, aat2965. “Measurement and implications of Saturn’s gravity field and ring mass”
- Čuk, M., Dones, L., & Nesvorný, D. 2016, *ApJ*, 820, 97. “Dynamical Evidence for a Late Formation of Saturn’s Moons”
- Yu, S.-Y., & Ho, L. C. 2020, *ApJ*, 900, doi:10.3847/1538-4357/abac5b. “The Statistical Properties of Spiral Arms in Nearby Disk Galaxies”
- Kormendy, J., & Ho, L. C. 2013, *ARA&A*, 51, 511. “Coevolution (Or Not) of Supermassive Black Holes and Host Galaxies”
- Reid, M. J., & Brunthaler, A. 2004, *ApJ*, 616, 872. “The Proper Motion of *Sagittarius A**”
- Gebhardt, K., Lauer, T. R., Kormendy, J., et al. 2001, *AJ*, 122, 2469. “M33: A Galaxy with No Supermassive Black Hole”
- Dressler, A. 1980, *ApJ*, 236, 351. “Galaxy morphology in rich clusters, Implications for the formation and evolution of galaxies”
- Nogueras-Lara, F., Schödel, R., Gallego-Calvente, A. T., et al. 2020, *Nature Astronomy*, 4, 377. “Early formation and recent starburst activity in the nuclear disk of the Milky Way”
- Su, M., Slatyer, T. R., & Finkbeiner, D. P. 2010, *ApJ*, 724, 1044. “Giant Gamma-ray Bubbles from Fermi-LAT: Active Galactic Nucleus Activity or Bipolar Galactic Wind?”
- Johansen, A., Youdin, A., & Klahr, H. 2009, *ApJ*, 697, 1269. “Zonal Flows and Long-lived Axisymmetric Pressure Bumps in Magnetorotational Turbulence”
- Bai, X.-N., & Stone, J. M. 2014, *ApJ*, 796, 31. “Magnetic Flux Concentration and Zonal Flows in Magnetorotational Instability Turbulence”
- Dittrich, K., Klahr, H., & Johansen, A. 2013, *ApJ*, 763, 117. “Gravoturbulent Planetesimal Formation: The Positive Effect of Long-lived Zonal Flows”
- Dullemond, C. P., Birnstiel, T., Huang, J., et al. 2018, *ApJL*, 869, L46. “The Disk Substructures at High Angular Resolution Project (DSHARP). VI. Dust Trapping in Thin-ringed Protoplanetary Disks”
- Lin, D. N. C., Bodenheimer, P., & Richardson, D. C. 1996, *Nature*, 380, 606. “Orbital migration of the planetary companion of 51 Pegasi to its present location”
- Masset, F. S., Morbidelli, A., Crida, A., & Ferreira, J. 2006, *ApJ*, 642, 478. “Disk Surface Density Transitions as Protoplanet Traps”
- ALMA Partnership, Brogan, C. L., Pérez, L. M., et al. 2015, *ApJL*, 808, L3. “The 2014 ALMA Long Baseline Campaign: First Results from High Angular Resolution Observations toward the HL Tau Region”
- Jutzi, M., & Asphaug, E. 2011, *Nature*, 476, 69. “Forming the lunar farside highlands by accretion of a companion moon”
- Canup, R. M., & Ward, W. R. 2006, *Nature*, 441, 834. “A common mass scaling for satellite systems of gaseous planets”
- Mestel, L. 1963, *MNRAS*, 126, 553. “On the galactic law of rotation”

- Mann, A. W., Feiden, G. A., Gaidos, E., Boyajian, T., & von Braun, K. 2015, *ApJ*, 804, 64. “*How to Constrain Your M Dwarf: Measuring Effective Temperature, Bolometric Luminosity, Mass, and Radius*”
- Díez Alonso, E., Caballero, J. A., Montes, D., et al. 2019, *A&A*, 621, A126. “*CARMENES input catalogue of M dwarfs. IV. New rotation periods from photometric time series*”
- Holmberg, J., Nordström, B., & Andersen, J. 2009, *A&A*, 501, 941. “*The Geneva-Copenhagen survey of the solar neighbourhood. III. Improved distances, ages, and kinematics*”
- Andrews-Hanna, J. C., Zuber, M. T., & Banerdt, W. B. 2008, *Nature*, 453, 1212. “*The Borealis basin and the origin of the martian crustal dichotomy*”
- Hesselbrock, A. J., & Minton, D. A. 2017, *Nature Geoscience*, 10, 266. “*An ongoing satellite–ring cycle of Mars and the origins of Phobos and Deimos*”
- Morel, T. 2018, *A&A*, 615, A172. “*The chemical composition of α Centauri AB revisited*”
- Zolotov, A., Dekel, A., Mandelker, N., et al. 2015, *MNRAS*, 450, 2327. “*Compaction and quenching of high- z galaxies in cosmological simulations: blue and red nuggets*”
- Huang, J., Andrews, S. M., Dullemond, C. P., et al. 2018, *ApJL*, 869, L42. “*The Disk Substructures at High Angular Resolution Project (DSHARP). II. Characteristics of Annular Substructures*”
- Paumard, T., Genzel, R., Martins, F., et al. 2006, *ApJ*, 643, 1011. “*The Two Young Star Disks in the Central Parsec of the Galaxy: Properties, Dynamics, and Formation*”
- Tegler, S. C., & Romanishin, W. 2000, *Nature*, 407, 979. “*Extremely red Kuiper-belt objects in near-circular orbits beyond 40 AU*”
- Brownlee, D., Tsou, P., Aléon, J., et al. 2006, *Science*, 314, 1711. “*Comet 81P/Wild 2 Under a Microscope*”
- Maiolino, R., Russell, H. R., Fabian, A. C., et al. 2017, *Nature*, 544, 202. “*Star formation inside a galactic outflow*”
- Gallagher, R., Maiolino, R., Belfiore, F., et al. 2019, *MNRAS*, 485, 3409. “*Widespread star formation inside galactic outflows*”
- Okamoto, Y. K., Kataza, H., Honda, M., et al. 2004, *Nature*, 431, 660. “*An early extrasolar planetary system revealed by planetesimal belts in β Pictoris*”
- Zhu, W., & Wu, Y. 2018, *AJ*, 156, 92. “*The Super Earth–Cold Jupiter Relations*”
- Meshkat, T., Mawet, D., Bryan, M. L., et al. 2017, *AJ*, 154, 245. “*A Direct Imaging Survey of Spitzer-detected Debris Disks: Occurrence of Giant Planets in Dusty Systems*”
- Rivera, E. J., Laughlin, G., Butler, R. P., et al. 2010, *ApJ*, 719, 890. “*The Lick-Carnegie Exoplanet Survey: A Uranus-mass Fourth Planet for GJ 876 in an Extrasolar Laplace Configuration*”
- Goździewski, K., & Migaszewski, C. 2014, *MNRAS*, 440, 3140. “*Multiple mean motion resonances in the HR 8799 planetary system*”
- Kervella, P., Thévenin, F., & Lovis, C. 2017, *A&A*, 598, L7. “*Proxima’s orbit around α Centauri*”
- Breslau, A., Steinhausen, M., Vincke, K., & Pfalzner, S. 2014, *A&A*, 565, A130. “*Sizes of protoplanetary discs after star-disc encounters*”
- Davis, L., Jr. 1955, *Phys. Rev.*, 100, 1440. “*Interplanetary Magnetic Fields and Cosmic Rays*”
- Parker, E. N. 1958, *ApJ*, 128, 664. “*Dynamics of the Interplanetary Gas and Magnetic Fields*”
- Luger, R., Sestovic, M., Kruse, E., et al. 2017, *Nature Astronomy*, 1, 0129. “*A seven-planet resonant chain in TRAPPIST-1*”
- Steffen, J. H., Ragozzine, D., Fabrycky, D. C., et al. 2012, *PNAS*, 109, 7982. “*Kepler constraints on planets near hot Jupiters*”

- Li, R., & Chiang, E. 2025, ApJ, in press (arXiv:2508.04776). “*In-Situ Formation of the Cold Classical Kuiper Belt*”
- Bovaird, T., & Lineweaver, C. H. 2013, MNRAS, 435, 1126. “*Exoplanet predictions based on the generalized Titius–Bode relation*”
- Raymond, S. N., & Izidoro, A. 2017, Icarus, 297, 134. “*Origin of water in the inner Solar System: Planetesimals scattered inward during Jupiter and Saturn’s rapid gas accretion*”
- Ward, W. R., & Hamilton, D. P. 2004, AJ, 128, 2501. “*Tilting Saturn. I. Analytic Model*”
- Kratter, K., & Lodato, G. 2016, ARA&A, 54, 271. “*Gravitational Instabilities in Circumstellar Disks*”
- Stern, S. A. 2003, Nature, 424, 639. “*The evolution of comets in the Oort cloud and Kuiper belt*”