Gene	eQTL colocalization, 35 Tier-1/2 GWAS SuSiE PP.H4, study	 ABF PP.H4, study	Fine-mapped coding variant	Bulk liver RNA 774 genetic-identity clusters (138 control, 636 disease)		Class
GLP1R	no Tier-1/2 SuSiE result	0.022 MVP NAFLD, EAS	none	not detected		Untested
	Route: gene level, acquire the missing decisive assay; molecular object, acquire the missing decisive measurement					
HKDC1	0.988, UK Biobank AST > 0.5 in 3 of 5 studies	0.989 UK Biobank AST	none		+1.07 q 8.0e-7	Both supported*
	Route: gene level, allele-aware edit plus gene perturbation; molecular object, gene perturbation + rescue					
MTARC1	no Tier-1/2 SuSiE result	0.046 FinnGen NASH	missense, PIP 0.76 PDFF GWAS PIP ≥ 0.5 in 3 studies		-0.15 q 1.2e-4	Unresolved
	Route: gene level, measure the single most decisive missing layer; molecular object, gene perturbation + rescue					
THRB	> 0.999, UK Biobank GGT > 0.5 in 1 of 4 studies	> 0.999 UK Biobank GGT	none		-0.20 q 4.2e-10	Genetically anchored
	Route: gene level, allele-aware regulatory editing; molecular object, gene perturbation + rescue					



RNA supported = BH q < 0.05 and |log2FC| > 0.5 (dashed lines). Genetic call = SuSiE PP.H4 > 0.5; ABF for reference only.
Coding variant: Catalog r9d field, not part of the genetic call.
* Requires a supported call in each domain; direction not compared.