
signed eQTL direction, best head, by |z| stratum
all rows; 111,426 rows / 55 blocks
|z| >= 1; 47,777 rows / 54 blocks
|z| >= 2; 17,113 rows / 49 blocks
|z| >= 3; 8,862 rows / 39 blocks
|z| >= 5; 3,847 rows / 31 blocks
0.450
0.475
0.500
0.525
sign AUC (95% LD−block bootstrap, uncorrected across 5 strata)
within−block permutation chance
sequence−attributable bound
achieved AUC
released expression instrument, severity transfer, no refit
GSE268273, 109 participants (chose transform)
GSE213621, 263 genetic identities
GSE276114 MASLD, 81 participants
GSE276114 chronic viral hepatitis, 82 participants
0.4
0.6
0.8
1.0
Spearman with stage (stage−stratified 95% bootstrap)
stage tie ceiling
achieved Spearman
LD−block join
own block left in; 644,556 rows / 7,576 genes
own block left in; 15,192 rows / 198 genes
own block held out; 15,192 rows / 198 genes
0.00
0.25
0.50
0.75
1.00
Pearson r, per−gene mean effect vs row effect
own block left in
own block held out
routing: 521 samples, 5 cohorts, no donor key
rule as fitted
oracle router
pre−declared floor
0.000
0.025
0.050
0.075
0.100
macro Spearman gain over the reference model
pre−declared floor
oracle router bound
rule as fitted