
supported
H3K27ac map, activity >= 3, women only; 36 vs 49 participants; post hoc
H3K27ac map, activity >= 3; 50 vs 49 participants; precommitted
H3K27ac map, activity >= 4; 35 vs 64 participants; post hoc
liver zone from expression; 19 participants; precommitted
−1
0
1
2
untestable
H3K27ac map, activity >= 4, women, normal livers out; 27 vs 34 participants; post hoc
ATAC map, any fibrosis; 15 vs 22 donors; precommitted
H3K27ac map, activity >= 4, normal livers out; 35 vs 40 participants; post hoc
ATAC map, steatosis >= 5%; 20 vs 18 donors; precommitted
program h2, liver fat MRI GWAS; h2 z 6.35
program h2, Ghouse cirrhosis GWAS; h2 z 6.29
program h2, MVP cirrhosis GWAS; h2 z 6.23
program h2, MVP NAFLD GWAS; h2 z 5.57
program h2, FinnGen NAFLD GWAS; h2 z 4.36
program h2, Ghodsian NAFLD GWAS; h2 z 2.83
program h2, FinnGen cirrhosis GWAS; h2 z 2.33
signed eQTL direction; 111,426 rows, 55 LD blocks
−1
0
1
2
tested negative
routing adds skill; 521 samples, 5 cohorts
treatment response, diet arm; 6 vs 16 participants
−1
0
1
2
(estimate − null) / (threshold − null): 0 = null, 1 = threshold
supported
untestable
tested negative
bound measured from the data