

# Genetic Characterization and Detection of *Angiostrongylus cantonensis* by Molecular Approaches

Muxin Chen,<sup>1,5,\*</sup> Dana Huang,<sup>1,\*</sup> Jiaxu Chen,<sup>2-6,\*</sup> Yalan Huang,<sup>1</sup> Huiwen Zheng,<sup>1</sup> Yijun Tang,<sup>1</sup> Qian Zhang,<sup>1</sup> Shaohong Chen,<sup>2-5</sup> Lin Ai,<sup>2-5,7</sup> Xiaonong Zhou,<sup>2-5,7</sup> and Renli Zhang<sup>1</sup>

## Abstract

*Angiostrongylus cantonensis* constitutes a major etiologic agent of eosinophilic meningoencephalitis. The detection methods for angiostrongyliasis mainly depend on morphology or immunology. A firmer diagnosis could be reached by directly detecting the parasite in the cerebrospinal fluid or through laboratory assays that are specific for *Angiostrongylus*-induced antibodies or the parasite's DNA. *A. cantonensis* detection could be carried out by larva release from the tissue upon pepsin digestion. However, the procedure requires live mollusks, which might complicate the analysis of large amounts of samples. Since morphological assays are limited, multiple molecular techniques have been put forward for detecting *A. cantonensis*, including PCR amplification of targets followed by fragment length or DNA sequence analysis. This allows rapid and accurate identification of *A. cantonensis* for efficient infection management and epidemiological purposes. In this study, we reviewed the current methods, concepts, and applications of molecular approaches to better understand the genetic characterization, molecular detection methods, and practical application of molecular detection in *A. cantonensis*.

**Keywords:** *Angiostrongylus cantonensis*, genetic characterization, molecular detection methods, detection

## Introduction

**A**NGIOSTRONGYLUS CANTONENSIS, also called rat lung-worm, represents an important pathogen responsible for a zoonotic disease affecting humans (accidental host), rats (definitive host), and freshwater snails (intermediate hosts) (Himsworth et al. 2013, Barratt et al. 2016, Huang et al. 2019, Lima et al. 2020). This nematode causes angiostrongyliasis with the prominent symptom of eosinophilic meningoencephalitis in humans, which involves elevated cerebrospinal fluid (CSF) eosinophil amounts (Sawanyawisuth et al. 2012, Tsai et al. 2012, Aghazadeh et al. 2015a, Wang et al. 2020, Zhou et al. 2020).

Angiostrongyliasis-related symptoms result from the presence of larvae or developing worms and host response in the brain. Dead or dying worms in the central nervous system (CNS) can induce inflammation and cause disease (Cross and Chen 2007, Barratt et al. 2016). Major signs of eosinophilic meningitis include serious headache, neck stiffness, paresthesia, and gastrointestinal tract symptoms such as vomiting and nausea (Wilkins et al. 2013, Ansdell and Wattanagoon 2018). Moreover, encephalitic angiostrongyliasis can be fatal (Sawanyawisuth et al. 2009, Wang et al. 2012, Cowie et al. 2017).

Clinical detection of angiostrongyliasis remains challenging, including both patient examination and laboratory

<sup>1</sup>Institute of Pathogenic Biology, Shenzhen Center for Disease Control and Prevention, Shenzhen, China.

<sup>2</sup>Health Education and Detection Center, National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention (Chinese Center for Tropical Diseases Research), Shanghai, China.

<sup>3</sup>Health Education and Detection Center, NHC Key Laboratory for Parasitology and Vector Biology, Shanghai, China.

<sup>4</sup>Health Education and Detection Center, WHO Collaborating Center for Tropical Diseases, Shanghai, China.

<sup>5</sup>Health Education and Detection Center, National Center for International Research on Tropical Diseases, Shanghai, China.

<sup>6</sup>Health Education and Detection Center, National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention, Shenzhen Center for Disease Control and Prevention, Joint Laboratory for Imported Tropical Disease Control, Shanghai, China.

<sup>7</sup>Department of One Health, School of Global Health, Chinese Center for Tropical Diseases Research, Shanghai Jiao Tong University School of Medicine, Shanghai, China.

\*These authors contributed equally to this work.

assessment of *Angiostrongylus*-specific antibodies or parasite DNA (Eamsobhana et al. 2018, Graeff-Teixeira et al. 2018, Somboonpatarakun et al. 2019, 2020). It is uncommon to detect *A. cantonensis* larvae in the CSF (Kuberski and Wallace 1979, Punyagupta et al. 1990, Prociv et al. 2000, Chye et al. 2004, Graeff-Teixeira et al. 2009). High circulating content of eosinophils is an indication of this disease, but is not definitive (Lv et al. 2017b). Computed tomography and magnetic resonance imaging could help detect brain lesions, but do not provide differential diagnosis (Kanpittaya et al. 2000, Tsai et al. 2003, Jin et al. 2005).

Researchers have examined the impact of *A. cantonensis* on human health by studying the epidemiology, morphology, pathology, diagnosis, and treatment of the associated disease (Kwon et al. 2013, Martin-Alonso et al. 2015, Aghazadeh et al. 2016, Chen et al. 2016, Waugh et al. 2016). In this review, we presented the current approaches and issues surrounding the genetic characterization of and molecular detection methods for *A. cantonensis*, as well as the practical application of molecular detection.

## Genotyping of *A. cantonensis*

### Ribosomal DNA sequences

A ribosomal DNA (rDNA) represents a DNA sequence encoding ribosomal RNA (rRNA). The rDNA in eukaryotic cells comprises a tandem repeat of a single operon encompassing NTS, ETS, 18S, ITS1, 5.8S, ITS2, and 28S tracts (Alvarez and Wendel 2003). The various coding regions of these rDNA repeats frequently indicate distinct evolutionary levels. This difference subsequently provides phylogenetic information of species that belong to multiple systematic levels (Hillis and Dixon 1991, Salim and Gerton 2019).

### Internal transcribed spacer sequences

Internal transcribed spacer (ITS) sequences refer to DNA spaces found between small-subunit (SSU) and large-subunit rRNA genes on DNA or the corresponding transcript (Lafontaine and Tollervey 2001, Coleman 2015). Previous reports have demonstrated that the first and second ITSs (ITS-1 and ITS-2) of rDNA could help identify *A. cantonensis* at the species level (Caldeira et al. 2003, Foronda 2010, Qvarnstrom et al. 2010, Liu et al. 2011, Carvalho et al. 2012, Fang et al. 2012, Eamsobhana et al. 2013, Lee et al. 2014, Guerino et al. 2017, Lv et al. 2017a).

Generally, to identify parasites, third instar larvae are obtained from naturally infected snails, and utilized for infecting vertebrate hosts by the oral route. Upon maturation of worms and migration to their habitat, parasite recovery is performed for morphological analysis. Such process is lengthy, and the vertebrate host must be sacrificed for worm recovery. ITS sequences, when used as biomarkers, allow the differentiation of larvae and adult worms in *Angiostrongylus costaricensis*, *A. cantonensis*, and *Angiostrongylus vasorum*. This in turn contributes to reducing morphological identification that generally requires ~40 days (Caldeira et al. 2003).

According to ITS sequences, Lv et al. (2017a) analyzed the genetics and evolution of internal species in *A. cantonensis* in China. Singleton variable sites make up 67.3% and 70.0% of all variable sites in ITS1 and ITS2, respectively; these rates are markedly elevated than found for mitochondrial (mt) genes. The study also ruled out the possibility of group II

constituting a distinct species, because there are no major differences in ITS sequences between this and other clades. Also striking intragenomic or within-individual heterogeneity was detected in ITS sequences and microsatellites, conflicting with the notion of concerted evolution (Ganley et al. 2007), but corroborating previously reported findings (Maslunka et al. 2014, Zhao et al. 2015).

Different local strains have been distinguished based on this gene. The DNA sequences of ITS-1 in *A. cantonensis* in China, United States, and Brazil have an intraspecific sequence variation of about 0.1–1.0%, whereas ITS-2 varies by 0.0–1.3% in isolates collected in China and Philippines (Eamsobhana et al. 2013). Moreover, the ITS gene of Hualien (H) *A. cantonensis* was shown to differ from those of counterparts in Pingtung (P) and three additional locations by 19%. H *A. cantonensis* was less infective and fecund in rats, with impaired development, causing a milder disease and reduced death rate in mice, compared with the P strain (Lee et al. 2014).

Furthermore, ITS has been utilized for the identification of the species of L3 stage *Angiostrongylus* larvae in snails (Foronda et al. 2010, Qvarnstrom et al. 2010, Liu et al. 2011, Carvalho et al. 2012, Guerino et al. 2017), as well as L1 stage larvae in mouse stool (Fang et al. 2012).

### The SSU rRNA gene

18S rRNA, as a rRNA, is considered a basic cell component in eukaryotes. 18S rRNA sequences are broadly utilized for reconstructing the evolutionary history of organisms (Liu et al. 2015, Cooper et al. 2016, Harl et al. 2019). Because it slowly evolves, it can be utilized for reconstructing ancient divergences (Meyer et al. 2010, Sulaiman et al. 2014, Zhang et al. 2017b, Vitari et al. 2019, Kenmotsu et al. 2020).

The identification of nematodes in the environment remains important for assessing their potential risks to humans. Most nematodes that are present in intermediate hosts are juvenile, making it difficult to differentiate species based on morphology. Nuclear 18S (SSU) rDNAs are used to differentiate multiple *Angiostrongylus* taxa and to conduct systematic and phylogenetic analyses of relationships among *Angiostrongylus* species and other metastrongyloid groups (Floyd et al. 2002, Fontanilla and Wade 2008, Tokiwa et al. 2012, Eamsobhana et al. 2015). It was suggested that two sequences are from a single species with 99.5–100% similarity for the first 450 bp of the SSU rDNA (Floyd et al. 2002).

Importantly, 18S (SSU) rDNAs precisely segregated the five *Angiostrongylus* spp., confirming that *A. cantonensis* and *Angiostrongylus malaysiensis* on the one hand, and *A. costaricensis* and *Angiostrongylus dujardini* on the other hand, are closely related (Eamsobhana et al. 2015). Thus, the identity of a species can be determined by matching the sequence to an accessible sequence database such as GenBank. It is important to evaluate the role of *Achatina fulica* in the spread of *A. cantonensis* and potentially other nematodes in Philippines, especially in the densely populated area of Metro Manila. This allows proper assessment of the epidemiology of the associated diseases.

Phylogenetic relationships of *A. cantonensis* from distinct geographical areas revealed multiple lineages. Two different SSU genotypes (G1 and G2) were detected in 17 individual *A. cantonensis* strains from 17 geographical areas in Japan, Mainland China, Taiwan, and Thailand (Tokiwa et al. 2012).

### The mt genome

Mitochondrial DNA (mtDNA or mDNA) is found in mitochondria, which produce adenosine triphosphate from food-associated energy for cell utilization (Iborra et al. 2004, Yan et al. 2019). Unlike nuclear DNA, mtDNA does not generally vary from parent to offspring. Since mtDNA copies can recombine (Zhang et al. 2017b), mtDNA mutates more than nuclear DNA in animals (Tatarenkov and Avise 2007, Zhang et al. 2017c). Therefore, mtDNA acts as a potent biomarker in tracking ancestry through females (matrilineage) for multiple millennia.

### Complete mt genome

The full mt genome of the rodent intra-arterial nematode *A. cantonensis* was reported in 2012 (Lv et al. 2012). It is 13,497 bp long, including 2 proteins, 22 transfer ribonucleic acid (tRNAs), and 2 rRNAs, all on the same strand. Meanwhile, *A. costaricensis*'s full mt genomes are 13,585 bp long, with 81.6% nucleotide similarity between both mt genomes, ranging between 77.7% and 87.1% in individual gene pairs. In 2015, Yong et al. (2015b) revealed the genetic diversity of *A. cantonensis* by analyzing the complete mt genome. The mt genomes of Thai (13,519 bp) and Chinese (13,497–13,502 bp) isolates differed in length, as well as five protein-coding genes, including *atp6*, *cox1*, *cox2*, *cob*, and *nad2*. Stop codons for four genes (*atp6*, *cob*, *nad2*, and *nad6*) differed between the Thai and Chinese isolates as well.

In addition, the Thai and Chinese isolates had 4 and 3 incomplete T stop codons, respectively. The Thai isolate had longer control (258 bp) in comparison with the Chinese (230–236 bp) and Taiwanese (237 bp) counterparts. The intergenic sequence between *nad4* and *cox1* in the Thai isolate was short 2 bp (indels) at 5' in comparison with the Chinese counterpart, also differing at 7 additional sites. In the Thai, Chinese, and Taiwanese isolates, 18, 17, and 16 tRNAs lacked the whole TΨC-arm, respectively.

A phylogenetic analysis of 36 mt genes, 12 polycomb group (PCGs), 2 rRNA genes, 22 tRNA genes, and control region revealed the Chinese isolate had closer genetic affinity with the Taiwanese counterpart in comparison with the Thai isolate. On the basis of 36 mt genes, the interisolate genetic distance varied from  $p=3.2\%$  between the Chinese and Taiwanese isolates to  $p=11.6\%$  between the Thai and Chinese isolates. Hence, the mt genome could help in population, phylogenetic, and phylogeographical analyses.

Furthermore, next-generation sequencing analysis of *A. cantonensis* may help better understand rat lung worm invasion and parasite detection. Novel *A. cantonensis* mt genomes in Sydney (Australia), Hawaii (United States), Canary Islands (Spain), and Fatu Hiva (French Polynesia) were comparatively assessed with those outside of Southeast Asia, showing low genetic diversity (0.02–1.03%). Both *cox1* and *cox2* are preferred biomarkers for detecting *A. cantonensis* haplotypes. Low-coverage whole-genome sequencing enables standardized identification of *A. cantonensis* (laboratory and field strains) (Červená et al. 2019).

Of the 13 *Angiostrongylus* species (Spratt 2015), *A. cantonensis* (Lv et al. 2012), *A. costaricensis* Brazil taxon (Lv et al. 2012), *A. costaricensis* Costa Rica taxon (Yong et al. 2015b), *A. mackerrasae* (Aghazadeh et al. 2015b, Valentyne et al. 2020), *A. malaysiensis* (Yong et al. 2016), and

*A. vasorum* (Gasser et al. 2012) have complete mt genomes available. The mitogenome is therefore useful for phylogenetic and systematic analyses of *Angiostrongylus* lung worms and other Metastrongyloid nematodes.

### Cytochrome c oxidase subunit I gene

Cytochrome c oxidase subunit I (COI) represents the major cytochrome c oxidase subunit. COI belongs to respiratory complex IV, which also includes cytochrome c oxidase 2 and 3. Complex IV represents the third and final enzyme of the electron transport chain during the process of mt oxidative phosphorylation. Cytochrome c oxidase constitutes a major enzyme involved in aerobic metabolism. Heme-copper oxidases (proton pumps) are terminal, energy-transfer enzymes of prokaryotic and eukaryotic respiratory chains (Kosakyan et al. 2012).

The *A. cantonensis* nematode was first described in China. Lee et al. (2014) analyzed the *cox1* gene of Taiwanese strains, where sequences of the H strain differed from those of the P and other three strains by 19% and 11%, respectively. A study conducted by Lv et al. (2017) reported the *cox1* genes of totally 130 *A. cantonensis* samples from 32 distinct sites covering the entire endemic area of angiostrongyliasis in China, which were successfully sequenced. There were 39 *cox1* haplotypes (Hd, 0.8114; Pi, 0.0284). Among the 843 sites of the complete nicotinamide adenine dinucleotide dehydrogenase subunit 1 (*nad1*) gene and 1577 sites of *cox1*, there were 171 variable sites per gene. Parsimony-informative sites were 94.7% in *cox1* (162/171).

No substitution saturation was detected in mt *cox1* gene in *A. cantonensis* (Wang and Lv 2014). Reports have examined the phylogenetic features of this nematode in other Asian countries, such as Japan, Lao PDR, Cambodia, Myanmar, and Thailand, besides China. Haplotypes *ac1* (Tokyo, Chiba, Kanagawa, Amamioshima Island, and Taichung), *ac2* (Ishikawa, Shenzhen, and Lianjiang), *ac5* (Okinawa and Ogasawara), and *ac7* (Miyagi, Aichi, and Kanagawa) were the commonest. The above regions are separated from one another by high mountains or seas (Tokiwa et al. 2012). Differences in the *cox1* genes of *A. cantonensis* and *A. malaysiensis* have also been identified. Larvae were obtained from land snails in Lao PDR, Cambodia, Myanmar, and Thailand.

The regional distributions of both species showed a broad overlap. Publicly available databases were queried to infer phylogenetic relationships in *Angiostrongylus* species. COI showed the highest intraspecific variation in study regions (five and four *A. cantonensis* and *A. malaysiensis* haplotypes, respectively) (Rodpai et al. 2016). This parasite was also reported in the Americas. *A. cantonensis* isolates from distinct geographical Brazilian regions were shown to be genetically variable, based on COI gene sequences. Phylogenetic analysis revealed that *A. cantonensis* obtained in Pará (Brazil) and the Japanese strain are similar. Also, the above specimens once had a single haplotype with high bootstrap support with the Rio de Janeiro's isolate (Moreira et al. 2013).

The Brazilian haplotype *ac5* and Japanese isolates cluster, whereas the Brazilian haplotype *ac8* (Rio de Janeiro, São Paulo, Pará, and Pernambuco) constitute a different clade. The Brazilian haplotype *ac9* has close relationships to the

Chinese and Japanese ac6 and ac7, respectively (Monte et al. 2012). There were also 7 Thai haplotypes, including AC4, AC10, AC11, AC13, AC14, and AC15 (Apichat et al. 2016, Eamsobhana et al. 2017). *A. cantonensis*, *A. costaricensis*, and *A. vasorum* COI sequences in GenBank were utilized for comparison. Two major clades were included: *A. cantonensis* and *A. malaysiensis* on one hand, and *A. costaricensis* and *A. vasorum* on the other hand.

The three *A. cantonensis* stains obtained in different geographical regions constituted a clade showing low-to-high bootstrap values, encompassing two subclades, namely the Chinese and Hawaiian isolates on one hand, and the Thai strain on the other hand. In the other main clade, European *A. vasorum* strains differ from Brazilian isolates. For *A. costaricensis*, the Costa Rican and Brazilian isolates differed, with an uncorrected (p) distance of 11.39%, suggesting the potential occurrence of cryptic species (Eamsobhana et al. 2010).

#### Nicotinamide adenine dinucleotide dehydrogenase subunit 1 (*nad1*) gene

Nad1 belongs to the NAD(P)H dehydrogenase (quinone) family that is translated into a cytoplasmic 2-electron reductase, which homodimerizes and catalyzes quinone reduction into hydroquinones and other redox substances. It preferentially transforms short-chain acceptor quinones, including ubiquinone, benzoquinone, juglone, and duroquinone (Sparla et al. 1996). Nad1 has a critical paralog, that is, NAD(P)H dehydrogenase quinone 2. This protein is cytosolic (Ross et al. 2000).

In China, the *nad1* genes of 130 *A. cantonensis* strains obtained in 32 distinct sites covering the entire reported endemic region for angiostrongyliasis were accumulated. There was significant variation in *nad1* sequences, with 75 haplotypes (Hd: 0.9260; Pi: 0.0314). Of the 843 *nad1* sites, 171 variable ones were identified. Parsimony-informative sites were most frequent, representing 60.2% (103/171). However, there were multiple singlets in *nad1*. In addition, three compound parsimony-informative sites (three or four nucleotide types concurrently found more than twice) were identified. The nicotinamide adenine dinucleotide 1 protein consisted of 281 amino acids, with 17 variable sites. The corresponding synonymous variations were 73. The rates of singleton variable sites markedly differed between nuclear and mt genes, as well as among mt genes (Lv et al. 2017).

#### Cytochrome b gene

Cytochrome b (Cytb), a mt protein in eukaryotes, is involved in the electron transport chain as the major subunit of transmembrane cytochrome bc1 and b6f complexes (Esposti et al. 1993). Cytb is an important constituent of respiratory chain complex III (bc1 complex or ubiquinol-cytochrome c reductase) found in mitochondria of eukaryotes as well as aerobic prokaryotes. Due to the sequence variability of Cytb, it is frequently utilized to determine phylogenetic relationships among organisms, notably at the family and genus levels. Comparative analyses of cytochrome b produced novel classification patterns and helped assign new species to a genus and provided insights into evolutionary relationships (Castresana 2001).

In phylogeographical researches, the Cytb gene has been used in studying *A. cantonensis* in southern China (including Taiwan) and some Southeast Asian regions (Thailand). Peng et al. (2017) analyzed 520 distinct *A. cantonensis* strains from 13 localities, which were sequenced and grouped into 42 haplotypes. Genetic variations among populations were prominent, but demographic expansion was not observed through neutrality assays or mismatch distribution analysis, indicating low local gene flow in various sample collection sites. Two distinct *A. cantonensis* lineages were found in Taiwan, indicating that the parasite came to the island from many locations. Hekou (China) and Laos populations had the highest genetic variations, as reflected by genetic diversity parameters and analysis of molecular variation.

The above findings jointly indicate that the region surrounding Thailand or Hekou in Yunnan province, China, might be the most plausible origin of *A. cantonensis*. In 2015, Dusitsittipon et al. (2015) and Yong et al. (2015a) used this gene to determine *A. cantonensis*' genetic structure in Thailand. Phylogenetic assessment revealed two major clades in Thailand, which comprised four subclades or clusters. Genetic variation of gene flow and geographical location were not overtly related. Such low genetic variation and geographical distribution of *A. cantonensis* in various locations indicated a founder effect, likely due to many independent origins, suggesting haplotype migration from endemic areas through anthropogenic factors (Dusitsittipon et al. 2015).

Moreover, Cytb haplotypes from different Thai regions have been established, with some apparently restricted to particular geographical areas. Incomplete Cytb DNA sequences could allow an unequivocal separation of *A. cantonensis* strains collected in Thailand, China, and Hawaii, as well as the congeners *A. malaysiensis*, *A. costaricensis*, and *A. vasorum*, with *A. malaysiensis* and *A. costaricensis* grouped with *A. cantonensis* and *A. vasorum*, respectively (Yong et al. 2015a). However, CYTB mtDNA contains pseudogenes (Dusitsittipon et al. 2018).

#### Molecular Identification and Diagnosis of *A. cantonensis*

To address the limitations of phenotypic assays, genotypic strategies have been utilized for identifying and differentiating *A. cantonensis* (Ash 1970, Lindo et al. 2002, Cowie 2013, Qvarnstrom et al. 2013, Heather et al. 2017). Beside the development of conventional PCR detection in *A. cantonensis* (Eamsobhana et al. 2010, Lv et al. 2012, Lee et al. 2014, Yong et al. 2015a), several PCR-based approaches, including PCR-linked restriction fragment length polymorphism, PCR-random amplified polymorphic DNA (PCR-RAPD), multiplex PCR, quantitative PCR (qPCR), and loop-mediated isothermal amplification (LAMP) assays, are currently available for the accurate identification of *A. cantonensis* (Caldeira et al. 2003, Wei et al. 2010, Chen et al. 2011, Liu et al. 2011, Cui et al. 2012, Jarvi et al. 2012, 2015, Thaenkhom et al. 2012, Qvarnstrom et al. 2016) (Table 1).

For example, according to the differentiation of COI and ITS2 regions of mtDNA and rDNA, respectively, in *A. cantonensis*, *A. costaricensis*, and *A. vasorum*, and subsequent fragment digestion by *RsaI*, *HapII*, *AluI*, *HaeIII*, *DdeI*, and *ClaI*, species can be differentiated and identified; *RsaI* and

TABLE 1. MOLECULAR ASSAYS FOR DETECTING AND/OR GENETICALLY DIFFERENTIATING *ANGIOSTRONGYLUS CANTONENSIS*

Molecular approaches	Source of samples					References
	Definitive host (rat)			Accidental host (human)		
	Intermediate host (snails and/or slugs)	Lung	Feces	Blood and peripheral tissues	Cerebrospinal fluid	
Conventional PCR	✓					ITS1
	✓					ITS2
		✓				ITS
	✓	✓		✓		ITS
						SSU rRNA
Multiplex PCR	✓					SSU rRNA
		✓				complete mitochondrial genome
						COI
PCR-RFLP	✓					COI
	✓	✓				COI
		✓				nad1
		✓				Cyt b
PCR-RAPD	✓					SSU rRNA
	✓					COI and ITS2
	✓					ITS2
	✓					Genomic DNA
				✓		ITS-2
Quantitative PCR	✓					ITS-1
					✓	ITS-1
						ITS-1
					✓	ITS-1
						ITS-1
LAMP	✓					18S rRNA
	✓				✓	ITS-1
	✓					18S rRNA and ITS-1
	✓					ITS1
	✓					ITS1
RAA	✓					ITS1
	✓					
	✓					
	✓					
	✓					
RPA-EXO	✓					
	✓					
	✓					
	✓					
	✓					
RPA-LFA	✓					
	✓					
	✓					
	✓					
	✓					

COI, cytochrome c oxidase subunit I; ITS, internal transcribed spacer; LAMP, loop-mediated isothermal amplification; RAA, recombinase-aided isothermal amplification; PCR-RAPD, PCR-random amplified polymorphic DNA; PCR-RFLP, PCR-linked restriction fragment length polymorphism; RPA-EXO, recombinase polymerase amplification Exo; RPA-LFA, recombinase polymerase amplification lateral flow assay; rRNA, ribosomal RNA; SSU, small subunit.

*Clal* were most potent in discriminating these two genes (Caldeira et al. 2003).

RAPD markers are 10-nt DNA fragments obtained by amplifying random segments of genomic DNA using a single primer of a random nucleotide sequence, which can distinguish genetically distinct individuals, although not reproducibly. They are utilized for assessing genetic diversity for an individual. In a study conducted by Thaenkham et al. (2012), *A. cantonensis* was collected in 8 Thai geographical areas. The results showed high genetic diversity and low gene flow among populations. Thai *A. cantonensis* was shown to comprise two groups with genetic differentiation. However, genotypic changes and haplotype relationships deserve further investigation utilizing additional markers.

qPCR could be applied in multiple research areas such as genotyping, and gene expression, microRNA, genetic variation and protein analyses. qPCR has higher efficiency and sensitivity compared with microscopy, providing a novel tool for quantifying *A. cantonensis* larvae in the intermediate host snails (Qvarnstrom et al. 2010) and potentially other sources (e.g., semi-slugs) as well (Jarvi et al. 2012).

*A. cantonensis* DNA could also be detected by qPCR in peripheral blood at various times during rat lungworm infection (Jarvi et al. 2015), as well as in human CSF specimens (Qvarnstrom et al. 2016, Sears et al. 2020). The AcanR3990 qPCR instrument has a limit of detection (LOD) of 1 fg (DNA equivalent of 1/100,000 dilution of 1 third instar larva), with no cross-reaction with CSF specimens from individuals with definite neurocysticercosis, toxocariasis, gnathostomiasis, and baylisascariasis (Sears et al. 2020). LAMP represents an isothermal single-tube method utilized for DNA amplification. Unlike PCR that uses alternating temperature steps and/or cycles, LAMP is operated at constant temperature, with no need of a thermal cycler (Notomi et al. 2000).

LAMP represents a suitable technique for routine *A. cantonensis* detection within the intermediate hosts, that is, *Pomacea canaliculata* and *A. fulica* due to its simplicity, sensitivity, and specificity. It has great prospect for monitoring *A. cantonensis* in endemic regions. Reactions amplified *A. cantonensis*' 18S rRNA and ITS-1 sequences, with high sensitivity, that is, only 1 fg of DNA detected in *P. canaliculata* specimens (Chen et al. 2011) and 10 times higher sensitivity versus conventional PCR in *A. fulica*. Furthermore, no cross-reactivity with other parasitic organism was detected (Liu et al. 2011).

The recombinase-aided isothermal amplification (RAA) assay was also applied in the nucleic acid detection of *A. cantonensis*. *A. cantonensis*' ITS1 gene served as the target sequence for detection. RAA was performed for successful nucleic acid detection in *A. cantonensis*, achieving real-time and specific amplification within 20 min at 37°C. The lowest LODs of the fluorescent RAA assay were 10 copies/ $\mu$ L and 100 pg/ $\mu$ L of recombinant plasmid and genomic DNA, respectively. This technique could not detect the genomic DNAs of *A. cantonensis*, *Schistosoma mansoni*, *Ascaris lumbricoides*, *Clonorchis sinensis*, *Echinococcus granulosus*, *Ancylostoma duodenale*, *P. canaliculata*, and *Biomphalaria straminea* in snail tissues (Zhang et al. 2020).

Currently, the recombinase polymerase amplification Exo (RPA-EXO) assay and recombinase polymerase amplifica-

tion lateral flow assay (RPA-LFA) with routine qPCR have been developed. These three methods have been applied to assess 35 slugs in Hawaii for *A. cantonensis* DNA, with consistent, discordant, and equivocal results in 23/35 (65.7%), 7/35 (20%, <0.01 larvae/mg tissue), and 5/35 (14.3%), respectively. They consistently detected 50–100 copies/ $\mu$ L versus ~13 copies/ $\mu$ L for qPCR. RPA-EXO could detect 25 copies/ $\mu$ L, whereas RPA-LFA only amplified consistently above 50 copies/ $\mu$ L. Although both RPA-EXO assay and RPA-LFA appear less sensitive compared with qPCR at low DNA levels, they are more convenient than qPCR (Jarvi et al. 2020).

### Application of Molecular Detection of *A. cantonensis* Infection

In *A. cantonensis*, first instar larval worms are expelled in the feces of rats (definitive host) (Alicata et al. 1965). The infected feces is then ingested by many species of snails and slugs (intermediate hosts); however, larvae might also enter the snail through the body wall or through the respiratory pore (Alicata 1991). These larvae develop into third instar larvae in snails, remaining at this stage until the snail is eaten or dies. After snails with third instar larvae are eaten by rats, the larvae move to the small intestine through the rat gut. They next penetrate the intestinal wall and enter the circulation, with some eventually entering the CNS and reaching the brain. Humans (incidental host) contract the infection mostly like rats through deliberate or accidental ingestion of infected snails or slugs (Lindo et al. 2002).

For molecular detection of *A. cantonensis* infection, blood and tissue specimens from rats (Jarvi et al. 2015), larvae in the stool of rats (Fang et al. 2012), L3 stage larvae in snails or slugs (Zhang et al. 2006, Qvarnstrom et al. 2007, 2010, Wei et al. 2010, Carvalho et al. 2012, Chan et al. 2015, Guerino et al. 2017), CSF samples from humans (Eamsobhana et al. 2013), and abdominal angiostrongyliasis in humans (da Silva et al. 2003) have been tested.

### Conclusions

In summary, most of the current knowledge of *A. cantonensis* epidemiology has stemmed from multiple observational and morphological analyses. Nevertheless, molecular epidemiologic studies reported in recent years have provided important insights into *Angiostrongylus* taxonomy and evolutionary biology. Importantly, these molecular detection methods have been widely used in diagnosing angiostrongyliasis. The application of such molecular methods, coupled with classical epidemiologic approaches, imparts greater resolution and accuracy in describing the epidemiology of *Angiostrongylus*.

### Author Disclosure Statement

There were no competing financial interests.

### Funding Information

This study was supported by the Natural Science Foundation of Shanghai (Grant No. 18ZR1443500), the General Program Shanghai Municipal Commission of Health and Family Planning of China (Grant Nos. 201840286 and 201640278), the Youth Science Foundation of Chinese

Center for Disease Control and Prevention (Grant No. 2018A105), the Fifth Round of Three-Year Public Health Action Plan of Shanghai (Grant No. GWV-10.1-XK13), the Open project of Key Laboratory of Tropical Disease Control (Sun Yat-Sen University), Ministry of Education (Grant No. 2019kfkt02), NHC Key Laboratory of Echinococcosis Prevention and Control (Project No. 2020WZK2002), and Shenzhen San-Ming Project for prevention and research on vector-borne diseases (Grant No. SZSM201611064).

## References

- Aghazadeh M, Harvie MC, Owen HC, Veríssimo C, et al. Comparative pathogenesis of eosinophilic meningitis caused by *Angiostrongylus mackerrasae* and *Angiostrongylus cantonensis* in murine and guinea pig models of human infection. *Parasitology* 2016; 143:1243–1251.
- Aghazadeh M, Jones MK, Aland KV, Reid SA, et al. Emergence of neural angiostrongyliasis in eastern Australia. *Vector Borne Zoonotic Dis* 2015a; 15:184–190.
- Aghazadeh M, Traub RJ, Mohandas N, Aland KV, et al. The mitochondrial genome of *Angiostrongylus mackerrasae* as a basis for molecular, epidemiological and population genetic studies. *Parasit Vectors* 2015b; 8:473.
- Alicata JE. Biology and distribution of the rat lungworm, *Angiostrongylus cantonensis*, and its relationship to eosinophilic meningoencephalitis and other neurological disorders of man and animals. *Adv Parasitol* 1965; 3:223–248.
- Alicata JE. The discovery of *Angiostrongylus cantonensis* as a cause of human eosinophilic meningitis. *Parasitol Today* 1991; 7:151–153.
- Alvarez I, Wendel JF. Ribosomal ITS sequences and plant phylogenetic inference. *Mol Phylogenet Evol* 2003; 29:417–434.
- Ansdell V, Wattanagoon Y. *Angiostrongylus cantonensis* in travelers: Clinical manifestations, diagnosis, and treatment. *Curr Opin Infect Dis* 2018; 31:399–408.
- Apichat V, Narongrit S, Jitranuch T, Anucha W, et al. Phylogeny of *Angiostrongylus cantonensis* in Thailand based on cytochrome C Oxidase subunit I gene sequence. *Southeast Asian J Trop Med Public Health* 2016; 47:377–386.
- Ash LR. Diagnostic morphology of the third-stage larvae of *Angiostrongylus cantonensis*, *Angiostrongylus vasorum*, *Aelurostrongylus abstrusus*, and *Anafilaroides rostratus* (Nematoda: Metastrongyloidea). *J Parasitol* 1970; 56:249–253.
- Barratt J, Chan D, Sandaradura I, Malik R, et al. *Angiostrongylus cantonensis*: A review of its distribution, molecular biology and clinical significance as a human pathogen. *Parasitology* 2016; 143:1087–1118.
- Caldeira RL, Carvalho OS, Mendonça CL, Graeff-Teixeira C, et al. Molecular differentiation of *Angiostrongylus costaricensis*, *A. cantonensis*, and *A. vasorum* by polymerase chain reaction-restriction fragment length polymorphism. *Mem Inst Oswaldo Cruz* 2003; 98:1039–1043.
- Carvalho Odo S, Scholte RG, Mendonça CL, et al. *Angiostrongylus cantonensis* (Nematode: Metastrongyloidea) in molluscs from harbour areas in Brazil. *Mem Inst Oswaldo Cruz* 2012; 107:740–746.
- Castresana J. Cytochrome b phylogeny and the taxonomy of great apes and mammals. *Mol Biol Evol* 2001; 18:465–471.
- Červená B, Modrý D, Feclová B, Hrazdilová K, et al. Low diversity of *Angiostrongylus cantonensis* complete mitochondrial DNA sequences from Australia, Hawaii, French Polynesia and the Canary Islands revealed using whole genome next-generation sequencing. *Parasit Vectors* 2019; 12:241.
- Chan D, Barratt J, Roberts T, Lee R, et al. The Prevalence of *Angiostrongylus cantonensis/mackerrasae* Complex in Molluscs from the Sydney Region. *PLoS One* 2015; 10:e0128128.
- Chen MX, Chen JX, Chen SH, Huang DN, et al. Development of lateral flow immunoassay for antigen detection in human *Angiostrongylus cantonensis* infection. *Korean J Parasitol* 2016; 54:375–380.
- Chen R, Tong Q, Zhang Y, Lou D, et al. Loop-mediated isothermal amplification: Rapid detection of *Angiostrongylus cantonensis* infection in *Pomacea canaliculata*. *Parasit Vectors* 2011; 4:204.
- Chye SM, Lin SR, Chen YL, Chung LY, et al. Immuno-PCR for detection of antigen to *Angiostrongylus cantonensis* circulating fifth-stage worms. *Clin Chemistry* 2004; 50:51–57.
- Coleman AW. Nuclear rRNA transcript processing versus internal transcribed spacer secondary structure. *Trends Genet* 2015; 31:157–163.
- Cooper MK, Phalen DN, Donahoe SL, Rose K, et al. The utility of diversity profiling using Illumina 18S rRNA gene amplicon deep sequencing to detect and discriminate *Toxoplasma gondii* among the cyst-forming coccidia. *Vet Parasitol* 2016; 216:38–45.
- Cowie RH. Biology, systematics, life cycle, and distribution of *Angiostrongylus cantonensis*, the cause of rat lungworm disease. *Hawaii J Med Public Health* 2013; 72(Suppl. 2):6–9.
- Cowie RH. *Angiostrongylus cantonensis*: Agent of a sometimes fatal globally emerging infectious disease (rat lungworm disease). *ACS Chem Neurosci* 2017; 8:2102–2104.
- Cross JH, Chen ER. Angiostrongyliasis. In: Murrel KD, Fried B, eds. *Food Borne Parasitic Zoonoses*. New York: Springer, 2007:263–290.
- Cui LY, Zhang XX, Li KH, Yang YM. Application of PCR and PCR-derived technologies in the detection of *Angiostrongylus cantonensis*. *Chin J Parasitol Parasit Dis* 2012; 30:321–324. (In Chinese)
- da Silva AC, Graeff-Teixeira C, Zaha A. Diagnosis of abdominal angiostrongyliasis by PCR from sera of patients. *Rev Inst Med Trop Sao Paulo* 2003; 45:295–297.
- Dusitsittipon S, Criscione CD, Morand S, Komalamisra C, et al. Hurdles in the evolutionary epidemiology of *Angiostrongylus cantonensis*: Pseudogenes, incongruence between taxonomy and DNA sequence variants, and cryptic lineages. *Evol Appl* 2018; 11:1257–1269.
- Dusitsittipon S, Thienkham U, Wathanakulpanich D, Adisakwattana P, et al. Genetic differences in the rat lungworm, *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae), in Thailand. *J Helminthol* 2015; 89:545–551.
- Eamsobhana P, Lim PE, Solano G, Zhang H, et al. Molecular differentiation of *Angiostrongylus taxa* (Nematoda: Angiostrongylidae) by cytochrome c oxidase subunit I (COI) gene sequences. *Acta Trop* 2010; 116:152–156.
- Eamsobhana P, Lim PE, Yong HS. Genetic diversity of the rat lungworm, *Angiostrongylus cantonensis*, the major cause of eosinophilic meningitis. *Hawaii J Med Public Health* 2013; 72(Suppl. 2):15–17.
- Eamsobhana P, Lim PE, Yong HS. Phylogenetics and systematics of *Angiostrongylus* lungworms and related taxa (Nematoda: Metastrongyloidea) inferred from the nuclear small subunit (SSU) ribosomal DNA sequences. *J Helminthol* 2015; 89:317–325.
- Eamsobhana P, Song SL, Yong HS, Prasartvit A, Boonyong S, et al. Cytochrome c oxidase subunit I haplotype diversity of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae). *Acta Trop* 2017; 171:141–145.

- Eamsobhana P, Tungtrongchitr A, Wanachiwanawin D, Yong HS. Immunochromatographic test for rapid serological diagnosis of human angiostrongyliasis. *Int J Infect Dis* 2018; 73: 69–71.
- Esposti MD, De Vries S, Crimi M, Ghelli A, et al. Mitochondrial cytochrome b: Evolution and structure of the protein. *Biochim Biophys Acta* 1993; 1143:243–271.
- Fang W, Wang J, Liu J, Xu C, et al. PCR assay for the cell-free copro-DNA detection of *Angiostrongylus cantonensis* in rat faeces. *Vet Parasitol* 2012; 183:299–304.
- Floyd R, Abebe E, Papert A, Blaxter M. Molecular barcodes for soil nematode identification. *Mol Ecol* 2002; 11:839–850.
- Fontanilla IK, Wade CM. The small subunit (SSU) ribosomal (r) RNA gene as a genetic marker for identifying infective 3 rdjuvenile stage *Angiostrongylus cantonensis*. *Acta Trop* 2008; 105:181–186.
- Foronda P, López-González M, Miquel J, Torres J, et al. Finding of *Parastrongylus cantonensis* (Chen, 1935) in *Rattus rattus* in Tenerife, Canary Islands (Spain). *Acta Trop* 2010; 114:123–127.
- Ganley AR, Kobayashi T. Highly efficient concerted evolution in the ribosomal DNA repeats: Total rDNA repeat variation revealed by wholegenome shotgun sequence data. *Genome Res* 2007; 17:184–191.
- Gasser RB, Jabbar A, Mohandas N, Schnyder M, et al. Mitochondrial genome of *Angiostrongylus vasorum*: Comparison with congeners and implications for studying the population genetics and epidemiology of this parasite. *Infect Genet Evol* 2012; 12:1884–1891.
- Graeff-Teixeira C, da Silva AC, Yoshimura K. Update on eosinophilic meningoencephalitis and its clinical relevance. *Clin Microbiol Rev* 2009; 2:322–348.
- Graeff-Teixeira C, Morassutti AL, Jones MK. Diagnosing and understanding angiostrongyliasis, a zoonotic cause of meningitis. *ACS Chem Neurosci* 2018; 9:393–394.
- Guerino LR, Pecora IL, Miranda MS, Aguiar-Silva C, et al. Prevalence and distribution of *Angiostrongylus cantonensis* (Nematoda, Angiostrongylidae) in *Achatina fulica* (Mollusca, Gastropoda) in Baixada Santista, São Paulo, Brazil. *Rev Soc Bras Med Trop* 2017; 50:92–98.
- Harl J, Himmel T, Valkiūnas G, Weissenböck H. The nuclear 18S ribosomal DNAs of avian haemosporidian parasites. *Malar J* 2019; 18:305.
- Hillis DM, Dixon MT. Ribosomal DNA: Molecular evolution and phylogenetic inference. *Q Rev Biol* 1991; 66:411–453.
- Himsworth CG, Parsons KL, Jardine C, Patrick DM. Rats, cities, people, and pathogens: A systematic review and narrative synthesis of literature regarding the ecology of rat-associated zoonoses in urban centers. *Vector Borne Zoonotic Dis* 2013; 13:349–359.
- Huang D, Huang Y, Tang Y, Zhang Q, et al. Survey of *Angiostrongylus cantonensis* infection status in host animals and populations in Shenzhen, 2016–2017. *Vector Borne Zoonotic Dis* 2019; 19:717–723.
- Iborra FJ, Kimura H, Cook PR. The functional organization of mitochondrial genomes in human cells. *BMC Biol* 2004; 2:9.
- Jarvi SI, Atkinson ES, Kaluna LM, Snook KA, et al. Development of a recombinase polymerase amplification (RPA-EXO) and lateral flow assay (RPA-LFA) based on the ITS1 gene for the detection of *Angiostrongylus cantonensis* in gastropod intermediate hosts. *Parasitology* 2020; 148:251–258.
- Jarvi SI, Farias ME, Howe K, Jacquier S, et al. Quantitative PCR estimates *Angiostrongylus cantonensis* (rat lungworm) infection levels in semi-slugs (*Parmarion martensi*). *Mol Biochem Parasitol* 2012; 185:174–176.
- Jarvi SI, Pitt WC, Farias ME, Shiels L, et al. Detection of *Angiostrongylus cantonensis* in the blood and peripheral tissues of wild Hawaiian rats (*Rattus rattus*) by a quantitative PCR (qPCR) assay. *PLoS One* 2015; 10:e0123064.
- Jin E, Ma D, Liang Y, Ji A, et al. MRI findings of eosinophilic myelomeningoencephalitis due to *Angiostrongylus cantonensis*. *Clin Radiol* 2005; 60:242–250.
- Kanpittaya J, Jitpimolmard S, Tiamkao S, Mairiang E. MR findings of eosinophilic meningoencephalitis attributed to *Angiostrongylus cantonensis*. *AJNR Am J Neuroradiol* 2000; 1:1090–1094.
- Kenmotsu H, Uchida K, Hirose Y, Eki T. Taxonomic profiling of individual nematodes isolated from copse soils using deep amplicon sequencing of four distinct regions of the 18S ribosomal RNA gene. *PLoS One* 2020; 15:e0240336.
- Kosakyan A, Heger TJ, Leander BS, Todorov M, et al. COI barcoding of Nebelid testate amoebae (Amoebozoa: Arcellinida): Extensive cryptic diversity and redefinition of the *Hyalospheniidae schultze*. *Protist* 2012; 163:415–434.
- Kuberski T, Wallace GD. Clinical manifestations of eosinophilic meningitis due to *Angiostrongylus cantonensis*. *Neurology* 1979; 29:1566–1570.
- Kwon E, Ferguson TM, Park SY, Manuzak A, et al. A severe case of *Angiostrongylus* eosinophilic meningitis with encephalitis and neurologic sequelae in Hawai'i. *Hawaii J Med Public Health* 2013; 72(Suppl. 2):41–45.
- Lafontaine DL, Tollervey D. The function and synthesis of ribosomes. *Nat Rev Mol Cell Biol* 2001; 2:514–520.
- Lee JD, Chung LY, Wang LC, Lin RJ, et al. Sequence analysis in partial genes of five isolates of *Angiostrongylus cantonensis* from Taiwan and biological comparison in infectivity and pathogenicity between two strains. *Acta Trop* 2014; 133:26–34.
- Lima MG, Augusto RC, Pinheiro J, Thiengo SC. Physiology and immunity of the invasive giant African snail, *Achatina* (Lissachatina) *fulica*, intermediate host of *Angiostrongylus cantonensis*. *Dev Comp Immunol* 2020; 105:103579.
- Lindo JF, Waugh C, Hall J, Cunningham-Myrie C, et al. Enzootic *Angiostrongylus cantonensis* in rats and snails after an outbreak of human *Eosinophilic meningitis*, Jamaica. *Emerg Infect Dis* 2002; 8:324–326.
- Liu CY, Song HQ, Zhang RL, Chen MX, et al. Specific detection of *Angiostrongylus cantonensis* in the snail *Achatina fulica* using a loop-mediated isothermal amplification (LAMP) assay. *Mol Cell Probes* 2011; 25:164–167.
- Liu J, Yu Y, Cai Z, Bartlam M, Wang Y. Comparison of ITS and 18S rDNA for estimating fungal diversity using PCR-DGGE. *World J Microbiol Biotechnol* 2015; 31:1387–1395.
- Lv S, Zhang Y, Steinmann P, Utzinger J, et al. The genetic variation of *Angiostrongylus cantonensis* in the People's Republic of China. *Infect Dis Poverty* 2017a; 6:125.
- Lv S, Zhang Y, Zhang L, Liu Q, et al. The complete mitochondrial genome of the rodent intra-arterial nematodes *Angiostrongylus cantonensis* and *Angiostrongylus costaricensis*. *Parasitol Res* 2012; 111:115–123.
- Lv S, Zhou XN, Andrews JR. *Eosinophilic meningitis* caused by *Angiostrongylus cantonensis*. *ACS Chem Neurosci* 2017b; 8: 1815–1816.
- Martin-Alonso A, Abreu-Yanes E, Feliu C, Mas-Coma S, et al. Intermediate hosts of *Angiostrongylus cantonensis* in Tenerife, Spain. *PLoS One* 2015; 10:e0120686.
- Maslunka C, Gifford B, Tucci J, Gurtler V, et al. Insertions or deletions (Indels) in the rrn 16S-23S rRNA gene inter-



- nal transcribed spacer region (ITS) compromise the typing and identification of strains within the *Acinetobacter calcoaceticus-baumannii* (Acb) complex and closely related members. *PLoS One* 2014; 9:e105390.
- Meyer A, Todt C, Mikkelsen NT, Lieb B. Fast evolving 18S rRNA sequences from Solenogastres (Mollusca) resist standard PCR amplification and give new insights into mollusk substitution rate heterogeneity. *BMC Evol Biol* 2010; 10:70.
- Monte TC, Simões RO, Oliveira AP, Novaes CF, et al. Phylogenetic relationship of the Brazilian isolates of the rat lungworm *Angiostrongylus cantonensis* (Nematoda: Metastrongylidae) employing mitochondrial COI gene sequence data. *Parasit Vectors* 2012; 5:248.
- Moreira VL, Giese EG, Melo FT, Simões RO, et al. Endemic angiostrongyliasis in the Brazilian Amazon: Natural parasitism of *Angiostrongylus cantonensis* in *Rattus rattus* and *R. norvegicus*, and sympatric giant African land snails, *Achatina fulica*. *Acta Trop* 2013; 125:90–97.
- Notomi T, Okayama H, Masubuchi H, Yonekawa T, et al. Loop-mediated isothermal amplification of DNA. *Nucleic Acids Res* 2000; 28:E63.
- Peng J, He ZP, Zhang S, Lun ZR, et al. Phylogeography of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae) in southern China and some surrounding areas. *PLoS Negl Trop Dis* 2017; 11:e0005776.
- Prociv P, Spratt DM, Carlisle MS. Neuro-angiostrongyliasis: Unresolved issues. *Int J Parasitol* 2000; 30:1295–1303.
- Punyagupta S, Bunnag T, Juttijudata P. Eosinophilic meningitis in Thailand. Clinical and epidemiological characteristics of 162 patients with myeloencephalitis probably caused by *Gnathostoma spinigerum*. *J Neurol Sci* 1990; 96:241–256.
- Qvarnstrom Y, Bishop HS, da Silva AJ. Detection of rat lungworm in intermediate, definitive, and paratenic hosts obtained from environmental sources. *Hawaii J Med Public Health* 2013; 72:63–69.
- Qvarnstrom Y, da Silva AC, Teem JL, Hollingsworth R, et al. Improved molecular detection of *Angiostrongylus cantonensis* in mollusks and other environmental samples with a species-specific internal transcribed spacer 1-based TaqMan assay. *Appl Environ Microbiol* 2010; 76:5287–5289.
- Qvarnstrom Y, Sullivan JJ, Bishop HS, Hollingsworth R, et al. PCR-based detection of *Angiostrongylus cantonensis* in tissue and mucus secretions from molluscan hosts. *Appl Environ Microbiol* 2007; 73:1415–1419.
- Qvarnstrom Y, Xayavong M, da Silva AC, Park SY, et al. Real-time polymerase chain reaction detection of *Angiostrongylus cantonensis* DNA in cerebrospinal fluid from patients with *Eosinophilic meningitis*. *Am J Trop Med Hyg* 2016; 94:176–181.
- Rodpai R, Intapan PM, Thanchomnang T, Sanpool O, et al. *Angiostrongylus cantonensis* and *A. malaysiensis* broadly overlap in Thailand, Lao PDR, Cambodia and Myanmar: A molecular survey of Larvae in Land Snails. *PLoS One* 2016; 11:e0161128.
- Ross D, Kepa JK, Winski SL, Beall HD, et al. NAD(P)H:quinone oxidoreductase 1 (NQO1): Chemoprotection, bioactivation, gene regulation and genetic polymorphisms. *Chem Biol Interact* 2000; 129:77–97.
- Salim D, Gerton JL. Ribosomal DNA instability and genome adaptability. *Chromosome Res* 2019; 27:73–87.
- Sawanyawisuth K, Sawanyawisuth K, Senthong V, Limpawattana P, et al. How can clinicians ensure the diagnosis of meningitic angiostrongyliasis? *Vector Borne Zoonotic Dis* 2012; 12:73–75.
- Sawanyawisuth K, Takahashi K, Hoshuyama T, Sawanyawisuth K, et al. Clinical factors predictive of encephalitis caused by *Angiostrongylus cantonensis*. *Am J Trop Med Hyg* 2009; 81:698–701.
- Sears WJ, Qvarnstrom Y, Dahlstrom E, Snook K, et al. AcanR3990 qPCR: A novel, highly sensitive, bioinformatically-informed assay to detect *Angiostrongylus cantonensis* infections. *Clin Infect Dis* 2020; 30:ciaa1791.
- Somboonpatarakun C, Intapan PM, Sadaow L, Rodpai R, et al. Development of an immunochromatographic device to detect antibodies for rapid diagnosis of human angiostrongyliasis. *Parasitology* 2020; 147:194–198.
- Somboonpatarakun C, Intapan PM, Sanpool O, Wongkham C, Maleewong W. Application of recombinant *Angiostrongylus cantonensis* galectin-2 protein for serodiagnosis of human angiostrongyliasis by immunoblotting. *Am J Trop Med Hyg* 2019; 101:851–858.
- Sparla F, Tedeschi G, Trost P. NAD(P)H:(quinone-acceptor) oxidoreductase of tobacco leaves is a flavin mononucleotide-containing flavoenzyme. *Plant Physiol* 1996; 112:249–258.
- Spratt DM. Species of *Angiostrongylus* (Nematoda: Metastrongyloidea) in wildlife: A review. *Int J Parasitol Parasites Wildl* 2015; 4:178–189.
- Stockdale Walden HD, Slapcinsky JD, et al. Geographic distribution of *Angiostrongylus cantonensis* in wild rats (*Rattus rattus*) and terrestrial snails in Florida, USA. *PLoS One* 2017; 12:e0177910.
- Sulaiman IM, Ortega Y, Simpson S, Kerdahi K. Genetic characterization of human-pathogenic *Cyclospora cayetanensis* parasites from three endemic regions at the 18S ribosomal RNA locus. *Infect Genet Evol* 2014; 22:229–234.
- Tatarenkov A, Avise JC. Rapid concerted evolution in animal mitochondrial DNA. *Proc Biol Sci* 2007; 274:1795–1798.
- Thaenkhom U, Pakdee W, Nuamtanong S, Maipanich W, et al. Population structure of *Angiostrongylus cantonensis* (Nematoda: Metastrongylidae) in Thailand based on PCR-RAPD markers. *Southeast Asian J Trop Med Public Health* 2012; 43:567–573.
- Tokiwa T, Harunari T, Tanikawa T, Komatsu N, et al. Phylogenetic relationships of rat lungworm, *Angiostrongylus cantonensis*, isolated from different geographical regions revealed widespread multiple lineages. *Parasitol Int* 2012; 61:431–436.
- Tsai HC, Liu YC, Kunin CM, Lai PH, et al. Eosinophilic meningitis caused by *Angiostrongylus cantonensis* associated with eating raw snails: Correlation of brain magnetic resonance imaging scans with clinical findings. *Am J Trop Med Hyg* 2003; 68:281–285.
- Tsai HC, Tseng YT, Yen CM, Chen ER, et al. Brain magnetic resonance imaging abnormalities in eosinophilic meningitis caused by *Angiostrongylus cantonensis* infection. *Vector Borne Zoonotic Dis* 2012; 12:161–166.
- Valentyne H, Spratt DM, Aghazadeh M, Jones MK, et al. The mitochondrial genome of *Angiostrongylus mackerrasae* is distinct from *A. cantonensis* and *A. malaysiensis*. *Parasitology* 2020; 147:681–688.
- Vitari GLV, Costa RL, Abreu APM, Peckle M, et al. Genetic diversity of *Theileria equi* from horses in different regions of Brazil based on the 18S rRNA Gene. *J Parasitol* 2019; 105:186–194.
- Wang JJ, Wu ZS, Chung LY, Lu CY, et al. Galectin-9-like from *Angiostrongylus cantonensis* young adult worms modulates eosinophil chemotaxis *in vitro*. *J Microbiol Immunol Infect* 2020; 53:604–611.

- Wang M, Lv S. Substitution saturation analysis of mitochondrial cytochrome C oxidase subunit 1 (COX1) gene of *Angiostrongylus cantonensis*. Chin J Parasitol Parasit Dis 2014; 32:205–209. (In Chinese)
- Wang QP, Wu ZD, Wei J, Owen RL, et al. Human *Angiostrongylus cantonensis*: An update. Eur J Clin Microbiol Infect Dis 2012; 31:389–395.
- Waugh CA, Lindo JF, Lorenzo-Morales J, Robinson RD. An epidemiological study of *A. cantonensis* in Jamaica subsequent to an outbreak of human cases of eosinophilic meningitis in 2000. Parasitology 2016; 143:1211–1217.
- Wei FR, Liu HX, Lv S, Hu L, et al. Multiplex PCR assay for the detection of *Angiostrongylus cantonensis* larvae in *Pomacea canaliculata*. Chin J Parasitol Parasit Dis 2010; 28:355–358. (In Chinese)
- Wilkins PP, Qvarnstrom Y, Whelen AC, Saucier C, et al. The current status of laboratory diagnosis of *Angiostrongylus cantonensis* infections in humans using serologic and molecular methods. Hawaii J Med Public Health 2013; 72: 55–57.
- Yan C, Duanmu X, Zeng L, Liu B, Song Z. Mitochondrial DNA: Distribution, mutations, and elimination. Cells 2019; 8:379.
- Yong HS, Eamsobhana P, Song SL, Prasartvit A, et al. Molecular phylogeography of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae) and genetic relationships with congeners using cytochrome b gene marker. Acta Trop 2015a; 148:66–71.
- Yong HS, Song SL, Eamsobhana P, Goh SY, et al. Complete mitochondrial genome reveals genetic diversity of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae). Acta Trop 2015b; 152:157–164.
- Yong HS, Song SL, Eamsobhana P, Lim PE. Complete mitochondrial genome of *Angiostrongylus malaysiensis* lungworm and molecular phylogeny of Metastrongyloid nematodes. Acta Trop 2016; 161:33–40.
- Zhang Q, Ding X, Liu YH, Liu JF, et al. Establishment of a recombinase-aided isothermal amplification assay for nucleic acid detection of *Angiostrongylus cantonensis*. Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi 2020; 32:350–354. (In Chinese)
- Zhang X, Duan JY, Shi YL, Jiang P, et al. Comparative mitochondrial genomics among Spirometra (Cestoda: Diphylobothriidae) and the molecular phylogeny of related tapeworms. Mol Phylogenet Evol 2017a; 117:75–82.
- Zhang X, Duan Y, Wang ZQ, Jiang P, et al. Using the small subunit of nuclear ribosomal DNA to reveal the phylogenetic position of the plerocercoid larvae of Spirometra tapeworms. Exp Parasitol 2017b; 175:1–7.
- Zhang X, Shi YL, Wang ZQ, Duan JY, et al. Morphological and mitochondrial genomic characterization of eyeworms (*Thelazia callipaeda*) from clinical cases in central China. Front Microbiol 2017c; 8:1335.
- Zhang Y, Zhou XN, Liu HX, Lv S, et al. Development of PCR assay for detection of *Angiostrongylus cantonensis* in *Pomacea canaliculata*. Chin J Parasitol Parasit Dis 2006; 24: 353–355. (In Chinese)
- Zhao Y, Tsang CC, Xiao M, Cheng J, et al. Intragenomic internal transcribed spacer region sequence heterogeneity and molecular diagnosis in clinical microbiology. Int J Mol Sci 2015; 16:25067–25079.
- Zhou H, Chen Z, Limpanont Y, Hu Y, et al. Necroptosis and Caspase-2-mediated apoptosis of astrocytes and neurons, but not microglia, of rat hippocampus and parenchyma caused by *Angiostrongylus cantonensis* infection. Front Microbiol 2020; 10:3126.

Address correspondence to:

Lin Ai

Department of One Health

School of Global Health

Chinese Center for Tropical Diseases Research

Shanghai Jiao Tong University School of Medicine

227 South Chongqing Road

Huangpu, Shanghai 200025

China

E-mail: ailin@nipd.chinacdc.cn

Xiaonong Zhou

National Institute of Parasitic Diseases

Chinese Center for Disease Control and Prevention

207 Rui Jin Er Road

Huangpu, Shanghai 200025

China

E-mail: zhouxn1@chinacdc.cn

Renli Zhang

Institute of Pathogenic Biology

Shenzhen Center for Disease Control and Prevention

8 Longyuan Road

Nanshan, Shenzhen 518055

China

E-mail: renlizhangszcdc@aliyun.com