

Assignment 4

● Graded

Student

James La Fontaine

Total Points

11 / 13 pts

Question 1

Question 2

11 / 13 pts

✓ + 1 pt 2a) M: use derivative substitution

✓ + 1 pt 2a) A: $-3 \cos(x^2) + c$

✓ + 1 pt 2b) M: Use a valid hyperbolic substitution. Expecting $x = \frac{1}{3} \cosh \theta$.

✓ + 1 pt 2b) J: Stated use of $\sinh \theta > 0$ to remove absolute value signs. ($\sinh \theta \geq 0$) is okay too.

✓ + 1 pt 2b) A: $\int \cosh^2(\theta) d\theta$. correct integrand after substitution up to a constant factor.

✓ + 1 pt 2b) M: Use a valid method such as a double angle formula to calculate the integral.

+ 1 pt 2b) A: $\frac{1}{54} (3x\sqrt{9x^2 - 1} + \operatorname{arccosh}(3x)) + c$. Must simplify compositions of hyperbolic functions and their inverses.

✓ + 1 pt 2c) M: Use integration by parts twice to obtain an equation for $\int e^{-x} \cos(x) dx$.

✓ + 1 pt 2c) A: $\frac{1}{2}e^{-x} (\sin(x) - \cos(x)) + c$

✓ + 1 pt 2d) M: Use complex exponential

✓ + 1 pt 2d) A: $\frac{1}{2}e^{-x} (\sin(x) - \cos(x)) + c$

✓ + 1 pt 2e) A: For a reason given that contains no mathematical errors.

+ 1 pt N: Notation correction throughout question 2 and no +c's missing in non-answer steps.

+ 0 pts Scored zero or not attempt

1 good

2 something wrong, missing a factor

3 dx

4 would be good to put big brackets around it

5 refer to solutions

6 integrate gives you 1/2 outside?

No questions assigned to the following page.

Calculus 2 Assignment 4

1. $f(t) = 20000 \operatorname{sech}^2(2t - 30)$, $t \geq 0$

a) y-intercept

$$y = 20000 \operatorname{sech}^2(2(0) - 30)$$

$$y = 20000 \operatorname{sech}^2(-30)$$

x-intercepts

$$20000 \operatorname{sech}^2(2t - 30) = 0$$

$$\operatorname{sech}^2(2t - 30) = 0$$

$$\operatorname{sech}(2t - 30) = 0$$

$$2t - 30 = \operatorname{arcsech}(0) \quad \operatorname{sech} > 0$$

= undefined over \mathbb{R} as
 $\operatorname{dom}(\operatorname{arcsech}) = (0, 1]$

horizontal asymptotes

$$y = \frac{20000}{\cosh^2(2t - 30)}$$

degree $n <$ degree d

$\Rightarrow y = 0$ is the horizontal asymptote

vertical asymptote

occurs when $\cosh^2(2t - 30) = 0$

$$\cosh(2t - 30) = 0, \quad \cosh > 0$$

$$2t - 30 = \operatorname{arccosh}(0)$$

= undefined over \mathbb{R}

as $\operatorname{dom}(\operatorname{arccosh}) = [1, \infty)$

No questions assigned to the following page.

● Stationary point

● occur when $f'(t) = 0$

$$f'(t) = \frac{d}{dt} (20000 \operatorname{sech}^2(2t-30))$$

$$= 20000 \frac{d}{dt} (\operatorname{sech}^2(2t-30))$$

$$= 20000 \cdot 2 \operatorname{sech}(2t-30) \cdot \frac{d}{dt} (\operatorname{sech}(2t-30))$$

$$= 40000 \operatorname{sech}(2t-30) \cdot -\operatorname{sech}(2t-30) \tanh(2t-30) \cdot 2$$

$$= -80000 \operatorname{sech}^2(2t-30) \tanh(2t-30)$$

$$\Rightarrow -80000 \operatorname{sech}^2(2t-30) \tanh(2t-30) = 0$$

$$\operatorname{sech}^2(2t-30) \tanh(2t-30) = 0$$

$$\tanh(2t-30) = 0$$

$$2t-30 = \operatorname{arctanh}(0)$$

$$2t-30 = 0$$

$$2t = 30$$

$$t = 15$$


sub $t=15$ into $f(t)$

$$f(15) = 20000 \operatorname{sech}^2(0) \\ = 20000$$

● turning point at $(15, 20000)$



No questions assigned to the following page.



b) $0.01 \cdot f(t)$ cases require hospitalisation

\Rightarrow the hospital will first be over capacity when $0.01 \cdot f(t)$ is first ~~equal to~~ greater than 10.

$$\Rightarrow 200 \operatorname{sech}^2(2t-30) = 10$$

$$\operatorname{sech}^2(2t-30) = \frac{1}{20}$$

$$|\operatorname{sech}(2t-30)| = \pm \frac{1}{\sqrt{20}}$$

$$\operatorname{sech}(2t-30) = \frac{1}{\sqrt{20}}, \quad \text{range}(\operatorname{sech}) > 0$$

$$\operatorname{sech}(2t-30) = \operatorname{sech}(-2t+30) \quad (\text{as sech is an even function})$$

$$\Rightarrow \operatorname{sech}(2t-30) = \frac{1}{\sqrt{20}} \quad \text{and} \quad \Rightarrow \operatorname{sech}(-2t+30) = \frac{1}{\sqrt{20}}$$

$$2t-30 = \operatorname{arcsech}\left(\frac{1}{\sqrt{20}}\right) \quad -2t+30 = \operatorname{arcsech}\left(\frac{1}{\sqrt{20}}\right)$$

$$t = \frac{\operatorname{arcsech}\left(\frac{1}{\sqrt{20}}\right) + 30}{2} \quad t = -\frac{\operatorname{arcsech}\left(\frac{1}{\sqrt{20}}\right) - 30}{2}$$

$$t \approx 16$$

$$t \approx 14$$

($t < 14$ so day 14 will

be the first day)

No questions assigned to the following page.

- Therefore, the hospital system is first expected to be over capacity 14 days after the start of the disease outbreak.

$$\begin{aligned} \text{c) } T(t) &= \int_0^t f(u) \, du \\ &= \int_0^t 20000 \operatorname{sech}^2(2u-30) \, du \end{aligned}$$

$$\text{Let } v = 2u - 30$$

$$\frac{dv}{du} = 2$$

$$\text{sub } u=t \Rightarrow v=2t-30$$

$$\frac{1}{2} \frac{dv}{du} = 1$$

$$\text{sub } u=0 \Rightarrow v=-30$$

$$\Rightarrow T(t) = 20000 \int_0^t \operatorname{sech}^2(v) \frac{1}{2} \frac{dv}{du} \, du$$

$$= 20000 \cdot \frac{1}{2} \int_{-30}^{2t-30} \operatorname{sech}^2(v) \, dv$$

$$= 10000 [\tanh(v)]_{-30}^{2t-30}$$

$$T(t) = 10000 (\tanh(2t-30) - \tanh(-30))$$

$$\text{or } T(t) = 10000 \left(\tanh(2t-30) - \frac{e^{-30} - e^{30}}{e^{-30} + e^{30}} \right)$$

No questions assigned to the following page.

d) Total number of expected cases $\approx 20,000$

$$\frac{20000}{1000000} = 0.02 \Rightarrow 2\% \text{ of the population are expected to be infected}$$

This model says that roughly 98% of the population will remain uninfected which is roughly 980,000 people

Question assigned to the following page: [1](#)

2.

$$a) \int 6x \sin(x^2) dx$$

$$\text{Let } u = x^2$$

$$\frac{du}{dx} = 2x \Rightarrow 3 \frac{du}{dx} = 6x$$

~~Derivative Substitution~~

Derivative
Substitution

$$\int 6x \sin(x^2) dx$$

$$= \int 3 \frac{du}{dx} \sin(u) dx$$

$$= 3 \int \sin(u) du$$

$$= -3 \cos(u) + C$$

$$= -3 \cos(x^2) + C$$

$$b) \int \frac{x^2}{\sqrt{9x^2 - 1}} dx$$

$$= \int \frac{x^2}{\sqrt{9(x^2 - \frac{1}{9})}} dx$$

$$= \frac{1}{3} \int \frac{x^2}{\sqrt{x^2 - \frac{1}{9}}} dx$$



Question assigned to the following page: [1](#)

$$\text{Let } x = \frac{1}{3} \cosh \theta$$

Hyperbolic
Substitution

$$\theta = \operatorname{arccosh}(3x)$$

This is valid when

$$3x \in \operatorname{dom}(\operatorname{arccosh}) \text{ and } \theta \in \operatorname{range}(\operatorname{arccosh})$$

$$\Rightarrow 3x \geq 1 \Rightarrow x \geq \frac{1}{3} \text{ and } \theta \geq 0$$

$$\text{Also need } \sqrt{9x^2 - 1} \neq 0$$

$$|9x^2 - 1| \neq 0$$

$$|9x^2 - 1| \neq 0$$

$$9x^2 \neq 1$$

$$x^2 \neq \frac{1}{9}$$

$$x \neq \frac{1}{3} \Rightarrow \theta \neq 0 \text{ as } \theta = 0 \text{ when } x = \frac{1}{3}$$

~~$$|9x^2 - 1| > 0 \text{ as } |0| = 0$$~~

$$\Rightarrow x > \frac{1}{3} \text{ and } \theta > 0$$

$$\bullet x = \frac{1}{3} \cosh \theta$$

$$\bullet \frac{dx}{d\theta} = \frac{1}{3} \sinh \theta$$

$$\bullet \frac{x^2}{\sqrt{x^2 - \frac{1}{9}}} = \frac{\frac{1}{9} \cosh^2 \theta}{\sqrt{\frac{1}{9} \cosh^2 \theta - \frac{1}{9}}}$$

$$= \frac{\frac{1}{9} \cosh^2 \theta}{\sqrt{\frac{1}{9} (\cosh^2 \theta - 1)}}$$

$$= \frac{\frac{1}{9} \cosh^2 \theta}{\sqrt{\frac{1}{9} \sinh^2 \theta}}$$

Question assigned to the following page: [1](#)

$$= \frac{1/9 \cosh^2 \theta}{1/3 |\sinh \theta|}$$

$$= \frac{1/9 \cosh^2 \theta}{1/3 \sinh \theta}$$

$$= \frac{1/9 \cosh^2 \theta}{1/3 \sinh \theta}$$

$$\sinh > 0 \text{ when } \theta > 0$$

$$\text{Therefore } \frac{1}{3} \int \frac{x^2}{\sqrt{x^2 - 1/9}} dx$$

$$= \frac{1}{3} \int \frac{1/9 \cosh^2 \theta}{1/3 \sinh \theta} \cdot \frac{1}{3} \sinh \theta d\theta$$

$$= \frac{1}{3} \int \frac{1}{9} \cosh^2 \theta d\theta$$

$$= \frac{1}{27} \int \cosh^2 \theta d\theta$$

$$= \frac{1}{27} \int \frac{1}{2} (\cosh(2\theta) + 1) d\theta, \text{ Double Angle formula}$$

$$= \frac{1}{54} [\sinh(2\theta) + \theta] + C$$

$$= \frac{1}{54} [2 \sinh \theta \cosh \theta + \theta] + C, \text{ Double Angle formula}$$

$$= \frac{1}{54} [2 \sqrt{\cosh^2 \theta - 1} \cosh \theta + \theta] + C$$

$$= \frac{1}{54} [2 \sqrt{9x^2 - 1} \cdot 3x + \operatorname{arccosh}(3x)] + C$$

$$= \frac{x}{9} \sqrt{9x^2 - 1} + \frac{1}{54} \operatorname{arccosh}(3x) + C$$

$$= \frac{x}{3} \sqrt{x^2 - \frac{1}{9}} + \frac{1}{54} \operatorname{arccosh}(3x) + C$$

Question assigned to the following page: [1](#)

$$c) \int e^{-x} \cos(x) dx$$

$$\text{Let } u = e^{-x} \quad \frac{dv}{dx} = \cos(x)$$

$$\frac{du}{dx} = -e^{-x} \quad v = \sin(x) \quad , \quad \text{Integration by parts}$$

$$\text{Therefore } \int e^{-x} \cos(x) dx$$

$$= e^{-x} \sin(x) + \int e^{-x} \sin(x) dx$$

$$\text{First integrate } \int e^{-x} \sin(x) dx:$$

$$\text{Let } u = e^{-x} \quad \frac{dv}{dx} = \sin(x) \quad , \quad \text{Integration by parts}$$

$$\frac{du}{dx} = -e^{-x} \quad v = -\cos(x)$$

$$\text{Therefore } \int e^{-x} \sin(x) dx$$

$$= -e^{-x} \cos(x) - \int e^{-x} \cos(x) dx$$

Putting everything together:

$$\Rightarrow \int e^{-x} \cos(x) dx$$

$$= e^{-x} \sin(x) + (-e^{-x} \cos(x) - \int e^{-x} \cos(x) dx)$$

$$\Rightarrow 2 \int e^{-x} \cos(x) dx = e^{-x} \sin(x) - e^{-x} \cos(x) + C$$

$$\Rightarrow \int e^{-x} \cos(x) dx = \frac{e^{-x} \sin(x) - e^{-x} \cos(x)}{2} + d, \quad d = \frac{1}{2} C$$

$$= e^{-x} \left(\frac{1}{2} \sin(x) - \frac{1}{2} \cos(x) \right) + d$$

Question assigned to the following page: [1](#)

d) $\int e^{-x} \cos(x) dx$, complex exponential

$$\begin{aligned} e^{-x} \cos(x) &= e^{-x} \operatorname{Re}(e^{ix}) \\ &= \operatorname{Re}(e^{-x} \cdot e^{ix}), \quad e^{-x} \in \mathbb{R} \\ &= \operatorname{Re}(e^{ix-x}) = \operatorname{Re}(e^{(-1+i)x}) \end{aligned}$$

$$\begin{aligned} \text{Therefore } \int e^{-x} \cos(x) dx &= \int \operatorname{Re}(e^{(-1+i)x}) dx \\ &= \operatorname{Re} \int e^{(-1+i)x} dx \end{aligned}$$

$$= \operatorname{Re} \left[\frac{1}{-1+i} e^{(-1+i)x} + C + di \right], \quad \begin{matrix} c, d \\ \in \mathbb{R} \end{matrix}$$

$$= \operatorname{Re} \left[\frac{-1-i}{(-1+i)(-1-i)} e^{-x} e^{ix} + C + di \right]$$

$$= \operatorname{Re} \left[\frac{-1-i}{2} e^{-x} (\cos(x) + i \sin(x)) + C + di \right]$$

$$= \operatorname{Re} \left[\frac{e^{-x}}{2} (-\cos(x) + \sin(x) - i \cos(x) - i \sin(x)) + C + di \right]$$

$$= \frac{e^{-x}}{2} (\sin(x) - \cos(x)) + C$$

$$= e^{-x} \left(\frac{1}{2} \sin(x) - \frac{1}{2} \cos(x) \right) + C$$

e) I found (d) simpler as using the complex exponential involves easier integration and simple algebra with complex numbers

